

Critical dynamics of Model H within the real-time fRG approach

Yong-rui Chen,¹ Yang-yang Tan,¹ and Wei-jie Fu¹

¹*School of Physics, Dalian University of Technology, Dalian, 116024, P.R. China*

The critical dynamics of Model H with a conserved order parameter coupled to a transverse momentum density which describes the gas-liquid or binary-fluid transitions is investigated within the functional renormalization group approach formulated on the closed time path. According to the dynamic scaling analysis, Model H and QCD critical end point belong to the same dynamic universality class in the critical region. The higher-order correction of the transport coefficient $\bar{\lambda}$ and shear viscosity $\bar{\eta}$ arising from mode-couplings are obtained by calculating the two-point correlation functions. The flow equation of a dimensionless coupling constant for nondissipative interactions is derived to look for the fixed-point solution of the system. The scaling relation between the critical exponent of the transport coefficient and that of the shear viscosity is estimated. Finally, the dynamic critical exponent z is obtained as a function of the spatial dimension d .

I. INTRODUCTION

Critical dynamics plays a crucial role in the studies of QCD phase structure that is related to the phase transition from the confinement hadronic phase to the deconfinement phase of quark-gluon plasma (QGP) [1–4]. It is expected that fluctuations of conserved charges can be used to search for the critical end point (CEP) in heavy-ion collision experiments [5–9]. Significant progresses have been made in the Beam Energy Scan (BES) program at Relativistic Heavy Ion Collider (RHIC) [10–13]. Notably, recent estimates of the location of CEP in the QCD phase diagram have arrived at convergent results from functional QCD [14–16], which are also consistent with the results obtained from finite-size scaling of proton cumulants in heavy-ion collisions [17] and from extrapolation of lattice simulations [18]. The correlation length tends to be divergent near the CEP, such that properties of the critical behavior are universal and independent of the detail of microscopic interactions. It is found that the chiral phase transition at vanishing chemical potential in the chiral limit of u and d quarks is classified into the universality of three dimensional ($3d$) $O(4)$ symmetry [19], and the CEP in the QCD phase diagram belongs to the $Z(2)$ universality class [20], see also e.g., [21] for relevant discussions.

The criticality is not only reflected in static equilibrium properties, but also in real-time nonequilibrium dynamics. When the critical end point is approached, the relaxation time from nonequilibrium to equilibrium is significantly increased, that is known as the effect of critical slowing down [22, 23]. The renormalization group technique by Wilson *et al.* have been extended to study the dynamic critical phenomena [24–27], where several characteristic dynamic universalities were investigated and classified, see [28, 29] for related reviews. It has been shown that the second-order chiral phase transition of QCD lies in the dynamic universality class of ‘Model G’ [22], and the dynamics of QCD critical end point belongs to ‘Model H’ [30]. A plethora of results about the real-time correlation functions and critical dynamics have been obtained within the classical-statistical lattice

simulations [31–36]. Recently, the functional renormalization group (fRG) within the Schwinger-Keldysh formalism [37–41] has been used to investigate the spectral functions [42–44], pseudo-Goldstone damping [45], critical dynamics and dynamic critical exponents in Model A [46–50], Model C [46, 51] and Model G [52] and so forth. This provides us with a powerful tool to explore the critical dynamics of fluctuations with nonperturbative interactions.

In this work we would like to study the properties of dynamic universality class ‘Model H’ using the real-time fRG approach. Model H is characterized by a conserved order parameter field, which can be treated as a combination of the chiral condensate and baryon density in QCD, and this order parameter couples with the energy-momentum density. That is related to the liquid-gas phase transition. Previous studies focus on the perturbative ϵ -expansion method, see [53, 54]. Recently stochastic fluid dynamics simulation has also been performed on Model H [55]. From the equation of motion, we construct the effective action in terms of two different kinds of field with the Keldysh rotation [56, 57]. The mode-couplings and dissipative dynamics are encoded through the calculations of correlation functions. By solving the fixed-point equations we obtain the anomalous dimensions and dynamic critical exponent.

This paper is organized as follows: In Sec. II we briefly introduce the Langevin equation of Model H. The formalism of Model H formulated on the closed time path within the fRG approach is specialized in Sec. III, where the flow equations of the effective potential and relevant parameters are shown. Numerical results are presented and discussed in Sec. IV. In Sec. V we give a summary and conclusion. The expressions of the propagators and Feynman rules of vertices are collected in App. A. In App. B, we show a specific calculation for loop diagram.

II. LANGEVIN EQUATION FOR MODEL H

The simplified model that describes the dynamics of the gas-liquid or binary-fluid transition was introduced

earlier in the mode-coupling theories, which was denoted as Model H [28, 53]. The equations of motion for the coupled modes in Model H are given by the Langevin equations

$$\frac{\partial \phi}{\partial t} = \lambda_0 \nabla^2 \frac{\delta \mathcal{H}}{\delta \phi} - g_0 \nabla \phi \cdot \frac{\delta \mathcal{H}}{\delta \mathbf{j}} + \theta(x, t), \quad (1)$$

$$\frac{\partial \mathbf{j}}{\partial t} = \Pi^\perp \left(\eta_0 \nabla^2 \frac{\delta \mathcal{H}}{\delta \mathbf{j}} + g_0 \nabla \phi \frac{\delta \mathcal{H}}{\delta \phi} + \zeta(x, t) \right), \quad (2)$$

with the Hamiltonian

$$\mathcal{H} = \int d^d x \left(\frac{1}{2} |\nabla \phi(x)|^2 + \frac{r}{2} \phi^2 + \frac{u}{4} \phi^4 + \frac{1}{2} \mathbf{j}^2 \right). \quad (3)$$

The correlations of Gaussian noises read

$$\begin{aligned} \langle \theta(x, t) \theta(x', t') \rangle &= -2\lambda_0 \nabla^2 \delta(x - x') \delta(t - t'), \\ \langle \zeta_\mu(x, t) \zeta_\nu(x', t') \rangle &= -2\eta_0 \nabla^2 \delta(x - x') \delta(t - t') \delta_{\mu\nu}. \end{aligned} \quad (4)$$

The scalar field ϕ of order parameter is a conserved quantity. The vector field $\mathbf{j}$ corresponds to the momentum density or velocity, that has only the transverse components, since the right hand side of equation of motion for $\mathbf{j}$ in Eq. (2) is multiplied by the transverse projection operator, i.e.,

$$\Pi_{\alpha\beta}^\perp = (\delta_{\alpha\beta} - \mathbf{q}_\alpha \mathbf{q}_\beta / \mathbf{q}^2). \quad (5)$$

In Eq. (1) and (2) λ_0 and η_0 are the transport coefficient and shear viscosity, respectively. Their dynamical scaling behaviors are described by the exponents x_λ and x_η in the critical regime, and these exponents can also be seen as dynamic anomalous dimensions. The Hamiltonian has the usual form except the term $\mathbf{j}^2/2$ included. Thus the static critical properties of ϕ are the same as the Ginzburg-Landau model with Ising symmetry. When the system is in thermal equilibrium, the equilibrium distribution function is given by $e^{-\mathcal{H}}$. The coupling constant g_0 describes the interaction between the two dynamical modes in model H. The Gaussian noises θ and ζ drive the system to thermal equilibrium and their correlation functions are shown in Eq. (4).

According to the dynamic scaling relation, the characteristic frequency of the order parameter has the following form [58]

$$\omega_\phi(k) = Dk^2 \Omega(k\xi) = D_0 \xi^{2-z} k^2 \Omega(k\xi), \quad (6)$$

in the limit $\xi \rightarrow \infty, k \rightarrow 0$. Here ξ is the correlation length, and the diffusion constant D satisfies the ‘Kawasaki-Stokes’ relation [59, 60]

$$D \equiv \frac{\lambda}{\chi_\phi} = \frac{Rk_B T}{\eta\xi}, \quad (7)$$

in $d = 3$ dimensions. Here λ and η stand for the renormalized transport coefficient and shear viscosity, respectively; χ_ϕ denotes the static susceptibility; T is the temperature and k_B the Boltzmann constant; R is a constant. Using the scaling relations as follows

$$\chi_\phi \sim \xi^{2-\eta_\phi}, \quad \lambda \sim \xi^{x_\lambda}, \quad \eta \sim \xi^{x_\eta}, \quad (8)$$

one arrives at the dynamic critical exponent z for the order parameter

$$z = 4 - \eta_\phi - x_\lambda, \quad (9)$$

where η_ϕ is the static anomalous dimension. Note that the Kawasaki-Stokes relation in Eq. (7) also leaves us with

$$x_\eta + x_\lambda = 1 - \eta_\phi, \quad (10)$$

in $d = 3$ dimensions. The results of the exponents x_λ and x_η in the ϵ -expansion up to the order of $O(\epsilon^2)$ are obtained as [53]

$$\begin{aligned} x_\lambda &= \frac{18}{19} \epsilon [1 - 0.033\epsilon + O(\epsilon^2)], \\ x_\eta &= \frac{1}{19} \epsilon [1 + 0.238\epsilon + O(\epsilon^2)]. \end{aligned} \quad (11)$$

In the next section, we will compute the aforementioned exponents using the functional renormalization group approach.

III. MODEL H WITHIN THE REAL-TIME FRG APPROACH

The crucial part of the functional renormalization group (fRG) approach is the scale-dependent effective action $\Gamma_k[\Phi]$ that interpolates smoothly between the short-distance and long-distance physics with the evolution of the exact renormalization group equation [61], for fRG reviews see [2, 3, 62]. At the ultraviolet cutoff $k = \Lambda$, the effective action $\Gamma_\Lambda[\Phi]$ coincides with the classical action $S[\Phi]$ and at $k = 0$ the effective action is the quantum action as fluctuations of different momentum modes are integrated in successively. To formulate the effective action in the closed time path, starting from the Langevin equation in Eq. (1) and (2), we perform the path integral about the noise variables that result in the Martin-Siggia-Rose response field [63]. The renormalization group (RG) scale k dependent effective action of Model H in the Schwinger-Keldysh formalism is

$$\begin{aligned} \Gamma_k[\Phi] &= i \int d^d x dt \left\{ \phi_q \left(\frac{\partial}{\partial t} \phi_c + \lambda_k \nabla^2 \nabla^2 \phi_c \right) \right. \\ &\quad - \lambda_k \phi_q \nabla^2 \frac{\delta V_k(\rho_c)}{\delta \phi_c} + g_k \phi_q \nabla \phi_c \cdot \mathbf{j}_c + 2\lambda_k \phi_q \nabla^2 \phi_q \\ &\quad + \mathbf{j}_{q,\alpha} \Pi_{\alpha\beta}^\perp \left[\left(\frac{\partial}{\partial t} \mathbf{j}_{c,\beta} - \eta_k \nabla^2 \mathbf{j}_{c,\beta} \right) \right. \\ &\quad \left. \left. - g_k \nabla \phi_c (-\nabla^2 \phi_c) + 2\eta_k \nabla^2 \mathbf{j}_{q,\beta} \right] \right\}. \end{aligned} \quad (12)$$

Here, (ϕ_c, ϕ_q) and $(\mathbf{j}_{c,\beta}, \mathbf{j}_{q,\alpha})$ are the order parameter fields and momentum densities with the subscripts ‘c’ and ‘q’ denoting classical and quantum fields, respectively, see

propagators, the final result on the power of external momenta should account for the contributions from both the vertices and propagators.

Here we also introduce the dimensionless quantities to convert the flow equations to fixed-point equations which is more convenient to investigate the scaling behavior, such as

$$\begin{cases} \bar{\lambda} = k^{4-z} \lambda_k \\ \bar{\eta} = k^{2-z} \eta_k \\ \bar{g} = k^{1+\frac{d}{2}-z} g_k \\ \bar{\rho} = k^{2-d} \rho \\ u(\bar{\rho}) = k^{-d} V_k(\rho) \end{cases} \quad (18)$$

In App. B, a specific example for the loop diagram calculation is presented for illustrative purpose, where we also show the expansion up to certain power of external momentum and the related approximation used there. Performing the integrals on the right hand side of Fig. 2 and Fig. 3, one arrives at the final form of the flow equations for the effective potential $u'(\bar{\rho})$, and the coefficients $\bar{\lambda}$, $\bar{\eta}$, $\bar{g}$ as follows

$$\begin{aligned} \partial_\tau u'(\bar{\rho}) &= (-2 + \eta_\phi) u'(\bar{\rho}) + (-2 + d + \eta_\phi) \bar{\rho} u^{(2)}(\bar{\rho}) \\ &\quad - \frac{2\nu_d}{d} \left(1 - \frac{\eta_\phi}{d+2}\right) \left(\frac{3u^{(2)}(\bar{\rho}) + 2\bar{\rho}u^{(3)}(\bar{\rho})}{(1 + u'(\bar{\rho}) + 2\bar{\rho}u^{(2)}(\bar{\rho}))^2} \right), \end{aligned} \quad (19)$$

$$\begin{aligned} \partial_\tau \bar{\lambda} &= -(z - 4 + \eta_\phi) \bar{\lambda} \\ &\quad - \frac{2\bar{g}^2}{\bar{\eta}} \frac{1}{(1 + \bar{m}^2)^2} \int d\Omega_d (-1 + \cos^2 \theta) \cos^2 \theta \\ &\quad + \frac{2}{d} \frac{2\bar{g}^2}{\bar{\eta}} \frac{1}{(1 + \bar{m}^2)^2} \int d\Omega_d (2 - 6 \cos^2 \theta + 4 \cos^4 \theta) \\ &\quad - \frac{2\bar{g}^2}{\bar{\eta}} \frac{1}{1 + \bar{m}^2} \int d\Omega_d (\cos^4 \theta - \cos^2 \theta) \\ &\quad + \frac{6\bar{g}^2 \bar{\lambda}}{\bar{\eta}^2} \int d\Omega_d (-1 + \cos^2 \theta) \cos^2 \theta \\ &\quad + \frac{2}{d+2} \frac{6\bar{g}^2 \bar{\lambda}}{\bar{\eta}^2} (1 + \bar{m}^2) \int d\Omega_d (-1 + \cos^2 \theta) \\ &\quad - \frac{3\bar{g}^2 \bar{\lambda}}{\bar{\eta}^3} \int d\Omega_d (4 \cos^2 \theta - 4 \cos^4 \theta) \\ &\quad + \frac{2}{d+4} \frac{32\bar{g}^2 \bar{\lambda}^2}{\bar{\eta}^3} (1 + \bar{m}^2) \int d\Omega_d (\cos^2 \theta - \cos^4 \theta), \end{aligned} \quad (20)$$

$$\begin{aligned} \partial_\tau \bar{\eta} &= -(z - 2) \bar{\eta} \\ &\quad + \frac{2}{d} \frac{\bar{g}^2}{\bar{\lambda}} \frac{1}{(1 + \bar{m}^2)^3} \int d\Omega_d (1 - 3 \cos^2 \theta + 2 \cos^4 \theta), \end{aligned} \quad (21)$$

$$\partial_\tau \bar{g} = -(z - 3 + (4 - d)/2) \bar{g}, \quad (22)$$

where $d\Omega_d$ represents the angular integration in d -dimension. The renormalized dimensionless mass square

of the order parameter reads

$$\bar{m}^2 = u'(\bar{\rho}) + 2\bar{\rho}u^{(2)}(\bar{\rho}). \quad (23)$$

As for the static anomalous dimension η_ϕ , we utilize the result with the modified local potential approximation (LPA')

$$\eta_\phi = \frac{8}{d} \frac{1}{2^d \pi^{d/2} \Gamma(d/2)} \frac{\bar{\rho}_0 (u^{(2)}(\bar{\rho}_0))^2}{(1 + 2\bar{\rho}_0 u^{(2)}(\bar{\rho}_0))^2}, \quad (24)$$

where $\bar{\rho}_0$ is the minimum of the effective potential, i.e., $u'(\bar{\rho}_0) = 0$.

Here we neglect the loop diagram correction for the coupling g_k . The flow for $\bar{g}$ only comes from the running of canonical dimension. It has been proved that there is no correction for the coupling to the one-loop order in the perturbative theory [53, 54]. Besides, the relevant symmetries of the system also protect the coupling constant from renormalization. It has been shown that it is the combination of $\bar{\lambda}\bar{\eta}$ that determines the scaling behavior, rather than the coefficients $\bar{\lambda}$ and $\bar{\eta}$ separately. Therefore we define a new dimensionless coupling constant

$$f = \nu_d \frac{\bar{g}^2}{\bar{\lambda}\bar{\eta}}, \quad (25)$$

with $\nu_d = 1/[2^d \pi^{d/2} \Gamma(d/2)]$ is the geometric factor corresponding to the angular integration. By combining the flow equations of (20) to (22), one arrives at

$$\partial_\tau f = (\eta_\phi - (4 - d))f - I_\lambda f^2 - I_\eta f^2, \quad (26)$$

where the functions I_λ, I_η read

$$\begin{aligned} I_\lambda &= - \frac{2}{(1 + \bar{m}^2)^2} \int d\Omega_d (-1 + \cos^2 \theta) \cos^2 \theta \\ &\quad + \frac{4}{d} \frac{1}{(1 + \bar{m}^2)^2} \int d\Omega_d (2 - 6 \cos^2 \theta + 4 \cos^4 \theta) \\ &\quad - \frac{2}{1 + \bar{m}^2} \int d\Omega_d (\cos^4 \theta - \cos^2 \theta), \end{aligned} \quad (27)$$

$$I_\eta = \frac{2}{d} \frac{1}{(1 + \bar{m}^2)^3} \int d\Omega_d (1 - 3 \cos^2 \theta + 2 \cos^4 \theta), \quad (28)$$

where the high-order terms in Eq. (27) from Eq. (20) have been neglected. The fixed-point equation of the coupling constant f as well as its solution is readily obtained from Eq. (26).

Once we find the fixed-point solution of f^* , the anomalous dimensions of transport coefficient λ_k and shear viscosity η_k can be easily obtained according to the scaling behavior in the critical regime

$$\lambda_k \sim k^{-x_\lambda}, \quad \eta_k \sim k^{-x_\eta}, \quad (29)$$

where the exponents x_λ and x_η are the anomalous dimensions, and they can be computed using the following relation:

$$x_\lambda = -I_\lambda f^*, \quad x_\eta = -I_\eta f^*. \quad (30)$$




FIG. 4: Left panel: Global solution for $u'(\bar{\rho})$ derived from the fixed-point equation Eq. (19). Right panel: Static anomalous dimension η_ϕ as a function of the spatial dimension d , in comparison with the ϵ -expansion result presented in Eq. (31).




FIG. 5: Dynamic anomalous dimensions x_λ (left panel) and x_η (right panel) as functions of the spatial dimensions d . Results from the fRG and ϵ -expansion in [53] are compared. Moreover, the result of x_η obtained by Ohta and Kawasaki in [64] is also shown in the right panel in blue dashed line, see text for more details.

Actually, Eq. (30) can be obtained from the right hand side of flow equations in Eq. (20) and Eq. (21). Finally, the dynamic critical exponent z of the order parameter is obtained based on Eq. (9). In the next section, we will show the numerical results of the fixed-point solution and various critical exponents.

IV. NUMERICAL RESULTS

In this work, the derivative of global effective potential $u'(\bar{\rho})$ is obtained by using the large field expansion method proposed in [65]. By directly solving the fixed-point equation starting from the large field asymptotic, we achieve the global effective potential with high numerical accuracy. The fixed-point value of the coupling

constant f^* is then determined from the global effective potential.

In the left panel of Fig. 4 we present the global solution of the first-order derivative of the effective potential $u'(\bar{\rho})$ as a function of $\bar{\rho}$ for three different spatial dimensions. The relevant static anomalous dimension η_ϕ in LPA' in Eq. (24) is shown in the right panel of Fig. 4. The function $u'(\bar{\rho})$ demonstrates monotonic behavior with respect to $\bar{\rho}$ and possesses a unique zero crossing point, indicating the presence of the Wilson-Fisher fixed point. As the spatial dimension d decreases, the position of zero point $\bar{\rho}_0$ increases, indicating the necessity of a global solution rather than a local Taylor expansion, as discussed in [65]. In the LPA' approximation, the static anomalous dimension follows the form given in Eq. (24), and we compare our results with the ϵ -expansion up to the third order




FIG. 6: Dynamical critical exponent z defined in Eq. (9) as a function of the spatial dimension d . The ϵ -expansion for η_ϕ in Eq. (31) and x_λ in Eq. (11) are also presented for comparison.

[66].

$$\eta = \frac{N+2}{2(N+8)^2} \epsilon^2 + \frac{N+2}{2(N+8)^2} \left[\frac{6(3N+14)}{(N+8)^2} - \frac{1}{4} \right] \epsilon^3. \quad (31)$$

The value of η_ϕ from fRG computation and ϵ -expansion are comparable when $d \geq 3.5$. We also present the exact value $\eta_\phi = 0.25$ for $d = 2$ with a star in this plot.

In Fig. 5, we investigate the dependence of the dynamic anomalous dimensions x_λ , x_η on the spatial dimension d . The ϵ -expansion results from Eq. (11) are also shown for comparison. The anomalous dimension x_λ , associated with the transport coefficient, shows excellent agreement with the perturbative expansion up to ϵ^2 , with deviations appearing for spatial dimensions $d < 2.8$. In contrast, the behavior of x_η differs. Results from the fRG calculations and ϵ -expansion are comparable only as the spatial dimension approaches $d = 4$. Additionally, our result for x_η exhibits non-monotonic behavior when $d \lesssim 2.5$. In [64], the exponent for x_η is evaluated within the formulation of self-consistent mode coupling, where it is found

$$\begin{aligned} x_\eta = & - (2 - \eta_\phi)^2 \Gamma(1 - \frac{\eta_\phi}{2}) \Gamma(d - 1 - \frac{x_\eta}{2} + \frac{\eta_\phi}{2}) \\ & \times \Gamma(2 + \frac{x_\eta}{2}) \left\{ 4(d-1) \Gamma(\frac{d}{2} + 2) \Gamma(2 - \frac{d}{2} + \frac{x_\eta}{2} - \frac{\eta_\phi}{2}) \right. \\ & \times \Gamma(\frac{d}{2} - 1 - \frac{x_\eta}{2}) \Gamma(\frac{d}{2} + \frac{\eta_\phi}{2}) \left. \right\}^{-1}. \end{aligned} \quad (32)$$

The relevant result is shown in the right panel of Fig. 5 in blue dashed line, obtained through numerical iterations for Eq. (32), where the anomalous dimension η_ϕ are used from fRG calculation. Our result is comparable with this result as $d \gtrsim 3$.

Finally, the dynamic critical exponent z as a function of spatial dimension d is shown in Fig. 6. The anomalous

dimensions η_ϕ and x_λ are obtained from Fig. 4 and Fig. 5 for our real-time fRG computation, respectively. The ϵ -expansion results are taken from Eq. (31) for η and from Eq. (11) for x_λ . Since the order parameter is a conserved quantity, the dynamic critical exponent $z = 4$ for spatial dimension $d = 4$. Both results are highly consistent, with only minor differences emerging as the spatial dimension approaches $d = 2$.

V. SUMMARY AND OUTLOOK

In this work, we investigate the critical dynamics of Model H, which describes a conserved order parameter field coupled with the transverse momentum density, using the real-time fRG approach. The Langevin equation of Model H incorporates kinetic terms and mode-coupling interactions. By integrating the noise terms and introducing the Martin-Siggia-Rose response field, i.e., the “q” type fields, which encode fluctuation effects in the Schwinger-Keldysh field theory, we obtain the RG scale-dependent effective action within the fRG approach. This formalism is well-suited for investigating the critical dynamics of Model H.

We solve the one-point and two-point correlation functions of the effective action to derive the flow equations for the effective potential, transport coefficient λ_k , and shear viscosity η_k . The contributions of mode couplings to the flow of λ_k and η_k are considered. We introduce the fixed-point equation for a coupling constant f , which determines the dynamical scaling behavior. The global fixed-point solution of the effective potential is obtained by integrating the flow equation from large to vanishing field. Subsequently, the anomalous dimensions for the transport coefficient x_λ and shear viscosity x_η are determined according to their scaling forms in the critical regime. Finally, the dynamic critical exponent for the order parameter is computed.

Our results for the anomalous dimension x_λ and the dynamic critical exponent show excellent agreement with the perturbative ϵ -expansion results in the range $2 \leq d \leq 4$. Only small deviations emerge as the spatial dimension approaches $d = 2$. However, result for anomalous dimension x_η from fRG computation and ϵ -expansion are comparable only in a narrow range of spatial dimensions close to $d = 4$. Additionally, x_η exhibits a non-monotonic behavior for $d \lesssim 2.5$. From the analysis in [30], the dynamics of QCD critical end point and Model H belong to the same dynamic universality class. Our results predict a dynamic critical exponent $z = 3.0507$ for $d = 3$.

After this work is finished, we note that the critical dynamics of Model H is also studied within the real-time fRG independently in [67].

ACKNOWLEDGMENTS

We thank Jan M. Pawłowski, Fabian Rennecke and Vladimir V. Skokov for insightful comments and discussions. This work is supported by the National Natural Science Foundation of China under Grant Nos. 12175030.

Appendix A: Propagators and mode-coupling vertices

As mentioned in Sec. III, in the real-time fRG framework, the propagators corresponding to the inverse of the two-point functions of the effective action, contain the retarded, advanced and Keldysh components. The general definition of the propagator is

$$G_{\Phi_i \Phi_j, k} = \left(\frac{\delta^2 \Gamma_k[\Phi]}{\delta \Phi_i \delta \Phi_j} + R_{\Phi, k} \right)^{-1}. \quad (\text{A1})$$

Taking the functional derivative with respect to the different types of fields, one arrives at the specific forms of the propagators as follows

$$\begin{aligned} iG_{\phi, k}^R(q) &= \frac{i}{q_0 + i\lambda_k \mathbf{q}^2 \left(\mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) + m_{\phi, k}^2 \right)}, \\ iG_{\phi, k}^A(q) &= \frac{i}{q_0 - i\lambda_k \mathbf{q}^2 \left(\mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) + m_{\phi, k}^2 \right)}, \\ iG_{\phi, k}^K(q) &= \frac{4\lambda_k \mathbf{q}^2}{q_0^2 + \left[\lambda_k \mathbf{q}^2 \left(\mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) + m_{\phi, k}^2 \right) \right]^2}, \end{aligned} \quad (\text{A2})$$

for the order parameter and

$$\begin{aligned} iG_{j_{\alpha\beta}, k}^R(q) &= \frac{i}{q_0 + i\eta_k \mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right)} \Pi_{\alpha\beta}^\perp, \\ iG_{j_{\alpha\beta}, k}^A(q) &= \frac{i}{q_0 - i\eta_k \mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right)} \Pi_{\alpha\beta}^\perp, \\ iG_{j_{\alpha\beta}, k}^K(q) &= \frac{4\eta_k \mathbf{q}^2}{q_0^2 + \left[\eta_k \mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) \right]^2} \Pi_{\alpha\beta}^\perp. \end{aligned} \quad (\text{A3})$$

for the transverse momentum density fields. Since the effective potential satisfy the Ising symmetry, the order parameter only has the radial component. The mass term in Eq. (A2) reads

$$m_{\phi, k}^2 = V'_k(\rho_c) + 2\rho_c V''_k(\rho_c). \quad (\text{A4})$$

The operator $\Pi^\perp$ in Eq. (A3) selects the transverse part of vector momentum density. The regulator functions are chosen as [68, 69]

$$R_{\phi, k}^{qc}(q) = i\lambda_k \mathbf{q}^4 r_B \left(\frac{\mathbf{q}^2}{k^2} \right), \quad R_{j, k}^{qc}(q) = i\eta_k \mathbf{q}^2 r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \delta_{\mu\nu}, \quad (\text{A5})$$

where $r_B(x)$ is a Heaviside step function $\Theta(x)$, that is

$$r_B(x) = \left(\frac{1}{x} - 1 \right) \Theta(1 - x). \quad (\text{A6})$$

For the flow equation of effective potential, the vertex $\Gamma_{\phi_q \phi_c \phi_c, k}^{(3)}$ emerging in the right hand of Eq. (16) is

$$i\Gamma_{\phi_q \phi_c \phi_c, k}^{(3)} = -3\rho_c^{1/2} V_k^{(2)}(\rho_c) - 2\rho_c^{3/2} V_k^{(3)}(\rho_c). \quad (\text{A7})$$

To evaluate the flow equation for the two-point correlation function in Fig. 3, higher-order vertices are needed. They can be easily obtained from the functional derivative of $\Gamma_k[\Phi]$ w.r.t the corresponding fields ϕ (or j). In this work, the three-point vertices arising from the coupling modes read


$$i\Gamma_{\phi_q \phi_c j_c, k}^{(3)}(q_1, q_2, q_3) = g_k \mathbf{q}_2 \cdot \mathbf{e}_{j_c}, \quad (\text{A8})$$

$$i\Gamma_{j_q \phi_c \phi_c, k}^{(3)}(q_1, q_2, q_3) = -g_k \mathbf{q}_3^2 \mathbf{e}_{j_q} \cdot \mathbf{q}_2 - g_k \mathbf{q}_2^2 \mathbf{e}_{j_q} \cdot \mathbf{q}_3, \quad (\text{A9})$$

where the explicit momentum dependence from the “classical” fields are labelled. And the $\mathbf{e}_j$ denotes the unit vector along the j field direction.

Appendix B: Loop diagram calculation

In this appendix, we present a specific example to show how the calculation are performed and to illustrate the approximations used. We employ a loop diagram for the two-point correlation function $\Gamma_{\phi_q \phi_c, k}^{(2)}$ with the regulator function inserted on the propagator of order parameter, that is,



$$= \int_q \partial_\tau R_{\phi, k} G_{\phi, k}^A(q) \Gamma_{\phi_q \phi_c, k}^{(3)} G_{j, k}^K(q-p) \Gamma_{\phi_q \phi_c, k}^{(3)} G_{\phi, k}^A(q). \quad (\text{B1})$$

Substituting the expressions of vertices and propagators in App. A into the equation above, one has

$$\begin{aligned} & \int_q \partial_\tau R_{\phi, k} \frac{1}{\left[i q_0 + \lambda_k \mathbf{q}^2 \left(\mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) + m_{\phi, k}^2 \right) \right]^2} \times \frac{4 \eta_k (\mathbf{q} - \mathbf{p})^2}{(q_0 - p_0)^2 + \left[\eta_k (\mathbf{q} - \mathbf{p})^2 \left(1 + r_B \left(\frac{(\mathbf{q} - \mathbf{p})^2}{k^2} \right) \right) \right]^2} \\ & g_k(i \mathbf{q}_\mu) \times \left(\delta^{\mu\nu} - \frac{(\mathbf{q} - \mathbf{p})_\mu (\mathbf{q} - \mathbf{p})_\nu}{(\mathbf{q} - \mathbf{p})^2} \right) \times g_k(-i \mathbf{p}_\nu) \\ & = \int_q \partial_\tau R_{\phi, k} g_k^2 (-1 + \cos^2 \theta) \mathbf{q}^2 \mathbf{p}^2 \frac{4 \eta_k}{q_0^2 + \left[\eta_k \mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) \right]^2} \frac{1}{\left[i q_0 + \lambda_k \mathbf{q}^2 \left(\mathbf{q}^2 \left(1 + r_B \left(\frac{\mathbf{q}^2}{k^2} \right) \right) + m_{\phi, k}^2 \right) \right]^2}. \end{aligned} \quad (\text{B2})$$

The vertices contribute to the external momentum of order p^2 . Since the flow equation for transport coefficient λ_k should be projected onto the p^4 term, we need to expand the $G_{j, k}^K$ propagator about the external momentum. The result is given by

$$-\frac{1}{2} \frac{1}{[q_0^2/k^2 + (\eta_k x(1 + r_B(x)))^2]^2} 2 \eta_k^2 \delta(1-x) \frac{4(\mathbf{q} \cdot \mathbf{p})^2}{k^4}, \quad (\text{B3})$$

where $x = \mathbf{q}^2/k^2$ is the ratio of internal momentum over RG scale.

Then we complete the integration in Eq. (B2),

$$\frac{1}{(2\pi)^d} \int d\mathbf{x} d\Omega_D k^d x^{\frac{d}{2}} 16 g_k^2 \lambda_k \eta_k^3 (-1 + \cos^2 \theta) \cos^2 \theta \mathbf{q}^4 \mathbf{p}^4 \frac{1}{k^{16}} \frac{k^6 (3\eta_k + \lambda_k k^2 x (1 + \bar{m}^2))}{4 \eta_k^3 (\eta_k + \lambda_k k^2 x (1 + \bar{m}^2))^3} \delta(1-x). \quad (\text{B4})$$

As the RG scale evolves towards the infrared, the shear viscosity η_k increases, and λ_k approaches to a finite value. Then we have $\eta_k \gg \lambda_k$, and the $\lambda_k k^2$ term in the denominator can be safely neglected, which leads us to

$$\frac{1}{(2\pi)^d} \int d\Omega_D \frac{6 \bar{g}^2}{\bar{\eta}^2} (-1 + \cos^2 \theta) \cos^2 \theta \lambda_k \mathbf{p}^4. \quad (\text{B5})$$

-
- [1] C. S. Fischer, QCD at finite temperature and chemical potential from Dyson-Schwinger equations, *Prog. Part. Nucl. Phys.* **105**, 1 (2019), [arXiv:1810.12938 \[hep-ph\]](#).
 - [2] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. M. Pawłowski, M. Tissier, and N. Wschebor, The nonperturbative functional renormalization group and its applications, *Phys. Rept.* **910**, 1 (2021), [arXiv:2006.04853 \[cond-mat.stat-mech\]](#).
 - [3] W.-j. Fu, QCD at finite temperature and density within the fRG approach: an overview, *Commun. Theor. Phys.* **74**, 097304 (2022), [arXiv:2205.00468 \[hep-ph\]](#).
 - [4] J. Braun *et al.*, Soft modes in hot QCD matter, (2023), [arXiv:2310.19853 \[hep-ph\]](#).
 - [5] M. Bluhm *et al.*, Dynamics of critical fluctuations: Theory – phenomenology – heavy-ion collisions, *Nucl. Phys. A* **1003**, 122016 (2020), [arXiv:2001.08831 \[nucl-th\]](#).
 - [6] X. Luo and N. Xu, Search for the QCD Critical Point with Fluctuations of Conserved Quantities in Relativistic Heavy-Ion Collisions at RHIC : An Overview, *Nucl. Sci. Tech.* **28**, 112 (2017), [arXiv:1701.02105 \[nucl-ex\]](#).
 - [7] M. Stephanov, Non-Gaussian fluctuations near the QCD critical point, *Phys. Rev. Lett.* **102**, 032301 (2009), [arXiv:0809.3450 \[hep-ph\]](#).
 - [8] W.-j. Fu, X. Luo, J. M. Pawłowski, F. Rennecke, R. Wen,

- and S. Yin, Hyper-order baryon number fluctuations at finite temperature and density, *Phys. Rev. D* **104**, 094047 (2021), [arXiv:2101.06035 \[hep-ph\]](#).
- [9] W.-j. Fu, X. Luo, J. M. Pawłowski, F. Rennecke, and S. Yin, Ripples of the QCD Critical Point, (2023), [arXiv:2308.15508 \[hep-ph\]](#).
- [10] L. Adamczyk *et al.* (STAR), Energy Dependence of Moments of Net-proton Multiplicity Distributions at RHIC, *Phys. Rev. Lett.* **112**, 032302 (2014), [arXiv:1309.5681 \[nucl-ex\]](#).
- [11] L. Adamczyk *et al.* (STAR), Bulk Properties of the Medium Produced in Relativistic Heavy-Ion Collisions from the Beam Energy Scan Program, *Phys. Rev. C* **96**, 044904 (2017), [arXiv:1701.07065 \[nucl-ex\]](#).
- [12] A. Bzdak, S. Esumi, V. Koch, J. Liao, M. Stephanov, and N. Xu, Mapping the Phases of Quantum Chromodynamics with Beam Energy Scan, *Phys. Rept.* **853**, 1 (2020), [arXiv:1906.00936 \[nucl-th\]](#).
- [13] J. Adam *et al.* (STAR), Nonmonotonic Energy Dependence of Net-Proton Number Fluctuations, *Phys. Rev. Lett.* **126**, 092301 (2021), [arXiv:2001.02852 \[nucl-ex\]](#).
- [14] W.-j. Fu, J. M. Pawłowski, and F. Rennecke, QCD phase structure at finite temperature and density, *Phys. Rev. D* **101**, 054032 (2020), [arXiv:1909.02991 \[hep-ph\]](#).
- [15] F. Gao and J. M. Pawłowski, Chiral phase structure and critical end point in QCD, *Phys. Lett. B* **820**, 136584 (2021), [arXiv:2010.13705 \[hep-ph\]](#).
- [16] P. J. Gunkel and C. S. Fischer, Locating the critical endpoint of QCD: Mesonic backcoupling effects, *Phys. Rev. D* **104**, 054022 (2021), [arXiv:2106.08356 \[hep-ph\]](#).
- [17] A. Sorensen and P. Sorensen, Locating the critical point for the hadron to quark-gluon plasma phase transition from finite-size scaling of proton cumulants in heavy-ion collisions, (2024), [arXiv:2405.10278 \[nucl-th\]](#).
- [18] D. A. Clarke, P. Dimopoulos, F. Di Renzo, J. Goswami, C. Schmidt, S. Singh, and K. Zambello, Searching for the QCD critical endpoint using multi-point Padé approximations, (2024), [arXiv:2405.10196 \[hep-lat\]](#).
- [19] R. D. Pisarski and F. Wilczek, Remarks on the Chiral Phase Transition in Chromodynamics, *Phys. Rev. D* **29**, 338 (1984).
- [20] A. M. Halasz, A. D. Jackson, R. E. Shrock, M. A. Stephanov, and J. J. M. Verbaarschot, On the phase diagram of QCD, *Phys. Rev. D* **58**, 096007 (1998), [arXiv:hep-ph/9804290](#).
- [21] Y.-r. Chen, R. Wen, and W.-j. Fu, Critical behaviors of the $O(4)$ and $Z(2)$ symmetries in the QCD phase diagram, *Phys. Rev. D* **104**, 054009 (2021), [arXiv:2101.08484 \[hep-ph\]](#).
- [22] K. Rajagopal and F. Wilczek, Static and dynamic critical phenomena at a second order QCD phase transition, *Nucl. Phys. B* **399**, 395 (1993), [arXiv:hep-ph/9210253](#).
- [23] B. Berdnikov and K. Rajagopal, Slowing out-of-equilibrium near the QCD critical point, *Phys. Rev. D* **61**, 105017 (2000), [arXiv:hep-ph/9912274](#).
- [24] B. I. Halperin, P. C. Hohenberg, and S.-k. Ma, Calculation of Dynamic Critical Properties Using Wilson's Expansion Methods, *Phys. Rev. Lett.* **29**, 1548 (1972).
- [25] B. I. Halperin, P. C. Hohenberg, and S.-k. Ma, Renormalization-group methods for critical dynamics: 1. Recursion relations and effects of energy conservation, *Phys. Rev. B* **10**, 139 (1974).
- [26] B. I. Halperin, P. C. Hohenberg, and S.-k. Ma, Renormalization-group methods for critical dynamics: 2. Detailed analysis of the relaxational models, *Phys. Rev. B* **13**, 4119 (1976).
- [27] B. I. Halperin, P. C. Hohenberg, and E. D. Siggia, Renormalization-group treatment of the critical dynamics of superfluid helium, the isotropic antiferromagnet, and the easy-plane ferromagnet, *Phys. Rev. B* **13**, 1299 (1976).
- [28] P. C. Hohenberg and B. I. Halperin, Theory of Dynamic Critical Phenomena, *Rev. Mod. Phys.* **49**, 435 (1977).
- [29] R. Folk and G. Moser, Critical dynamics: a field-theoretical approach, *Journal of Physics A: Mathematical and General* **39**, R207 (2006).
- [30] D. T. Son and M. A. Stephanov, Dynamic universality class of the QCD critical point, *Phys. Rev. D* **70**, 056001 (2004), [arXiv:hep-ph/0401052](#).
- [31] J. Berges, S. Schlichting, and D. Sexty, Dynamic critical phenomena from spectral functions on the lattice, *Nucl. Phys. B* **832**, 228 (2010), [arXiv:0912.3135 \[hep-lat\]](#).
- [32] S. Schlichting, D. Smith, and L. von Smekal, Spectral functions and critical dynamics of the $O(4)$ model from classical-statistical lattice simulations, *Nucl. Phys. B* **950**, 114868 (2020), [arXiv:1908.00912 \[hep-lat\]](#).
- [33] D. Schweitzer, S. Schlichting, and L. von Smekal, Spectral functions and dynamic critical behavior of relativistic Z_2 theories, *Nucl. Phys. B* **960**, 115165 (2020), [arXiv:2007.03374 \[hep-lat\]](#).
- [34] D. Schweitzer, S. Schlichting, and L. von Smekal, Critical dynamics of relativistic diffusion, *Nucl. Phys. B* **984**, 115944 (2022), [arXiv:2110.01696 \[hep-lat\]](#).
- [35] A. Florio, E. Grossi, A. Soloviev, and D. Teaney, Dynamics of the $O(4)$ critical point in QCD, *Phys. Rev. D* **105**, 054512 (2022), [arXiv:2111.03640 \[hep-lat\]](#).
- [36] A. Florio, E. Grossi, and D. Teaney, Dynamics of the $O(4)$ critical point in QCD: critical pions and diffusion in Model G, (2023), [arXiv:2306.06887 \[hep-lat\]](#).
- [37] J. Berges and G. Hoffmeister, Nonthermal fixed points and the functional renormalization group, *Nucl. Phys. B* **813**, 383 (2009), [arXiv:0809.5208 \[hep-th\]](#).
- [38] T. Gasenzer and J. M. Pawłowski, Towards far-from-equilibrium quantum field dynamics: A functional renormalisation-group approach, *Phys. Lett. B* **670**, 135 (2008), [arXiv:0710.4627 \[cond-mat.other\]](#).
- [39] J. M. Pawłowski and N. Strodthoff, Real time correlation functions and the functional renormalization group, *Phys. Rev. D* **92**, 094009 (2015), [arXiv:1508.01160 \[hep-ph\]](#).
- [40] L. Corell, A. K. Cyrol, M. Heller, and J. M. Pawłowski, Flowing with the temporal renormalization group, *Phys. Rev. D* **104**, 025005 (2021), [arXiv:1910.09369 \[hep-th\]](#).
- [41] J. Braun *et al.*, Renormalised spectral flows, *SciPost Phys. Core* **6**, 061 (2023), [arXiv:2206.10232 \[hep-th\]](#).
- [42] S. Huelsmann, S. Schlichting, and P. Scior, Spectral functions from the real-time functional renormalization group, *Phys. Rev. D* **102**, 096004 (2020), [arXiv:2009.04194 \[hep-ph\]](#).
- [43] Y.-y. Tan, Y.-r. Chen, and W.-j. Fu, Real-time dynamics of the $O(4)$ scalar theory within the fRG approach, *SciPost Phys.* **12**, 026 (2022), [arXiv:2107.06482 \[hep-ph\]](#).
- [44] J. V. Roth, D. Schweitzer, L. J. Sieke, and L. von Smekal, Real-time methods for spectral functions, *Phys. Rev. D* **105**, 116017 (2022), [arXiv:2112.12568 \[hep-ph\]](#).
- [45] Y.-y. Tan, Y.-r. Chen, W.-j. Fu, and W.-J. Li, Universality of pseudo-Goldstone damping near critical points, (2024), [arXiv:2403.03503 \[hep-th\]](#).

- [46] J. V. Roth and L. von Smekal, Critical dynamics in a real-time formulation of the functional renormalization group, *JHEP* **10**, 065, [arXiv:2303.11817 \[hep-ph\]](#).
- [47] Y.-r. Chen, Y.-y. Tan, and W.-j. Fu, Critical dynamics within the real-time FRG approach, *Phys. Rev. D* **109**, 094044 (2024), [arXiv:2312.05870 \[hep-ph\]](#).
- [48] L. Canet and H. Chaté, A non-perturbative approach to critical dynamics, *Journal of Physics A: Mathematical and Theoretical* **40**, 1937 (2007), [arXiv:cond-mat/0610468](#).
- [49] L. Canet, H. Chate, and B. Delamotte, General framework of the non-perturbative renormalization group for non-equilibrium steady states, *J. Phys. A* **44**, 495001 (2011), [arXiv:1106.4129 \[cond-mat.stat-mech\]](#).
- [50] L. Batini, E. Grossi, and N. Wink, Dissipation dynamics of a scalar field, *Phys. Rev. D* **108**, 125021 (2023), [arXiv:2309.06586 \[hep-th\]](#).
- [51] D. Mesterházy, J. H. Stockemer, L. F. Palhares, and J. Berges, Dynamic universality class of Model C from the functional renormalization group, *Phys. Rev. B* **88**, 174301 (2013), [arXiv:1307.1700 \[cond-mat.stat-mech\]](#).
- [52] J. V. Roth, Y. Ye, S. Schlichting, and L. von Smekal, Dynamic critical behavior of the chiral phase transition from the real-time functional renormalization group, (2024), [arXiv:2403.04573 \[hep-ph\]](#).
- [53] E. D. Siggia, B. I. Halperin, and P. C. Hohenberg, Renormalization-group treatment of the critical dynamics of the binary-fluid and gas-liquid transitions, *Phys. Rev. B* **13**, 2110 (1976).
- [54] H.-U. Yee, Dynamic universality class of model H with frustrated diffusion: ϵ expansion, *Phys. Rev. D* **97**, 016003 (2018), [arXiv:1707.08560 \[hep-ph\]](#).
- [55] C. Chattopadhyay, J. Ott, T. Schaefer, and V. V. Skokov, Simulations of stochastic fluid dynamics near a critical point in the phase diagram, (2024), [arXiv:2403.10608 \[nucl-th\]](#).
- [56] J. S. Schwinger, Brownian motion of a quantum oscillator, *J. Math. Phys.* **2**, 407 (1961).
- [57] L. V. Keldysh, Diagram technique for nonequilibrium processes, *Zh. Eksp. Teor. Fiz.* **47**, 1515 (1964).
- [58] B. I. Halperin and P. C. Hohenberg, Scaling Laws for Dynamic Critical Phenomena, *Phys. Rev.* **177**, 952 (1969).
- [59] G. Arcovito, C. Faloci, M. Roberti, and L. Mistura, Shear viscosity of the binary system aniline-cyclohexane near the critical point, *Phys. Rev. Lett.* **22**, 1040 (1969).
- [60] S.-M. Lo and K. Kawasaki, Frequency-dependence correction to the order-parameter decay rates near the critical point of fluids, *Phys. Rev. A* **8**, 2176 (1973).
- [61] C. Wetterich, Exact evolution equation for the effective potential, *Phys. Lett. B* **301**, 90 (1993), [arXiv:1710.05815 \[hep-th\]](#).
- [62] J. M. Pawłowski, Aspects of the functional renormalisation group, *Annals Phys.* **322**, 2831 (2007), [arXiv:hep-th/0512261 \[hep-th\]](#).
- [63] P. C. Martin, E. D. Siggia, and H. A. Rose, Statistical Dynamics of Classical Systems, *Phys. Rev. A* **8**, 423 (1973).
- [64] T. Ohta, Selfconsistent Calculation of Dynamic Critical Exponents for Classical Liquid, *Progress of Theoretical Physics* **54**, 1566 (1975), <https://academic.oup.com/ptp/article-pdf/54/5/1566/5399799/54-5-1566.pdf>.
- [65] Y.-y. Tan, C. Huang, Y.-r. Chen, and W.-j. Fu, Criticality of the $O(N)$ universality via global solutions to nonperturbative fixed-point equations, (2022), [arXiv:2211.10249 \[hep-ph\]](#).
- [66] K. G. Wilson, Feynman graph expansion for critical exponents, *Phys. Rev. Lett.* **28**, 548 (1972).
- [67] J. V. Roth, Y. Ye, S. Schlichting, and L. von Smekal, Universal critical dynamics near the chiral phase transition and the QCD critical point, (2024), [arXiv:2409.14470 \[hep-ph\]](#).
- [68] D. F. Litim, Optimization of the exact renormalization group, *Phys. Lett. B* **486**, 92 (2000), [arXiv:hep-th/0005245 \[hep-th\]](#).
- [69] D. F. Litim, Optimized renormalization group flows, *Phys. Rev. D* **64**, 105007 (2001), [arXiv:hep-th/0103195 \[hep-th\]](#).