A class of Taub-NUT metrics in the presence of a scalar field

Ali Derekeh * Behrouz Mirza † Pouya Heidari † Fatemeh Sadeghi [§]and Reza Bahani [¶]

Department of Physics, Isfahan University of Technology, Isfahan 84156-83111, Iran

Department of Physics and Astronomy, University of Calgary, Calgary Alberta T2N 1N4 Canada

We derive a class of Taub-NUT metrics in the presence of a scalar field (TNS) by applying the Ehlers transformation on the exact solutions that was recently introduced in [1, 2]. Furthermore, we investigate the effective potential, geodesics, topological charge, and quasinormal modes (QNMs) for the obtained TNS metrics. We also use conformal transformations to generate a new class of exact solutions of the Einstein-conformalscalar theory by using the obtained TNS solutions as seed metrics. Finally we compare QNMs of the class of exact solutions.

1. Introduction

Einstein's equations consist of ten second-order nonlinear differential equations. As a consequence, there is no general method to solve them. Therefore, it is necessary to consider special symmetries that will minimize the number of equations, making it easier to solve them. The more symmetries there are, the simpler it is to solve these equations. A method was proposed by Ernst in 1968 in a series of papers [3, 4]. Ernst demonstrated that for any axially-symmetric and stationary space-time (without considering the cosmological constant), the Einstein-Maxwell equations can be rewritten in terms of a complex function. It is known that some important symmetry transformations are hidden in the Ernst method, called Ehlers or Harrison transformations [5–7]. By using Ehlers transformations, it is possible to add the NUT parameter [8, 9] to axially symmetric metrics, which is a type of pseudomagnetism for the gravitational field.

One of the most important results of the theory of general relativity is predicting the existence of black holes and, as a result, singularities. The defining character-

^{*}a.derekeh@ph.iut.ac.ir

[†]b.mirza@iut.ac.ir

[‡]pouya.heidari@ucalgary.ca

[§]fatemeh.sadeghi96@ph.iut.ac.ir

 $^{{}^{\}P}$ bahanireza@gmail.com

istic of a black hole is that it is concealed from an external observer by an event horizon. The cosmic censorship conjecture (CCC) [10, 11] suggest that wherever there is a singularity in curved space-time, this singularity is covered by a special hypersurface called the event horizon.

Two solutions to the Einstein's field equations are the JNW (Janis-Newman-Winicour) [12, 14?] and the γ -metric (Zipoy-Voorhees metric) [15–17], both of which are asymptotically flat and axially symmetric. These two solutions have naked singularities, meaning that there is no event horizon covering the singularities arising from them. However, these types of exact solutions can be candidates for the mass distribution of the center of the galaxies [18, 19]. It should be noted that the γ -metric is an exact solution of the Einstein field equations in vacuum. The JNW metric is an exact solution of the scalar field, the JNW metric is the same as the Schwarzschild metric. The JNW space-time and its geodesics are studied in [20, 21]. Also, gravitational lensing has been investigated in [22, 23].

In a recent article [1], a three-parameter class of metrics in the presence of a scaler field was introduced. By choosing certain values of the parameters one can derive both JNW and γ -metrics. Therefore, JNW metrics and γ -metric are only special cases of the three-parameter class of metrics. Rotating form of the three-parameter class of metrics of the Einstein field equations was obtained in [2].

In this paper, we consider the three-parameter class of metrics and by using the Ehlers transformation, obtain a new class of Taub-NUT metrics. Then, following the Φ -mapping theory of Duran's topological current [24], a topological current and charge for light rings is obtained for the new Taub-NUT metric and it is proved that there is at least one light ring whose topological number is -1.

By using conformal transformations and the new class of Taub-NUT metrics in the presence of a scalar field and using the Bekenstein method [25], we obtain a new class of exact solutions of the Einstein-conformal-scalar theory. The conformal transformations used here include a redefinition of the massless scalar field and a conformal rescaling of the metric. Additionally, we investigated the curvature invariants, singularities, and quasinormal modes (QNMs) for the obtained metrics.

This article is organized as follows. In Sec. 2, we go through a quick preliminary review of the new class of metrics in the presence of a massless scalar field. Then, in Sec. 3, we obtain the TNS metric by using the class of three-parameter metrics and the Ehlers transformation. Then, we investigate various properties of the TNS metric, such as geodesics, effective potential, and topological charge, and represent the Klenian form of the metric. In Sec. 4, we consider the Bekenstein

 $\mathbf{2}$

approach to find the proper conformal transformation to derive a new class of exact solutions of the Einstein-conformal-scalar theory that are counterpart of the class of three parameter metrics. Thereafter, in Sec. 5, we propose the conformal transformation of the TNS metric, and obtain another new class of exact solutions of the Einstein-conformal-scalar theory. We also investigate the geodesics and the effective potential. Finally, in Sec. 6, we derive the QNMs of TNS and conformal TNS (CTNS) metrics using the light ring method.

2. A CLASS OF METRICS IN THE PRESENCE OF A MASSLESS SCALAR FIELD

Recently, a class of exact solutions of Einstein's equations and their rotating forms were introduced in [1, 2]. The class of solutions depends on three parameters, i.e., mass and two other free parameters and one may obtain the well-known γ -metric and the JNW metric at some values of the parameters. In the following, we will consider the Einstein-Hilbert action in the presence of a massless scalar field

$$S = \int d^4x \sqrt{-g} \left(\mathcal{R} - \partial_\alpha \psi(r) \partial^\alpha \psi(r) \right) , \qquad (1)$$

where \mathcal{R} , g, and ψ are the Ricci scalar, determinant of metric, and scalar field, respectively. By varying the above action with respect to the metric tensor $g_{\mu\nu}$ and the massless scalar field ψ , respectively, the equations of motion will be obtained as follows

$$\mathcal{R}_{\alpha\beta} = \partial_{\alpha}\psi(r)\partial_{\beta}\psi(r) , \qquad (2)$$

$$\nabla_{\alpha} \nabla^{\alpha} \psi(r) = 0 . \tag{3}$$

Now, we consider the following exact solutions of the Einstein's equations that recently was obtained in [1]

$$ds^{2} = -f(r)^{\gamma} dt^{2} + f(r)^{\mu} k(r,\theta)^{\nu} \left(\frac{dr^{2}}{f(r)} + r^{2} d\theta^{2}\right) + f(r)^{1-\gamma} r^{2} \sin^{2}(\theta) d\phi^{2} , \qquad (4)$$

where the functions f(r) and $k(r, \theta)$ are given as below

$$f(r) = 1 - \frac{2m}{r} , \qquad (5)$$

$$k(r,\theta) = 1 - \frac{2m}{r} + \frac{m^2}{r^2} \sin^2(\theta) .$$
 (6)

The metric in (4) depends on three independent parameters. The parameters are mass m, and three other parameters γ , μ , and ν that only two of them are independent as we have a constraint, i.e. $\mu + \nu = 1 - \gamma$. The expression for the scalar field ψ can be written as

$$\psi(r) = \sqrt{\frac{1 - \gamma^2 - \nu}{2}} \ln(1 - \frac{2m}{r}) .$$
(7)

Assuming that the scalar field is a real function, the following conditions on the parameters μ and ν can be obtained

$$\mu \ge \gamma^2 - \gamma ,$$

$$1 - \gamma^2 \ge \nu .$$
(8)

Overall, this three parameter solution is important. Choosing the parameters as $\mu = \gamma^2 - \gamma$ and $\nu = 1 - \gamma^2$, the metric represents the γ -metric, and by choosing $\nu = 0$ will give us the JNW metric. We will define the conformal transformation of metric (4) and hence a new metric in section 4 by using the scalar field (7). Various futures of the three-parameter metrics have been investigated in [1, 2].

3. TAUB-NUT-SCALAR METRIC VIA EHLERS TRANSFORMATION

We use the class of three parameter metrics (4) and apply the Ehlers transformation to drive a new class of TNS metrics. We also consider that the parameter $\gamma = 1$ in (4) and $\mu = -\nu$. Therefore, the metric can be written as follows

$$ds^{2} = -f(r)dt^{2} + f(r)^{\mu}k(r,\theta)^{\nu}\left(\frac{dr^{2}}{f(r)} + r^{2}d\theta^{2}\right) + r^{2}\sin^{2}(\theta)d\phi^{2} .$$
(9)

The metric in (9) has at least one time-like killing vector representing the isometry. Accordingly, following the Ehlers transformation approach, we define the 4-dimensional space-time as follows [26-28]

$$ds^{2} = -e^{2U}(dx^{0} + A_{i}dx^{i})^{2} + dl^{2} , \qquad (10)$$

where

$$A_i = -\frac{g_{0i}}{g_{00}}, \ e^{2U} = -g_{00}, \ dl^2 = \gamma_{ij} dx^i dx^j , \qquad (11)$$

$$\gamma_{\alpha\beta} = -g_{\alpha\beta} + \frac{g_{0\alpha}g_{0\beta}}{g_{00}} \ . \tag{12}$$

The Ehlers transformation states that if the space-time can be written in the form

$$g_{\mu\nu} dx^{\mu} dx^{\nu} = -e^{2U} (dx^0)^2 + e^{-2U} d\tilde{l}^2, \qquad (13)$$

where $d\tilde{l}^2 = e^{2U} dl^2$, we can find a new static solution that satisfies Einstein's field equations. Here we have

June 11, 2024 0:27

$$\bar{g}_{\mu\nu} \, dx^{\mu} \, dx^{\nu} = -\left(\alpha \cosh(2U)\right)^{-1} \left(dx^{0} + A_{i} dx^{i}\right)^{2} - \alpha \cosh(2U) d\tilde{l}^{2} \,, \qquad (14)$$

where α has a constant and positive value and the functions $U = U(x^i)$ and $A_i(x^j)$ satisfy the Ehlers equation

$$-\alpha \sqrt{\tilde{\gamma}} \epsilon_{ijk} U^{,k} = A_{[i,j]} .$$
⁽¹⁵⁾

where $\bar{\gamma}$ is determinant of the spatial part of the conformal metric and $A_{[i,j]} = \frac{\partial A_i}{\partial x^j} - \frac{\partial A_j}{\partial x^i}$. According to the Ehlers approach, for the metric (4), the function above can be defined as

$$U_c = \frac{1}{2} \ln\left(1 - \frac{2m}{r}\right) + \frac{1}{2} \ln(c) , \qquad (16)$$

where c is a constant. As a result, for the three-parameter metric with $\gamma=1,$ we have

$$d\tilde{l}^{2} = \left(1 + \frac{m^{2}\sin^{2}(\theta)}{r^{2} - 2mr}\right)^{\nu} dr^{2} + r^{2} \left(1 - \frac{2m}{r}\right) \left[\left(1 + \frac{m^{2}\sin^{2}(\theta)}{r^{2} - 2mr}\right)^{\nu} d\theta^{2} + \sin^{2}(\theta) d\phi^{2}\right]$$
(17)

and the determinant of the metric is

$$\tilde{\gamma} = r^4 \left(1 - \frac{2m}{r} \right)^2 \left(1 + \frac{m^2 \sin^2(\theta)}{r^2 - 2mr} \right)^{2\nu} \sin^2(\theta) .$$
(18)

If we choose the field A_i in such a way that it is only a function of θ , by solving equation (15) and using metric (17), we get

$$A_{\phi}(\theta) = -2\alpha \ m\cos(\theta) \ . \tag{19}$$

By substituting the field (19) in (14), it is just a matter of calculation to find

$$ds^{2} = \frac{2c \left(r^{2} - 2mr\right)}{\alpha \left[r^{2} + c^{2} \left(r - 2m\right)^{2}\right]} dt^{2} - \frac{8cm \left(r^{2} - 2mr\right) \cos(\theta)}{r^{2} + c^{2} \left(r - 2m\right)^{2}} dt d\phi$$

$$+ \frac{\alpha \left(1 + c^{2} \left(1 - \frac{2m}{r}\right)^{2}\right) \left(1 + \frac{m^{2} \sin^{2}(\theta)}{r^{2} - 2mr}\right)^{\nu}}{2c \left(1 - \frac{2m}{r}\right)} dr^{2}$$

$$+ \frac{\alpha \left[r^{2} + c^{2} \left(r - 2m\right)^{2}\right] \left(1 + \frac{m^{2} \sin^{2}(\theta)}{r^{2} - 2mr}\right)^{\nu}}{2c} d\theta^{2}$$

$$+ \alpha \left(-\frac{8cm^{2}r \left(r - 2m\right) \cos^{2}(\theta)}{r^{2} + c^{2} \left(r - 2m\right)^{2}} + \frac{\left[r^{2} + c^{2} \left(r - 2m\right)^{2}\right] \sin^{2}(\theta)}{2c}\right) d\phi^{2} .$$
(20)

,

Now, if we make the following substitutions

$$r = R - M + \sqrt{M^2 + n^2}, \quad \alpha = \frac{n}{\sqrt{M^2 + n^2}},$$

$$m = \sqrt{M^2 + n^2}, \quad c = \frac{\sqrt{M^2 + n^2} - M}{n},$$
 (21)

in (20) where n is the NUT charge, and then using the following replacements $M \to m$ and $R \to r$, we will find the class of TNS metrics as below

$$ds^{2} = -\left(1 - \frac{2\left(n^{2} + mr\right)}{r^{2} + n^{2}}\right)dt^{2} + \frac{\left(r^{2} + n^{2}\right)\left(1 + \frac{\left(m^{2} + n^{2}\right)\sin^{2}(\theta)}{r^{2} - 2mr - n^{2}}\right)^{\nu}}{r^{2} - 2mr - n^{2}}dr^{2} + \left(r^{2} + n^{2}\right)\left(1 + \frac{\left(m^{2} + n^{2}\right)\sin^{2}(\theta)}{r^{2} - 2mr - n^{2}}\right)^{\nu}d\theta^{2} + \left(-\frac{4n^{2}\left(r^{2} - 2mr - n^{2}\right)\cos^{2}(\theta)}{r^{2} + n^{2}} + \left(r^{2} + n^{2}\right)\sin^{2}(\theta)\right)d\phi^{2} - \frac{4n\left(r^{2} - 2mr - n^{2}\right)\cos(\theta)}{r^{2} + n^{2}}dtd\phi .$$
(22)

Now, considering replacements in (21), the scalar field (7) for the metric (22) is obtained as below

$$\psi(r) = \sqrt{\frac{-\nu}{2}} \ln\left(\frac{r - m - \sqrt{m^2 + n^2}}{r - m + \sqrt{m^2 + n^2}}\right) .$$
(23)

To find the space-time singularities, one must calculate the Kretschmann or Ricci scalar. The Kretschmann Scalar is too complex to report here, however the Ricci scalar has a simple form as below

$$\mathcal{R} = -\frac{2\nu \left(m^2 + n^2\right) \left(1 + \frac{\left(m^2 + n^2\right)\sin^2(\theta)}{r^2 - 2mr - n^2}\right)^{\nu}}{\left(r^2 - 2mr - n^2\right)\left(r^2 + n^2\right)} .$$
(24)

It should be noted that the Ricci scalar is not equal to zero in the presence of a massless scalar field. The Ricci scalar diverges at the following point

$$r = m + \sqrt{m^2 + n^2} , \qquad (25)$$

which indicates the naked singularity of the class of TNS metrics. As is depicted in Fig. 1, for constant values n = 1, m = 1, $\nu = -1$, and $\theta = \pi/4$, the Ricci scalar diverges; hence, there is a naked singularity for the class of TNS metrics.



Fig. 1. Ricci scalar with respect to r for a TNS metric for constant values $n = 1, m = 1, \nu = -1$, and $\theta = \pi/4$.

3.1. KLEINIAN CLASS OF TNS METRICS

The generalization of Lorentzian signature (1, 3) to Kleinian split-signature (2, 2) is useful in the study of quantum field theory and quantum gravity. Therefore, studying Kleinian geometry and its geometric features with a split-signature is important. Recently, the study of black holes has been investigated in the Kleinian signature, and the solutions are found in the Kleinian space through a simple change of coordinates [29–31].

Using the class of TNS metrics (22) and the following coordinate transformations

$$t \to it, \quad \theta \to i\theta$$
, (26)

7

and NUT charge transformation $n \to in$, class of TNS metrics in Klein space can be written in the following form

$$ds^{2} = \left(1 + \frac{2(n^{2} - mr)}{r^{2} - n^{2}}\right) dt^{2} + \frac{\left(r^{2} - n^{2}\right)\left(1 + \frac{\left(m^{2} - n^{2}\right)\sinh^{2}(\theta)}{r^{2} - 2mr + n^{2}}\right)^{\nu}}{r^{2} - 2mr + n^{2}} dr^{2}$$

$$- \left(r^{2} - n^{2}\right)\left(1 + \frac{\left(m^{2} - n^{2}\right)\sinh^{2}(\theta)}{r^{2} - 2mr + n^{2}}\right)^{\nu} d\theta^{2}$$

$$+ \left(\frac{4n^{2}\left(r^{2} - 2mr + n^{2}\right)\cosh^{2}(\theta)}{r^{2} - n^{2}} + \left(r^{2} - n^{2}\right)\sinh^{2}(\theta)\right) d\phi^{2}$$

$$+ \frac{4n\left[r^{2} - 2mr + n^{2}\right]\cosh(\theta)}{r^{2} - n^{2}} dt d\phi ,$$

$$(27)$$

and the scalar field (23) corresponding to the metric (27) with the replacement $n \rightarrow in$ finds as

$$\Psi(r) = \sqrt{\frac{-\nu}{2}} \ln\left(\frac{r - m - \sqrt{m^2 - n^2}}{r - m + \sqrt{m^2 - n^2}}\right) .$$
(28)

It is worth mentioning that in the limit $r \to \infty$, all the components of the Riemann curvature tensor and the curvature invariants and the scalar field become zero.

3.2. GEODESICS IN TNS SPACE-TIME

We use the Lagrangian and the effective potential method to investigate geodesics in the TNS space-time. It should be noted that on the equatorial plane $\theta = \pi/2$, the movement of the test particle is in a circular orbit. Here, the metric components are not a function of ϕ and t, therefore, two Killing vector fields are as follows

$$\xi^{\mu} = \left(\frac{\partial}{\partial t}, 0, 0, 0\right) , \qquad (29)$$

$$\zeta^{\mu} = \left(0, 0, 0, \frac{\partial}{\partial \phi}\right) \,. \tag{30}$$

When the observer moves with four-velocity u^{μ} in the direction of these two vector fields, the metric components will not change, and as a result, we are dealing with two constants here, ξ^{μ} associated with the constant energy E and ζ^{μ} is related to the constant quantity of angular momentum L of the test particles.

$$-E = g_{\mu\nu}u^{\mu}\xi^{\nu} = g_{tt}\dot{t} + g_{t\phi}\dot{\phi} , \qquad (31)$$

$$L = g_{\mu\nu} u^{\mu} \zeta^{\nu} = g_{t\phi} \dot{t} + g_{\phi\phi} \dot{\phi} . \tag{32}$$

Solving the two Eqs. (31) and (32) simultaneously, we have

$$\dot{t} = \frac{1}{B} \left(E g_{\phi\phi} + L g_{t\phi} \right) , \qquad (33)$$

$$\dot{\phi} = -\frac{1}{B} \left(E g_{t\phi} + L g_{tt} \right) , \qquad (34)$$

where

$$B = g_{t\phi}^2 - g_{tt}g_{\phi\phi} \ . \tag{35}$$

Therefore, geodesics related to coordinates can be obtained by solving Eqs. (33) and (34). It is possible to obtain the geodesic related to the radial coordinate by defining a Lagrangian as follows

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} \dot{x}^{\mu} \dot{x}^{\nu} = -\frac{1}{2} \kappa , \qquad (36)$$

where $\kappa = 1, 0, -1$ are affine parameters for time-like, light-like, and space-like paths, respectively. Also, conjugate momentum is defined as

June 11, 2024 0:27

$$\pi_{\mu} = \frac{\partial \mathcal{L}}{\partial \dot{x}^{\mu}} = g_{\mu\nu} \dot{x}^{\nu} . \qquad (37)$$

The Hamiltonian for the test particle for the TNS metric is

$$\mathcal{H} = \pi_{\mu} \dot{x}^{\mu} - \mathcal{L} = \frac{1}{2} \left(g_{tt} \dot{t}^2 + g_{rr} \dot{r}^2 + g_{\theta\theta} \dot{\theta}^2 + g_{\phi\phi} \dot{\phi}^2 + 2g_{t\phi} \dot{t} \dot{\phi} \right) = -\frac{1}{2} \kappa , \qquad (38)$$

or

$$g_{tt}\dot{t}^2 + g_{rr}\dot{r}^2 + g_{\theta\theta}\dot{\theta}^2 + g_{\phi\phi}\dot{\phi}^2 + 2g_{t\phi}\dot{t}\dot{\phi} + \kappa = 0.$$
(39)

The kinetic energy K and potential energy \mathcal{V} for particles are defined as below

$$K = g_{rr}\dot{r}^2 + g_{\theta\theta}\dot{\theta}^2 , \qquad (40)$$

$$\mathcal{V} = g_{tt}\dot{t}^2 + g_{\phi\phi}\dot{\phi}^2 + 2g_{t\phi}\dot{t}\dot{\phi} + \kappa \ . \tag{41}$$

Therefore, the trajectory of the test particle corresponds to the following relation

$$K + \mathcal{V} = 0 \ . \tag{42}$$

From the Eqs. (33), (34), and (41), the effective potential can be written as

$$\mathcal{V}_{eff} = -\frac{1}{B} \left(E^2 g_{\phi\phi} + 2ELg_{t\phi} + L^2 g_{tt} \right) + \kappa .$$
(43)

Therefore, by using metric in Eq. (22), we can find the effective potential for the TNS metric. In Fig. 2, the effective potentials are depicted for different values of the NUT charge and m = 0.8, E = 0.5, L = 4, and $\kappa = 0$.



Fig. 2. The effective potential with respect to r for different values of NUT charge and the constants m = 0.8, E = 0.5, L = 4, and $\kappa = 0$.

The stable paths for TNS metric can be found using the conditions

9

$$\partial_r^2 \mathcal{V}_{eff} \partial_\theta^2 \mathcal{V}_{eff} - \left(\partial_r \partial_\theta \mathcal{V}_{eff}\right)^2 > 0, \quad \frac{1}{g_{rr}} \partial_r^2 \mathcal{V}_{eff} + \frac{1}{g_{\theta\theta}} \partial_\theta^2 \mathcal{V}_{eff} > 0.$$
(44)

Since we examine the light rings, we must deal with the ones in which $\kappa = 0$. As a result, the effective potential can be written as

$$\mathcal{V}_{eff} = -\frac{g_{\phi\phi}}{B} \left(E - E_1 \right) \left(E - E_2 \right) , \qquad (45)$$

where

$$E_1 = \frac{-Lg_{t\phi} + L\sqrt{B}}{g_{\phi\phi}} , \qquad (46)$$

$$E_2 = \frac{-Lg_{t\phi} - L\sqrt{B}}{g_{\phi\phi}} \ . \tag{47}$$

Considering that the time-like circular orbits for a test particle satisfies $\mathcal{V}_{eff} = 0$ and $\partial_r \mathcal{V}_{eff} = 0$, using Eqs. (45) and (46), light ring radius r_{LR} obtains as [32–36]

$$\Phi^r = \frac{\partial_r E_1}{\sqrt{g_{rr}}} = 0 , \quad \Phi^\theta = \frac{\partial_\theta E_1}{\sqrt{g_{\theta\theta}}} = 0 , \qquad (48)$$

where (r, θ) represent the light ring coordinates. Using E_1 in the two dimensional space-time (r, θ) we introduce a vector field as

$$\Phi^r = \frac{\partial_r E_1}{\sqrt{g_{rr}}} , \quad \Phi^\theta = \frac{\partial_\theta E_1}{\sqrt{g_{\theta\theta}}} .$$
(49)

As a result, from the topological current point of view, these points form a topological number so that a topological current can be attributed to it. In the following, by introducing the topological current and the topological number, the vector field will be constructed for the TNS space-time, and the light ring will be determined using zero points of the vector field.

3.3. TOPOLOGICAL CURRENT AND CHARGE

It is known that light rings around black holes can be studied by using topological charges corresponding to them [34, 37–39]. The topological current is defined as

$$j^{\mu} = \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \epsilon_{ab} \partial_{\nu} n^a \partial_{\rho} n^b .$$
 (50)

where a normal vector field can be constructed using the vector field Φ as simply show that j^{μ} is a conserved current June 11, 2024 0:27

$$n^r = \frac{\Phi^r}{\sqrt{\Phi_r^2 + \Phi_\theta^2}} , \qquad (51)$$

$$n^{\theta} = \frac{\Phi^{\theta}}{\sqrt{\Phi_r^2 + \Phi_{\theta}^2}} , \qquad (52)$$

$$\partial_{\mu}j^{\mu} = 0 . (53)$$

As a result, a constant quantity can be related to this conserved current, which is the topological charge. Using the following integral over the parameter region Σ , the constant charge is obtained as

$$Q = \int_{\Sigma} j^0 d^2 x \ . \tag{54}$$

Since this current is caused by point-like topological charges that are placed on the light ring, it can be obtained using the definition of Dirac's delta function as

$$j^{\mu} = \int_{\Sigma} \delta^2 \left(\Phi \right) j^{\mu} \left(\frac{\Phi}{x} \right) \ . \tag{55}$$

This equation is completely in agreement with the concept of topological current because the topological current is non-zero only on the light ring. At the topological charge points, the vector field Φ is 0, and according to the definition, the Dirac's delta function (55) is non-zero, as a result, we have topological current. It should be noted that $\Phi = (X^i, t)$ are the vector field components obtained from the effective potential. As a result, the topological charge is simplified as

$$Q = \int_{S} j^{0} d^{2}x = \sum_{i=1}^{N} \beta_{i} \eta_{i} = \sum_{i=1}^{N} w_{i} , \qquad (56)$$

in which β_i is the Hopf index, $\eta = \pm 1$ is the Brouwer degree, and the w_i is the winding index for the zero points of the vector field Φ^{α} in S area. In Eq. (56), at the points where the vector field Φ is zero, the topological charge is non-zero. Note that these points are located on the light ring so that each light ring will correspond to one topological charge. If S covers zero points of Φ , then topological charge Q corresponds to the sum of the winding index of these points on the light ring. On the other hand, if only one zero point of the vector field Φ is covered by the S, then the topological charge Q will be directly proportional to the winding number.

As it can be seen in Fig. 3, for specific values of the parameters of the TNS metric, the topological charge Q is -1, which indicates that there is an unstable orbit for massless particles in TNS space-time.

11



Fig. 3. The unit vector field for m = 1, $\mu = 0$, l = 1.9, $\nu = -1$, and n = 0.5.

4. NEW CLASS OF EXACT SOLUTIONS OF THE EINSTEIN-CONFORMAL-SCALAR THEORY

Bekenstein has shown that the Einstein-scalar theory and the Einstein-conformalscalar theory are coformally related and obtained its exact solutions in [25], which is the conformal transformation of the JNW metric [40]. In this article, we take the three-parameter metric in Eq. (4) as a seed metric and by introducing a conformal transformation derive a new class of exact solutions of the Einstein-conformal-scalar theory.

Consider the following action of the Einstein-conformal-scalar theory that is related to the action of the Einstein-scalar theory in Eq. (1), by a conformal transformation as follows

$$S = \int d^4x \sqrt{-\bar{g}} \left(\bar{\mathcal{R}} - \bar{\nabla}_\mu \bar{\psi} \bar{\nabla}^\mu \bar{\psi} - \frac{1}{6} \bar{\mathcal{R}} \bar{\psi}^2 \right) , \qquad (57)$$

where

June 11, 2024 0:27

$$\bar{g}_{\mu\nu} = \cosh^2\left(\sqrt{\frac{1}{6}}\psi\right)g_{\mu\nu} , \qquad (58)$$

$$\bar{\psi} = \sqrt{6} \tanh\left(\sqrt{\frac{1}{6}}\psi\right) \ . \tag{59}$$

Varying the action (57) with respect to scalar field $\bar{\psi}$, and spacetime metric we derive the following field equations

$$\left(\bar{\nabla}_{\mu}\bar{\nabla}^{\mu} - \frac{1}{6}\bar{\mathcal{R}}\right)\bar{\psi} = 0 , \qquad (60)$$

$$\bar{G}_{\mu\nu} = \bar{T}_{\mu\nu} \ , \tag{61}$$

where $\bar{G}_{\mu\nu}$ is the conformal Einstein tensor and the energy-momentum tensor is

$$\bar{T}_{\mu\nu} = \bar{\nabla}_{\mu}\bar{\psi}\bar{\nabla}_{\nu}\bar{\psi} - \frac{1}{2}\bar{g}_{\mu\nu}\bar{\nabla}_{\mu}\bar{\psi}\bar{\nabla}^{\mu} + \frac{1}{6}\left(\bar{g}_{\mu\nu}\bar{\nabla}_{\mu}\bar{\nabla}^{\mu} - \bar{\nabla}_{\mu}\bar{\nabla}_{\nu} + \bar{G}_{\mu\nu}\right)\bar{\psi}^{2} .$$
(62)

The new solution should solve the transformed field equations. It is straightforward to show that for a conformal non-minimal coupling, the Ricci scalar vanishes ($\bar{\mathcal{R}} = 0$) and the equation of motion for $\bar{\psi}$ in Eq. (60) reduces to

$$\bar{\nabla}_{\mu}\bar{\nabla}^{\mu}\bar{\psi} = 0.$$
(63)

We can obtain exact class of solutions that satisfies the field Eqs. (60) and (61) by using conformal transformations (58) and (59) of the class of three-parameter (4) and TNS metrics (22). Using the two transformations in Eqs. (58) and (59) with the scalar field (7) and renaming m = 2M, and using the coordinates $(t, \tilde{r}, \theta, \phi)$ with

$$r = \frac{\tilde{r}^2}{\tilde{r} - M} , \qquad (64)$$

and using Eqs. (4), (7), the rescaled metric $\bar{g}_{\mu\nu}$ and the scalar field $\bar{\psi}$ are obtained as follows

$$\bar{g}_{\mu\nu} = \frac{\left(\left(1 - \frac{2M}{\tilde{r}}\right)^{2\sqrt{\frac{1 - \gamma^2 - \nu}{3}}} + 1\right)^2}{4\left(1 - \frac{2M}{\tilde{r}}\right)^{2\sqrt{\frac{1 - \gamma^2 - \nu}{3}}}} g_{\mu\nu} , \qquad (65)$$

$$\bar{\psi} = \sqrt{6} \frac{\left(1 - \frac{2M}{\tilde{r}}\right)^{2\sqrt{\frac{1 - \gamma^2 - \nu}{3}}} - 1}{\left(1 - \frac{2M}{\tilde{r}}\right)^{2\sqrt{\frac{1 - \gamma^2 - \nu}{3}}} + 1}$$
(66)

Therefore, the new exact class of three parameter solutions of equation (61) can be written as follows

$$ds^{2} = \frac{1}{4} \left(\left(1 - \frac{2M}{\tilde{r}} \right)^{-\sqrt{\frac{\gamma - \gamma^{2} + \mu}{3}}} + \left(1 - \frac{2M}{\tilde{r}} \right)^{\sqrt{\frac{\gamma - \gamma^{2} + \mu}{3}}} \right)^{2} \left[- \left(1 - \frac{2M}{\tilde{r}} \right)^{2\gamma} dt^{2} + \frac{\left(1 - \frac{2M}{\tilde{r}} \right)^{2\mu} \tilde{k}^{1 - \gamma - \mu}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} \left(\frac{dr^{2}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} + \tilde{r}^{2} d\theta^{2} \right) + \frac{\tilde{r}^{2} \left(1 - \frac{2M}{\tilde{r}} \right)^{2 - 2\gamma}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} \sin^{2}(\theta) d\phi^{2} \right],$$
(67)

where

$$\tilde{k} = \frac{\tilde{r}^2 \left(1 - \frac{2M}{\tilde{r}}\right)^2 + 4M^2 \left(1 - \frac{M}{\tilde{r}}\right)^2 \sin^2(\theta)}{\tilde{r}^2}.$$
(68)

It should be noted that the metric in Eq. (67) represents a class of exact solutions of the Einstein-conformal-scalar theory with three parameters mass M, γ , and μ with the condition $\mu + \nu = 1 - \gamma$. A special case is when $\gamma = 1$ and $\mu = -\nu$, which can be written as follows

$$ds^{2} = \frac{1}{4} \left(\left(1 - \frac{2M}{\tilde{r}} \right)^{-\sqrt{\frac{\mu}{3}}} + \left(1 - \frac{2M}{\tilde{r}} \right)^{\sqrt{\frac{\mu}{3}}} \right)^{2} \left[-\left(1 - \frac{2M}{\tilde{r}} \right)^{2} dt^{2} + \frac{\left(1 - \frac{2M}{\tilde{r}} \right)^{2\mu} \tilde{k}^{-\mu}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} \left(\frac{dr^{2}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} + \tilde{r}^{2} d\theta^{2} \right) + \frac{\tilde{r}^{2}}{\left(1 - \frac{M}{\tilde{r}} \right)^{2}} \sin^{2}(\theta) d\phi^{2} \right].$$
(69)

For certain circular paths r_0 located in the plane $\theta = \frac{\pi}{2}$ considering the conformal metric (69) and using the effective potential in (41), we have

$$\bar{g}_{tt} + \bar{g}_{\phi\phi} \left(\frac{d\phi}{dt}\right)^2 = 0 , \qquad (70)$$

$$\bar{\Gamma}^{r}_{tt} + \bar{\Gamma}^{r}_{\phi\phi} \left(\frac{d\phi}{dt}\right)^{2} = 0 .$$
(71)

By substituting the components of the conformal metric (69) and the related Christoffel symbols into Eqs. (70) and (71), one obtain the following relations

$$\frac{d\phi}{dt} = \frac{\left(1 - \frac{2M}{\tilde{r}}\right)\left(1 - \frac{M}{\tilde{r}}\right)}{\tilde{r}} , \qquad (72)$$

$$\frac{d\phi}{dt} = \sqrt{-\frac{\bar{\Gamma}^r_{tt}}{\bar{\Gamma}^r_{\phi\phi}}} \ . \tag{73}$$

Now, as the right sides of Eqs. (72) and (73), are equal, the radius of the light ring can be obtained as

$$\tilde{r} = \left(3 + \sqrt{3}\right)M\tag{74}$$

5. CONFORMAL-TAUB-NUT-SCALAR SOLUTION

In section 3, we used Ehlers transformation to establish the TNS metric (22)in Einstein-Scalar theory. Now, we tend to find the Conformal-Taub-NUT-Scalar (CTNS) metric using the conformal transformations (58) and (59) that is a class of exact solutions ($\gamma = 1, \mu = -\nu$) of Eq. (61). For this, we replace m = 2M and n = 2N and use the coordinates $(t, \tilde{r}, \theta, \phi)$ with

$$r = \frac{\tilde{r}^2}{\tilde{r} - M} , \qquad (75)$$

$$\bar{\rho}^2 = M^2 + N^2 , \qquad (76)$$

$$\bar{\Delta} = \tilde{r}^2 - 2M\tilde{r} + M^2 . \tag{77}$$

Using Bekenstein's approach and considering the scalar field (23) and the metric (22) with transformations (65) and (66), the conformal scalar field and conformal metric can be obtained as follows

$$\bar{g}_{\mu\nu} = \cosh^2\left(\sqrt{-\frac{\nu}{12}}\ln\left(\frac{\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}{\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}\right)\right)g_{\mu\nu} , \qquad (78)$$

$$\bar{\psi} = \sqrt{6} \frac{\left(\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2\right)^{\sqrt{-\frac{\nu}{3}}} - \left(\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2\right)^{\sqrt{-\frac{\nu}{3}}}}{\left(\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2\right)^{\sqrt{-\frac{\nu}{3}}} + \left(\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2\right)^{\sqrt{-\frac{\nu}{3}}}}, \quad (79)$$

so the CTNS metric components can be derived as follows

$$\begin{split} \bar{g}_{tt} &= -\frac{W - V^2 U}{4(1 + V^2 U)} H , \\ \bar{g}_{rr} &= \frac{W^{-1} \left(1 + V^2 U\right)}{4(W - V^2 U)} \left(1 + \frac{4\bar{\rho}^2 \sin^2(\theta)}{\tilde{r}^2 \left(U^{-1} W - V^2\right)}\right)^{-\mu} H , \\ \bar{g}_{\theta\theta} &= \frac{\tilde{r}^2}{4} \left(U^{-1} + V^2\right) \left(1 + \frac{4\bar{\rho}^2 \sin^2(\theta)}{\tilde{r}^2 \left(U^{-1} W - V^2\right)}\right)^{-\mu} H , \\ \bar{g}_{\phi\phi} &= \frac{1}{4} \left\{ \frac{16N^2 \left[(1 - V) - \frac{2M}{\tilde{r}} \left(1 - \frac{N}{\tilde{r}}\right)\right] \left[\frac{2M}{\tilde{r}} \left(1 + \frac{N}{\tilde{r}}\right) - \left(1 + V\right)\right] \cos^2(\theta)}{1 + V^2 U} + \tilde{r}^2 \left(U^{-1} + V^2\right) \sin^2(\theta) \right\} H, \\ \bar{g}_{t\phi} &= \frac{N \left[\left(1 - V\right) - \frac{2M}{\tilde{r}} \left(1 - \frac{N}{\tilde{r}}\right)\right] \left[\frac{2M}{\tilde{r}} \left(1 + \frac{N}{\tilde{r}}\right) - \left(1 + V\right)\right]}{1 + V^2 U} H \cos(\theta) , \end{split}$$
(80)

were H, W, U, and V are

15

(80)

$$H = \left(\frac{\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}{\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}\right)^{-\sqrt{\frac{\mu}{3}}} \left(1 + \left(\frac{\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}{\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}\right)^{\sqrt{\frac{\mu}{3}}}\right)^2 ,$$
$$W = \left(1 - \frac{2M}{\tilde{r}}\right)^2 ,$$
$$U = \left(1 - \frac{M}{\tilde{r}}\right)^2 ,$$
$$V = \frac{2N}{\tilde{r}} ,$$
(81)

which satisfies the field equation in (61). The Ricci scalar for metric in (80) is too complicated to be written here, however, we have derived its singularities as follows

$$\tilde{r} = M + N \pm \sqrt{M^2 + N^2}$$
,
 $\tilde{r} = M - N + \sqrt{M^2 + N^2}$.
(82)

It should be noted that the curvature singularities of the CTNS metric differs from the TNS metric in Eq. (25).

5.1. GEODESICS IN CTNS SPACE-TIME

In order to investigate the geometric features of the CTNS space-time, we determine the equations of motion of test particles in this space-time. Considering the CTNS metric (80), the geodesic equation is written as

$$\frac{d^2 x^{\mu}}{ds^2} + \bar{\Gamma}^{\mu}_{\ \nu\rho} \frac{dx^{\nu}}{ds} \frac{dx^{\rho}}{ds} = 0 , \qquad (83)$$

where $x^{\mu} = x^{\mu}(s)$ is the test particle paths, s is the affine parameter, and $\bar{\Gamma}^{\mu}_{\nu\rho}$ is the Christoffel symbol for the CTNS metric $\bar{g}_{\mu\nu} = \Omega(x)g_{\mu\nu}$ whic is defined as

$$\bar{\Gamma}^{\mu}_{\ \nu\rho} = \Gamma^{\mu}_{\ \nu\rho} + \frac{1}{2} \left(\delta^{\mu}_{\nu} \partial_{\rho} \ln \Omega + \delta^{\mu}_{\rho} \partial_{\nu} \ln \Omega - g_{\nu\rho} g^{\mu\lambda} \partial_{\lambda} \ln \Omega \right) , \qquad (84)$$

where

$$\Omega(x) = \Omega(\tilde{r}, \theta) = \cosh^2\left(\sqrt{-\frac{\nu}{12}}\ln\left(\frac{\bar{\Delta} - 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}{\bar{\Delta} + 2\bar{\rho}\left(\tilde{r} - M\right) + M^2}\right)\right) .$$
(85)

Substituting Eq. (84) in Eq. (83) and using $d\bar{s} = \Omega^{-1} ds$ we have

$$\frac{d^2 x^{\mu}}{d\bar{s}^2} + \Gamma^{\mu}_{\ \nu\rho} \frac{dx^{\nu}}{d\bar{s}} \frac{dx^{\rho}}{d\bar{s}} = F^{\mu} , \qquad (86)$$

where $F^{\mu} = \frac{\kappa}{2} g^{\mu\lambda} \partial_{\lambda} \ln \Omega$ is the external effective force due to the CTNS metric and $\kappa = g_{\nu\rho} \frac{dx^{\nu}}{d\bar{s}} \frac{dx^{\rho}}{d\bar{s}}$. To investigate the time-like orbits in CTNS space-time, we will use

the following effective potential

$$\mathcal{V}(\bar{r},\theta) = 1 + \frac{L^2 \bar{g}_{tt} + 2EL \bar{g}_{t\phi} + E^2 \bar{g}_{\phi\phi}}{\bar{g}_{tt} \bar{g}_{\phi\phi} - \bar{g}_{t\phi}^2} .$$
(87)

In CTNS space-time, E and L are defined as

$$E = -\frac{\bar{g}_{tt} + \bar{g}_{t\phi}\omega}{\sqrt{-(\bar{g}_{tt} + 2\bar{g}_{t\phi}\omega + \bar{g}_{\phi\phi}\omega^2)}} , \qquad (88)$$

$$L = \frac{g_{t\phi} + g_{\phi\phi}\omega}{\sqrt{-(\bar{g}_{tt} + 2\bar{g}_{t\phi}\omega + \bar{g}_{\phi\phi}\omega^2)}} , \qquad (89)$$

where $\omega = \frac{d\phi}{dt}$ is the angular velocity. Now, it is a matter of calculation to find the effective potential for the CTNS metric. Fig. 4 represents the effective potential for CTNS metric for different values of NUT charge and M = 0.1, E = 0.5, L = 4, $\nu = -1$, and $\kappa = 0$.



Fig. 4. The effective potential for CTNS metric (80) with respect to r for different values of NUT charge and the constants M = 0.1, E = 0.5, L = 4, $\nu = -1$, and $\kappa = 0$.

Here, we would like to compare the time-like geodesics of TNS and CTNS spacetimes for $\kappa = 1$. For this purpose, we find the effective potential for these two metrics. Fig. 5 is the comparison diagram for E = 0.5, L = 4, $\nu = -1$, and $\kappa = 1$. We have used m = 2M and n = 2N to compare the metrics, so each color belongs to a specific value for mass and NUT charge.



Fig. 5. The effective potentials for TNS and CTNS metrics for different values of NUT charge and constant values E = 0.5, L = 4, $\nu = -1$, m = 0.8, and M = 0.4. The straight and dashed lines correspond to TNS and CTNS space-times, respectively.

6. QNMs OF TNS AND CTNS METRICS

In this section, we will derive the QNMs of TNS and CTNS metrics using the light ring method in the eikonal limit [41–43].

6.1. QNMs OF TNS METRIC

The QNMs of the TNS metric can be obtained by assuming $\theta = \pi/2$ in metric (22) and then expand the metric components up to the second order of n and the first order of ν and omit the terms $n\nu$ (we use these approximations until the end of the QNMs calculations)

$$ds^{2} = -\left(1 - \frac{2m}{r} + \frac{2n^{2}(m-r)}{r^{3}}\right)dt^{2} + \frac{1}{r^{3}(1-2m/r)^{2}}\left[r^{3} + 2rn^{2} - 2mr(r + n^{2}) + r^{3}\nu(1-2m/r)\ln\left(1 + \frac{m^{2}}{r^{2}(1-2m/r)}\right)\right]dr^{2} + \left(n^{2} + r^{2}\left[1 + \nu\ln\left(1 + \frac{m^{2}}{r^{2}(1-2m/r)}\right)\right]\right)d\theta^{2} + (n^{2} + r^{2})d\phi^{2}.$$
(90)

The QNMs are represented by $Q = \Omega + i \Gamma$, which in the light ring method and in the eikonal limit, Ω is defined by the following equation

$$\Omega = \pm j \,\Omega_{\pm},\tag{91}$$

where j is the member of the integers and Ω_{\pm} is the frequency of the circular orbits of the null rays. Also, wave perturbations can be considered using the superposition of eigenmodes as follows

$$e^{i\left(\Omega t - \ell \phi\right)} S_{\Omega j \ell s}(r, \theta), \tag{92}$$

where, S and Ω are the spin and wave's frequency respectively. Also, ℓ and j are angular momentum with the following condition

$$|\ell| \le j. \tag{93}$$

We assume that $\Omega \gg 1/M$ and $|\ell| = j \gg 1$ in the eikonal limit. Moreover, from the numerical calculations of black holes QNMs [41–43] oscillation frequency can be written as follow

$$\Omega = \ell \, \frac{d\phi}{dt} = \pm \, j \, \Omega_{\pm}, \tag{94}$$

which is equal to the QNMs oscillations. To calculate QNMs, we derive the radius of null circular ($\theta = \pi/2$). In every light ring, we have

$$g_{tt} + g_{\phi\phi} \left(\frac{d\phi}{dt}\right)^2 = 0.$$
(95)

The radial component of the geodesics is as follows

$$\Gamma_{tt}^r + \Gamma_{\phi\phi}^r \left(\frac{d\phi}{dt}\right)^2 = 0.$$
(96)

By placing the metric components (90) in Eqs. (95) and (96), we have

$$\frac{d\phi}{dt} = \frac{2\,r^3 - 3\,r\,n^2 + 4\,m\,(n^2 - r^2)}{2\,r^4\,\sqrt{1 - 2\,m/r}},\tag{97}$$

$$\frac{d\phi}{dt} = \frac{2\,m\,r^2 - n^2\,(3\,m - 2\,r)}{2\,r^3\,\sqrt{m\,r}}.\tag{98}$$

By equating Eqs. (97) and (98) and assuming the following perturbation

$$r = 3\,m + \bar{\epsilon},\tag{99}$$

where $\bar{\epsilon}$ is a very small parameter that indicates a perturbation, and 3m is the light ring radius for the Schwarzschild black hole and $r \equiv r_{0_{TNS}}$. By keeping the perturbation terms up to first order of $\bar{\epsilon}$, one has

$$\bar{\epsilon} = \frac{8\,n^2}{9\,m}, \quad \Rightarrow \quad r_{0_{TNS}} = 3\,m + \frac{8\,n^2}{9\,m}.\tag{100}$$

By placing $r_{0_{TNS}}$ in Eq. (97) or Eq. (98), we have

$$\Omega_{\pm} = \frac{d\phi}{dt} = \frac{1}{3\sqrt{3}m} \left(1 \pm \frac{5}{18} \left(\frac{n}{m}\right)^2\right).$$
(101)

Now we have to calculate the other term of QNMs (Γ). Γ represents the decay rate in the amplitude of QNMs. To calculate Γ , we perturb the null equatorial circular orbit. On the other hand, we assume a perturbation in coordinates $\bar{x}^{\mu} = (t, r, \theta, \phi) = (t, r_0, \frac{\pi}{2}, \Omega_{\pm} t)$ as follows

$$r = r_0 \left[1 + \epsilon \,\delta_r(t) \right], \quad \phi = \Omega_{\pm} \left[t + \epsilon \,\delta_{\phi}(t) \right], \quad \ell = t + \epsilon \,\delta_{\ell}(t), \tag{102}$$

where, ϵ is perturbation parameter and following conditions for $\delta_r(t)$, $\delta_{\phi}(t)$ and $\delta_{\ell}(t)$ are applied

$$\delta_r(0) = \delta_\phi(0) = \delta_\ell(0) = 0, \tag{103}$$

from Eqs. (94) and (102), we have $\delta_{\phi}(t) = 0$. The propagation vector is defined as follows (up to $O(e^2)$)

$$K^{\mu} = \frac{dx^{\mu}}{d\ell} = \left(1 - \epsilon \,\delta'_{\ell}, \, \epsilon \, r_0 \,\delta'_r, \, 0, \, \Omega_{\pm} \left(1 - \epsilon \,\delta'_{\ell}\right)\right). \tag{104}$$

Here, the prime represents the derivative with respect to t. The conservation law for the congruence of the null rays can be represented as follows

$$\nabla_{\mu} \left(\varrho_n \, K^{\mu} \right) = 0,$$

$$\Rightarrow \quad \frac{1}{\varrho_n} \frac{d\varrho_n}{d\ell} = -\nabla_{\mu} K^{\mu} = -\frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^{\alpha}} \left(\sqrt{-g} \, K^{\alpha} \right),$$
(105)

where ρ_n is the density of null rays. We consider the terms up to the first order of ϵ , so we have

$$\frac{1}{\varrho_n} \frac{d\varrho_n}{dt} = -\frac{\delta_r''(t)}{\delta_r'(t)}.$$
(106)

To obtain ρ_n , we must obtain $\delta_r(t)$. To calculate δ_r , we use the following equation, which is the radial component of geodesic

$$\frac{d^2r}{d\ell^2} + \Gamma_{tt}^r \left(\frac{dt}{d\ell}\right)^2 + \Gamma_{\phi\phi}^r \left(\frac{d\phi}{d\ell}\right)^2 + \Gamma_{\theta\theta}^r \left(\frac{d\theta}{d\ell}\right)^2 = 0.$$
(107)

By putting $\theta = \pi/2$ and expanding the above equation up to the first order of ϵ , we have

$$9m^{2}\left(27m^{2}+8n^{2}\right)\delta_{r}^{\prime\prime}(t) + \left[n^{2}-9m^{2}\left(1-\nu\ln(\frac{4}{3})\right)\right]\delta_{r}^{\prime}(t) = 0.$$
(108)

By solving the differential Eq. (108), $\delta_r(t)$ is obtained as follows

$$\delta_r(t) = \sinh(\zeta t),\tag{109}$$

where ζ is

$$\zeta = \frac{1}{3\sqrt{3}m} \Big[1 - \frac{11n^2}{54m^2} - \nu \ln 2 + \frac{\nu \ln 3}{2} \Big].$$
(110)

Using Eq. (109) in Eq. (106), ρ_n is obtained as follows

$$\varrho_n(t) = \varrho_n(0) \frac{1}{\cosh(\zeta)} \simeq 2 \, \varrho_n(0) \, (e^{-\zeta \, t} - e^{-3 \, \zeta \, t} + e^{-5 \, \zeta \, t} - \dots). \tag{111}$$

Therefore, the imaginary parts related to the QNMs can be written as below

$$\Gamma = \left(\tilde{n} + \frac{1}{2}\right)\zeta. \tag{112}$$

20

QNMs for the TNS metric are given by

$$Q_{TNS} = \left(\Omega + i\,\Gamma\right)_{TNS} = j\,\left[\frac{1}{3\sqrt{3}\,m}\left(1 \pm \frac{5}{18}\,(\frac{n}{m})^2\right)\right] + i\,\left(\tilde{n} + \frac{1}{2}\right)\left\{\frac{1}{3\sqrt{3}\,m}\left[1 - \frac{11\,n^2}{54\,m^2}\right] - \nu\,\ln 2 + \frac{\nu\,\ln 3}{2}\right]\right\}.$$
(113)

6.2. QNMs OF THE CTNS METRIC

Now, we derive the QNMs of the CTNS metric with the same method introduced in the previous section. To derive the QNMs, we first expand the metric in Eq. (80) to the second order of N and the first order of ν .

$$ds^{2} = \left\{ \frac{(\tilde{r} - 2M)^{2}}{3\tilde{r}^{2}} \left[\nu \ln(\tilde{r} - 2M) - 3 \right] + \frac{8N^{2}}{\tilde{r}^{6}} \left[(M - \tilde{r})^{2} \left(\tilde{r}^{2} - 2M(\tilde{r} - M) \right) \right] \right\} dt^{2} \\ + \left\{ \frac{1}{(M - \tilde{r})^{4}} \left[\tilde{r}^{4} + \frac{8N^{2} \left(M - \tilde{r} \right)^{2} \left(\tilde{r}^{2} + 2M(M - \tilde{r}) \right)}{(\tilde{r} - 2M)^{2}} \right] - \frac{\nu \tilde{r}^{4}}{12 \left(M - \tilde{r} \right)^{4}} \\ \times \left[6 \ln \left(\tilde{r} - 2M \right) - \ln \left(\frac{\tilde{r}^{2} \left(\tilde{r} - 2M \right)^{2} + 4M^{2} \left(\tilde{r} - M \right)^{2}}{\tilde{r}^{2}} \right) \right] \right\} dr^{2} \\ + \frac{1}{(\tilde{r} - M)^{2}} \left\{ \tilde{r}^{4} + 4N^{2} \left(\tilde{r} - M \right)^{2} + \nu \tilde{r}^{4} \left[\ln \left(\frac{\tilde{r}^{2} \left(\tilde{r} - 2M \right)^{2} + 4M^{2} \left(\tilde{r} - M \right)^{2}}{\tilde{r}^{2}} \right) \\ - \ln(\tilde{r} - 2M) \right] \right\} d\theta^{2} + \left\{ 4N^{2} + \frac{\tilde{r}^{4}}{3 \left(\tilde{r} - M \right)^{2}} \left[3 - \nu \ln(\tilde{r} - 2M) \right] \right\} d\phi^{2}.$$

$$(114)$$

We perform all the calculations of the previous part for this metric, with the difference that the radius of the light ring for the conformal Schwarzschild metric is as follows

$$r_{CS} = (3 + \sqrt{3}) M. \tag{115}$$

Finally, we obtain the QNMs of the CTNS metric as follows

$$Q_{CTNS} = \left(\Omega + i\,\Gamma\right)_{CTNS} = j\left[\frac{1}{108\,\sqrt{3}\,M^3}\,(18\,M^2 \pm 5\,N^2)\right] + i\left(n + \frac{1}{2}\right)\left\{-\frac{11N^2}{54M^3} + \frac{1 - 0.144\nu}{M}\right\}.$$
(116)

According to Eqs. (113) and (116) and the data in Table 1, for all values of (n = N), (m = M) and ν , the ratio of Γ_{TNS} to Γ_{CTNS} is always constant and smaller than one. Therefore, the perturbations created in the metric Γ_{CTNS} disappear faster than the perturbations created in the metric Γ_{TNS} .

CONCLUSION

Using the Ehlers method, we obtained a new class of Taub-NUT metrics in the presence of a massless scalar field, which is a new class of exact solutions for Einstein's

22 REFERENCES

Table 1. Numerica	l values of the	imaginary p	part of the	QNMs of	TNS and	CTNS metric	cs
-------------------	-----------------	-------------	-------------	---------	---------	-------------	----

n, N	m, M	ν	$\frac{\Gamma_{TNS}}{\Gamma_{CTNS}}$
1	1	-1	0.19

field equations. The calculation of the Ricci scalar for the TNS metric showed that this metric has a naked singularity. Using the topological charge method, the light ring in the TNS geometry were investigated in such a way that the presence of the topological charge indicates the existence of the light ring in this new geometry, and the topological charge location, with a good approximation, was the extreme point of the effective potential. Furthermore, by using the appropriate conformal transformation, we obtained the exact solutions of the CTNS theory. It should be noted that these conformal solutions are new class of metrics in the presence of a massless scalar field. We also compared the effective potentials of TNS and CTNS spece-times. Lastly, by examination and comparing the QNMs of the TNS and CTNS metrics, we concluded that small perturbations in the TNS space-time are more stable than the small perturbations in the CTNS space-time.

References

- A. Azizallahi, B. Mirza, A. Hajibarat, H. Anjomshoa, Three parameter metrics in the presence of a scalar field in four and higher dimensions, Nuclear Physics B. 998, 116414 (2023), arXiv:2307.09328 [gr-qc].
- [2] B. Mirza, P. K. Kangazi, F. Sadeghi, A class of rotating metrics in the presence of a scalar field, Eur. Phys. J. C. 83, 1161 (2023), arXiv:2307.13588 [gr-qc].
- [3] F. J. Ernst, New formulation of the axially symmetric gravitational field problem, Phys. Rev. 167, 1175 (1968).
- [4] F. J. Ernst, New formulation of the axially symmetric gravitational field problem. II, Phys. Rev. 168, 1415 (1968).
- [5] J. Ehlers, Konstruktionen und Charakterisierung von Lösungen der Einsteinschen Gravitationsfeld Gleichungen, dissertation, Hamburg. (1957).
- B. Kent Harrison; New Solutions of the Einstein-Maxwell Equations from Old, J. Math. Phys. 9, 1744-1752 (1968).
- [7] F. J. Ernst; Black holes in a magnetic universe, J. Math. Phys. 17, 54-56 (1976).
- [8] C. W. Misner. "Taub-NUT Space as a Counterexample to almost anything", Relativity Theory and Astrophysics. 1, 167 (1967).
- [9] B. Chng, R. Mann, and C. Stelea. "Accelerating Taub-NUT and Eguchi-Hanson solitons in four dimensions", Physical Review D 74. 084031 (2006), arXiv:gr-qc/0608092.
- [10] R. Penrose, Gravitational collapse: The role of general relativity, Nuovo Cimento Rivista Serie. 1, 252 (1969).
- [11] S. Hawking, W. Israel, and D. Liebscher, General relativity: an Einstein centenary survey, Astron. Nachrichten. 301, 331 (1980).

REFERENCES 23

- [12] I. Fisher, Scalar mesostatic field with regard for gravitational effects, Zh. Eksp. Teor. Fiz. 18, 636-640 (1948).
- [13] A. I. Janis, E. T. Newman, and J. Winicour, Reality of the Schwarzschild singularity, Phys. Rev. Lett. 20, 878 (1968).
- [14] M. Wyman, Static spherically symmetric scalar fields in general relativity, Phys. Rev. D. 24, 839 (1981).
- [15] G. Darmois, M'emorial des sciences math'ematiques, Gauthier-Villars. (1927).
- [16] D. M. Zipoy, Topology of some spheroidal metrics, J. Math. Phys. 7, 1137 (1966).
- [17] B. Voorhees, Static axially symmetric gravitational fields, Phys. Rev. D. 2, 2119 (1970).
- [18] K. Destounis, G. Huez, and K.D. Kokkotas, Geodesics and gravitational waves in chaotic extreme-mass-ratio inspirals: the curious case of Zipoy-Voorhees black-hole mimickers, Gen. Relativ. Gravit. 55, 71 (2023), arXiv:2301.11483 [gr-qc].
- [19] F. D. Lora-Clavijo, G. D. Prada-Méndez, L. M. Becerra, and E. A. Becerra-Vergara, The q-metric naked singularity: a viable explanation for the nature of the central object in the Milky Way, Class. Quantum Grav. 40, 245012 (2023), arXiv:2311.06653 [gr-qc].
- [20] A. N. Chowdhury, M. Patil, D. Malafarina, and P. S. Joshi, Circular geodesics and accretion disks in the Janis-Newman-Winicour and gamma metric spacetimes, Phys. Rev. D. 85, 104031 (2012), arXiv:1112.2522 [gr-qc].
- [21] B. Turimov, B. Ahmedov, M. Kološ, and Z. Stuchlík, Axially symmetric and static solutions of Einstein equations with self-gravitating scalar field, Phys. Rev. D. 98, 084039 (2018), arXiv:1810.01460 [gr-qc].
- [22] K. Virbhadra, D. Narasimha, and S. Chitre, Role of the scalar field in gravitational lensing, Astron. Astrophys. 337, 1-8 (1998), arXiv:astro-ph/9801174.
- [23] T. K. Dey and S. Sen, Gravitational lensing by wormholes, Mod. Phys. Lett. 23, 953 (2008), arXiv:gr-qc/0602062.
- [24] Y. Duan, L. Fu, G. Jia; Topological tensor current of p̃-branes in the Φ-mapping theory, J. Math. Phys. 41, 4379-4386 (2000), arXiv:hep-th/9904123.
- [25] J. Bekenstein, "Exact solution of Einstein conformal scalar equations", Ann. Phys. (N.Y.) 82, 535 (1974).
- [26] J. Ehlers, "Transformations of static exterior solutions of Einstein's gravitational field equations into different solutions by means of conformal mapping," Colloq. Int. 91, 275–284 (1962).
- [27] D. Momeni, M. Nouri-Zonoz, and R. Ramezani-Arani, "Morgan-Morgan-NUT disk space via Ehlers transformation," Phys. Rev. D. 72, 064023 (2005), arXiv:gr-qc/0508036.
- [28] R. Alawadhi, D.S. Berman, B. Spence, et al. S-duality and the double copy, J. High Energ. Phys. 2020, 1-27 (2020), arXiv:1911.06797 [hep-th].
- [29] N. Arkani-Hamed, Y. Huang, and D. O'Connell, "Kerr black holes as ele-

24 REFERENCES

mentary particles," J. High Energ. Phys. 2020, 1-12 (2020), arXiv:1906.10100 [hep-th].

- [30] E. Crawley, A. Guevara, N. Miller, and A. Strominger," Black holes in Klein space," J. High Energ. Phys. 2022, 1-41 (2022), arXiv:2112.03954 [hep-th].
- [31] E. Crawley, A. Guevara, E. Himwich, and A. Strominger," Self-dual black holes in celestial holography," J. High Energ. Phys. 2023, 1-33 (2023), arXiv:2302.06661 [hep-th].
- [32] X. Ye, and S. Wei, Distinct topological configurations of equatorial timelike circular orbit for spherically symmetric (hairy) black holes, Journal of Cosmology and Astroparticle Physics 2023, 049 (2023), arXiv:2301.04786 [gr-qc].
- [33] P. V. P. Cunha, J. Grover, C. Herdeiro, E. Radu, H. Runarsson and A. Wittig, Chaotic lensing around boson stars and Kerr black holes with scalar hair, Phys. Rev. D. 94, 104023 (2016), arXiv:1609.01340 [gr-qc].
- [34] P. V. P. Cunha, and C. A. R. Herdeiro," Stationary black holes and light rings", Phys. Rev. Lett. 124, 181101 (2020), arXiv:2003.06445[gr-qc].
- [35] S. W. Wei," Topological Charge and Black Hole Photon Spheres", Phys. Rev. D. 102, 064039 (2020), arXiv:2006.02112[gr-qc].
- [36] M. Guo and S. Gao," Universal Properties of Light Rings for Stationary Axisymmetric Spacetime", Phys. Rev. D. 103, 104031 (2021), arXiv:2011.02211[gr-qc].
- [37] Y. S. Duan," The structure of the topological current", Phys. Rev. D. (1984).
- [38] P. V. P. Cunha, E. Berti, and C. A. R. Herdeiro, Light ring stability in ultracompact objects, Phys. Rev. Lett. 119, 251102 (2017), arXiv:1708.04211 [gr-qc].
- [39] S. W. Wei and Y. X. Liu," Topology of equatorial timelike circular orbits around stationary black holes", Phys. Rev. D. 107, 064006 (2023), arXiv:2207.08397[gr-qc].
- [40] A. Cisterna, K. Müller, K. Pallikaris, and A. Viganò, Exact rotating wormholes via Ehlers transformations, Phys. Rev. D. 108, 024066 (2023), arXiv:2306.14541 [gr-qc].
- [41] V. Ferrari and B. Mashhoon, Oscillations of a black hole, Phys. Rev. Lett. 52, 1361 (1984).
- [42] V. Ferrari and B. Mashhoon, New approach to the quasinormal modes of a black hole, Phys. Rev. D. 30, 295 (1984).
- [43] B. Mashhoon, Stability of charged rotating black holes in the eikonal approximation, Phys. Rev. D. 31, 290 (1985).