






















ODIN: Identifying Protoclusters and Cosmic Filaments Traced by Ly α -emitting Galaxies

VANDANA RAMAKRISHNAN ¹, KYOUNG-SOO LEE ¹, MARIA CELESTE ARTALE ², ERIC GAWISER ³, YUJIN YANG ⁴,
CHANGBOM PARK ⁵, ROBIN CIARDULLO ^{6,7}, LUCIA GUAITA ², SANG HYEOK IM ⁸, SEONGJAE KIM ⁴,
ANKIT KUMAR ², JAEHYUN LEE ^{4,5}, SEONG-KOOK LEE ^{9,10}, BYEONGHA MOON ⁴, NELSON PADILLA ¹¹,
ALEXANDRA POPE ¹², ROXANA POPESCU ¹³, HYUNMI SONG ¹⁴, PAULINA TRONCOSO ¹⁵, FRANCISCO VALDES ¹⁶, AND
ANN ZABLUDOFF ¹⁷

¹Department of Physics and Astronomy, Purdue University, 525 Northwestern Avenue, West Lafayette, IN 47907, USA

²Departamento de Ciencias Fisicas, Universidad Andres Bello, Fernandez Concha 700, Las Condes, Santiago, Chile

³Physics and Astronomy Department, Rutgers, The State University, Piscataway, NJ 08854

⁴Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

⁵Korea Institute for Advanced Study, 85 Hoegi-ro, Dongdaemun-gu, Seoul 02455, Republic of Korea

⁶Department of Astronomy & Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

⁷Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

⁸Department of Physics and Astronomy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

⁹SNU Astronomy Research Center, Department of Physics and Astronomy, Seoul National University, Seoul, Korea

¹⁰Astronomy Program, Department of Physics and Astronomy, Seoul National University, Seoul, Korea

¹¹Instituto de Astronomía Teórica y Experimental (IATE), CONICET-UNC, Laprida 854, X500BGR, Córdoba, Argentina

¹²Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA

¹³Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA

¹⁴Department of Astronomy and Space Science, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon, 34134, Republic of Korea

¹⁵Escuela de Ingeniería, Universidad Central de Chile, Avenida Francisco de Aguirre 0405, 171-0614 La Serena, Coquimbo, Chile

¹⁶NSF's National Optical-Infrared Astronomy Research Laboratory, 950 N. Cherry Avenue, Tucson, AZ 85719, USA

¹⁷Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

ABSTRACT

To understand the formation and evolution of massive cosmic structures, studying them at high redshift, in the epoch when they formed the majority of their mass is essential. The One-hundred-deg² DECam Imaging in Narrowbands (ODIN) survey is undertaking the widest-area narrowband program to date, to use Ly α -emitting galaxies (LAEs) to trace the large-scale structure (LSS) of the Universe at three cosmic epochs. In this work, we present results at $z = 3.1$ based on early ODIN data in the COSMOS field. We identify and characterize protoclusters and cosmic filaments using multiple methods and discuss their strengths and weaknesses. We then compare our observations against the IllustrisTNG suite of cosmological hydrodynamical simulations. The two are in excellent agreement, with a similar number and angular size of structures identified above a specified density threshold. We are able to recover the simulated protoclusters with $\log(M_{z=0}/M_{\odot}) \gtrsim 14.4$ in $\sim 60\%$ of the cases. With these objects we show that the descendant masses of the protoclusters in our sample can be estimated purely based on our 2D measurements, finding a median $z = 0$ mass of $\sim 10^{14.5} M_{\odot}$. The lack of information on the radial extent of each protocluster introduces a ~ 0.4 dex uncertainty in its descendant mass. Finally, we show that the recovery of the cosmic web in the vicinity of protoclusters is both efficient and accurate. The similarity of our observations and the simulations imply that our structure selection is likewise robust and efficient, demonstrating that LAEs are reliable tracers of the LSS.

1. INTRODUCTION

According to the hierarchical theory of structure formation, matter is organized into a *cosmic web*, comprised of linear filaments intersecting at nodes of high density surrounded by vast voids (Bond et al. 1996;

Springel et al. 2005; Boylan-Kolchin et al. 2009). This large-scale structure (LSS) determines how much cold gas is available to a galaxy and the likelihood of a merger or interaction with another galaxy, thereby acting as one of the fundamental drivers of galaxy evolution.

Out to $z \approx 1.5$, redshift surveys and other observational techniques have enabled the selection of samples of galaxies inhabiting clusters, groups, and filaments (e.g., Eisenhardt et al. 2008; Rykoff et al. 2014; Tempel et al. 2014; Bleem et al. 2015; Rykoff et al. 2016; Malavasi et al. 2017; Gonzalez et al. 2019). These studies show that galaxies in cluster or group environments tend to be older and more massive, and are more likely to have ceased star formation than those in the field (e.g., Peng et al. 2010; Quadri et al. 2012; van der Burg et al. 2013). Filaments may have a weaker but similar effect and may be responsible for pre-processing galaxies that are falling into cluster- or group environments (e.g., Sarron et al. 2019; Salerno et al. 2019).

At Cosmic Noon ($z \gtrsim 2$), when the global star formation rate reached its peak (Madau & Dickinson 2014), these environmental effects are predicted to be even more dramatic. In the high-density regions within the LSS, the accretion rates of infalling gas and the incidence of galaxy interactions are expected to be greatest, fostering both enhanced in-situ star formation and black hole activity. A popular hypothesis is that highly dissipative gas-rich mergers help the efficient feeding of gas into the central black hole and trigger an active galactic nuclei (AGNs), which may ultimately quench the star formation activity (e.g., Hopkins et al. 2006). These expectations are indeed in line with the heightened SF and AGN activities found in a handful of *protocluster* systems (e.g., Casey et al. 2015; Wang et al. 2016; Umemoto et al. 2015; Oteo et al. 2018; Harikane et al. 2019; Lemaux et al. 2022) as well as the emergence of quenched galaxies in such environments (e.g., Shi et al. 2021; Ito et al. 2023).

Yet, the role that LSS environment plays in galaxy formation at Cosmic Noon remains under-explored. Our limited knowledge is due to a combination of factors, including the observational limitations of measuring precise redshifts of faint, high-redshift galaxies, the inherent scarcity of massive cosmic structures, and our incomplete grasp of the indicators for the locations of these structures. The lack of readily identifiable signatures—such as a hot intracluster medium and/or a concentration of quiescent galaxies—in young, yet-to-be-virialized structures of mostly star-forming galaxies leads to a strong reliance on spectroscopy for finding protoclusters. While the lack of large, statistical samples prevent us from disentangling the effects of cosmic variance from general properties of protocluster galaxies, studies from heterogeneously selected samples can lead to seemingly conflicting conclusions. Although cosmic filaments connected to these protoclusters likely play a vital role in replenishing fresh gas for sustained star for-

mation, such medium-density features are even more difficult to discern than dense protocluster cores.

The One-hundred-deg² DECam Imaging in Narrow-bands (ODIN, Lee et al. 2024) survey is designed to obtain large and *uniformly selected* samples of protoclusters and filaments at three cosmic epochs ($z = 2.4, 3.1$ and 4.5) using Ly α -emitting galaxies (LAEs) as tracers of underlying matter distribution. As the most common electron transition in the universe, Ly α emission traces ionized and/or excited gas from star formation, black hole activity, and the gravitational collapse of dark matter halos. A large fraction of low-luminosity star-forming galaxies (which dominate the cosmic star formation rate density: Reddy & Steidel 2009) show Ly α emission (Stark et al. 2010). These LAEs tend to have lower stellar masses, younger population ages, less internal extinction than systems selected via their broadband colors, and low galaxy bias (Gawiser et al. 2007; Kusakabe et al. 2018; Weiss et al. 2021; White et al. 2024). These traits make LAEs a most efficient tracer of the underlying dark matter distribution, one which can be used to constrain cosmology (e.g., Gebhardt et al. 2021; White et al. 2024) and most massive cosmic structures (Dey et al. 2016; Huang et al. 2022). Upon completion, ≈ 600 protoclusters are expected to be discovered by ODIN, which will facilitate robust statistical investigations of the cosmic evolution of protoclusters and their galaxy inhabitants.

In this paper, we use the early ODIN data in the COSMOS field to present our selection of protoclusters and cosmic filaments. Building on the results presented in Ramakrishnan et al. (2023), we calibrate and fine-tune our LSS detection methods by carrying out careful comparisons with cosmological hydrodynamical simulations. The outline of this paper is as follows. In Section 2, we give details of the observational and simulation data. In Sections 3 and 4, we describe how we construct LAE surface density maps and how we use them to detect protoclusters and filaments. We validate our procedures and interpret the results using the mock data created from the simulations in Section 5. Finally, the properties of our observationally selected structures are discussed in Section 6 followed by a summary of our findings in Section 7. Throughout this paper, we assume a Planck cosmology (Planck Collaboration et al. 2016): $\Omega_\Lambda = 0.6911$, $\Omega_b = 0.0486$, $\Omega_m = 0.3089$, $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $h = 0.6774$. Distances are given in units of comoving Mpc (cMpc) unless noted otherwise.

2. OBSERVATIONAL AND SIMULATION DATA

2.1. The ODIN survey

The ODIN survey is conducting the widest-area deep narrowband imaging program to date, using three custom narrowband filters ($N419$, $N501$ and $N673$) to identify redshifted Ly α emission. In this work, we make use of the Year 1 ODIN data taken with the $N501$ filter ($\lambda_C/\Delta\lambda = 5014/75$ Å $\bar{z}/\Delta z = 3.12/0.06$) in the extended COSMOS field. For more details about the survey fields and observing strategy, we refer interested readers to Lee et al. (2024).

The method for selecting ODIN LAEs is detailed in Firestone et al. (2023). Briefly, we identify LAEs as $N501$ -detected sources exhibiting a narrowband excess over the continuum, corresponding to a rest-frame equivalent width of 20 Å. The continuum magnitude is calculated as a weighted combination of the magnitude in two broadband filters (g and r for $N501$ LAEs). We exclude sources flagged for saturated pixels or other image defects and those close to bright stars. The continuum is estimated using broadband data from the Hyper Suprime-Cam Subaru Strategic Program (HSC SSP: Aihara et al. 2018a,b) second data release (Aihara et al. 2019). Our resulting LAE sample comprises 6,056 sources over ≈ 7.5 deg². While dedicated spectroscopic follow-up programs with Keck, Gemini, and DESI are ongoing, more than 25% of our $N501$ -selected LAEs have been targeted to date. Of the sources that yielded a redshift, $\approx 97\%$ are confirmed as LAEs (Ramakrishnan et al. in prep.).

2.2. Building Mock ODIN Observations with TNG

To build a concrete framework in which we can interpret our observations, we use the IllustrisTNG300-1 simulation (hereafter TNG300: Nelson et al. 2019; Pillepich et al. 2018a,b) and define our mock LAE samples. TNG300 provides the largest simulation volume (302.6 cMpc on a side) of all the IllustrisTNG simulations. Given the rarity of massive galaxy (proto)clusters, this is especially crucial for our work. All TNG simulations assume the Planck cosmology (Planck Collaboration et al. 2016). The surface area viewed along the X, Y, or Z direction is $\approx 90,000$ cMpc², well matched to the angular extent of the ODIN data in the COSMOS field, $\approx 95,500$ cMpc², and the simulation box is several times larger than the ODIN radial extent of ~ 60 cMpc at $z = 3.1$. TNG300 has a baryon mass resolution of $1.1 \times 10^7 M_\odot$ and a dark matter mass resolution of $5.9 \times 10^7 M_\odot$. In our analysis we employ galaxies with stellar mass greater than $10^7 M_\odot$, corresponding to a halo mass of $\gtrsim 10^9 M_\odot$; as do not employ any stellar or gas physics in our analysis, the resolution is sufficient for our purposes.

In constructing mock LAE samples, our primary goal is to mimic the spatial distribution of the ODIN LAEs as closely as possible so that we can evaluate their utility as tracers of the LSS. Understanding the complex behavior of Ly α radiative transfer requires vastly higher resolution simulations and thus is outside the scope of this work. In what follows, we describe how we select mock LAEs from the TNG galaxies.

First, we match the 75 Å (60 cMpc at $z = 3.1$) full-width-at-half-maximum of $N501$ by creating cosmic ‘slices’ from the TNG $z = 3$ snapshot with ≈ 80 cMpc in line-of-sight thickness. The filter and thus the window function are not a perfect top hat in shape. To emulate this effect, we assign the LAE selection probability to match the shape of the filter transmission function. In all things being equal, the probability of being selected as a mock LAE is 1, 0.5, and 0 at the distance of 0, 30, and 40 cMpc from the center of the slice, respectively. In practice, the probability of an LAE being detected at a given position along the redshift direction also depends on its line luminosity, as bright LAEs are more likely to be detected when they fall on the wings of the filter than fainter ones, and hence are detected over a larger volume. However, since we are interested only in the average number density of *all* LAEs, irrespective of line luminosity, this does not significantly affect our analysis.

Second, we aim to reproduce the small- and large-scale clustering of LAEs. The galaxy bias for $z \sim 3$ LAEs is relatively low at $b \lesssim 2$ (Gawiser et al. 2007; Ouchi et al. 2008; White et al. 2024), suggesting that LAEs have low stellar mass content and are hosted by low-mass halos (e.g., Guaita et al. 2010; Kusakabe et al. 2018). To match the small-scale clustering, we must simultaneously match the LAE overdensity distribution across the field; the details of this measurement are presented in Section 3. Motivated by the findings of Hagen et al. (2014), we model the stellar masses of the mock LAEs as a lognormal distribution: i.e., $\log M_*/M_\odot$ is a Gaussian function, with mean and standard deviation (μ, σ). Lowering μ values would shift the host halos to lower masses (thus lower bias). A larger scatter σ (while fixing μ) would permit both higher- and lower-mass halos to host LAEs, thereby changing how LAEs trace these halos. We find that $\mu = 8.75$ and $\sigma = 0.75$ yield the best fit to our surface density measurements. The left panel of Figure 5 shows the resultant LAE overdensity map relative to the data. Our best-fit parameters are fully consistent with the stellar mass distribution of LAEs based on SED fitting reported by Hagen et al. (2014), $\log(M_*/M_\odot) = 8.97^{+0.60}_{-0.71}$. They are also similar to the

result of Vargas et al. (2014), who find $\log(M_*/M_\odot) = 8.45^{+0.72}_{-0.67}$ for LAEs at $z = 2.1$.

Finally, we match the measured sky density of LAEs. The $N501$ LAE surface density is 0.22 arcmin^{-2} . In comparison, the number of TNG galaxies selected based on the above criteria is typically ten times greater. Additionally, we assume 10% of our LAEs may be contaminants. This number is based on the fraction of our spectroscopic targets in the two faintest bins ($N501=24.5\text{--}25.0$ and $25.0\text{--}25.5$ AB) that yielded a redshift, which is 94.3% and 85.9%, respectively (Ramakrishnan et al. in prep.). By doing so, we are assuming that all sources that did not result in a redshift are interlopers. Our spectroscopic success rate (i.e., the fraction of DESI-confirmed sources that yield a redshift for which it is within the expected range) is 97%, so our estimate is conservative. Changing it to 5% does not change our results. From the selected TNG galaxies, we randomly choose a subset whose number is equal to 90% of the LAE density; assuming that LAE sample contaminants are unclustered, the remaining 10% is drawn as a purely random distribution within the same region.

To compare directly with our observations, we utilize the progenitors of 30 of the most massive galaxy clusters in the final simulation box of TNG300, ranked based on their total stellar mass. The stellar mass, halo mass and Group ID of these clusters at $z = 0$ are found in Andrews et al. (2024). We identify the protoclusters by tracing the main progenitor branch of the merger trees of the $z = 0$ clusters to their $z = 3$ predecessors. The descendant masses of the protoclusters ranges from $1.5 \times 10^{15} M_\odot$ for the most massive system to $2 \times 10^{14} M_\odot$ for the least massive. We generate cosmic slices, aligning the mid-point of each with that of the protocluster using the periodic boundary conditions of the simulation. Slicing the TNG volume along the X, Y and Z axes produces 90 distinct slices for comparison. In a given slice, typically a few other protoclusters from the top 30 most massive sample are included although, depending on their positions in the redshift direction, they may be only partially represented.

3. TRACING LARGE-SCALE STRUCTURE

This section aims to delineate the large-scale structure in our observations by constructing the LAE surface density maps. We employ two methods; first, smoothing over the LAE positions with a fixed Gaussian kernel; second, constructing the Voronoi diagram of the LAEs. These two methods are summarized below but are discussed in more detail in Ramakrishnan et al. (2023).

3.1. Gaussian smoothing

The simplest approach to measure the surface density across a field involves smoothing over the positions of the galaxies within it using a fixed-size kernel. We employ a two-dimensional Gaussian kernel with FWHM 10 cMpc. This FWHM is decided following the methodology of Bădescu et al. (2017), also utilized in Ramakrishnan et al. (2023). The kernel size is chosen such that the resultant surface density map maximizes the total probability at the positions of the LAEs in the real data. This is achieved through a leave-one-out cross-validation, where the likelihood of finding a point at the location \vec{r}_j of the j th data point is estimated as:

$$p(\vec{r}_j) = \sum_{i \neq j} \frac{1}{\sqrt{2\pi}\sigma} \exp \frac{-(\vec{r}_i - \vec{r}_j)^2}{2\sigma^2} \quad (1)$$

The optimum σ value is the one which maximizes $\prod_j p(\vec{r}_j)$.

Before creating the LAE surface density map, we fill in the holes left by removing the sources near bright stars and image defects with uniformly distributed random points with surface density matched to that of the LAEs. After convolving with the kernel, the overdensity map is computed by dividing the Σ_{LAE} map by the mean surface density, $\bar{\Sigma}_{LAE}$:

$$(1 + \delta_{LAE}) = \frac{\Sigma_{LAE}}{\bar{\Sigma}_{LAE}} \quad (2)$$

The mean and standard deviation are determined by fitting the LAE surface density distribution with a Gaussian function, $\exp[-(\Sigma_{LAE} - \bar{\Sigma}_{LAE})^2/2\sigma^2]$. The fit is restricted to within $\pm 1.5\sigma$ (after iterative sigma-clipping) to ensure that our estimate is not biased by the presence of multiple high LAE overdensities, which show up at the high $(1 + \delta_{LAE})$ end of the distribution.

The left panel of Figure 1 shows the resultant $(1 + \delta_{LAE})$ map, which we will refer to as the GS map, hereafter. It shows overdense regions that are both strongly clustered and highly irregular in shape, consistent with expectations from the hierarchical theory of structure formation, in which smaller structures continuously merge to form larger ones, a phenomenon supported by hydrodynamical simulations (e.g., Boylan-Kolchin et al. 2009). The three most prominent overdensity complexes, highlighted by red boxes and labeled as complexes A, B, and C, were discussed in detail in Ramakrishnan et al. (2023).

Smoothing with a fixed kernel is most effective at identifying structures with a size and shape similar to that of the kernel itself. Thus, the Gaussian smoothing method may not adequately capture non-isotropic features, potentially resulting in an underestimation of the significance of many observed structures. To address this, we explore tessellation-based methods in Section 3.2.




Figure 1. *Left:* Gaussian smoothed (GS) LAE surface density map. Black, green, white, and cyan contours indicate LAE overdensities of 2 , 3 , 4 , and 5σ significance respectively; the corresponding values of $(1 + \delta_{LAE})$ are marked on the color bar. The three largest overdensity complexes are highlighted in red. *Right:* VT-based density map. White contours indicate overdensities of 3σ significance. The LSS revealed by the surface density maps is highly clumpy and irregular. Non-spherical/irregular overdensities are better delineated at fixed significance in the VT map than the GS map. The clover shape of the surface density maps arise from the HSC-SSP data, which consists of four Deep and one central UltraDeep pointing in the COSMOS field.

3.2. Voronoi tessellation

Tessellation-based density estimates offer the advantage of being scale-independent and do not assume any specific shape or size for the underlying structures. The two algorithms most commonly used in the literature are the Voronoi tessellation (e.g., Dey et al. 2016; Cucciati et al. 2018; Lemaux et al. 2018; Hung et al. 2020) and the Delaunay tessellation (e.g., Malavasi et al. 2021; Sarron & Conselice 2021). Darvish et al. (2015) find, through analysis of a simulated dataset, that the Delaunay tessellation fares more poorly at estimating the ‘true’ surface density value at a given point as compared to the Voronoi tessellation. Additionally, the method tends to overestimate the surface density in overdense regions. Considering these results, we opt for the Voronoi tessellation algorithm as outlined below.

In a two-dimensional case, the Voronoi tessellation (VT) divides a plane into distinct cells based on the positions of a set of generating points e.g., the sky locations of LAEs. The Voronoi cell of each generating point includes all regions in the plane that are closer to it than to any other generating point. Consequently, the area of each cell is a measure of surface density: cells associated with LAEs in overdense regions will be small due to the proximity of numerous other LAEs, whereas those in underdense regions will be larger. Since each

cell contains a single LAE, its LAE surface density Σ_i is given by,

$$\Sigma_i = \frac{1}{A_i} \quad (3)$$

where A_i is the area of the i th cell. As before, the masked regions are filled in before constructing the Voronoi diagram.

In Figure 1, we show the VT $(1 + \delta_{LAE})$ map together with the GS map. In both panels, white contours delineate 3σ overdensities, with σ denoting the field density fluctuations computed from the $(1 + \delta_{LAE})$ distribution. While both maps detect the most significant structures, the VT method detects them at higher significance. As expected, the method also identifies a greater number of overdensities by capturing irregular or anisotropic features, as expected.

4. FEATURES OF THE LSS IN OBSERVATIONS: PROTOCLUSTERS AND FILAMENTS

Galaxy protoclusters represent some of the most striking features of large-scale structure at high redshift and are expected to be observed as significant galaxy overdensities spanning ≈ 10 cMpc in scale (Chiang et al. 2013). According to the hierarchical theory of structure formation, massive halos hosting protoclusters are connected to filaments of the cosmic web (e.g., Kuchner et al. 2022) along which pristine gas is being ac-

creted to feed star formation. Indeed, these predictions align qualitatively with the features observed in our $(1 + \delta_{LAE})$ map. Multiple overdensities with angular scales of 5–10 cMpc are evident in Figure 1 (right), frequently clustered together to form *complexes* comprising 2 to 5 adjacent overdensities. Additionally, regions of high overdensity, where $(1 + \delta_{LAE}) > 3$, are interconnected by more moderate ‘bridges’ with $(1 + \delta_{LAE}) = 2$ –3. Several features show a distinctly linear morphology, reminiscent of cosmic filaments (for example, the northwestern corner of “Complex A” and the southeastern corner of “Complex C”, see Ramakrishnan et al. 2023, for more detail).

Motivated by these observations, our next objective is to pinpoint the locations of protoclusters and filaments of the cosmic web using LAEs as tracers. By directly comparing our observations with TNG300 predictions, we will also test how robust our protocluster candidates are and measure their key properties.

4.1. Selecting protoclusters from density maps

Protoclusters are generally defined as regions that will evolve into a virialized structure with masses $\gtrsim 10^{14} M_{\odot}$ by $z = 0$ (e.g., Overzier 2016). In distant look-back times, protoclusters are loosely bound regions comprised of multiple dark matter halos. This poses a challenge in measuring their physical extent. In this study, we define a protocluster as a region exhibiting significant size and overdensity, unlikely to occur by chance alone, and containing sufficient mass that could plausibly collapse to form a single halo with mass greater than or equal to $10^{14} M_{\odot}$ by $z = 0$.

According to this operational definition, selecting protoclusters entails identifying contiguous regions above a pre-set overdensity threshold and minimum area. This procedure mirrors source detection in pixelated astrophysical images. Therefore, adopting approaches similar to those outlined by Lemaux et al. (2018); Hung et al. (2020); Sarron & Conselice (2021), we utilize source detection software on our density maps for protocluster selection.

We begin by pixelating the surface density maps by interpolating them over a two-dimensional grid of positions with pixel size $3''6$. Our grid size, corresponding to 115 kpc at $z = 3.1$, is small enough to clearly delineate the boundaries of the structures, as it is two orders of magnitude smaller than the typical size of a protocluster. For source detection, we use SEP (Barbary 2016), a Python implementation of the SExtractor (Bertin & Arnouts 1996) software. The number of detected structures depends strongly on the detection threshold (DETECT_THRESH) and minimum area (DETECT_MINAREA).

Our goal is to optimize these parameters to maximize the identification of robust candidates while minimizing the inclusion of spurious objects resulting from chance alignments of LAEs.

To assess contamination, we randomly select a subset of *all* continuum and line-emitting sources detected within the *N501* image, matching the number of LAEs in the field. We then generate GS and VT maps for these random points following the same procedure as for the LAE surface density maps. Since these points are distributed across a wide range of redshift, any overdensity observed in these ‘random maps’ is unlikely to be genuine but rather the result of chance alignments. Hence, the contamination fraction for a specific set of detection parameters can be approximated by dividing the number of sources detected in the random map by the number detected in the LAE surface density map. This evaluation is depicted in the top and bottom right panels of Figure 2 for the VT and GS map, respectively, with the contamination fraction averaged over 30 iterations.

For the GS-selected protoclusters, the contamination fraction is low for all sets of detection parameters used. Our fiducial setup, DETECT_THRESH of 3σ and DETECT_MINAREA of 4600 pixels ($\simeq 60$ cMpc²), yields a contamination fraction of $< 0.1\%$ where σ is the (sigma-clipped) standard deviation of δ_{LAE} . Compared to the GS map, the pixel-to-pixel density fluctuations are much greater in the VT map, introducing much higher noise in structure detection. We mitigate this effect by smoothing the map with a 5 cMpc FWHM Gaussian kernel, large enough to reduce the noise but small enough to avoid adjacent structures blending into one considering that a typical diameter of a protocluster at $z \sim 3$ is 10 cMpc. We choose DETECT_THRESH of 4.5σ and DETECT_MINAREA of 3000 pixels ($\simeq 40$ cMpc²), yielding the contamination rate of 20%.

Figure 3 (top and middle rows) shows the protoclusters identified from the GS and VT-based maps within the highlighted Complexes A, B, and C. The GS map predominantly captures the largest structures, with smaller and more irregular formations remaining undetected; nonetheless, the identified structures exhibit high robustness. In contrast, the VT map reveals numerous structures overlooked by the GS map but is also subject to higher noise levels and is more susceptible to chance alignments of LAEs. The detailed comparison of different detection methods is presented in Section 6.

4.2. Selecting protoclusters with HDBSCAN

Density-based clustering techniques, as a subset of unsupervised machine learning algorithms (Kriegel et al. 2011), discern clusters by identifying regions of high




Figure 2. Number of candidates and contamination fraction for protoclouds selected from the VT map (top) and GS map (bottom), for different density thresholds and minimum areas. The density threshold is expressed in units of $N\sigma$ above the field surface density. White stars indicate the detection settings adopted for this work. While the VT map suffers from higher contamination than the GS map, it also detects far more candidates.

point density in data space, set apart by areas of low density. These methods can be applied to pinpoint protoclouds from the clustering of the LAEs. Here, we employ the Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN) algorithm (Campello et al. 2013). HDBSCAN offers the advantage of detecting clusters with varying densities, regardless of their shape, provided they exceed a specified size threshold. Briefly, this algorithm measures density at each location, finding clusters as peaks in the density distribution separated by troughs. Density estimation is based on the distance to the K th nearest neighbor.

In HDBSCAN, each point is classified into *clusters* or *noise* based on two parameters: the K th nearest neighbor distance and the minimum cluster size. The latter dictates the minimum number of data points constituting a cluster. In the Python implementation of the HDBSCAN algorithm (`hdbscan`: McInnes et al. 2017), these parameters are denoted as `min_samples`

and `min_cluster_size`, respectively. In our setup, we opt for values of 15 and 10, respectively (i.e., a minimum cluster size of 15 LAEs and the 10th nearest neighbour distance; see Appendix A). Additionally, we require that each cluster exhibit a median surface density above a specified threshold. This surface density is calculated as the number of LAEs divided by the enclosed area, computed as the sum of the areas of all Voronoi cells encompassing the cluster members. We set the threshold at 0.23 arcmin^{-2} (i.e., $\gtrsim 1\sigma$ above the field mean), which yields a contamination rate of $\approx 20\%$ similar to that for the VT sample. The HDBSCAN-detected protoclouds are illustrated in the bottom row of Figure 3. Symbols of the same color denote membership within the same structure, while gray circles represent LAEs outside of protoclouds. They are also shown in Figure 4.

4.3. Cosmic filaments with DisPerSE




Figure 3. Protocluster candidates selected from the GS map (top row), the VT map (middle) and with HDBSCAN (bottom). Red crosses indicate the geometric centers of the selected overdensities. Yellow swathes indicate the extent of the protoclusters in the upper two rows. In the bottom row, colored points indicate LAEs classified as belonging to a cluster by HDBSCAN, enclosed by an ellipse for clarity, while gray points are classified as noise.

The cosmic web at low redshift ($z \lesssim 1$) has been investigated in detail through multiple surveys (e.g. [Alpaslan et al. 2014](#); [Tempel et al. 2014](#); [Malavasi et al. 2017](#)). These studies have yielded valuable insight into the complex interplay of filaments and the clusters at the nodes of the cosmic web. Notably, [Sarron et al. \(2019\)](#) found evidence that galaxies accreted onto clusters along filaments show signs of pre-processing. Moreover, several studies have observed that galaxies within filaments show higher quenched fractions than those within the field ([Malavasi et al. 2016](#); [Kraljic et al. 2017](#); [Laigle](#)

[et al. 2018](#)). At higher redshift, the role of the cosmic web on galaxy formation remains largely observationally unconstrained. Motivated by the discernible presence of filamentary structures in the LAE surface density map, we aim to detect cosmic web filaments in our observations in this section. We explore the reliability of our filament detection in Section 5.3.

To identify cosmic filaments, we use the Discrete Persistent Structure Extractor (DisPerSE; [Sousbie 2011](#); [Sousbie et al. 2011](#)), a widely utilized tool in both observational (e.g., [Sarron et al. 2019](#); [Malavasi et al.](#)




Figure 4. Protoclusters selected from the GS map (top left), VT map (top right), with HDBSCAN (bottom left) and filaments (bottom right) overlaid on the VT map. The red dashed region in the bottom left panel highlights a structure which is missed by the VT and GS maps but is detected with HDBSCAN. The filaments correspond to regions of moderate ($\sim 2\text{--}3\sigma$) overdensity, and connect protoclusters together.

2017, 2020) and simulation studies (e.g., Byrohl & Nelson 2023). DisPerSE utilizes the Delaunay Tessellation Field Estimator (Schaap & van de Weygaert 2000; van de Weygaert & Schaap 2009; Cautun & van de Weygaert 2011) to create a discrete representation of the density field based on a given set of points. It then locates critical points within this density field approximation and defines filaments as arcs connecting saddle points to maxima. The filaments are constructed by creating short segments tangent to the gradient of the density field at each point. To account for data noise, DisPerSE

employs the concept of persistence, which measures the density contrast between critical points defining a filament. Persistence represents the range of density thresholds over which a filament connecting two critical points remains significant relative to the noise. This significance is expressed in units of σ , where an $N\sigma$ filament corresponds to a probability under a 1D Gaussian distribution. We stress that this method of measuring the persistence level does not imply the use of a Gaussian distribution within DisPerSE but rather is purely an expression of the likelihood of the filaments arising from

noise. For example, a persistence level of 3σ (2σ) means that the extracted filament has a 99.7 (95.5)% probability of being a true feature.

After filling in the voids left by star masks, we run DisPerSE with a persistence of 2σ . The resultant filament network is illustrated in the bottom right panel of Figure 4 as red curves superimposed on the VT density map in greyscale. The figure reveals a complex web extending over the entire field. Several previously noted linear overdensities (e.g., within complexes A and C) are part of this network. Moreover, the filaments appear to converge at the locations of the extended complexes, interconnecting the individual protoclusters within. Indeed, every protocluster is connected to at least one filament, with several positioned at the convergence of multiple filaments. Our findings are fully consistent with the hierarchical picture wherein protoclusters occupy the nodes of the cosmic web.

Given that the line-of-sight window function set by the $N501$ filter is substantially larger than the dimensions of individual protoclusters and cosmic filaments, our 2D-based detection algorithm is expected to include false positives arising from random noise fluctuations. In Section 5.3, we show that while false detection - in particular, of filaments - does indeed occur, all filaments around overdense protoclusters are robustly identified. Protocluster detection is even more secure.

5. COSMIC STRUCTURES IN SIMULATIONS: BUILDING EXPECTATIONS WITH TNG

Cosmological hydrodynamical simulations, such as TNG300, offer invaluable insights not readily accessible through direct observations. By facilitating connections between observable traits of protoclusters and cosmic filaments with more fundamental attributes and allowing us to track their evolution across cosmic time, simulations provide a crucial context in which we understand our data. In this section, we compare the ODIN protocluster and filament samples with those derived from the TNG300 simulation to gain an understanding of the physical properties of the structures detected in Section 4.

5.1. Comparison of observations and simulations

In the left panel of Figure 5, we compare the LAE surface density distribution of the real data with the TNG slices. As detailed in Section 2.2, the latter are constructed from TNG300 and are designed to match the ODIN filter transmission as well as the LAE surface density and clustering bias. Multiple realizations are averaged over to result in the surface density distribution shown for the simulation. The high-end tail of the LAE

surface density distribution, which represents the field's highest density regions corresponding to protoclusters and filaments, is nearly perfectly reproduced.

In Figure 6, we show the VT map of both data and one $z = 3$ TNG slice side by side. The displayed slice is chosen at random and is centered on a massive protocluster with descendant mass $5.5 \times 10^{14} M_\odot$ (Group ID=13 at $z = 0$: cyan cross). It also contains five additional structures (yellow crosses) that will evolve into a cluster with mass greater than 2×10^{14} by $z = 0$, as well as several objects that will evolve into less massive clusters (green crosses). Indeed the majority of the largest overdensities seen within the slice are associated with protoclusters. The two maps are remarkably similar in that both display structures of similar sizes and irregular morphologies, arranged into extended complexes. We highlight three such similar regions (Complexes A1, B1, and C1 in the real data, and A2, B2, and C2 in the simulations, shown by dashed-line boxes). These mock structures are also connected by cosmic web filaments.

The distributions of the total transverse area and median LAE surface density of protoclusters detected in real data and simulation span a similar range, demonstrating good agreement given the constraints of modeling. This is illustrated in Figure 5. The median number of protoclusters (averaged over 90 TNG slices) is 37 compared to 33 detected in our data. They correspond to the protocluster surface density of $(4.1 \pm 0.6) \times 10^{-4}$ and $(3.3 \pm 0.6) \times 10^{-4} \text{ cMpc}^{-2}$, respectively. The uncertainties represent the shot noise only and thus should be considered as a lower limit. The excellent agreement between our data and TNG simulations suggests that the massive cosmic structures identified with current and future ODIN data are robust.

5.2. Calibration of descendant mass estimate

Estimations of the descendant mass allow us to establish direct connections between high-redshift protoclusters and present-day galaxy clusters, thereby tracing the evolution of massive cosmic structures and their galaxy populations across cosmic time. Following the methodology outlined by Steidel et al. (1998), we compute the descendant mass of ODIN protoclusters by assuming that the mass contained within the overdensity region will collapse into a cluster-sized halo by $z = 0$, with a total mass $M_{z=0}$ given by

$$M_{z=0} = \rho_m V_{PC} = \rho_{0,z} (1 + \delta_m) V_{PC} \quad (4)$$

where ρ_m and δ_m are density and overdensity of matter, respectively. V_{PC} is the protocluster volume, and $\rho_{0,z}$ is the mean density of the Universe at redshift z . If the




Figure 5. *Left:* LAE surface density distribution of the TNG300 slices, in comparison to that of the observed LAEs. The red line shows the median distribution across all 90 slices while the shaded region shows the spread. The observed and simulated LAE distributions are very similar, particularly at the high end which represents massive structures. *Middle and right:* Area and median LAE overdensity (δ_{LAE}) for the protocluster candidates detected in the 90 TNG300 slices compared to those of the observationally detected protocluster candidates. Blue/red vertical lines indicate the median value for the ODIN data/TNG300. The properties of the structures are similar in the observations and simulation.

galaxy bias, b_g , is known, Equation 4 becomes:

$$M_{z=0} = \left(1 + \frac{\delta_g}{b_g}\right) \rho_{0,z} V_{PC} \quad (5)$$

where δ_g is galaxy overdensity. We assume $b_g = 1.8$ (Ga-wiser et al. 2007; White et al. 2024) and fix δ_g to the median LAE overdensity within the protocluster region. Since we do not have any redshift information for the majority of our LAEs, we assume that the size of a protocluster in the line-of-sight direction is similar to that in the transverse direction and approximate the volume of a protocluster from its area (A_{PC}) as $V_{PC} = A_{PC}^{1.5}$: i.e., the shape of a protocluster is approximated as a cube. $M_{z=0}$ is then estimated as,

$$M_{z=0} = \left(1 + \frac{\delta_g}{b_g}\right) \rho_{0,z} A_{PC}^{1.5} \quad (6)$$

The relation $V_{PC} = A_{PC}^{1.5}$ is a good estimate for an isotropic structure. As we have already seen, many of our protoclusters are strikingly non-spherical in shape, and thus the above relation may be a poor approximation. If a protocluster is extended in the transverse dimension, it would be observed as a more modest overdensity with a larger angular extent. Conversely, a structure stretched along the line-of-sight direction is expected to be more compact with a larger overdensity. Using TNG-detected structures, we quantify the detection rate and the uncertainty in mass estimates due to the lack of information in the third dimension.

Figure 5 shows that there is a good agreement between TNG-selected and observational protoclusters for both angular sizes (middle) and median LAE overdensity (right). We also assess the likelihood of the most

30 massive cluster progenitors to be selected as a protocluster candidate when viewed along the X-, Y-, and Z directions. To this end, we only consider the structures for which: i) the measured (angular) position center lies within 10 cMpc of the true center; and ii) the line-of-sight position center is within 25 cMpc of the center of a given TNG slice. For reference, the FWHM of all slices is 60 cMpc and a typical size of a protocluster is ≈ 10 cMpc.

Based on this definition, the median recovery rate of the protoclusters is 60%. The progenitor of the most massive cluster is always detected while the progenitors of the next 4 most massive clusters are detected $> 90\%$ of the time. As expected, the likelihood of finding smaller structures depends on the sightlines being favorable or adverse to robust detection. If we lower the threshold for our VT-based protocluster selection from 4.5σ to 3.5σ , the recovery rate of the protoclusters increases to 80%. The higher success rate comes at the price of a much greater contamination of $\sim 45\%$, compared to $\approx 20\%$ for our fiducial setup.

In the left panel of Figure 7, we compare the *estimated* and true descendant mass, $M_{est,z=0}$ and $M_{200,z=0}$ for the recovered clusters in the 90 slices. The figure illustrates that $M_{est,z=0}$ tends to underestimate the true mass about 80% of the time. The median $M_{est,z=0}/M_{true,z=0}$ ratio is 0.35, as indicated by the solid black line. This implies that the cosmic volume that ends up in a galaxy cluster by $z = 0$ is much greater than what we identify in the data as significant galaxy overdensities likely associated with a protocluster.




Figure 6. VT-based surface density map of our observed LAEs versus that of a 60 cMpc thick slice of the TNG300 simulation box at $z = 3$. Red lines in the top panels represent filaments selected with DisPerSE. The cyan cross in the top right panel indicates the position of the massive protocluster on which the slice is centered (Group ID = 13 at $z = 0$), while yellow crosses show the other massive protoclusters which fall within the slice (Group IDs = 1, 2, 7, 11, 23 at $z = 0$). Green crosses show the progenitors of less massive clusters which fall within the slice. Some overdensity complexes (A1, B1, C1; A2, B2, C2) are highlighted and zoomed in on in the bottom panels. The two maps show visible similarities.




Figure 7. $M_{z=0}$ values estimated from Equation 6 compared to the true values for the progenitors of the 30 most massive clusters in the TNG300 box. The left panel (blue points) indicate the masses estimated from the 2D projection of the LAE overdensity corresponding to the protocluster, with the LAEs matched in number density to observations. The dashed line indicates a one-to-one relation between estimated and true descendant masses, while the solid line indicates a relation of $M_{est,z=0} = 0.35 M_{200,z=0}$. Highlighted points show cases where the protocluster was merged with another, more massive object by the detection. The right panel (red points) indicate the masses estimated from the 3D positions of the cluster members (see Section 5.2 for more details). Shaded regions indicate the range of the 1 and 2σ scatter of the estimated masses. While the 2D mass estimate is subject to considerable scatter, primarily arising from the loss of information in the z direction, it correlates well with the true descendant mass.

Our findings are consistent with the results from Chiang et al. (2013), who, based on the semianalytical models implemented in the Millennium simulations, found that about 40% of the total mass at $z = 0$, M_{today} , is enclosed within a 6 (8) cMpc-radius sphere for Virgo- and Coma-sized protoclusters with $M_{\text{today}} = (3 - 10) \times 10^{14} M_{\odot}$ and $> 10^{15} M_{\odot}$, respectively. In comparison, the median effective radius of our simulated protoclusters (computed as $(A_{PC}/\pi)^{0.5}$) is ~ 5 cMpc. To account for this effect, we correct the estimated masses by a factor of 0.35.

Figure 7 shows that the 2D-based mass estimates yield a considerable scatter of 0.49 dex. The most extreme outliers arise from cases where two or more protoclusters are merged together during detection, particularly for the protoclusters with $M_{200,z=0} \leq 10^{14.5} M_{\odot}$; these points are highlighted with a black outline in Figure 7. Even excluding these cases, the scatter remains high at ~ 0.44 dex. We ascertain that the scatter is a result of the loss of information in the z direction as follows. We perform the 3D Voronoi tessellation of the 3D volume centered on the 30 massive clusters, this time, using *all* the galaxies with $M_{\text{star}} > 10^7 M_{\odot}$. Using the subhalo merger trees of the $z = 0$ cluster members, we also identify the ‘member galaxies’ in the $z = 3$ snapshot. The LAE overdensity and the protocluster region are com-

puted similarly to the 2D case but Voronoi cells are now 3D polyhedra instead of 2D polygons, while galaxy overdensity is measured as the median LAE overdensity of the Voronoi cells containing member galaxies. V_{PC} is obtained by summing over these Voronoi cells.

The resultant mass estimate is shown in the right panel of Figure 7. As expected, the scatter of the 3D $z = 0$ mass estimates is considerably smaller, at ~ 0.15 dex, than in the 2D case, once we have eliminated all uncertainties regarding the cluster membership and the information on the third dimension. The estimated descendant mass is also significantly less underestimated than in the 2D case, being $\sim 85\%$ of the true value. It is noteworthy that the *intrinsic* scatter of 0.15 dex is comparable to 0.2 dex estimated by Chiang et al. (2013) in converting galaxy overdensity measured within a $(15 \text{ cMpc})^3$ cubic window into descendant mass.

When we compare the ‘true’ volume found with the 3D Voronoi tessellation to that derived from the projected protocluster area, we find that the scatter of the latter (≈ 0.52 dex) is sufficient to account for the entirety of the uncertainty in the mass estimate. This suggests that the uncertainty in $M_{z=0}$ is dominated by the volume estimation and that the isotropic assumption is less than ideal for protoclusters at high redshift. This result is not surprising, but it does quantitatively demonstrate

the need for large-scale spectroscopic follow-up of protocluster candidates. However, we also emphasize that the 2D mass estimation is correct in an *average* sense, and statistical analyses for a large sample of protocluster candidates which make use of this mass estimate will be robust.

5.3. The fidelity of filament recovery

At high redshift, active star formation and galaxy growth taking place in protoclusters are likely supported by cool gas transported along filaments of the cosmic web (e.g., Dekel et al. 2009; Daddi et al. 2021). The wide area coverage of ODIN presents a unique opportunity to study protoclusters in concert with the surrounding cosmic web, provided that we can accurately recover the filaments. As they are even more extended structures than protoclusters, the effect of projection along the line-of-sight on filament recovery must be considered separately for filaments. In this section, we use the TNG300 slices to examine how well the observed 2D filament network corresponds to the true underlying 3D network. We focus specifically on filaments in the vicinity of protoclusters, to understand how likely it is for a filament that appears near a protocluster in projection to be physically connected to the protocluster.

We make use of the 90 slices of the TNG300 full box at $z = 3$ described in Section 2.2. For each slice, we use DisPerSE to extract filaments using the positions of the mock LAEs matched to ODIN, projected along the slice axis. This is henceforth referred to as the ‘2D skeleton’ or ‘2D filaments’. In order to create a comparison set of filaments which represents the ‘true’ cosmic web, we also run DisPerSE on the 3D positions of the slice galaxies with $10^7 M_\odot < M_* < 10^{12} M_\odot$, henceforth referred to as the ‘3D skeleton’ or ‘3D filaments’. We use a persistence threshold of 2σ to extract the 2D skeleton, identical to that used on the observed LAEs. For the 3D skeleton, we use multiple persistence thresholds of 5, 6, and 7σ in order to compare the 2D skeleton against 3D features of varying significance.

In Figure 8, we show the dark matter distribution in one of the TNG300 slices overlaid with the 2D and 3D skeletons. The similarity between the DisPerSE filaments (both 2D and 3D) and the underlying dark matter distribution is clear, as is the fact that the 2D and 3D skeletons mirror each other closely. However, the 2D skeleton does not perfectly reproduce the true 3D cosmic web. To quantify the extent to which the cosmic web can be recovered by our observational data, we compare the 2D filaments to the 3D skeleton. We do this following a similar procedure to Sarron et al. (2019) (see also Laigle et al. 2018; Kuchner et al. 2022),

by matching the individual segments out of which DisPerSE constructs the 2D and 3D skeletons (see Section 4.3) to each other. We measure the (projected) separation between the midpoints of each segment of the 2D skeleton and those of the 3D skeleton. We denote the minimum distance *from* a segment of the 2D filament network *to* one of the 3D network as $d_{2d \rightarrow 3d}$, and the inverse as $d_{3d \rightarrow 2d}$. This mapping is illustrated in the left panel of Figure 9. The former measurement enables us to determine which of the 2D filaments represent true features of the 3D cosmic web, while the latter allows us to determine which features of the underlying filament network are successfully recovered by observations.

In the top right panel of Figure 9, we show histograms of $d_{2d \rightarrow 3d}$ for the three persistence thresholds. Hydrodynamical simulations have found that the typical radius of a filament at $z \sim 3$ is 2–3 pMpc (8–12 cMpc; Zhu et al. 2021). The median values of $d_{2d \rightarrow 3d}$ are well below this range, being 2.0, 2.7, and 4 cMpc respectively for the 3D skeletons found with persistence 5, 6, and 7σ . The values are less than 5 cMpc (1.25 pMpc) for the majority of the 2D segments. We refer to the segments with $d_{2d \rightarrow 3d} < 5$ cMpc as ‘matched 2D segments’, meaning that they have a ‘match’ in the 3D skeleton. The fraction of matched segments decreases with increasing persistence threshold, from $\sim 80\%$ for a persistence threshold of 5σ to $\sim 60\%$ when the persistence threshold is 7σ . This is reasonable; as the persistence threshold is raised, the extracted network is increasingly restricted to only the most significant filaments and finer features are lost, leading to a greater number of 2D segments going unmatched. Irrespective of the adopted persistence value, 55% of the 3D segments have 2D matches within $d_{3d \rightarrow 2d} < 5$ cMpc. We conclude that 2D skeleton serves as a good representation of the underlying cosmic web even though it faithfully recovers the individual features $\approx 55\%$ of the time.

Next, we examine whether the fraction of matched 2D segments depends on the persistence threshold used to extract the 2D skeleton. If the matched fraction increases substantially with increasing persistence threshold, it would indicate that our chosen persistence threshold is too generous. Reassuringly, we find that the matched fraction only rises marginally with the persistence threshold, increasing by $\lesssim 5\%$ from a persistence threshold of 2σ to 5σ . This is visualized in Figure 10, where we show the 2D skeletons extracted with persistence thresholds of 2, 3, and 5σ in comparison to the dark matter distribution and the 3D skeletons. The majority of the 2D filaments extracted with a lower persistence threshold, which do not appear with a higher one,




Figure 8. *Top left:* The dark matter distribution for a randomly selected slice, centered on a protocluster with descendant mass $\log(M_{200}/M_{\odot}) = 15.18$ (red cross; Group ID = 0 within the $z = 0$ TNG300 simulation box). *Top right:* The filaments (pink lines) extracted by DisPerSE for this slice based on the mock LAE positions, which are shown by black dots. This network is comparable to the cosmic web selected with ODIN data. *Bottom:* The filaments extracted by DisPerSE for the same slice using the 3D positions of the slice galaxies with $10^7 M_{\odot} < M_{*} < 10^{12} M_{\odot}$, with varying persistence thresholds. The filaments are color-coded by their separation from the center of the slice along the line-of-sight direction. At large distances from the slice center, the filaments are truncated by the slice boundaries. The number of such truncated filaments reduces at higher persistence thresholds.




Figure 9. *Left:* An example comparison of the 2D and 3D skeletons, illustrating how $d_{2d \rightarrow 3d}$ and $d_{3d \rightarrow 2d}$ are measured. *Right:* Comparison of the 2D and 3D skeleton using the separation of the segments, $d_{2d \rightarrow 3d}$ (top) and $d_{3d \rightarrow 2d}$ (bottom). The separations are calculated across all 90 slices of 80 cMpc thickness from the $z = 3$ TNG300 simulation. In the majority of cases, both $d_{2d \rightarrow 3d}$ and $d_{3d \rightarrow 2d}$ are less than 5 cMpc, regardless of the specific persistence threshold.

can be visually matched to the true features of the 3D skeleton.

Intriguingly, the fraction of matched 2D segments increases with increasing surface density for all cases, as shown in the top panel of Figure 11, suggesting that the higher the observed LAE surface density is, the more likely the detected filament is to be an accurate representation of the 3D cosmic web. We also examine the converse, i.e. the probability that a filament within an overdense region is recovered by observations. On considering the 3D skeleton within 10 cMpc of a massive protocluster, we find that the fraction of matched 3D segments is considerably higher than in average regions, increasing from a median of $\sim 55\%$ to a median of $\sim 86\%$ percent in all three persistence cases. The matched fraction is even higher if we restrict the region under consideration to within 5 cMpc of the protoclusters, being 100% in the majority of slices.

In the bottom panel of Figure 11, we show $d_{3d \rightarrow 2d}$ as a function of the (2D) LAE surface density in the slices. To avoid confusion, we only show the result for the 5σ threshold skeleton, as the remaining two persistence thresholds give nearly identical results. With increasing LAE surface density, $d_{3d \rightarrow 2d}$ decreases rapidly; i.e., in dense protocluster regions, not only is the recovery rate of the 3D skeleton considerably higher but also

the 2D skeleton resembles the 3D skeleton more closely. This is illustrated in Figure 12 where we show the 2D and 3D skeletons for the same slice as in Figure 8 but zoom in on the massive protocluster. These results show that the ODIN data will be capable of studying protoclusters within the context of the surrounding cosmic web.

6. DISCUSSION

In Section 4, we based our protocluster selection on the GS and VT maps and the HDBSCAN algorithm. These methods resulted in 9, 33, and 47 structures, respectively. We now wish to evaluate the similarities and differences between these methods.

The false positive rate in the GS method is very low for all sets of detection parameters and is ~ 0 for our fiducial criteria. The 9 GS-based protoclusters may thus be considered the most secure candidates. All GS structures are present in the VT sample with similar morphologies. In all cases, the separation between the centers determined by the GS and VT method is within 2.5 cMpc. For all but one that is present in both GS and VT samples, the measured area is slightly greater (by a factor of $\sim 1.1 - 1.8$) in the latter. This is because irregular overdensities are generally detected at higher significance in the VT map.




Figure 10. Similar to Figure 8, but showing the 2D skeletons extracted with multiple persistence thresholds.

All structures in the GS sample are also present in the HDBSCAN sample. One is split into two structures by HDBSCAN as shown in the middle column of Figure 3. For the remaining 8, the structures have similar morphologies in both the GS and HDBSCAN analyses; however, the measured area in HDBSCAN is larger than that estimated from the GS method in by a factor of $\sim 2 - 5$.

More interesting is the comparison between the structures detected from the VT map and using HDBSCAN. Figure 4 shows the locations of the VT (upper right) and HDBSCAN samples (lower left). Of the 33 protocluster candidates in the VT sample, only two are not detected by HDBSCAN. As for the remaining 31, one is recovered as two separate objects and two are merged into a single structure. For all the protoclusters detected by both the VT and HDBSCAN methods, the latter yields larger measured areas than the former, typically by more than a factor of 2.

Why are the VT-based structures smaller and fewer in number compared to the HDBSCAN-selected ones? Visual inspection of Figure 4 suggests that many HDBSCAN-selected structures with no VT counterpart are more elongated and filamentary. In particular, the filamentary arm previously highlighted in Complex A is identified by HDBSCAN (highlighted in red in the figure) but not from the VT map. If we relax the overdensity threshold for the VT detection from 4.5σ to 3.5σ , the number of HDBSCAN-selected protoclusters that overlap one or more VT-based ones increases from 30 to all 47. The areas of the two sets of protoclusters become more comparable. However, the downside of lowering the detection threshold for the VT map is an increase in the contamination rate. We conclude that HDBSCAN

fares better in recovering lower surface density features with a more elongated morphology, but with the downside that it is more difficult to ascribe a physical meaning to the detection parameters for this method (K th nearest neighbour and minimum cluster size) than for those for the GS and VT methods (minimum area and density threshold).

We now estimate the descendant masses of our protocluster candidates. We showed in Section 5.2 that for our choice of protocluster detection parameters, Equation 6 underestimates the descendant masses of the structures selected from the VT map by a factor of ~ 3 . A similar analysis for the GS and HDBSCAN methods (see Appendix B) shows that the masses of the former are underestimated by a similar factor of ~ 3 , and the latter by a factor of ~ 1.15 . In Figure 13, we show histograms of the descendant masses after making the appropriate corrections. The median masses are $\log(M_{z=0}/M_{\odot}) = 14.35, 14.75$, and 14.52 for the VT, HDBSCAN, and GS protoclusters, respectively, suggesting that our protocluster candidates will evolve into moderately massive ‘Virgo-type’ clusters (Chiang et al. 2013).

7. CONCLUSIONS

The ODIN survey is the largest-area deep field narrow-band survey undertaken to date. By enabling the selection of LAEs, which are low-mass, star-forming galaxies and are well-localized in redshift space over a wide contiguous area, ODIN makes it possible to comprehensively trace the large-scale structure at high redshift. In this paper, we have used the early ODIN data taken in the COSMOS field with the $N501$ filter to compile a sample of protoclusters and cosmic filaments.




Figure 11. *Top:* The fraction of 2D filament segments with a match with the 3D skeleton (detected with various persistence thresholds), as a function of the LAE surface density at the position of the filament segment. Regardless of the persistence threshold used to select the 3D skeleton, the fraction of matched 2D segments increases with increasing surface density. *Bottom:* $d_{3d \rightarrow 2d}$ as a function of the (2D) LAE surface density for the 5σ threshold 3D skeleton; the other two skeletons give virtually identical results. The solid line indicates the median value while the shaded region indicates the spread as measured from the 16-84 percentile scatter.

We create LAE surface density maps using two methods - by smoothing over the LAE positions with a fixed-size Gaussian kernel (GS map, Section 3.1) and by constructing the Voronoi diagram of the LAEs (VT map, Section 3.2). We select protocluster candidates from the surface density maps (Section 4.1) and by applying the density-based clustering algorithm HDBSCAN to the LAE positions (Section 4.2). We also select filaments of the cosmic web (Section 4.3). We assess the reliability of our structure detection by comparing our observations against the results obtained with a carefully created mock sample of LAEs from the IllustrisTNG300-1 hydrodynamical simulation (Sec-

tion 5). Our main results and conclusions are as follows:

1. The large-scale structure revealed by our surface density maps is distinctly clumpy and irregular, with overdense regions clustered together in extended complexes. The VT map identifies filaments as regions of moderate LAE overdensity ($(1 + \delta_{LAE}) = 2-3$) connecting the regions of highest density ($(1 + \delta_{LAE}) > 3$). These features are in line with the expectations of hierarchical structure formation.

2. The three methods of identifying protoclusters which we explore all have their own strengths and weaknesses. The VT map identifies a greater number of objects but also suffers from a higher contamination rate than the GS map. The GS map recovers only the most significant overdensities but is almost free of interlopers. The HDBSCAN map recovers the maximum number of structures, however the input parameters are less straightforward to interpret or assign physical meaning to than for the surface density maps.

3. The surface density maps of the $z = 3$ TNG300 simulation box display remarkably similar features to those seen in the observations. The number and size of the overdensities selected with our fiducial detection parameters are likewise similar in the two. As shown in Figure 6, many of the prominent overdensities in the TNG300 surface density maps correspond to the progenitors of $z = 0$ clusters.

4. The simulation shows that we can successfully recover $\sim 60\%$ of the protoclusters with descendant mass $\gtrsim 2 \times 10^{14} M_{\odot}$. We find that the descendant mass of the simulated protoclusters can be estimated purely based on their area and LAE overdensity measured in 2D. The lack of information in the redshift direction introduces a scatter of ~ 0.4 dex on the measurement.

5. By comparing the filaments identified in 2D and 3D in TNG300, we show that despite projection effects, the cosmic web recovered in 2D is a close representation of the true LSS in regions with high LAE surface density. In the vicinity of protoclusters, the 3D filament network is almost perfectly recovered.

6. On estimating the descendant masses of our observed protocluster samples, we find that they span the range $\log(M_{z=0}/M_{\odot}) \sim 14.0 - 15.0$, with the median of the GS, VT and HDBSCAN samples being 14.35, 14.52 and 14.75 respectively. The majority of our protoclusters are thus likely to evolve into intermediate-mass ‘Virgo-




Figure 12. As Figure 8, but zoomed in on the region surrounding a massive protocluster

type' clusters.

Our results establish the robustness of our protocluster and filament samples and demonstrate that LAEs are reliable and efficient tracers of large-scale structures at high redshift. Upon completion, the ODIN survey will allow us to select secure samples of massive structures that are nearly ten times larger than those presented here and span three cosmic epochs. ODIN will thus be well placed to grant insight into the formation and evolution of cosmic structures near Cosmic Noon.

1 The authors acknowledge financial support from the
 2 National Science Foundation under Grant Nos. AST-
 3 2206705 and AST-2206222 and from the Ross-Lynn Pur-
 4 due Research Foundation Grant. This material is based
 5 upon work supported by the National Science Foun-
 6 dation Graduate Research Fel- lowship Program un-
 7 der Grant No. DGE-2233066 to NF. The Institute
 8 for Gravitation and the Cosmos is supported by the
 9 Eberly College of Science and the Office of the Se-
 10 nior Vice President for Research at the Pennsylvania
 11 State University. J.L. is supported by the National Re-
 12 search Foundation of Korea (NRF-2021R1C1C2011626).
 13 SL is supported by the National Research Foun-
 14 dation of Korea (NRF) grant funded by the Ko-
 15 rean government (MSIT; Nos. 2020R1I1A1A01060310,
 16 2020R1A2C3011091, 2021M3F7A1084525). Based on
 17 observations at Cerro Tololo Inter-American Observa-
 18 tory, NSF's NOIRLab (Prop. ID 2020B-0201; PI: K.-S.
 19 Lee), which is managed by the Association of Univer-
 20 sities for Research in Astronomy under a cooperative
 21 agreement with the National Science Foundation.




Figure 13. Descendant masses of the individual protocusters for all three samples, after applying appropriate corrections to the mass calculated with Equation 6.

Facility: Blanco

Software: SExtractor (Bertin & Arnouts 1996), SEP (Barbary 2016), hdbscan (McInnes et al. 2017), Astropy (Astropy Collaboration et al. 2013, 2018, 2022)

APPENDIX

A. PARAMETER SELECTION FOR HDBSCAN

As stated in Section 4.2, HDBSCAN is a density-based clustering algorithm which separates points into ‘clusters’ and ‘noise’. HDBSCAN performs this classification by estimating the density at each data point from its K th nearest neighbor distance, referred to as the ‘core distance’. Points in dense(sparse) regions will have relatively small(large) core distances, and the core distance of a data point is thus inversely proportional to its density.

HDBSCAN is an extension of the DBSCAN algorithm. DBSCAN selects clusters by imposing a user-defined density threshold. This is implemented by placing a ceiling on the core distance/ K th nearest neighbour distance, that is, by requiring that there be K points within a distance ϵ or less of a given point. DBSCAN begins with a point which passes the requirement of having a core distance less than or equal to ϵ , and classifies it and its neighbours within ϵ as a single cluster. If any of these neighbours also pass the density criterion, their neighbours within ϵ are assigned to the same cluster as well. This continues until there are no more points within the cluster which have a core distance less than ϵ , at which point the cluster is closed. DBSCAN then selects an unassigned point with K or more neighbours within ϵ and repeats the process. This continues until no points which pass the density threshold remain, at which juncture all remaining data points are classified as noise.

The approach of DBSCAN, of applying a constant density/core distance threshold, is restrictive. The optimal value of ϵ is difficult to identify, and can vary from region to region within the data. HDBSCAN circumvents the difficulty of selecting a single density threshold by making use of *hierarchical clustering*; that is, it creates a hierarchy of clusters by applying successively higher density thresholds, and then selects the optimal set of clusters from this hierarchy. This makes it ideal for selecting clusters of varying density from a single dataset, as in our case.

As it constructs a cluster hierarchy, HDBSCAN does not require the user to supply the parameter ϵ . Instead, the parameters which govern the cluster selection are K , as before (which determines how density is measured) and an additional parameter, the minimum cluster size. HDBSCAN uses the minimum cluster size to condense the cluster hierarchy - any cluster in the hierarchy which has fewer points than this size is discarded. This is shown in Figure 14 for a sample dataset. From this condensed hierarchy HDBSCAN selects the final set of clusters.

The HDBSCAN library provides two ways of selecting the final set of clusters. The first method, the ‘excess-of-mass’ method, selects the most stable set of clusters, defined as those which persist for the greatest range of density




Figure 14. The cluster hierarchy (middle panel) and condensed cluster hierarchy (right panel) constructed by HDBSCAN for a sample dataset (left panel). The sample dataset contains three regions of densely clustered points along with random noise. The cluster hierarchy is constructed by applying successive density cuts. Each branch of the tree represents a cluster of points selected above some density cut. The condensed cluster hierarchy shows the cluster hierarchy after clusters smaller than the minimum cluster size are discarded.

thresholds. The second method, the ‘leaf’ method, selects the smallest surviving clusters in the hierarchy. These two methods are illustrated in Figure 15 for the same sample dataset.

Based on the above description, there are two main parameters which govern the clustering based on HDBSCAN, namely K and the minimum cluster size. By determining which nearest-neighbour distance will be used for estimating the density, K controls how finely the density distribution is sampled, in essence acting as a kind of smoothing scale. Changing the minimum cluster size affects the condensed cluster hierarchy by changing the number of points required to be considered a cluster, that is, more points are discarded as noise and the cluster selection is more conservative. Setting either of these parameters to too low a value will result in many noise peaks being erroneously selected as clusters, whereas setting them to be too large may result in structures being blended together. Our key consideration in choosing the parameters is ensuring that nearby structures are separated to the extent possible, which we enforce as described below.

We run HDBSCAN on the positions of the LAEs with `min_samples` and `min_cluster_size` both in the range 10 – 30. We observe that the excess-of-mass method tends to blend structures together irrespective of the specific parameter values used, thus we choose to use the leaf method in our cluster selection. As discussed in the main text, we additionally require that the clusters be above a surface density threshold determined such that the contamination fraction is $\lesssim 0.2$. We estimate the mean area of the structures selected for each combination of parameters, as shown Figure 16. With increasing values of `min_samples` and `min_cluster_size`, the mean area of the detected structures increases, indicating that smaller structures are indeed being blended together. We select our final parameters to be `min_cluster_size` = 10 and `min_samples` = 15, which as shown in Figure 16 yields a protocluster sample with a mean area of $\sim 200 \text{ cMpc}^2$. This choice is based on the physical motivation that the scale of a protocluster is expected to be $\sim 10 - 15 \text{ cMpc}$ (Chiang et al. 2013), corresponding to a projected area (for an isotropic structure) of $\sim 100 - 225 \text{ cMpc}^2$. We make use of a value of `min_samples` on the larger end of the allowed range because while over-smoothing the density distribution is undesirable, it is equally undesirable for the distribution to be too finely sampled as it becomes very noisy.

B. CALIBRATION OF THE MASS ESTIMATION FOR GS MAP AND HDBSCAN

In Section 5.2, we evaluated the reliability of the estimated descendant masses for individual protocluster candidates selected from the VT map. We found that the descendant masses were on average underestimated by a factor of ~ 3 , which we attributed to the underestimation of the protocluster volume by our detection algorithm. Here we similarly examine the accuracy of the descendant mass estimates for the protocluster candidates selected from the GS map and with HDBSCAN.

We once again use the 90 slices constructed in Section 2.2. Analogously to the middle and right panels of Figure 5, we compare the properties of the structures selected with HDBSCAN (selected from the GS map) from these




Figure 15. The selection of clusters from the condensed cluster hierarchy with the excess-of-mass (EOM, top row) and leaf (bottom row) methods. Selected branches of the cluster hierarchy are circled in red in the left panels and the corresponding set of cluster points are shown in color in the right panel, while noise points are gray. The EOM method selects the clusters which persist for the greatest range of density cuts, while the leaf method selects the ‘leaves’ of the cluster hierarchy, i.e. the smallest surviving clusters.

slices to those of the observed protoclusters in the top (bottom) panel of Figure 17. The structures remain similar in projected area and median LAE overdensity between theory and simulations, reaffirming the fact that our observational protocluster candidates are reasonable.

As in Section 5.2, for each slice we consider those of the 30 massive cluster progenitors lying within the slice which are successfully recovered. The recovery rate of the HDBSCAN is $\sim 70\%$, slightly higher than that of the VT map. This is in accordance with the fact that the HDBSCAN candidates are the most numerous. By contrast, the recovery rate of the GS map is very low, $\sim 20\%$, in line with our observation that only the most prominent structures are recovered.

We compare the estimated and true descendant masses for recovered clusters for both methods in Figure 18. We find that the descendant masses of the protoclusters are underestimated by a factor of ~ 2 with the GS method, and by a factor of ~ 1.15 with HDBSCAN. In both cases the scatter is similar to that observed with the VT map, ~ 0.48 dex.

REFERENCES

Aihara, H., Arimoto, N., Armstrong, R., et al. 2018a,

PASJ, 70, S4, doi: [10.1093/pasj/psx066](https://doi.org/10.1093/pasj/psx066)

Aihara, H., Armstrong, R., Bickerton, S., et al. 2018b,

PASJ, 70, S8, doi: [10.1093/pasj/psx081](https://doi.org/10.1093/pasj/psx081)




Figure 16. Average area of the structures detected by HDBSCAN, for various combinations of the K th nearest neighbour used to measure density (`min_samples`) and minimum number of points to constitute a cluster (`min_cluster_size`). The expected area of a protocluster is $\sim 100 - 200 \text{ cMpc}^2$; areas much larger than this suggest that multiple structures are being blended together by the algorithm.

- Aihara, H., AlSayyad, Y., Ando, M., et al. 2019, PASJ, 71, 114, doi: [10.1093/pasj/psz103](https://doi.org/10.1093/pasj/psz103)
- Alpaslan, M., Robotham, A. S. G., Driver, S., et al. 2014, MNRAS, 438, 177, doi: [10.1093/mnras/stt2136](https://doi.org/10.1093/mnras/stt2136)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Barbary, K. 2016, The Journal of Open Source Software, 1, 58, doi: [10.21105/joss.00058](https://doi.org/10.21105/joss.00058)
- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393, doi: [10.1051/aas:1996164](https://doi.org/10.1051/aas:1996164)
- Bleem, L. E., Stalder, B., de Haan, T., et al. 2015, ApJS, 216, 27, doi: [10.1088/0067-0049/216/2/27](https://doi.org/10.1088/0067-0049/216/2/27)
- Bond, J. R., Kofman, L., & Pogosyan, D. 1996, Nature, 380, 603, doi: [10.1038/380603a0](https://doi.org/10.1038/380603a0)
- Boylan-Kolchin, M., Springel, V., White, S. D. M., Jenkins, A., & Lemson, G. 2009, MNRAS, 398, 1150, doi: [10.1111/j.1365-2966.2009.15191.x](https://doi.org/10.1111/j.1365-2966.2009.15191.x)
- Bădescu, T., Yang, Y., Bertoldi, F., et al. 2017, ApJ, 845, 172, doi: [10.3847/1538-4357/aa8220](https://doi.org/10.3847/1538-4357/aa8220)
- Byrohl, C., & Nelson, D. 2023, MNRAS, 523, 5248, doi: [10.1093/mnras/stad1779](https://doi.org/10.1093/mnras/stad1779)
- Campello, R. J. G. B., Moulavi, D., & Sander, J. 2013, in Advances in Knowledge Discovery and Data Mining, ed. J. Pei, V. S. Tseng, L. Cao, H. Motoda, & G. Xu (Berlin, Heidelberg: Springer Berlin Heidelberg), 160–172
- Casey, C. M., Cooray, A., Capak, P., et al. 2015, ApJL, 808, L33, doi: [10.1088/2041-8205/808/2/L33](https://doi.org/10.1088/2041-8205/808/2/L33)
- Cautun, M. C., & van de Weygaert, R. 2011, arXiv e-prints, arXiv:1105.0370, doi: [10.48550/arXiv.1105.0370](https://doi.org/10.48550/arXiv.1105.0370)
- Chiang, Y.-K., Overzier, R., & Gebhardt, K. 2013, ApJ, 779, 127, doi: [10.1088/0004-637X/779/2/127](https://doi.org/10.1088/0004-637X/779/2/127)
- Cucciati, O., Lemaux, B. C., Zamorani, G., et al. 2018, A&A, 619, A49, doi: [10.1051/0004-6361/201833655](https://doi.org/10.1051/0004-6361/201833655)
- Daddi, E., Valentino, F., Rich, R. M., et al. 2021, A&A, 649, A78, doi: [10.1051/0004-6361/202038700](https://doi.org/10.1051/0004-6361/202038700)
- Darvish, B., Mobasher, B., Sobral, D., Scoville, N., & Aragon-Calvo, M. 2015, ApJ, 805, 121, doi: [10.1088/0004-637X/805/2/121](https://doi.org/10.1088/0004-637X/805/2/121)
- Dekel, A., Birnboim, Y., Engel, G., et al. 2009, Nature, 457, 451, doi: [10.1038/nature07648](https://doi.org/10.1038/nature07648)
- Dey, A., Lee, K.-S., Reddy, N., et al. 2016, ApJ, 823, 11, doi: [10.3847/0004-637X/823/1/11](https://doi.org/10.3847/0004-637X/823/1/11)
- Eisenhardt, P. R. M., Brodwin, M., Gonzalez, A. H., et al. 2008, ApJ, 684, 905, doi: [10.1086/590105](https://doi.org/10.1086/590105)
- Firestone, N. M., Gawiser, E., Ramakrishnan, V., et al. 2023, arXiv e-prints, arXiv:2312.16075, doi: [10.48550/arXiv.2312.16075](https://doi.org/10.48550/arXiv.2312.16075)
- Gawiser, E., Francke, H., Lai, K., et al. 2007, ApJ, 671, 278, doi: [10.1086/522955](https://doi.org/10.1086/522955)
- Gebhardt, K., Mentuch Cooper, E., Ciardullo, R., et al. 2021, ApJ, 923, 217, doi: [10.3847/1538-4357/ac2e03](https://doi.org/10.3847/1538-4357/ac2e03)
- Gonzalez, A. H., Gettings, D. P., Brodwin, M., et al. 2019, ApJS, 240, 33, doi: [10.3847/1538-4365/aafad2](https://doi.org/10.3847/1538-4365/aafad2)
- Guaita, L., Gawiser, E., Padilla, N., et al. 2010, ApJ, 714, 255, doi: [10.1088/0004-637X/714/1/255](https://doi.org/10.1088/0004-637X/714/1/255)




Figure 17. Area and median LAE overdensity (δ_{LAE}) for the structures selected with HDBSCAN (top) and from the GS map (bottom) in the TNG300 slices compared to those in observations. Solid lines indicate the median values of each property. As with the structures selected from the VT map, the properties of the objects in the simulation and observations are similar.

Hagen, A., Ciardullo, R., Gronwall, C., et al. 2014, *ApJ*, 786, 59, doi: [10.1088/0004-637X/786/1/59](https://doi.org/10.1088/0004-637X/786/1/59)
 Harikane, Y., Ouchi, M., Ono, Y., et al. 2019, *ApJ*, 883, 142, doi: [10.3847/1538-4357/ab2cd5](https://doi.org/10.3847/1538-4357/ab2cd5)
 Hopkins, P. F., Hernquist, L., Cox, T. J., et al. 2006, *ApJS*, 163, 1, doi: [10.1086/499298](https://doi.org/10.1086/499298)
 Huang, Y., Lee, K.-S., Cucciati, O., et al. 2022, *ApJ*, 941, 134, doi: [10.3847/1538-4357/ac9ea4](https://doi.org/10.3847/1538-4357/ac9ea4)
 Hung, D., Lemaux, B. C., Gal, R. R., et al. 2020, *MNRAS*, 491, 5524, doi: [10.1093/mnras/stz3164](https://doi.org/10.1093/mnras/stz3164)
 Ito, K., Tanaka, M., Valentino, F., et al. 2023, *ApJL*, 945, L9, doi: [10.3847/2041-8213/acb49b](https://doi.org/10.3847/2041-8213/acb49b)
 Kraljic, K., Arnouts, S., Pichon, C., et al. 2017, *Monthly Notices of the Royal Astronomical Society*, 474, 547, doi: [10.1093/mnras/stx2638](https://doi.org/10.1093/mnras/stx2638)
 Kriegel, H.-P., Kröger, P., Sander, J., & Zimek, A. 2011, *Wiley interdisciplinary reviews: data mining and knowledge discovery*, 1, 231

Kuchner, U., Haggard, R., Aragón-Salamanca, A., et al. 2022, *MNRAS*, 510, 581, doi: [10.1093/mnras/stab3419](https://doi.org/10.1093/mnras/stab3419)
 Kusakabe, H., Shimasaku, K., Ouchi, M., et al. 2018, *PASJ*, 70, 4, doi: [10.1093/pasj/psx148](https://doi.org/10.1093/pasj/psx148)
 Laigle, C., Pichon, C., Arnouts, S., et al. 2018, *MNRAS*, 474, 5437, doi: [10.1093/mnras/stx3055](https://doi.org/10.1093/mnras/stx3055)
 Lee, K.-S., Gawiser, E., Park, C., et al. 2024, *ApJ*, 962, 36, doi: [10.3847/1538-4357/ad165e](https://doi.org/10.3847/1538-4357/ad165e)
 Lemaux, B. C., Le Fèvre, O., Cucciati, O., et al. 2018, *A&A*, 615, A77, doi: [10.1051/0004-6361/201730870](https://doi.org/10.1051/0004-6361/201730870)
 Lemaux, B. C., Cucciati, O., Le Fèvre, O., et al. 2022, *A&A*, 662, A33, doi: [10.1051/0004-6361/202039346](https://doi.org/10.1051/0004-6361/202039346)
 Madau, P., & Dickinson, M. 2014, *ARA&A*, 52, 415, doi: [10.1146/annurev-astro-081811-125615](https://doi.org/10.1146/annurev-astro-081811-125615)
 Malavasi, N., Aghanim, N., Douspis, M., Tanimura, H., & Bonjean, V. 2020, *A&A*, 642, A19, doi: [10.1051/0004-6361/202037647](https://doi.org/10.1051/0004-6361/202037647)




Figure 18. Descendant mass estimated according to Equation 6 compared to the true descendant mass for structures selected with HDBSCAN (left) and from the GS map (right). The dashed lines represent a 1-to-1 relation of estimated and true mass whereas the solid lines represent the median ratio of the two.

Malavasi, N., Lee, K.-S., Dey, A., et al. 2021, *ApJ*, 921, 103, doi: [10.3847/1538-4357/ac1c6e](https://doi.org/10.3847/1538-4357/ac1c6e)

Malavasi, N., Arnouts, S., Vibert, D., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 465, 3817, doi: [10.1093/mnras/stw2864](https://doi.org/10.1093/mnras/stw2864)

Malavasi, N., Arnouts, S., Vibert, D., et al. 2017, *MNRAS*, 465, 3817, doi: [10.1093/mnras/stw2864](https://doi.org/10.1093/mnras/stw2864)

McInnes, L., Healy, J., & Astels, S. 2017, *The Journal of Open Source Software*, 2, doi: [10.21105/joss.00205](https://doi.org/10.21105/joss.00205)

Nelson, D., Springel, V., Pillepich, A., et al. 2019, *Computational Astrophysics and Cosmology*, 6, 2, doi: [10.1186/s40668-019-0028-x](https://doi.org/10.1186/s40668-019-0028-x)

Oteo, I., Ivison, R. J., Dunne, L., et al. 2018, *ApJ*, 856, 72, doi: [10.3847/1538-4357/aaaf1](https://doi.org/10.3847/1538-4357/aaaf1)

Ouchi, M., Shimasaku, K., Akiyama, M., et al. 2008, *ApJS*, 176, 301, doi: [10.1086/527673](https://doi.org/10.1086/527673)

Overzier, R. A. 2016, *A&A Rv*, 24, 14, doi: [10.1007/s00159-016-0100-3](https://doi.org/10.1007/s00159-016-0100-3)

Peng, Y.-j., Lilly, S. J., Kovač, K., et al. 2010, *ApJ*, 721, 193, doi: [10.1088/0004-637X/721/1/193](https://doi.org/10.1088/0004-637X/721/1/193)

Pillepich, A., Springel, V., Nelson, D., et al. 2018a, *MNRAS*, 473, 4077, doi: [10.1093/mnras/stx2656](https://doi.org/10.1093/mnras/stx2656)

Pillepich, A., Nelson, D., Hernquist, L., et al. 2018b, *MNRAS*, 475, 648, doi: [10.1093/mnras/stx3112](https://doi.org/10.1093/mnras/stx3112)

Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13, doi: [10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830)

Quadri, R. F., Williams, R. J., Franx, M., & Hildebrandt, H. 2012, *ApJ*, 744, 88, doi: [10.1088/0004-637X/744/2/88](https://doi.org/10.1088/0004-637X/744/2/88)

Ramakrishnan, V., Moon, B., Im, S. H., et al. 2023, *The Astrophysical Journal*, 951, 119, doi: [10.3847/1538-4357/acd341](https://doi.org/10.3847/1538-4357/acd341)

Reddy, N. A., & Steidel, C. C. 2009, *ApJ*, 692, 778, doi: [10.1088/0004-637X/692/1/778](https://doi.org/10.1088/0004-637X/692/1/778)

Rykoff, E. S., Rozo, E., Busha, M. T., et al. 2014, *ApJ*, 785, 104, doi: [10.1088/0004-637X/785/2/104](https://doi.org/10.1088/0004-637X/785/2/104)

Rykoff, E. S., Rozo, E., Hollowood, D., et al. 2016, *ApJS*, 224, 1, doi: [10.3847/0067-0049/224/1/1](https://doi.org/10.3847/0067-0049/224/1/1)

Salerno, J. M., Martínez, H. J., & Muriel, H. 2019, *MNRAS*, 484, 2, doi: [10.1093/mnras/sty3456](https://doi.org/10.1093/mnras/sty3456)

Sarron, F., Adami, C., Durret, F., & Laigle, C. 2019, *A&A*, 632, A49, doi: [10.1051/0004-6361/201935394](https://doi.org/10.1051/0004-6361/201935394)

Sarron, F., & Conselice, C. J. 2021, *MNRAS*, 506, 2136, doi: [10.1093/mnras/stab1844](https://doi.org/10.1093/mnras/stab1844)

Schaap, W. E., & van de Weygaert, R. 2000, *A&A*, 363, L29, doi: [10.48550/arXiv.astro-ph/0011007](https://doi.org/10.48550/arXiv.astro-ph/0011007)

Shi, K., Toshikawa, J., Lee, K.-S., et al. 2021, *ApJ*, 911, 46, doi: [10.3847/1538-4357/abe62e](https://doi.org/10.3847/1538-4357/abe62e)

Sousbie, T. 2011, *Monthly Notices of the Royal Astronomical Society*, 414, 350, doi: [10.1111/j.1365-2966.2011.18394.x](https://doi.org/10.1111/j.1365-2966.2011.18394.x)

Sousbie, T., Pichon, C., & Kawahara, H. 2011, *Monthly Notices of the Royal Astronomical Society*, 414, 384, doi: [10.1111/j.1365-2966.2011.18395.x](https://doi.org/10.1111/j.1365-2966.2011.18395.x)

Springel, V., White, S. D. M., Jenkins, A., et al. 2005, *Nature*, 435, 629, doi: [10.1038/nature03597](https://doi.org/10.1038/nature03597)

Stark, D. P., Ellis, R. S., Chiu, K., Ouchi, M., & Bunker, A. 2010, *MNRAS*, 408, 1628, doi: [10.1111/j.1365-2966.2010.17227.x](https://doi.org/10.1111/j.1365-2966.2010.17227.x)

- Steidel, C. C., Adelberger, K. L., Dickinson, M., et al. 1998, ApJ, 492, 428, doi: [10.1086/305073](https://doi.org/10.1086/305073)
- Tempel, E., Stoica, R. S., Martínez, V. J., et al. 2014, MNRAS, 438, 3465, doi: [10.1093/mnras/stt2454](https://doi.org/10.1093/mnras/stt2454)
- Umehata, H., Tamura, Y., Kohno, K., et al. 2015, ApJL, 815, L8, doi: [10.1088/2041-8205/815/1/L8](https://doi.org/10.1088/2041-8205/815/1/L8)
- van de Weygaert, R., & Schaap, W. 2009, in Data Analysis in Cosmology, ed. V. J. Martínez, E. Saar, E. Martínez-González, & M. J. Pons-Bordería, Vol. 665, 291–413, doi: [10.1007/978-3-540-44767-2_11](https://doi.org/10.1007/978-3-540-44767-2_11)
- van der Burg, R. F. J., Muzzin, A., Hoekstra, H., et al. 2013, A&A, 557, A15, doi: [10.1051/0004-6361/201321237](https://doi.org/10.1051/0004-6361/201321237)
- Vargas, C. J., Bish, H., Acquaviva, V., et al. 2014, ApJ, 783, 26, doi: [10.1088/0004-637X/783/1/26](https://doi.org/10.1088/0004-637X/783/1/26)
- Wang, T., Elbaz, D., Daddi, E., et al. 2016, ApJ, 828, 56, doi: [10.3847/0004-637X/828/1/56](https://doi.org/10.3847/0004-637X/828/1/56)
- Weiss, L. H., Bowman, W. P., Ciardullo, R., et al. 2021, ApJ, 912, 100, doi: [10.3847/1538-4357/abedb9](https://doi.org/10.3847/1538-4357/abedb9)
- White, M., Raichoor, A., Dey, A., et al. 2024, arXiv e-prints, arXiv:2406.01803, doi: [10.48550/arXiv.2406.01803](https://doi.org/10.48550/arXiv.2406.01803)
- Zhu, W., Zhang, F., & Feng, L.-L. 2021, ApJ, 920, 2, doi: [10.3847/1538-4357/ac15f1](https://doi.org/10.3847/1538-4357/ac15f1)