On the Expressibility of the Reconstructional Color Refinement

V. Arvind, Johannes Köbler, Oleg Verbitsky[†]

Abstract

One of the most basic facts related to the famous Ulam reconstruction conjecture is that the connectedness of a graph can be determined by the deck of its vertex-deleted subgraphs, which are considered up to isomorphism. We strengthen this result by proving that connectedness can still be determined when the subgraphs in the deck are given up to equivalence under the color refinement isomorphism test. Consequently, this implies that connectedness is recognizable by Reconstruction Graph Neural Networks, a recently introduced GNN architecture inspired by the reconstruction conjecture (Cotta, Morris, Ribeiro 2021).

1 Introduction

For a vertex v of a graph G, the vertex-deleted subgraph $G \setminus v$ is obtained by removing v along with all incident edges from G. The isomorphism type of $G \setminus v$ is sometimes referred to as a card, and then the multiset of all cards is called the deck of G. The famous Ulam reconstruction conjecture [19] says that every graph G with more than 2 vertices is, up to isomorphism, reconstructible from its deck. Though published in 1960, the problem collection [19] is a descendant of the much earlier famous Scottish Book [12]. According to Harary [9], the conjecture has already been discussed since 1929 and, about a hundred years later, it remains a notoriously hard open problem in graph theory.

The subject has a rich literature, and one of the most frequently taken approaches is examination of how much information about a graph G can be retrieved from its deck. One of the earliest results in this direction is referred to in [9] as a "little theorem": G is connected if and only if at least two subgraphs in its deck are connected. This implies the following fact (see also [15, Theorem 2.2])

Theorem 1 (Harary [8]). The connectedness of a graph G is determined by the multiset of all vertex-deleted subgraphs $G \setminus v$ of G.

^{*}The Institute of Mathematical Sciences (HBNI) and Chennai Mathematical Institute, Chennai, India.

[†]Institut für Informatik, Humboldt-Universität zu Berlin, Germany.

[‡]Supported by DFG grant KO 1053/8–2. On leave from the IAPMM, Lviv, Ukraine.

Recall that the subgraphs $G \setminus v$ are considered up to isomorphism, which means that the full structural information about them is available. It is natural to ask which partial information suffices to decide the connectedness of G. For example, the aforementioned "little theorem" of Harary shows that it would be just enough to know the connectedness of each card in the deck. The question addressed in this note is whether connectedness can still be decided when the subgraphs $G \setminus v$ are considered up to an important relaxation of the isomorphism relation, namely up to equivalence under the color refinement test (also known as one-dimensional Weisfeiler-Leman equivalence).

Color refinement iteratively computes an isomorphism-invariant coloring of the vertex set of an input graph G. Let CR(G) denote the mutisets of colors assigned by the algorithm to the vertices of G (see Section 2 for definitions). Color refinement distinguishes two graphs G and H if $CR(G) \neq CR(H)$. If CR(G) = CR(H), the graphs are called *CR-equivalent*. Since CR(G) is a graph invariant, CR-equivalence is a coarser equivalence relation than isomorphism of graphs.

CR-equivalence has natural characterizations discovered in different research contexts. Two graphs are CR-equivalent exactly when they are equivalent in the twovariable first-order logic with counting quantifiers [10], when they have equally many vertices and share a common covering [1], when they are fractionally isomorphic [17], and when the number of homomorphisms from every tree to these graphs is the same [5].

Color refinement is a versatile tool with applications in diverse fields [1, 7, 16]. Most prominently, it is a practical method of isomorphism testing. The graph invariant CR(G) can be computed efficiently [10] and identifies the isomorphism type of G for almost all graphs [2]. In machine learning, color refinement is used for analysis and comparison of graph-structured data [18] and for understanding the expressiveness of graph neural networks (GNNs) [13].

Our main result is the following strengthening of Theorem 1.

Theorem 2. The connectedness of a graph G is determined by the multiset of the invariants $CR(G \setminus v)$ for all vertex-deleted subgraphs of G.

Note that the invariant CR(G) is not powerful enough to decide whether or not G is connected. Indeed, $CR(C_6) = CR(2C_3)$, where C_n denotes the cycle graph of length n, and $2C_3$ stands for the vertex-disjoint union of two copies of C_3 . This cuts off the easiest route to Theorem 2 through the potential use of the "little theorem" of Harary and shows that running color refinement on the deck of a graph can be more beneficial than running it on the graph alone. Theorem 2 is interesting in several respects, which we discuss now.

Consequences for GNNs. The expressive power of a representative class of GNNs is characterizable in terms of color refinement [14]. In the search for a more expressive GNN architecture, Cotta, Morris, Ribeiro [4] introduce the model of k-*Reconstruction GNNs*, whose conceptual novelty consists in enhancing the standard GNNs by running them on the deck of k-vertex subgraphs of a graph G rather than on G itself. The simplest instructive example, highlighted in [4], is given by considering the 6-cycle graph C_6 . As we just mentioned, C_6 is not identified by

 $CR(C_6)$. However, C_6 is reconstructible from its deck, which consists of six copies of the 5-path graph P_5 . Since P_5 is identified by $CR(P_5)$, the cycle graph C_6 is identified by color refinement applied to its deck. We state an immediate consequence of Theorem 2, providing further evidence that k-Reconstruction GNNs can often be more powerful.

Corollary 3. The connecteness of an n-vertex graph is recognizable by (n-1)-Reconstruction GNNs.

Graph reconstruction. Theorem 2 shows that a prototypical reconstructible graph property can be recognized even from an incomplete information about the deck. The research on the reconstruction conjecture provides many results of this kind, and we here mention two of them which concern connectedness. One can determine the connectedness of an *n*-vertex graph from any $\lfloor n/2 \rfloor + 2$ of its cards [3]. This can also be done using the *k*-deck of a graph whenever $k \ge 9n/10$ [6]. Note in this respect that the *k*-deck of a graph determines its k'-deck whenever $k' \le k$ and, therefore, the *k*-deck with a smaller *k* potentially provides less information. Turning back to Theorem 2, it is natural to combine it with the logical characterization of color refinement [10]. As a consequence, the connectedness of a graph can be determined based on the information about its deck expressible in the two-variable logic with counting quantifiers.

Isomorphism invariants of graphs. Let D-CR(G) denote the multiset of CR(G)v) over all vertices v of G. This graph invariant is interesting to compare not only with with $\mathsf{CR}(G)$ but as well with other related invariants. Color refinement is also commonly known as the one-dimensional Weisfeiler-Leman algorithm. Weisfeiler and Leman [20] actually introduced the two-dimensional version, exploiting the same idea as color refinement but coloring pairs of vertices instead of single vertices. Let WL(G) denote the graph invariant computed by this algorithm on input G. A higher dimension results in a more powerful isomorphism test and, in particular, CR(G) is determined by WL(G). It is not hard to show that D-CR(G) is also determined by WL(G). Whether CR(G), D-CR(G), and WL(G) form a hierarchy of graph invariants linearly ordered by their strength is an open question. More precisely, it is unknown whether or not CR(G) is determined by D-CR(G). This question is discussed in [17, Section 6.8] and [4, Section 4.1]. As we mentioned above, CR(G) does not allow us to see whether G is connected. On the other hand, it is well known that the connectedness of a graph G can be determined from WL(G). Theorem 2 provides a formal strengthening of the last fact by showing that the connectedness of G is determined even by $\mathsf{D}\text{-}\mathsf{CR}(G)$.

2 Color refinement

The vertex set of a graph G is denoted by V(G). The *neighborhood* $N_G(x)$ of a vertex $x \in V(G)$ consists of all vertices adjacent to x. We write $\deg_G x = |N_G(x)|$ to denote the degree of x.

On an input graph G, the *color-refinement* algorithm (to be abbreviated as CR) iteratively computes a sequence of colorings C_G^r of V(G). The initial coloring C_G^0 is monochromatic, i.e., $C_G^0(x) = C_G^0(x')$ for all $x, x' \in V(G)$, where the same initial color is used for all G. Each subsequent color of a vertex x is obtained from the preceding coloring by the rule

$$C_G^{r+1}(x) = \{\!\!\{C_G^r(y)\}\!\!\}_{y \in N_G(x)},$$

where $\{\!\!\{\}\!\!\}\$ denotes a multiset. Note that $C_G^1(x) = C_G^1(x')$ exactly when $\deg_G(x) = \deg_G(x')$, that is, the color $C_G^1(x)$ can just be seen as the degree of the vertex x.

We define $C_G(x) = (C_G^0(x), C_G^1(x), C_G^2(x), \ldots)$ as the sequence of $C_G^r(x)$ for all rand set $CR(G) = \{\!\!\{C_G(x)\}\!\!\}_{x \in V(G)}$. We say that CR *distinguishes* two graphs G and H if $CR(G) \neq CR(H)$. If G and H are indistinguishable by CR, i.e., CR(G) = CR(H), they are referred to as CR-equivalent.

Defining $C_G(x)$ as an infinite sequence facilitates proving Theorem 2 in the subsequent sections. For example, this definition is needed for the equivalences in Lemma 8 below. Color refinement is, nevertheless, an absolutely practical procedure. Indeed, if *n*-vertex graphs G and H are distinguishable by CR, then they are distinguishable after the *n*-th refinement round at latest in the sense that $\{\!\!\{C_G^n(x)\}\!\!\}_{x \in V(G)} \neq \{\!\!\{C_H^n(x)\}\!\!\}_{x \in V(H)}$. Note in this respect that the color names of $C_G^r(x)$ increase exponentially with r if they are encoded in a straightforward way. This can be avoided by renaming the colors synchronously on G and H after each round.

3 Useful lemmas

As $C_G^1(x) = \deg_G x$, we have $C_G^2(x) = \{\!\!\{\deg_G(y) : y \in N_G(x)\}\!\!\}$, which can be referred to as the *iterated degree* of x. We begin with a particularly useful fact about the reconstructibility of iterated degrees.

Lemma 4 (Nash-Williams [15, Lemma 3.3]). Suppose that graphs G and H have the same deck up to CR-equivalence, that is,

$$\{\!\{\mathsf{CR}(G \setminus x)\}\!\}_{x \in V(G)} = \{\!\{\mathsf{CR}(H \setminus y)\}\!\}_{y \in V(H)}$$

If $\mathsf{CR}(G \setminus u) = \mathsf{CR}(H \setminus v)$ for a vertex $u \in V(G)$ and a vertex $v \in V(H)$, then $C^2_G(u) = C^2_H(v)$.

If two graphs have different number of vertices, then they are straightforwardly distinguished by CR. Nevertheless, such a pair of graphs might still be similar in the sense that the CR colors occurring in them are the same, and the difference is only in the color multiplicities. In the proof of Theorem 2, this similarity concept will play an important role.

Definition 5. We say that graphs A and B are *CR-similar* and write $A \equiv B$ if $\{C_A(x)\}_{x \in V(A)} = \{C_B(x)\}_{x \in V(B)}$. That is, we require that the *sets* of the colors produced by CR on A and B be equal (although the *multisets* of these colors might be unequal).

Note that two graphs with different numbers of vertices can be CR-similar, for example, $C_3 \equiv C_4$.

Lemma 6.

- **1.** If A' is a proper subgraph of a connected graph A, then $A \not\equiv A'$.
- **2.** Let A' be a proper subgraph of A and B' be a proper subgraph of B. Suppose that both A and B are connected. Then it is impossible that simultaneously $A \equiv B'$ and $B \equiv A'$.

The proof is based on a characterization of the CR-color $C_G(x)$ in terms of the universal cover of the graph G rooted at the vertex x. We first give the relevant definitions.

Let G and K be connected graphs, where G is finite graph, while K can be either finite or infinite. Let α be a surjective homomorphism from K onto G. We call α a covering map if its restriction to the neighborhood of each vertex in K is bijective. If such an α exists, we say that K covers G. A graph U is called a universal cover of G if U covers every graph covering G. A universal cover $U = U_G$ of G is unique up to isomorphism. It can be seen as an unfolding of G into a (possibly infinite) tree starting from an arbitrarily chosen vertex x of G. Denote this unfolding by $U_{G,x}$ and note that $U_{G,x} \cong U_G$ whatever x. If G is a tree, then $U_{G,x} \cong G$. If G contains a cycle, then $U_{G,x}$ is an infinite tree.

The unfolding $U_{G,x}$ can formally be described as follows. A sequence of vertices $x_0x_1 \ldots x_k$ in G is called a *walk starting at* x_0 if every two successive vertices x_i and x_{i+1} are adjacent. If $x_{i+1} \neq x_{i-1}$ for all 0 < i < k, then the walk is *non-backtracking*. Now, U_x is the graph whose vertices are all non-backtracking walks in G starting at $x_0 = x$, with any two walks of the form $x_0x_1 \ldots x_k$ and $x_0x_1 \ldots x_kx_{k+1}$ being adjacent.

A straightforward inductive argument shows that a covering map α preserves the coloring produced by CR, that is, $C_K^r(u) = C_G^r(\alpha(u))$ for all r. This has the following consequence. Let x be a vertex of a connected graph G and y be a vertex of a connected graph H. If $U_{G,x}$ and $U_{H,y}$ are isomorphic as rooted trees, then $C_G(x) = C_H(y)$. The converse implication is also true; see, e.g., Lemmas 2.3 and 2.4 in [11]. We state this equivalence in a more precise form. Let $U_{G,x}^r$ denote the rooted tree $U_{G,x}$ truncated at level r.

Lemma 7. $C_G^r(x) = C_H^r(y)$ if and only if $U_{G,x}^r \cong U_{H,y}^r$, where \cong denotes the isomorphism of rooted trees.

The characterization of CR-colors in terms of universal covers readily implies the following equivalences.

Lemma 8. The following conditions are equivalent for connected graphs G and H:

- 1. $G \equiv H$;
- 2. $U_G \cong U_H;$

3. $\{C_G(x)\}_{x \in V(G)} \cap \{C_H(y)\}_{y \in V(H)} \neq \emptyset.$

Lemma 8 has a simple consequence about the CR-similarity of not necessarily connected graphs. For graphs G_1, \ldots, G_s , we write $G_1 + \cdots + G_s$ to denote the vertex-disjoint union of these graphs.

Lemma 9.

- **1.** If $G \equiv H$, then for every connected component G' of G there is a connected component H' of H such that $G' \equiv H'$ (and vice versa).
- **2.** If $G \equiv H$ where G is connected and H = H' + H'', then $G \equiv H' \equiv H''$.

Proof of Lemma 6. Let us extend the notation $U_{G,x}^r$ to not necessarily connected graphs by setting $U_{G,x}^r = U_{G',x}^r$ where G' is the connected component of G containing the vertex x.

1. Let the positive integer D be one more than the diameter of A. Fix a vertex u of the subgraph A' maximizing the number of vertices in the truncated tree $U^{D}_{A',u}$. The description of a universal cover in terms of non-backtracking walks readily implies that $U^{D}_{A,u}$ has strictly more vertices than $U^{D}_{A',u}$ and, hence, than $U^{D}_{A',w}$ for any vertex w of A'. Indeed, our choice of D ensures that there is a walk of length at most D starting from u and passing through an edge of A absent in A'. Lemma 7, therefore, implies that $C_A(u) \neq C_{A'}(w)$ for any $w \in V(A')$.

2. Towards a contradiction, assume that $A \equiv B'$ and $B \equiv A'$. Let D be the positive integer that is one more than the larger of the diameters of A and B. Choose a vertex u of the subgraph A' maximizing the number of vertices in the truncated tree $U^{D}_{A',u}$. As in the proof of the first part, note that $U^{D}_{A,u}$ has strictly more vertices than $U^{D}_{A',u}$ and, hence, than $U^{D}_{A',w}$ for any vertex w of A'. The assumption $B \equiv A'$ implies by Lemma 7 that every tree $U^{D}_{A',w}$ for $w \in V(A')$ has an isomorphic mate $U^{D}_{B,w'}$ for some $w' \in V(B)$ and vice versa. Therefore, $U^{D}_{A,u}$ has strictly more vertices than $U^{D}_{B,w}$ for any vertex w of B. The same argument with the roles of A and B interchanged provides us with a vertex v in B which has strictly more vertices than $U^{D}_{A,w}$ for any vertex w of A, in particular, than $U^{D}_{A,u}$. This contradiction completes the proof.

4 Proof of Theorem 2

Let G be a connected graph and H any disconnected graph. We have to prove that the multisets $\{\!\{ \mathsf{CR}(G \setminus x) \}\!\}_{x \in V(G)}$ and $\{\!\{ \mathsf{CR}(H \setminus y) \}\!\}_{y \in V(H)}$ are unequal. We will assume that these multisets are equal and derive a contradiction.

Let H_1, \ldots, H_s , where $s \ge 2$, be the connected components of H. Let V = V(H)and $V_i = V(H_i)$ for $i \le s$. Identify V(G) with V so that

$$\mathsf{CR}(G \setminus v) = \mathsf{CR}(H \setminus v) \tag{1}$$

for all $v \in V$. By Lemma 4, this yields

$$C_G^2(v) = C_H^2(v)$$

for all $v \in V$. Taking into account the decomposition of H into connected components, we have

$$C_G^2(v) = C_{H_i}^2(v) \text{ if } v \in V_i.$$
 (2)

Denote the set of articulation (or cut) vertices of G by A, and set $B = V \setminus A$. Let $B_i = B \cap V_i$. If $v \in B_i$, then $G \setminus v$ is connected and $H \setminus v = H_1 + \ldots + H_i \setminus v + \ldots + H_s$. By Part 2 of Lemma 9, Equality (1) implies that

$$G \setminus v \equiv H_i \setminus v \equiv H_j \text{ if } v \in B_i \text{ and } j \neq i.$$
 (3)

Applying the similarity $G \setminus v \equiv H_j$ to any two vertices in B_i , we derive the similarity $G \setminus v \equiv G \setminus v'$ whenever $v, v' \in B_i$. Applying Lemma 4 in the particular case of G = H, we conclude that

$$C_G^2(v) = C_G^2(v') \text{ for all } v, v' \in B_i.$$

$$\tag{4}$$

Let $i \neq j$ and assume that $B_i \neq \emptyset$ and $B_j \neq \emptyset$. Let $x \in B_i$ and $y \in B_j$. By (3), we obtain $H_i \setminus x \equiv H_j$ and $H_j \setminus y \equiv H_i$, which contradicts Part 2 of Lemma 6. Therefore, only one of the sets B_i is not empty, that is, $B = B_i$ for some *i*. Without loss of generality, we suppose that $B = B_1$.

Claim A. $B_1 = V_1$.

Proof of Claim A. Towards a contradiction, assume that V_1 contains a cut vertex v of G. By Part 1 of Lemma 9, Equality (1) implies that the disconnected graph $G \setminus v$ has a connected component G' such that $G' \equiv H_2$. Choose a vertex $x \in B$ in a connected component of $G \setminus v$ different from G'. Since $x \in B_1$, the similarity (3) implies that $G \setminus x \equiv H_2$ and, therefore, $G \setminus x \equiv G'$. However, G' is a subgraph of the connected graph $G \setminus x$, contrary to Part 1 of Lemma 6.

By Claim A and Equality (4), we have

$$C_G^2(x) = C_G^2(x')$$
 for all $x, x' \in V_1$.

Taking into account Equality (2), this implies that

$$C_{H_1}^2(x) = C_{H_1}^2(x') \text{ for all } x, x' \in V_i.$$
(5)

In particular, H_1 is a regular graph, say, of degree d.

Consider the block-cut tree of G. Recall that it describes the decomposition of G into biconnected components, that is, maximal biconnected subgraphs, which are also called blocks. The vertex set of the tree consists of the blocks and the cut vertices of G. A block M is adjacent to a cut vertex c if $c \in M$.

Let L be a block that is a leaf in the block-cut tree of G. Let a be the single cut vertex of G belonging to L. By Claim A, a belongs to $A = V \setminus B = V_2 \cup \ldots \cup V_s$, and the other vertices of L belong to $B = B_1 = V_1$. Without loss of generality, suppose that $a \in V_2$. Let x be a vertex in L adjacent to a. Since $x \in B_1$, the similarity (3) implies that

$$G \setminus x \equiv H_1 \setminus x \equiv H_2. \tag{6}$$

Since H_1 is a *d*-regular graph, every vertex of $H_1 \setminus x$ has degree d - 1 or d. By the similarity (6), the neighbors of a in H_2 are of degree d - 1 or d. By Equality

(2), the same holds true for a in G. Therefore, every connected component G' of $G \setminus a$ has a vertex of degree d-2 or d-1 (namely a neighbor of a in G). As a consequence, $G' \not\equiv H_1$ for every G'. On the other hand, Equality (1) implies that

$$G \setminus a \equiv H_1 + H_2 \setminus a + \dots + H_s.$$

By Part 1 of Lemma 9, we conclude that $H_1 \equiv G'$ for some component G' of $G \setminus a$. This contradiction completes the proof of Theorem 2.

References

- D. Angluin. Local and global properties in networks of processors (extended abstract). In Proc. of the 12th Annual ACM Symposium on Theory of Computing (STOC'80), pages 82–93. ACM, 1980.
- [2] L. Babai, P. Erdös, and S. M. Selkow. Random graph isomorphism. SIAM J. Comput., 9(3):628–635, 1980.
- [3] A. Bowler, P. Brown, T. Fenner, and W. Myrvold. Recognizing connectedness from vertex-deleted subgraphs. *Journal of Graph Theory*, 67(4):285–299, 2011.
- [4] L. Cotta, C. Morris, and B. Ribeiro. Reconstruction for powerful graph representations. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, editors, Advances in Neural Information Processing Systems (NeurIPS 2021), volume 34, pages 1713–1726. Curran Associates, Inc., 2021.
- [5] Z. Dvořák. On recognizing graphs by numbers of homomorphisms. Journal of Graph Theory, 64(4):330–342, 2010.
- [6] C. Groenland, T. Johnston, A. Scott, and J. Tan. Reconstruction from smaller cards. Technical report, arxiv.org/abs/2103.13359, 2023.
- [7] M. Grohe, K. Kersting, M. Mladenov, and P. Schweitzer. Color refinement and its applications. In An Introduction to Lifted Probabilistic Inference. The MIT Press, 2021.
- [8] F. Harary. On the reconstruction of a graph from a collection of subgraphs. In *Theory of Graphs and its Applications*, pages 47–52. Academic Press: New York, 1964.
- [9] F. Harary. A survey of the reconstruction conjecture. In Graphs and Combinatorics, volume 406 of Lect. Notes Math, pages 18–28. Springer: Berlin, Heidelberg, 1974.
- [10] N. Immerman and E. Lander. Describing graphs: A first-order approach to graph canonization. In A. Selman, editor, *Complexity Theory Retrospective*, pages 59–81. Springer-Verlag, 1990.

- [11] A. Krebs and O. Verbitsky. Universal covers, color refinement, and two-variable counting logic: Lower bounds for the depth. In 30th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS'15), pages 689–700. IEEE Computer Society, 2015.
- [12] R. D. Mauldin, editor. The Scottish Book. Birkhäuser, Boston, MA, 1981. Mathematics from the Scottish Café, Including selected papers presented at the Scottish Book Conference held at North Texas State University, Denton, Tex., May 1979.
- [13] C. Morris, Y. Lipman, H. Maron, B. Rieck, N. M. Kriege, M. Grohe, M. Fey, and K. Borgwardt. Weisfeiler and Leman go machine learning: The story so far. *Journal of Machine Learning Research*, 24(333):1–59, 2023.
- [14] C. Morris, M. Ritzert, M. Fey, W. L. Hamilton, J. E. Lenssen, G. Rattan, and M. Grohe. Weisfeiler and Leman go neural: Higher-order graph neural networks. In *The 33-rd Conference on Artificial Intelligence (AAAI'19)*, pages 4602–4609. AAAI Press, 2019.
- [15] C. S. J. A. Nash-Williams. The reconstruction problem. In Selected topics in graph theory, pages 205–236. Academic Press: New York, 1978.
- [16] C. Riveros, B. Scheidt, and N. Schweikardt. Using color refinement to boost enumeration and counting for acyclic CQs of binary schemas. Technical report, arxiv.org/abs/2405.12358, 2024.
- [17] E. R. Scheinerman and D. H. Ullman. Fractional graph theory. A rational approach to the theory of graphs. Wiley: John Wiley & Sons, 1997.
- [18] N. Shervashidze, P. Schweitzer, E. J. van Leeuwen, K. Mehlhorn, and K. M. Borgwardt. Weisfeiler-Lehman graph kernels. *Journal of Machine Learning Research*, 12:2539–2561, 2011.
- [19] S. M. Ulam. A collection of mathematical problems, volume 8 of Intersci. Tracts Pure Appl. Math. Interscience Publishers, New York, NY, 1960.
- [20] B. Weisfeiler and А. Leman. The reduction of a graph tocanonical and the algebra which NTI. form appears therein. English 2. 9:12-16,1968. translation Ser. is available athttps://www.iti.zcu.cz/wl2018/pdf/wl_paper_translation.pdf.