

The gauge coupling evolutions of an SU(8) theory with the maximally symmetry breaking pattern

Ning Chen[♡] , Zhanpeng Hou[♠], Ying-nan Mao[◇] , Zhaolong Teng[♣] 

^{♡ ♠ ♣} *School of Physics, Nankai University, Tianjin, 300071, China*

[◇] *Department of Physics, School of Science, Wuhan University of Technology, Wuhan, 430070, Hubei, China*

Abstract

We study the renormalizable group equations (RGEs) of the extended strong and weak gauge couplings in an SU(8) theory, where three-generational SM fermions are non-trivially embedded. This framework was previously found to generate the observed SM quark/lepton mass hierarchies and the Cabibbo-Kobayashi-Maskawa mixing pattern through its maximally breaking pattern. The field theoretical two-loop RGEs can not achieve the gauge coupling unification with the minimal setup, unless additional adjoint Higgs fields as well as the gravity-induced $d = 5$ term to the SU(8) field strength term are included.

Emails:

[♡] chenning_symmetry@nankai.edu.cn,
[♠] houzhanpeng@mail.nankai.edu.cn,
[◇] ynmao@whut.edu.cn,
[♣] tengcl@mail.nankai.edu.cn

Contents

1	Introduction	1
2	The SU(8) theory and its symmetry breaking pattern	3
2.1	The setup of the SU(8) theory	3
2.2	The SWW symmetry breaking pattern of the SU(8) theory	5
2.3	Decompositions of the SU(8) fermions	5
2.4	Decompositions of the SU(8) Higgs fields	7
2.5	A summary of the vectorlike fermions of the SWW symmetry breaking pattern	8
3	The gauge sector of the SU(8) theory and the flavor non-universality	9
3.1	The $SU(4)_s \otimes U(1)_{X_0}$ gauge bosons and gauge couplings	9
3.1.1	Covariant derivatives and gauge boson masses	9
3.1.2	Gauge couplings of the \mathcal{G}_{441} fermions	11
3.2	The $SU(4)_W \otimes U(1)_{X_1}$ gauge bosons and gauge couplings	12
3.2.1	Covariant derivatives and gauge boson masses	12
3.2.2	Gauge couplings of the \mathcal{G}_{341} fermions	16
3.3	The $SU(3)_W \otimes U(1)_{X_2}$ gauge bosons	18
3.3.1	Covariant derivatives and gauge boson masses	18
3.3.2	Gauge couplings of the \mathcal{G}_{331} fermions	21
3.4	The $SU(2)_W \otimes U(1)_Y$ gauge bosons	23
3.4.1	Covariant derivatives and gauge boson masses	23
3.4.2	Fermion \mathcal{G}_{SM} gauge couplings	24
4	The gauge coupling unification in the SU(8) theory	24
4.1	RGEs of the minimal setup	25
4.2	The gauge coupling unification with additional adjoint Higgs fields	30
5	Summary and outlook	31
5.1	The main results	31
5.2	On the SUSY extension and the Kač-Moody extension	31
A	Conventions, decomposition rules and charge quantizations in the SU(8) theory	32
B	Note added in proof of the SU(8) maximal breaking pattern	34

1 Introduction

Grand Unified Theories (GUTs), with their original formulations based on the gauge groups of SU(5) [1] and SO(10) [2], were proposed to not only unify all three fundamental symmetries into one simple Lie group of \mathcal{G}_U , but to unify all Standard Model (SM) fermions into some anomaly-free irreps of the \mathcal{G}_U . The well-known minimal setup include the Georgi-Glashow SU(5) GUT [1] with $3 \times [\overline{\mathbf{5}}_{\mathbf{F}} \oplus \mathbf{10}_{\mathbf{F}}]$, and the Fritzsch-Minkowski SO(10) GUT [2] with $3 \times \mathbf{16}_{\mathbf{F}}$. Meanwhile, these SU(5) and/or SO(10) GUTs, with or without their supersymmetric (SUSY) extensions, are far from satisfactory, given that three-generational SM fermion masses and the Cabibbo-Kobayashi-Maskawa (CKM) mixing pattern [3, 4] were not ultimately explained.

In a pioneering work by Georgi [5], an extension to the gauge group of $SU(N > 5)$ was proposed to embed three-generational SM fermions non-trivially. In his original proposal, any anti-symmetric rank- k fermion irrep can appear at most once with the anomaly-free condition. Physically, this leads to a non-repetitive structure for both left-handed and right-handed components of chiral fermions based on an $SU(11)$ gauge group. Given the current LHC measurements of one single SM Higgs boson so far, which have already confirmed the hierarchical Yukawa couplings of the third-generational SM quarks/leptons and displayed the evidence for the muon [6, 7], Georgi’s original proposal is quite insightful and can be further relaxed by requiring distinctive symmetry properties of three generations in the UV theory. A minimal $SU(8)$ theory [8–10] was suggested to embed three-generational SM quarks/leptons non-trivially¹. The $SU(8)$ symmetry breaking pattern includes three intermediate scales above the EW scale. By analyzing the emergent non-anomalous global $\tilde{U}(1)_T$ symmetry in the UV, we derived the non-anomalous global $B-L$ symmetry in the SM, and one unique SM Higgs doublet from the $\mathbf{70}_H$ of the minimal setup was conjectured based on the requirement of the vanishing global $B-L$ charge [9]. This SM Higgs doublet was found to give natural top quark mass with the renormalizable Yukawa coupling at the tree level. Due to the distinctive symmetry properties of three-generational SM fermions in the UV, we found in Ref. [10] that all SM quark/lepton masses as well as the CKM mixing pattern can be explained with (i) both $d = 5$ direct Yukawa couplings and indirect Yukawa couplings to the $d = 5$ irreducible Higgs mixing operators induced by the inevitable gravitational effects, (ii) three reasonable intermediate symmetry-breaking scales, and (iii) precise identifications of the SM flavors in their UV irreps.

As a unified gauge theory, a central issue is to achieve the gauge coupling unification in terms of their renormalization group equations (RGEs)². A strong-weak-weak (SWW) symmetry breaking pattern of

$$\begin{aligned} SU(8) &\xrightarrow{v_U} \mathcal{G}_{441} \xrightarrow{v_{441}} \mathcal{G}_{341} \xrightarrow{v_{341}} \mathcal{G}_{331} \xrightarrow{v_{331}} \mathcal{G}_{SM} \xrightarrow{v_{EW}} SU(3)_c \otimes U(1)_{EM}, \\ \mathcal{G}_{441} &\equiv SU(4)_s \otimes SU(4)_W \otimes U(1)_{X_0}, \quad \mathcal{G}_{341} \equiv SU(3)_c \otimes SU(4)_W \otimes U(1)_{X_1}, \\ \mathcal{G}_{331} &\equiv SU(3)_c \otimes SU(3)_W \otimes U(1)_{X_2}, \quad \mathcal{G}_{SM} \equiv SU(3)_c \otimes SU(2)_W \otimes U(1)_Y, \end{aligned} \quad (1)$$

was previously considered in Ref. [10], and three intermediate scales of

$$v_{441} \simeq 1.4 \times 10^{17} \text{ GeV}, \quad v_{341} \simeq 4.8 \times 10^{15} \text{ GeV}, \quad v_{331} \simeq 4.8 \times 10^{13} \text{ GeV}, \quad (2)$$

were predicted with the reduced Planck scale of $M_{\text{pl}} = (8\pi G_N)^{-1/2} = 2.4 \times 10^{18} \text{ GeV}$. The maximally broken subgroup of \mathcal{G}_{441} further admits the \mathcal{G}_{SM} as its subgroup³. These scales were obtained by the reasonable predictions of three-generational SM quark/lepton masses and the CKM mixing pattern, together with the precise identifications of the SM flavors in their UV irreps. In this paper, we perform the RG analysis of the gauge couplings in the extended strong and weak sectors based on the benchmark point in Eq. (2).

The rest of the paper is organized as follows. We review the $SU(8)$ theory in Sec. 2, with the focus on the symmetry breaking pattern. In Sec. 3, we define the gauge sector of the $SU(8)$ theory along its symmetry breaking pattern defined in Sec. 2. The gauge couplings at each symmetry breaking stage will be defined. Due to the intrinsic symmetry properties, we explicitly display the flavor non-universalities of the \mathcal{G}_{441} and \mathcal{G}_{341} theories as well as the flavor universality of the \mathcal{G}_{331} theory through

¹By decomposing the $SU(8)$ fermionic irreps into the $SU(5)$ irreps, one finds three identical SM $\overline{\mathbf{5}}_F$ ’s and three distinctive $\mathbf{10}_F$ ’s. Hence, it is sufficient to obtain three distinctive SM generations in the UV.

²Some early studies of the RGEs based on the $SU(5)$ and $SO(10)$ groups include Refs. [11–13].

³Other different symmetry breaking pattern of the $SU(8)$ group was previously described in Ref. [14].

their neutral currents. In Sec. 4, we study gauge coupling unification in terms of the two-loop RGEs. The SU(8) theory with the minimal fermion/Higgs contents as well as the intermediate symmetry breaking scales in Refs. [9, 10] cannot achieve the gauge coupling unification in terms of the conventional field theoretical analyses. Instead, inclusion of additional SU(8) adjoint Higgs fields and the $d = 5$ Hill-Shafi-Wetterich (HSW) operator [15, 16] are suggested to achieve the gauge coupling unification within the field theoretical framework. We summarize and make future perspective in Sec. 5, such as the possible issues in the supersymmetric extensions. An App. A is given to define the gauge group indices and the decomposition rules according to the symmetry breaking pattern in the SU(8) theory. App. B is added to prove a unique feature of the U(1) charge definition in the maximally symmetry breaking pattern in Eq. (1).

2 The SU(8) theory and its symmetry breaking pattern

2.1 The setup of the SU(8) theory

The SU(8) theory contains two following chiral IRAFFSs at the GUT scale

$$\begin{aligned} \{f_L\}_{\text{SU}(8)}^{n_g=3} &= \left[\overline{\mathbf{8}}_{\mathbf{F}}^\omega \oplus \mathbf{28}_{\mathbf{F}} \right] \oplus \left[\overline{\mathbf{8}}_{\mathbf{F}}^{\dot{\omega}} \oplus \mathbf{56}_{\mathbf{F}} \right], \quad \dim_{\mathbf{F}} = 156, \\ \Omega \equiv (\omega, \dot{\omega}), \quad \omega &= (\mathbf{3}, \text{IV}, \text{V}, \text{VI}), \quad \dot{\omega} = (\dot{\mathbf{1}}, \dot{\mathbf{2}}, \text{VII}, \text{VIII}, \text{IX}), \end{aligned} \quad (3)$$

with undotted/dotted indices for the $\overline{\mathbf{8}}_{\mathbf{F}}$'s in the rank-2 IRAFFS and the rank-3 IRAFFS, respectively. The Roman numbers and the Arabic numbers are used for the heavy partner fermions and the SM fermions, respectively. These three $\mathbf{10}_{\mathbf{F}}$'s are transforming differently, while three identical $\overline{\mathbf{5}}_{\mathbf{F}}$'s can be identified from nine $\overline{\mathbf{8}}_{\mathbf{F}}$'s, which are sufficient to guarantee that three-generational SM quarks and leptons must transform differently in the UV theory.

The non-anomalous global DRS symmetries from fermions in Eq. (3) are

$$\tilde{\mathcal{G}}_{\text{DRS}} [\text{SU}(8), n_g = 3] = \left[\widetilde{\text{SU}}(4)_\omega \otimes \tilde{\text{U}}(1)_{T_2} \right] \otimes \left[\widetilde{\text{SU}}(5)_{\dot{\omega}} \otimes \tilde{\text{U}}(1)_{T_3} \right], \quad (4)$$

and we also denote the anomalous global Peccei-Quinn symmetries [17] as

$$\tilde{\mathcal{G}}_{\text{PQ}} [\text{SU}(8), n_g = 3] = \tilde{\text{U}}(1)_{\text{PQ}_2} \otimes \tilde{\text{U}}(1)_{\text{PQ}_3}. \quad (5)$$

Both $\tilde{\text{U}}(1)_{T_2}$ and $\tilde{\text{U}}(1)_{\text{PQ}_2}$ can be thought as the linear combinations of two global $\tilde{\text{U}}(1)$'s of the $\overline{\mathbf{8}}_{\mathbf{F}}^\omega$ and the $\mathbf{28}_{\mathbf{F}}$ in the rank-2 IRAFFS, and similarly for the $\tilde{\text{U}}(1)_{T_3}$ and $\tilde{\text{U}}(1)_{\text{PQ}_3}$ symmetries in the rank-3 IRAFFS. Since the global $B - L$ symmetry should be identical for all three generations, we further require a common non-anomalous $\tilde{\text{U}}(1)_T \equiv \tilde{\text{U}}(1)_{T_2} = \tilde{\text{U}}(1)_{T_3}$ between two IRAFFSs. Likewise, we also assume a common anomalous $\tilde{\text{U}}(1)_{\text{PQ}} \equiv \tilde{\text{U}}(1)_{\text{PQ}_2} = \tilde{\text{U}}(1)_{\text{PQ}_3}$ between two IRAFFSs in Eq. (5). Accordingly, the non-anomalous global $\tilde{\text{U}}(1)_T$ charges and the anomalous global $\tilde{\text{U}}(1)_{\text{PQ}}$ charges for fermions are assigned in Tab. 1.

Fermions	$\overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}$	$\mathbf{28}_{\mathbf{F}}$	$\mathbf{56}_{\mathbf{F}}$	
$\widetilde{\mathbf{U}}(1)_T$	$-3t$	$+2t$	$+t$	
$\widetilde{\mathbf{U}}(1)_{\text{PQ}}$	p	q_2	q_3	
Higgs	$\overline{\mathbf{8}}_{\mathbf{H},\omega}$	$\mathbf{28}_{\mathbf{H},\dot{\omega}}$	$\mathbf{70}_{\mathbf{H}}$	$\mathbf{63}_{\mathbf{H}}$
$\widetilde{\mathbf{U}}(1)_T$	$+t$	$+2t$	$-4t$	0
$\widetilde{\mathbf{U}}(1)_{\text{PQ}}$	$-(p+q_2)$	$-(p+q_3)$	$-2q_2$	0

Table 1: The non-anomalous $\widetilde{\mathbf{U}}(1)_T$ charges and the anomalous global $\widetilde{\mathbf{U}}(1)_{\text{PQ}}$ charges for the SU(8) fermions and Higgs fields.

The most general gauge-invariant Yukawa couplings at least include the following renormalizable and non-renormalizable terms ⁴

$$\begin{aligned}
-\mathcal{L}_Y &= Y_{\mathcal{B}} \overline{\mathbf{8}}_{\mathbf{F}}^{\omega} \mathbf{28}_{\mathbf{F}} \overline{\mathbf{8}}_{\mathbf{H},\omega} + Y_{\mathcal{T}} \mathbf{28}_{\mathbf{F}} \mathbf{28}_{\mathbf{F}} \mathbf{70}_{\mathbf{H}} \\
&+ Y_{\mathcal{D}} \overline{\mathbf{8}}_{\mathbf{F}}^{\dot{\omega}} \mathbf{56}_{\mathbf{F}} \overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}} + \frac{c_4}{M_{\text{pl}}} \mathbf{56}_{\mathbf{F}} \mathbf{56}_{\mathbf{F}} \overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}}^{\dagger} \mathbf{63}_{\mathbf{H}} + H.c..
\end{aligned} \tag{6}$$

All renormalizable Yukawa couplings are assumed to be $(Y_{\mathcal{B}}, Y_{\mathcal{T}}, Y_{\mathcal{D}}) \sim \mathcal{O}(1)$, and the Wilson coefficient c_4 of the non-renormalizable term was assigned in Ref. [10]. Altogether, we collect the SU(8) Higgs fields as follows

$$\{H\}_{\text{SU}(8)}^{n_g=3} = \overline{\mathbf{8}}_{\mathbf{H},\omega} \oplus \overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}} \oplus \mathbf{70}_{\mathbf{H}} \oplus \underline{\mathbf{63}}_{\mathbf{H}}, \quad \dim_{\mathbf{H}} = 547, \tag{7}$$

where the adjoint Higgs field of $\mathbf{63}_{\mathbf{H}}$ is real while all others are complex. The adjoint Higgs field of $\underline{\mathbf{63}}_{\mathbf{H}}$ will maximally break the GUT symmetry as $\text{SU}(8) \rightarrow \mathcal{G}_{441}$ through its VEV of

$$\langle \underline{\mathbf{63}}_{\mathbf{H}} \rangle = \frac{1}{4} \text{diag}(-\mathbb{I}_{4 \times 4}, +\mathbb{I}_{4 \times 4}) v_U. \tag{8}$$

The $\underline{\mathbf{63}}_{\mathbf{H}}$ can be decomposed under the \mathcal{G}_{441} as follows

$$\underline{\mathbf{63}}_{\mathbf{H}} = (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{H}} \oplus (\mathbf{15}, \mathbf{1}, 0)_{\mathbf{H}} \oplus (\mathbf{1}, \mathbf{15}, 0)_{\mathbf{H}} \oplus (\overline{\mathbf{4}}, \mathbf{4}, +\frac{1}{2})_{\mathbf{H}} \oplus (\mathbf{4}, \overline{\mathbf{4}}, -\frac{1}{2})_{\mathbf{H}}, \tag{9}$$

where the VEV is developed by the $(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{H}}$ component, and the $(\overline{\mathbf{4}}, \mathbf{4}, +\frac{1}{2})_{\mathbf{H}} \oplus (\mathbf{4}, \overline{\mathbf{4}}, -\frac{1}{2})_{\mathbf{H}}$ represent the Nambu-Goldstone bosons. We assign the non-anomalous global $\widetilde{\mathbf{U}}(1)_T$ charges and the anomalous global $\widetilde{\mathbf{U}}(1)_{\text{PQ}}$ charges for Higgs fields in Tab. 1 according to the renormalizable Yukawa couplings in Eq. (6). The anomalous global $\widetilde{\mathbf{U}}(1)_{\text{PQ}}$ charges are assigned such that

$$p : q_2 \neq -3 : +2, \quad p : q_3 \neq -3 : +1. \tag{10}$$

Obviously, the only non-renormalizable term in Eq. (6) also has a non-vanishing global $\widetilde{\mathbf{U}}(1)_{\text{PQ}}$ charge according to Tab. 1 and Eq. (10). Thus, it can be explicitly broken by the gravitational effect.

⁴The term of $\mathbf{56}_{\mathbf{F}} \mathbf{56}_{\mathbf{F}} \mathbf{28}_{\mathbf{H}} + H.c.$ vanishes due to the anti-symmetric property [8]. Instead, only a $d = 5$ non-renormalizable term of $\frac{1}{M_{\text{pl}}} \mathbf{56}_{\mathbf{F}} \mathbf{56}_{\mathbf{F}} \overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}}^{\dagger} \mathbf{63}_{\mathbf{H}}$ is possible to generate masses for vectorlike fermions in the $\mathbf{56}_{\mathbf{F}}$. Since it transforms as an $\widetilde{\text{SU}}(5)_{\dot{\omega}}$ vector and carries non-vanishing $\widetilde{\mathbf{U}}(1)_{\text{PQ}}$ charge of $p + 3q_3 \neq 0$ from Eq. (10), it is only possible due to the gravitational effect.

2.2 The SWW symmetry breaking pattern of the SU(8) theory

According to Ref. [18], only the adjoint Higgs field can achieve the maximal symmetry breaking pattern of $SU(N) \rightarrow SU(N-k) \otimes SU(k) \otimes U(1)$ (with $k = \lfloor \frac{N}{2} \rfloor$). The SWW symmetry breaking pattern in Eq. (1) follows from Ref. [10] will be further studied in the following discussions. The GUT scale symmetry breaking will be due to the VEV of the SU(8) adjoint Higgs field. In the subgroup of \mathcal{G}_{441} , both the strong and the weak symmetries are extended beyond the SM gauge symmetries. All intermediate symmetry breaking stages will be due to the VEVs of the Higgs fields from the gauge-invariant Yukawa couplings of the SU(8) theory. After the first stage of symmetry breaking pattern of $\mathcal{G}_{441} \rightarrow \mathcal{G}_{341}$ in Eq. (1), the sequential symmetry breaking stages are uniquely determined. Along this symmetry breaking pattern, the non-anomalous global $U(1)_T$ symmetry will evolve to the global $\tilde{U}(1)_{B-L}$ at the EW scale according to the following sequence [9]

$$\begin{aligned} \mathcal{G}_{441} &: \mathcal{T}' \equiv \mathcal{T} - 4t\mathcal{X}_0, & \mathcal{G}_{341} &: \mathcal{T}'' \equiv \mathcal{T}' + 8t\mathcal{X}_1, \\ \mathcal{G}_{331} &: \mathcal{T}''' \equiv \mathcal{T}'', & \mathcal{G}_{\text{SM}} &: \mathcal{B} - \mathcal{L} \equiv \mathcal{T}'''. \end{aligned} \quad (11)$$

With these definitions, we have also checked that all Higgs components that can develop the VEVs at each symmetry breaking stage are neutral under the corresponding global $\tilde{U}(1)_T$ symmetries [9].

2.3 Decompositions of the SU(8) fermions

SU(8)	\mathcal{G}_{441}	\mathcal{G}_{341}	\mathcal{G}_{331}	\mathcal{G}_{SM}
$\overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}$	$(\overline{\mathbf{4}}, \mathbf{1}, +\frac{1}{4})_{\mathbf{F}}^{\Omega}$	$(\overline{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega}$	$(\overline{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega}$	$(\overline{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega} : \mathcal{D}_R^{\Omega c}$
		$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega}$	$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega}$	$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega} : \tilde{\mathcal{N}}_L^{\Omega}$
	$(\mathbf{1}, \overline{\mathbf{4}}, -\frac{1}{4})_{\mathbf{F}}^{\Omega}$	$(\mathbf{1}, \overline{\mathbf{4}}, -\frac{1}{4})_{\mathbf{F}}^{\Omega}$	$(\mathbf{1}, \overline{\mathbf{3}}, -\frac{1}{3})_{\mathbf{F}}^{\Omega}$	$(\mathbf{1}, \overline{\mathbf{2}}, -\frac{1}{2})_{\mathbf{F}}^{\Omega} : \mathcal{L}_L^{\Omega} = (\mathcal{E}_L^{\Omega}, -\mathcal{N}_L^{\Omega})^T$
			$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega'}$	$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega'} : \tilde{\mathcal{N}}_L^{\Omega'}$
			$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega''}$	$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega''} : \tilde{\mathcal{N}}_L^{\Omega''}$

Table 2: The SU(8) fermion representation of $\overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}$ under the $\mathcal{G}_{441}, \mathcal{G}_{341}, \mathcal{G}_{331}, \mathcal{G}_{\text{SM}}$ subgroups for the three-generational SU(8) model, with $\Omega \equiv (\omega, \dot{\omega})$. Here, we denote $\mathcal{D}_R^{\Omega c} = d_R^{\Omega c}$ for the SM right-handed down-type quarks, and $\mathcal{D}_R^{\Omega c} = \mathfrak{D}_R^{\Omega c}$ for the right-handed down-type heavy partner quarks. Similarly, we denote $\mathcal{L}_L^{\Omega} = (\ell_L^{\Omega}, -\nu_L^{\Omega})^T$ for the left-handed SM lepton doublets, and $\mathcal{L}_L^{\Omega} = (\mathbf{e}_L^{\Omega}, -\mathbf{n}_L^{\Omega})^T$ for the left-handed heavy lepton doublets. All left-handed neutrinos of $\tilde{\mathcal{N}}_L$ are sterile neutrinos, which are \mathcal{G}_{SM} -singlets and do not couple to the EW gauge bosons.

By following the symmetry breaking pattern in Eq. (1), we tabulate the fermion representations at various stages of the SU(8) model in Tabs. 2, 3, and 4. All U(1) charges at different stages are obtained according to Eqs. (108a), (108b), (108c), and (108d). According to the rule by Georgi [5], it is straightforward to find $n_g = 3$ in the current setup. All chiral fermions are named by their \mathcal{G}_{SM} irreps. For the right-handed down-type quarks of $\mathcal{D}_R^{\Omega c}$, they are named as follows

$$\begin{aligned} \mathcal{D}_R^{i c} &\equiv d_R^c, \quad \mathcal{D}_R^{j c} \equiv s_R^c, \quad \mathcal{D}_R^{\text{VII} c} \equiv \mathfrak{D}_R^{\text{VII} c}, \quad \mathcal{D}_R^{\text{VIII} c} \equiv \mathfrak{D}_R^{\text{VIII} c}, \quad \mathcal{D}_R^{\text{IX} c} \equiv \mathfrak{D}_R^{\text{IX} c}, \\ \mathcal{D}_R^{3 c} &\equiv b_R^c, \quad \mathcal{D}_R^{\text{IV} c} \equiv \mathfrak{D}_R^{\text{IV} c}, \quad \mathcal{D}_R^{\text{V} c} \equiv \mathfrak{D}_R^{\text{V} c}, \quad \mathcal{D}_R^{\text{VI} c} \equiv \mathfrak{D}_R^{\text{VI} c}. \end{aligned} \quad (12)$$

SU(8)	\mathcal{G}_{441}	\mathcal{G}_{341}	\mathcal{G}_{331}	\mathcal{G}_{SM}
28_F	$(\mathbf{6}, \mathbf{1}, -\frac{1}{2})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}} : \mathcal{D}_L$
		$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}} : t_R^c$
	$(\mathbf{1}, \mathbf{6}, +\frac{1}{2})_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{6}, +\frac{1}{2})_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{3}, +\frac{1}{3})_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{2}, +\frac{1}{2})_{\mathbf{F}} : (\epsilon_R^c, \mathbf{n}_R^c)^T$
				$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}} : \check{\mathbf{n}}_R^c$
	$(\mathbf{4}, \mathbf{4}, 0)_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{4}, -\frac{1}{12})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{3}, 0)_{\mathbf{F}}$	$(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'_{\mathbf{F}} : (\mathbf{n}'_R{}^c, -\epsilon'_R{}^c)^T$
				$(\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}} : \tau_R^c$
		$(\mathbf{1}, \mathbf{4}, +\frac{1}{4})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{2}, +\frac{1}{6})_{\mathbf{F}} : (t_L, b_L)^T$
			$(\mathbf{1}, \mathbf{3}, +\frac{1}{3})''_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})'_{\mathbf{F}} : \mathcal{D}'_L$
				$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''_{\mathbf{F}} : \mathcal{D}''_L$
			$(\mathbf{1}, \mathbf{1}, 0)''_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{2}, +\frac{1}{2})_{\mathbf{F}} : (\epsilon''_R{}^c, \mathbf{n}''_R{}^c)^T$
			$(\mathbf{1}, \mathbf{1}, 0)'_{\mathbf{F}} : \check{\mathbf{n}}'_R{}^c$	
			$(\mathbf{1}, \mathbf{1}, 0)''_{\mathbf{F}} : \check{\mathbf{n}}''_R{}^c$	

Table 3: The SU(8) fermion representation of **28_F** under the $\mathcal{G}_{441}, \mathcal{G}_{341}, \mathcal{G}_{331}, \mathcal{G}_{\text{SM}}$ subgroups for the three-generational SU(8) model. All SM fermions are marked with underlines.

SU(8)	\mathcal{G}_{441}	\mathcal{G}_{341}	\mathcal{G}_{331}	\mathcal{G}_{SM}	
56_F	$(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})_{\mathbf{F}}$	$(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})_{\mathbf{F}}$	$(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})'_{\mathbf{F}}$	$(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'''_{\mathbf{F}} : (\mathbf{n}'''_R{}^c, -\epsilon'''_R{}^c)^T$	
				$(\mathbf{1}, \mathbf{1}, +1)''_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{1}, +1)'_{\mathbf{F}} : \mu_R^c$
	$(\bar{\mathbf{4}}, \mathbf{1}, -\frac{3}{4})_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{1}, +1)''_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}} : \mathfrak{E}_R^c$
		$(\mathbf{1}, \mathbf{1}, -1)_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{1}, -1)_{\mathbf{F}}$		$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}} : u_R^c$
	$(\mathbf{4}, \mathbf{6}, +\frac{1}{4})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{6}, +\frac{1}{6})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{3}, 0)'_{\mathbf{F}}$		$(\mathbf{1}, \mathbf{1}, -1)_{\mathbf{F}} : \mathfrak{E}_L$
				$(\mathbf{3}, \bar{\mathbf{3}}, +\frac{1}{3})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{2}, +\frac{1}{6})'_{\mathbf{F}} : (c_L, s_L)^T$
	$(\mathbf{6}, \mathbf{4}, -\frac{1}{4})_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{4}, -\frac{1}{12})'_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''_{\mathbf{F}}$		$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})'''_{\mathbf{F}} : \mathcal{D}''_L$
				$(\mathbf{1}, \mathbf{3}, +\frac{1}{3})'_{\mathbf{F}}$	$(\mathbf{3}, \bar{\mathbf{2}}, +\frac{1}{6})''_{\mathbf{F}} : (\mathfrak{d}_L, -u_L)^T$
		$(\mathbf{1}, \mathbf{6}, +\frac{1}{2})'_{\mathbf{F}}$	$(\mathbf{1}, \mathbf{3}, +\frac{1}{3})'_{\mathbf{F}}$		$(\mathbf{3}, \mathbf{1}, +\frac{2}{3})_{\mathbf{F}} : \mathfrak{U}_L$
			$(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})''_{\mathbf{F}}$		$(\mathbf{1}, \mathbf{2}, +\frac{1}{2})'''_{\mathbf{F}} : (\epsilon'''_R{}^c, \mathbf{n}'''_R{}^c)^T$
					$(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}} : \check{\mathbf{n}}_R^c$
			$(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})''_{\mathbf{F}}$		$(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})''''_{\mathbf{F}} : (\mathbf{n}''''_R{}^c, -\epsilon''''_R{}^c)^T$
				$(\mathbf{1}, \mathbf{1}, +1)'''_{\mathbf{F}} : e_R^c$	
		$(\mathbf{3}, \mathbf{4}, -\frac{1}{12})'_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{3}, 0)''_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{2}, +\frac{1}{6})'''_{\mathbf{F}} : (u_L, d_L)^T$	
			$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''''_{\mathbf{F}}$	$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''''_{\mathbf{F}} : \mathcal{D}'''_L$	
	$(\bar{\mathbf{3}}, \mathbf{4}, -\frac{5}{12})_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{3}, -\frac{1}{3})_{\mathbf{F}}$		$(\mathbf{3}, \mathbf{1}, -\frac{1}{3})''''_{\mathbf{F}} : \mathcal{D}''''_L$	
			$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'''_{\mathbf{F}}$	$(\bar{\mathbf{3}}, \mathbf{2}, -\frac{1}{6})_{\mathbf{F}} : (\mathfrak{d}_R^c, u_R^c)^T$	
				$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})''_{\mathbf{F}} : \mathfrak{U}_R^c$	
				$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}} : c_R^c$	

Table 4: The SU(8) fermion representation of **56_F** under the $\mathcal{G}_{441}, \mathcal{G}_{341}, \mathcal{G}_{331}, \mathcal{G}_{\text{SM}}$ subgroups for the three-generational SU(8) model. All SM fermions are marked with underlines.

For the left-handed $SU(2)_W$ lepton doublets of $(\mathcal{E}_L^\Omega, -\mathcal{N}_L^\Omega)$, they are named as follows

$$\begin{aligned}
(\mathcal{E}_L^{\text{I}}, -\mathcal{N}_L^{\text{I}}) &\equiv (e_L, -\nu_{eL}), & (\mathcal{E}_L^{\text{II}}, -\mathcal{N}_L^{\text{II}}) &\equiv (\mu_L, -\nu_{\mu L}), \\
(\mathcal{E}_L^{\text{VII}}, -\mathcal{N}_L^{\text{VII}}) &\equiv (\mathbf{e}'', -\mathbf{n}''), & (\mathcal{E}_L^{\text{VIII}}, -\mathcal{N}_L^{\text{VIII}}) &\equiv (\mathbf{e}''', -\mathbf{n}'''), & (\mathcal{E}_L^{\text{IX}}, -\mathcal{N}_L^{\text{IX}}) &\equiv (\mathbf{e}'''' , -\mathbf{n}''''), \\
(\mathcal{E}_L^{\text{III}}, -\mathcal{N}_L^{\text{III}}) &\equiv (\tau_L, -\nu_{\tau L}), & (\mathcal{E}_L^{\text{IV}}, -\mathcal{N}_L^{\text{IV}}) &\equiv (\mathbf{e}'', -\mathbf{n}''), \\
(\mathcal{E}_L^{\text{V}}, -\mathcal{N}_L^{\text{V}}) &\equiv (\mathbf{e}_L, -\mathbf{n}_L), & (\mathcal{E}_L^{\text{VI}}, -\mathcal{N}_L^{\text{VI}}) &\equiv (\mathbf{e}'_L, -\mathbf{n}'_L).
\end{aligned} \tag{13}$$

Through the analysis in Sec. 2, we shall see that all heavy $(\mathcal{D}^\Omega, \mathcal{E}^\Omega, \mathcal{N}^\Omega)$ (with $\Omega = \text{IV}, \dots, \text{IX}$) acquire vectorlike masses during the intermediate symmetry breaking stages. For the remaining left-handed sterile neutrinos of $(\tilde{\mathcal{N}}_L^\Omega, \tilde{\mathcal{N}}_L^{\Omega'}, \tilde{\mathcal{N}}_L^{\Omega''})$, several of them are massive and they are named as follows

$$\tilde{\mathcal{N}}_L^{\text{IV}'} \equiv \tilde{\mathbf{n}}'_L, \quad \tilde{\mathcal{N}}_L^{\text{IV}''} \equiv \tilde{\mathbf{n}}''_L, \quad \tilde{\mathcal{N}}_L^{\text{V}'} \equiv \tilde{\mathbf{n}}_L, \quad \tilde{\mathcal{N}}_L^{\text{VII}'} \equiv \tilde{\mathbf{n}}'''_L. \tag{14}$$

The flavor indices in Eqs. (12) and (13) are chosen in accordance with the symmetry breaking pattern depicted in Sec. 2. The SM flavor names in Tabs. 2, 3, and 4 have been determined according to the consistent mass hierarchies and the CKM mixing pattern predicted in Ref. [10].

Through Tabs. 2, 3, and 4, it is straightforward to find three-generational SM fermions transform differently in the UV theories. For example, three left-handed quark doublets belong to $(\mathbf{3}, \mathbf{2}, +\frac{1}{6})_{\mathbf{F}} \subset (\mathbf{4}, \mathbf{4}, 0)_{\mathbf{F}}, (\mathbf{3}, \mathbf{2}, +\frac{1}{6})'_{\mathbf{F}} \subset (\mathbf{4}, \mathbf{6}, +\frac{1}{4})_{\mathbf{F}}$, and $(\mathbf{3}, \mathbf{2}, +\frac{1}{6})''_{\mathbf{F}} \subset (\mathbf{6}, \mathbf{4}, -\frac{1}{4})_{\mathbf{F}}$, when one traces back their irreps in the \mathcal{G}_{441} effective theory. Likewise, three $(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}}$'s and three $(\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}}$'s all stem from distinctive \mathcal{G}_{441} irreps. Thus, the flavor non-universality under the flavor-conserving neutral currents of the \mathcal{G}_{441} and \mathcal{G}_{341} theories can be expected. When the theory flows to the \mathcal{G}_{331} along the symmetry breaking pattern in Eq. (1), all three-generational SM fermions together with their heavy partner fermions, have already transformed the same. Therefore, the flavor universality under the flavor-conserving neutral currents of the \mathcal{G}_{331} theory can be expected. These features will be explicitly demonstrated in terms of the neutral currents at each symmetry breaking stage in Sec. 3.

2.4 Decompositions of the $SU(8)$ Higgs fields

We decompose the Higgs fields in Eq. (6) into components that can be responsible for the sequential symmetry breaking pattern in Eq. (1). For Higgs fields of $\overline{\mathbf{8}}_{\mathbf{H},\omega}$, they read

$$\begin{aligned}
\overline{\mathbf{8}}_{\mathbf{H},\omega} &\supset \boxed{(\bar{\mathbf{4}}, \mathbf{1}, +\frac{1}{4})_{\mathbf{H},\omega}} \oplus \boxed{(\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\omega}} \\
&\supset \boxed{(\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\omega}} \supset \boxed{(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},\omega}} \supset \boxed{(\mathbf{1}, \bar{\mathbf{2}}, -\frac{1}{2})_{\mathbf{H},\omega}}.
\end{aligned} \tag{15}$$

For Higgs fields of $\overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}}$, they read

$$\begin{aligned}
\overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}} &\supset (\mathbf{6}, \mathbf{1}, +\frac{1}{2})_{\mathbf{H},\dot{\omega}} \oplus (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{\omega}} \oplus \overline{(\bar{\mathbf{4}}, \bar{\mathbf{4}}, 0)_{\mathbf{H},\dot{\omega}}} \\
&\supset (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{\omega}} \oplus \overline{(\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\dot{\omega}}} \\
&\supset \left[\boxed{(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})'_{\mathbf{H},\dot{\omega}}} \oplus \overline{(\mathbf{1}, \mathbf{3}, -\frac{2}{3})_{\mathbf{H},\dot{\omega}}} \right] \oplus \boxed{(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},\dot{\omega}}} \\
&\supset \left[\boxed{(\mathbf{1}, \bar{\mathbf{2}}, -\frac{1}{2})'_{\mathbf{H},\dot{\omega}}} \oplus \overline{(\mathbf{1}, \mathbf{2}, -\frac{1}{2})_{\mathbf{H},\dot{\omega}}} \right] \oplus \boxed{(\mathbf{1}, \bar{\mathbf{2}}, -\frac{1}{2})_{\mathbf{H},\dot{\omega}}}.
\end{aligned} \tag{16}$$

For Higgs field of $\mathbf{70}_H$, they read

$$\begin{aligned}
\mathbf{70}_H &\supset (\mathbf{1}, \mathbf{1}, -1)''_H \oplus (\mathbf{1}, \mathbf{1}, +1)'''_H \oplus (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_H \oplus (\bar{\mathbf{4}}, \mathbf{4}, -\frac{1}{2})_H \oplus (\mathbf{6}, \mathbf{6}, 0)_H \\
&\supset \underline{(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})'_H} \supset \underline{(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})'''_H} \supset \boxed{(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'''_H}.
\end{aligned} \tag{17}$$

Higgs	$\mathcal{G}_{441} \rightarrow \mathcal{G}_{341}$	$\mathcal{G}_{341} \rightarrow \mathcal{G}_{331}$	$\mathcal{G}_{331} \rightarrow \mathcal{G}_{SM}$	$\mathcal{G}_{SM} \rightarrow \text{SU}(3)_c \otimes \text{U}(1)_{EM}$
$\bar{\mathbf{8}}_{H,\omega}$	$\checkmark (\omega = \text{IV})$	$\checkmark (\omega = \text{V})$	$\checkmark (\omega = \mathbf{3}, \text{VI})$	\checkmark
$\mathbf{28}_{H,\dot{\omega}}$	\times	$\checkmark (\dot{\omega} = \dot{\mathbf{1}}, \text{VII})$	$\checkmark (\dot{\omega} = \dot{\mathbf{2}}, \text{VIII}, \text{IX})$	\checkmark
$\mathbf{70}_H$	\times	\times	\times	\checkmark

Table 5: The Higgs fields and their symmetry breaking directions in the SU(8) theory. The \checkmark and \times represent possible and impossible symmetry breaking directions for a given Higgs field. The flavor indices for the Higgs fields that develop VEVs at different stages are also specified in the parentheses.

All components that are likely to develop VEVs for the sequential symmetry breaking stages are framed with boxes, and other $\text{SU}(3)_c \otimes \text{U}(1)_{EM}$ -exact components are neglected. Accordingly, we mark the allowed/disallowed symmetry breaking directions for all Higgs fields in Tab. 5. Schematically, we assign the Higgs VEVs according to the symmetry breaking pattern in Eq. (1) as follows

$$\mathcal{G}_{441} \rightarrow \mathcal{G}_{341} : \langle (\bar{\mathbf{4}}, \mathbf{1}, +\frac{1}{4})_{H,\text{IV}} \rangle \equiv \frac{1}{\sqrt{2}} W_{\bar{\mathbf{4}},\text{IV}}, \tag{18a}$$

$$\mathcal{G}_{341} \rightarrow \mathcal{G}_{331} : \langle (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{H,\text{V}} \rangle \equiv \frac{1}{\sqrt{2}} w_{\bar{\mathbf{4}},\text{V}}, \langle (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{H,\dot{\mathbf{1}},\text{VII}} \rangle \equiv \frac{1}{\sqrt{2}} w_{\bar{\mathbf{4}},\dot{\mathbf{1}},\text{VII}}, \tag{18b}$$

$$\begin{aligned}
\mathcal{G}_{331} \rightarrow \mathcal{G}_{SM} : \langle (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{H,\mathbf{3},\text{VI}} \rangle &\equiv \frac{1}{\sqrt{2}} V_{\bar{\mathbf{3}},\mathbf{3},\text{VI}} \\
\langle (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{H,\dot{\mathbf{3}},\text{IX}} \rangle &\equiv \frac{1}{\sqrt{2}} V_{\bar{\mathbf{3}},\dot{\mathbf{3}},\text{IX}}, \langle (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})'_{H,\dot{\mathbf{2}},\text{VIII}} \rangle \equiv \frac{1}{\sqrt{2}} V'_{\bar{\mathbf{3}},\dot{\mathbf{2}},\text{VIII}},
\end{aligned} \tag{18c}$$

$$\text{EWSB} : \langle (\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'''_H \rangle \equiv \frac{1}{\sqrt{2}} v_{\text{EW}}. \tag{18d}$$

These VEV assignments are made such that the three-generational SM fermions are likely to generate the observed mass hierarchies and the CKM mixing pattern [10]. Given that the $\mathbf{70}_H$ is self-conjugate, one may speculate two EWSB components of $(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'''_H \subset (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_H$ and $(\mathbf{1}, \mathbf{2}, -\frac{1}{2})'''_H \subset (\bar{\mathbf{4}}, \mathbf{4}, -\frac{1}{2})_H$ from the $\mathbf{70}_H$. In Ref. [9], we have proved that only one component of $(\mathbf{1}, \mathbf{2}, +\frac{1}{2})'''_H \subset \mathbf{70}_H$ is $\text{U}(1)_{B-L}$ -neutral according to the definition in Eq. (11), and thus can develop the EWSB VEV. Furthermore, this component also gives the top quark mass through the Yukawa coupling of $Y_{\mathcal{T}} \mathbf{28}_F \mathbf{28}_F \mathbf{70}_H + H.c.$. An immediate consequence of this result is that the third-generational SM $\mathbf{10}_F$ must reside in the $\mathbf{28}_F$, while the first- and second-generational SM $\mathbf{10}_F$'s must reside in the $\mathbf{56}_F$. The same observation was also pointed out in Ref. [8].

2.5 A summary of the vectorlike fermions of the SWW symmetry breaking pattern

Here, we summarize all vectorlike fermions that acquire masses above the EWSB scale in Tab. 6, with the conventions of flavor indices displayed explicitly.

stages	$Q_e = -\frac{1}{3}$	$Q_e = +\frac{2}{3}$	$Q_e = -1$	$Q_e = 0$
v_{441} { Ω }	\mathfrak{D} IV	-	$(\mathbf{e}'', \mathbf{n}'')$ IV	$\{\check{\mathbf{n}}', \check{\mathbf{n}}''\}$ {IV', IV''}
v_{341} { Ω }	$\mathfrak{d}, \{\mathfrak{D}'', \mathfrak{D}''''\}$ {V, VII}	\mathbf{u}, \mathfrak{U}	$\mathfrak{E}, (\mathbf{e}, \mathbf{n}), (\mathbf{e}''''', \mathbf{n}''''')$ {V, VII}	$\{\check{\mathbf{n}}, \check{\mathbf{n}}'''\}$ {V', VII'}
v_{331} { Ω }	$\{\mathfrak{D}', \mathfrak{D}''', \mathfrak{D}'''''\}$ {VI, VIII, IX}	-	$(\mathbf{e}', \mathbf{n}'), (\mathbf{e}''''', \mathbf{n}''''')$ {VI, VIII, IX}	-

Table 6: The vectorlike fermions at different intermediate symmetry breaking scales in the SU(8) theory.

3 The gauge sector of the SU(8) theory and the flavor non-universality

In this section, we describe the gauge sector according to the SWW symmetry breaking pattern in Eq. (1). The gauge couplings at each symmetry breaking stage will be defined for the derivation of RGEs. Due to the intrinsic symmetry properties, the flavor non-universality of three-generational SM fermions in the \mathcal{G}_{441} and the \mathcal{G}_{341} theories will be explicitly displayed. Some of the earlier and recent references addressing the flavor non-universality scenario with product gauge groups in the UV setup include [19–29].

3.1 The SU(4)_s ⊗ U(1)_{X₀} gauge bosons and gauge couplings

3.1.1 Covariant derivatives and gauge boson masses

We express the SU(4)_s ⊗ U(1)_{X₀} covariant derivatives as follows

$$iD_\mu \psi_4 \equiv i\partial_\mu \psi_4 + (g_{4s} G_\mu^{\bar{A}} T_{\text{SU}(4)}^{\bar{A}} + g_{X_0} \mathcal{X}_0 \mathbb{I}_4 X_{0\mu}) \cdot \psi_4, \quad (19)$$

for the SU(4)_s fundamental representation of ψ_4 , and

$$iD_\mu \psi_{\bar{4}} \equiv i\partial_\mu \psi_{\bar{4}} + (-g_{4s} G_\mu^{\bar{A}} (T_{\text{SU}(4)}^{\bar{A}})^T + g_{X_0} \mathcal{X}_0 \mathbb{I}_4 X_{0\mu}) \cdot \psi_{\bar{4}}, \quad (20)$$

for the SU(4)_s anti-fundamental representation of $\psi_{\bar{4}}$. For the SU(4)_s rank-2 anti-symmetric field of ψ_6 , the covariant derivative acts as follows

$$iD_\mu \psi_6 \equiv i\partial_\mu \psi_6 + g_{4s} G_\mu^{\bar{A}} \left(T_{\text{SU}(4)}^{\bar{A}} \cdot \psi_6 + \psi_6 \cdot (T_{\text{SU}(4)}^{\bar{A}})^T \right) + g_{X_0} \mathcal{X}_0 \mathbb{I}_4 X_{0\mu} \cdot \psi_6. \quad (21)$$

The SU(4)_s generators of $T_{\text{SU}(4)}^{\bar{A}}$ are normalized such that $\text{Tr} \left(T_{\text{SU}(4)}^{\bar{A}} T_{\text{SU}(4)}^{\bar{B}} \right) = \frac{1}{2} \delta^{\bar{A}\bar{B}}$, with $\bar{A}, \bar{B} = 1, \dots, 15$. Specifically, we split the SU(4)_s adjoint indices into $\bar{A} = \underbrace{1, \dots, 8}_A, \underbrace{9, \dots, 15}_{\bar{A}}$. All SU(4)_s gluons

of $G_\mu^{\bar{A}}$ are decomposed into eight massless SU(3)_c adjoint gluons, three massive SU(3)_c fundamental and anti-fundamental vector bosons of $(\tilde{C}_\mu^a, \tilde{C}_{\mu,a})$, and one massive SU(3)_c-singlet vector boson of M_μ . It is useful to define a strong mixing angle of

$$t_{\vartheta_C} \equiv \tan \vartheta_C = \frac{g_{X_0}}{\sqrt{6}g_{4s}}. \quad (22)$$

The explicit form for the gauge fields of $g_{4s}G_\mu^{\bar{A}}T_{\text{SU}(4)}^{\bar{A}} + g_{X_0}\mathcal{X}_0\mathbb{I}_4X_{0\mu}$ can be expressed in terms of a 4×4 matrix in the gauge eigenbasis

$$g_{4s}G_\mu^{\bar{A}}T_{\text{SU}(4)}^{\bar{A}} + g_{X_0}\mathcal{X}_0\mathbb{I}_4X_{0\mu} = \frac{g_{4s}}{2} \begin{pmatrix} & & & G_\mu^9 - iG_\mu^{10} \\ & G_\mu^A T_{\text{SU}(3)}^A & & G_\mu^{11} - iG_\mu^{12} \\ & & & G_\mu^{13} - iG_\mu^{14} \\ G_\mu^9 + iG_\mu^{10} & G_\mu^{11} + iG_\mu^{12} & G_\mu^{13} + iG_\mu^{14} & 0 \end{pmatrix} + \frac{g_{4s}}{2} \text{diag}\left(\frac{1}{\sqrt{6}}G_\mu^{15} + 2\sqrt{6}t_{\vartheta_C}\mathcal{X}_0X_{0\mu}\right)\mathbb{I}_{3 \times 3}, -\frac{3}{\sqrt{6}}G_\mu^{15} + 2\sqrt{6}t_{\vartheta_C}\mathcal{X}_0X_{0\mu}\right). \quad (23)$$

By using the $U(1)_{X_1}$ charge matrix for the $\text{SU}(4)_s$ adjoint in Eq. (109a), we find that

$$\mathcal{X}_1(G_\mu^9 \mp iG_\mu^{10}) = \mathcal{X}_1(G_\mu^{11} \mp iG_\mu^{12}) = \mathcal{X}_1(G_\mu^{13} \mp iG_\mu^{14}) = \mp \frac{1}{3}, \quad (24)$$

through the relation of $[\hat{X}_1(\mathbf{15}_s), G_\mu^{\bar{A}}T_{\text{SU}(4)}^{\bar{A}}] = \mathcal{X}_1 G_\mu^{\bar{A}}T_{\text{SU}(4)}^{\bar{A}}$. Apparently, they are $\text{SU}(3)_c$ fundamental and anti-fundamental vector bosons with electric charges of $\mp \frac{1}{3}$, which are vectorial leptoquarks (LQs). Accordingly, we name these vectorial LQs as

$$\begin{aligned} \frac{1}{\sqrt{2}}(G_\mu^9 - iG_\mu^{10}) &\equiv \tilde{C}_\mu^\clubsuit, & \frac{1}{\sqrt{2}}(G_\mu^{11} - iG_\mu^{12}) &\equiv \tilde{C}_\mu^\diamond, & \frac{1}{\sqrt{2}}(G_\mu^{13} - iG_\mu^{14}) &\equiv \tilde{C}_\mu^\heartsuit, \\ \frac{1}{\sqrt{2}}(G_\mu^9 + iG_\mu^{10}) &\equiv \tilde{C}_{\mu,\clubsuit}, & \frac{1}{\sqrt{2}}(G_\mu^{11} + iG_\mu^{12}) &\equiv \tilde{C}_{\mu,\diamond}, & \frac{1}{\sqrt{2}}(G_\mu^{13} + iG_\mu^{14}) &\equiv \tilde{C}_{\mu,\heartsuit}. \end{aligned} \quad (25)$$

The first symmetry breaking stage is achieved by the Higgs VEV given in Eq. (18a), and the Higgs kinematic terms lead to the gauge boson masses of

$$\frac{1}{4}g_{4s}^2v_{441}^2 \cdot \left[\sum_{a=\clubsuit,\diamond,\heartsuit} \tilde{C}_\mu^a \tilde{C}_{\mu,a} + \frac{3}{4} \left(G_\mu^{15} + t_{\vartheta_C} X_{0\mu} \right)^2 \right], \quad v_{441}^2 = W_{4,\text{IV}}^2. \quad (26)$$

The mass eigenstates are diagonalized in terms of the mixing angle ϑ_C in Eq. (22) as follows

$$\begin{pmatrix} M_\mu \\ X_{1\mu} \end{pmatrix} = \begin{pmatrix} c_{\vartheta_C} & s_{\vartheta_C} \\ -s_{\vartheta_C} & c_{\vartheta_C} \end{pmatrix} \cdot \begin{pmatrix} G_\mu^{15} \\ X_{0\mu} \end{pmatrix}, \quad (27)$$

with $(c_{\vartheta_C}, s_{\vartheta_C}) \equiv (\cos \vartheta_C, \sin \vartheta_C)$.

The $\text{SU}(3)_c \otimes U(1)_{X_1}$ couplings of $(\alpha_{3c}, \alpha_{X_1})$ are related to the $\text{SU}(4)_s \otimes U(1)_{X_0}$ couplings of $(\alpha_{4s}, \alpha_{X_0})$ as

$$\begin{aligned} \alpha_{3c}^{-1}(v_{441}) &= \alpha_{4s}^{-1}(v_{441}), & \alpha_{X_1}^{-1}(v_{441}) &= \frac{1}{6}\alpha_{4s}^{-1}(v_{441}) + \alpha_{X_0}^{-1}(v_{441}), \\ \frac{1}{6}\alpha_{4s}^{-1} &= \alpha_{X_1}^{-1}s_{\vartheta_C}^2, & \alpha_{X_0}^{-1} &= \alpha_{X_1}^{-1}c_{\vartheta_C}^2. \end{aligned} \quad (28)$$

Obviously, three LQs of \tilde{C}_μ^a and one neutral M_μ obtain masses of

$$m_{\tilde{C}_\mu^a}^2 = \frac{g_{X_1}^2}{24s_{\vartheta_C}^2}v_{441}^2, \quad m_{M_\mu}^2 = \frac{g_{X_1}^2}{16s_{\vartheta_C}^2c_{\vartheta_C}^2}v_{441}^2, \quad (29)$$

at this stage of symmetry breaking. The remaining massless gauge boson of $X_{1\mu}$ will further mix with the $\text{SU}(4)_W$ gauge bosons afterwards.

Expressed in terms of mass eigenstates, the gauge fields in Eq. (23) become

$$\begin{aligned}
& g_{4s} G_\mu^{\bar{A}} T_{\text{SU}(4)}^{\bar{A}} + g_{X_0} \mathcal{X}_0 \mathbb{I}_4 X_{0\mu} = \\
& \frac{g_{4s}}{2} \begin{pmatrix} G_\mu^A T_{\text{SU}(3)}^A & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} + \frac{g_{4s}}{\sqrt{2}} \begin{pmatrix} & & & \tilde{C}_\mu^\clubsuit \\ & 0_{3 \times 3} & & \tilde{C}_\mu^\diamond \\ & & & \tilde{C}_\mu^\heartsuit \\ \tilde{C}_\mu^\clubsuit & \tilde{C}_\mu^\diamond & \tilde{C}_\mu^\heartsuit & 0 \end{pmatrix} \\
& + g_{X_1} \text{diag} \left(\left(-\frac{1}{12} + \mathcal{X}_0 \right) \mathbb{I}_{3 \times 3}, \frac{1}{4} + \mathcal{X}_0 \right) X_{1\mu} \\
& + \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \text{diag} \left(\left[\frac{1}{12} + \left(-\frac{1}{12} + \mathcal{X}_0 \right) s_{\vartheta_C}^2 \right] \mathbb{I}_{3 \times 3}, -\frac{1}{4} + \left(\frac{1}{4} + \mathcal{X}_0 \right) s_{\vartheta_C}^2 \right) M_\mu, \tag{30}
\end{aligned}$$

for the $\text{SU}(4)_s$ fundamental representation. It is straightforward to find

$$\begin{aligned}
& -g_{4s} G_\mu^{\bar{A}} (T_{\text{SU}(4)}^{\bar{A}})^T + g_{X_0} \mathcal{X}_0 \mathbb{I}_4 X_{0\mu} \\
& = -\frac{g_{4s}}{2} \begin{pmatrix} G_\mu^A (T_{\text{SU}(3)}^A)^T & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} - \frac{g_{4s}}{\sqrt{2}} \begin{pmatrix} & & & \tilde{C}_\mu^\clubsuit \\ & 0_{3 \times 3} & & \tilde{C}_\mu^\diamond \\ & & & \tilde{C}_\mu^\heartsuit \\ \tilde{C}_\mu^\clubsuit & \tilde{C}_\mu^\diamond & \tilde{C}_\mu^\heartsuit & 0 \end{pmatrix} \\
& + g_{X_1} \text{diag} \left(\left(\frac{1}{12} + \mathcal{X}_0 \right) \mathbb{I}_{3 \times 3}, -\frac{1}{4} + \mathcal{X}_0 \right) X_{1\mu} \\
& + \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \text{diag} \left(\left[-\frac{1}{12} + \left(\frac{1}{12} + \mathcal{X}_0 \right) s_{\vartheta_C}^2 \right] \mathbb{I}_{3 \times 3}, \frac{1}{4} + \left(-\frac{1}{4} + \mathcal{X}_0 \right) s_{\vartheta_C}^2 \right) M_\mu, \tag{31}
\end{aligned}$$

for the $\text{SU}(4)_s$ anti-fundamental representation. For the $\text{SU}(4)_s$ -singlet representations, we have

$$g_{X_0} \mathcal{X}_0 X_{0\mu} = g_{X_1} (+\mathcal{X}_0) X_{1\mu} + \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} (+\mathcal{X}_0 s_{\vartheta_C}^2) M_\mu. \tag{32}$$

3.1.2 Gauge couplings of the \mathcal{G}_{441} fermions

We proceed to obtain the neutral currents mediated by the $\text{SU}(3)_c$ -singlet vector boson of M_μ . For the anti-fundamental fermions of $\bar{\mathbf{8}}_F^\Omega$, we have the following covariant derivatives of

$$i\bar{\psi}_{(\bar{4}, \mathbf{1}, +\frac{1}{4})^\Omega} \not{D} \psi_{(\bar{4}, \mathbf{1}, +\frac{1}{4})^\Omega} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(\frac{1}{12} - \frac{1}{3} s_{\vartheta_C}^2 \right) \overline{\mathcal{D}}_R^\Omega \gamma^\mu \mathcal{D}_R^\Omega + \left(+\frac{1}{4} \right) \overline{\mathcal{N}}_L^\Omega \gamma^\mu \check{\mathcal{N}}_L^\Omega \right] \cdot M_\mu, \tag{33a}$$

$$\begin{aligned}
& i\bar{\psi}_{(\mathbf{1}, \bar{4}, -\frac{1}{4})^\Omega} \not{D} \psi_{(\mathbf{1}, \bar{4}, -\frac{1}{4})^\Omega} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left(-\frac{1}{4} s_{\vartheta_C}^2 \right) \left[\overline{\mathcal{E}}_L^\Omega \gamma^\mu \mathcal{E}_L^\Omega + \overline{\mathcal{N}}_L^\Omega \gamma^\mu \mathcal{N}_L^\Omega \right. \\
& + \left. \overline{\check{\mathcal{N}}}_L^{\Omega'} \gamma^\mu \check{\mathcal{N}}_L^{\Omega'} + \overline{\check{\mathcal{N}}}_L^{\Omega''} \gamma^\mu \check{\mathcal{N}}_L^{\Omega''} \right] \cdot M_\mu. \tag{33b}
\end{aligned}$$

For the rank-2 anti-symmetric fermions of $\mathbf{28}_F$, we have the covariant derivatives of

$$\begin{aligned} & \text{Tr} \left[i\bar{\psi}_{(\mathbf{6}, \mathbf{1}, -\frac{1}{2})} \not{D}\psi_{(\mathbf{6}, \mathbf{1}, -\frac{1}{2})} \right] \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(-\frac{1}{6} + \frac{2}{3} s_{\vartheta_C}^2 \right) \bar{t}_R \gamma^\mu t_R \right. \\ & + \left. \left(-\frac{1}{6} - \frac{1}{3} s_{\vartheta_C}^2 \right) \bar{\mathcal{D}}_L \gamma^\mu \mathcal{D}_L \right] \cdot M_\mu, \end{aligned} \quad (34a)$$

$$\begin{aligned} & i\bar{\psi}_{(\mathbf{1}, \mathbf{6}, +\frac{1}{2})} \not{D}\psi_{(\mathbf{1}, \mathbf{6}, +\frac{1}{2})} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left(-\frac{1}{2} s_{\vartheta_C}^2 \right) \left[\bar{\mathbf{e}}_R \gamma^\mu \mathbf{e}_R + \bar{\mathbf{n}}_R \gamma^\mu \mathbf{n}_R + \bar{\check{\mathbf{n}}}_R \gamma^\mu \check{\mathbf{n}}_R \right. \\ & + \left. \bar{\mathbf{n}}'_R \gamma^\mu \mathbf{n}'_R + \bar{\mathbf{e}}'_R \gamma^\mu \mathbf{e}'_R + \bar{\tau}_R \gamma^\mu \tau_R \right] \cdot M_\mu, \end{aligned} \quad (34b)$$

$$\begin{aligned} & i\bar{\psi}_{(\mathbf{4}, \mathbf{4}, 0)} \not{D}\psi_{(\mathbf{4}, \mathbf{4}, 0)} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(\frac{1}{12} - \frac{1}{12} s_{\vartheta_C}^2 \right) (\bar{t}_L \gamma^\mu t_L + \bar{b}_L \gamma^\mu b_L + \bar{\mathcal{D}}'_L \gamma^\mu \mathcal{D}'_L + \bar{\mathcal{D}}''_L \gamma^\mu \mathcal{D}''_L) \right. \\ & + \left. \left(\frac{1}{4} - \frac{1}{4} s_{\vartheta_C}^2 \right) (\bar{\mathbf{e}}''_R \gamma^\mu \mathbf{e}''_R + \bar{\mathbf{n}}''_R \gamma^\mu \mathbf{n}''_R + \bar{\check{\mathbf{n}}}'_R \gamma^\mu \check{\mathbf{n}}'_R + \bar{\check{\mathbf{n}}}_R \gamma^\mu \check{\mathbf{n}}_R) \right] \cdot M_\mu. \end{aligned} \quad (34c)$$

For the rank-3 anti-symmetric fermions of $\mathbf{56}_F$, we have the covariant derivatives of

$$\begin{aligned} & i\bar{\psi}_{(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})} \not{D}\psi_{(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left(-\frac{3}{4} s_{\vartheta_C}^2 \right) \left[\bar{\mathbf{n}}'''_R \gamma^\mu \mathbf{n}'''_R + \bar{\mathbf{e}}'''_R \gamma^\mu \mathbf{e}'''_R + \bar{\mu}_R \gamma^\mu \mu_R \right. \\ & + \left. \bar{\mathbf{e}}_R \gamma^\mu \mathbf{e}_R \right] \cdot M_\mu, \end{aligned} \quad (35a)$$

$$i\bar{\psi}_{(\bar{\mathbf{4}}, \mathbf{1}, -\frac{3}{4})} \not{D}\psi_{(\bar{\mathbf{4}}, \mathbf{1}, -\frac{3}{4})} \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(\frac{1}{12} + \frac{2}{3} s_{\vartheta_C}^2 \right) \bar{u}_R \gamma^\mu u_R + \left(\frac{1}{4} - s_{\vartheta_C}^2 \right) \bar{\mathbf{e}}_L \gamma^\mu \mathbf{e}_L \right] \cdot M_\mu, \quad (35b)$$

$$\begin{aligned} & i\bar{\psi}_{(\mathbf{4}, \mathbf{6}, +\frac{1}{4})} \not{D}\psi_{(\mathbf{4}, \mathbf{6}, +\frac{1}{4})} \\ & \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(\frac{1}{12} + \frac{1}{6} s_{\vartheta_C}^2 \right) (\bar{c}_L \gamma^\mu c_L + \bar{s}_L \gamma^\mu s_L + \bar{\mathcal{D}}'''_L \gamma^\mu \mathcal{D}'''_L + \bar{\mathcal{D}}_L \gamma^\mu \mathcal{D}_L + \bar{u}_L \gamma^\mu u_L + \bar{\mathfrak{U}}_L \gamma^\mu \mathfrak{U}_L) \right. \\ & + \left. \left(\frac{1}{4} - \frac{1}{2} s_{\vartheta_C}^2 \right) (\bar{\mathbf{e}}''''_R \gamma^\mu \mathbf{e}''''_R + \bar{\mathbf{n}}''''_R \gamma^\mu \mathbf{n}''''_R + \bar{\check{\mathbf{n}}}'''_R \gamma^\mu \check{\mathbf{n}}'''_R + \bar{\mathbf{n}}''''_R \gamma^\mu \mathbf{n}''''_R + \bar{\mathbf{e}}''''_R \gamma^\mu \mathbf{e}''''_R + \bar{e}_R \gamma^\mu e_R) \right] \cdot M_\mu, \end{aligned} \quad (35c)$$

$$\begin{aligned} & \text{Tr} \left[i\bar{\psi}_{(\mathbf{6}, \mathbf{4}, -\frac{1}{4})} \not{D}\psi_{(\mathbf{6}, \mathbf{4}, -\frac{1}{4})} \right] \supset \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} \left[\left(-\frac{1}{6} + \frac{5}{12} s_{\vartheta_C}^2 \right) (\bar{\mathfrak{D}}_R \gamma^\mu \mathfrak{D}_R + \bar{u}_R \gamma^\mu u_R + \bar{\mathfrak{U}}_R \gamma^\mu \mathfrak{U}_R + \bar{c}_R \gamma^\mu c_R) \right. \\ & + \left. \left(-\frac{1}{6} - \frac{1}{12} s_{\vartheta_C}^2 \right) (\bar{u}_L \gamma^\mu u_L + \bar{d}_L \gamma^\mu d_L + \bar{\mathcal{D}}''''_L \gamma^\mu \mathcal{D}''''_L + \bar{\mathcal{D}}'''_L \gamma^\mu \mathcal{D}'''_L) \right] \cdot M_\mu. \end{aligned} \quad (35d)$$

The flavor-conserving neutral currents mediated by M_μ are expressed as follows in the $V - A$ basis

$$\mathcal{L}_{\text{SU}(4)_s}^{\text{NC}, \text{F}} = \frac{g_{X_1}}{s_{\vartheta_C} c_{\vartheta_C}} (g_f^V \mathcal{M} \bar{f} \gamma^\mu f + g_f^A \mathcal{M} \bar{f} \gamma^\mu \gamma_5 f) M_\mu, \quad (36)$$

and we tabulate the vectorial and axial couplings of $(g_f^V \mathcal{M}, g_f^A \mathcal{M})$ in Tab. 7. The flavor non-universality of the neutral currents mediated by M_μ among three generational SM quarks and leptons are displayed explicitly.

3.2 The $\text{SU}(4)_W \otimes \text{U}(1)_{X_1}$ gauge bosons and gauge couplings

3.2.1 Covariant derivatives and gauge boson masses

We express the $\text{SU}(4)_W \otimes \text{U}(1)_{X_1}$ covariant derivatives as follows

$$iD_\mu \psi_4 \equiv i\partial_\mu \psi_4 + (g_{4W} A_\mu^{\bar{I}} T_{\text{SU}(4)}^{\bar{I}} + g_{X_1} \mathcal{X}_1 \mathbb{I}_4 X_{1\mu}) \cdot \psi_4, \quad (37)$$

	u	c	t	u	\mathfrak{U}	
g_f^{VM}	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$	
g_f^{AM}	$\frac{1}{8} + \frac{3}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{3}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{1}{8}s_{\vartheta_C}^2$	
	d	s	b	\mathfrak{D}		
g_f^{VM}	$-\frac{1}{24} - \frac{5}{24}s_{\vartheta_C}^2$	$\frac{1}{12} - \frac{1}{12}s_{\vartheta_C}^2$	$\frac{1}{12} - \frac{5}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} + \frac{7}{24}s_{\vartheta_C}^2$		
g_f^{AM}	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{4}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{1}{8}s_{\vartheta_C}^2$		
	\mathfrak{D}	\mathfrak{D}'	\mathfrak{D}''	\mathfrak{D}'''	\mathfrak{D}''''	\mathfrak{D}'''''
g_f^{VM}	$-\frac{1}{24} - \frac{1}{3}s_{\vartheta_C}^2$	$\frac{1}{12} - \frac{5}{24}s_{\vartheta_C}^2$	$\frac{1}{12} - \frac{5}{24}s_{\vartheta_C}^2$	$\frac{1}{12} - \frac{1}{12}s_{\vartheta_C}^2$	$-\frac{1}{24} - \frac{5}{24}s_{\vartheta_C}^2$	$-\frac{1}{24} - \frac{5}{24}s_{\vartheta_C}^2$
g_f^{AM}	$\frac{1}{8}$	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{4}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$
	e	μ	τ	\mathfrak{E}		
g_f^{VM}	$\frac{1}{8} - \frac{3}{8}s_{\vartheta_C}^2$	$-\frac{1}{2}s_{\vartheta_C}^2$	$-\frac{3}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{7}{8}s_{\vartheta_C}^2$		
g_f^{AM}	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{4}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8} + \frac{1}{8}s_{\vartheta_C}^2$		
	$(\mathfrak{e}, \mathfrak{n})$	$(\mathfrak{e}', \mathfrak{n}')$	$(\mathfrak{e}'', \mathfrak{n}'')$	$(\mathfrak{e}''', \mathfrak{n}''')$	$(\mathfrak{e}'''', \mathfrak{n}'''')$	$(\mathfrak{e}''''', \mathfrak{n}''''')$
g_f^{VM}	$-\frac{3}{8}s_{\vartheta_C}^2$	$-\frac{3}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{4}s_{\vartheta_C}^2$	$-\frac{1}{2}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{3}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{3}{8}s_{\vartheta_C}^2$
g_f^{AM}	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$\frac{1}{8}$	$-\frac{1}{4}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$
	$\nu_{eL}/\nu_{\mu L}/\nu_{\tau L}$	$\tilde{\mathcal{N}}_L^{\Omega, \Omega', \Omega''}$	$\tilde{\mathfrak{n}}$	$\tilde{\mathfrak{n}}'$	$\tilde{\mathfrak{n}}''$	$\tilde{\mathfrak{n}}'''$
g_f^{VM}	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{3}{8}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{4}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{1}{4}s_{\vartheta_C}^2$	$\frac{1}{8} - \frac{3}{8}s_{\vartheta_C}^2$
g_f^{AM}	$\frac{1}{8}s_{\vartheta_C}^2$	$\frac{1}{8}s_{\vartheta_C}^2$	$-\frac{1}{8}s_{\vartheta_C}^2$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8} - \frac{1}{8}s_{\vartheta_C}^2$

Table 7: The couplings of the neutral currents mediated by M_μ in the $V - A$ basis. The $\tilde{\mathcal{N}}_L^{\Omega, \Omega', \Omega''}$ represent all 23 left-handed massless sterile neutrinos except four massive sterile neutrinos of $(\tilde{\mathfrak{n}}, \tilde{\mathfrak{n}}', \tilde{\mathfrak{n}}'', \tilde{\mathfrak{n}}''')$.

for the $SU(4)_W$ fundamental representation, and

$$iD_\mu \psi_{\bar{\mathbf{4}}} \equiv i\partial_\mu \psi_{\bar{\mathbf{4}}} + (-g_{4W} A_\mu^{\bar{I}} (T_{SU(4)}^{\bar{I}})^T + g_{X_1} \mathcal{X}_1 \mathbb{1}_4 X_{1\mu}) \cdot \psi_{\bar{\mathbf{4}}}, \quad (38)$$

for the $SU(4)_W$ anti-fundamental representation. The $SU(4)_W$ generators of $T_{SU(4)}^a$ are normalized such that $\text{Tr} \left(T_{SU(4)}^{\bar{I}} T_{SU(4)}^{\bar{J}} \right) = \frac{1}{2} \delta^{\bar{I}\bar{J}}$. For the $SU(4)_W$ rank-2 anti-symmetric field of $\psi_{\mathbf{6}}$, the covariant derivative acts as follows

$$iD_\mu \psi_{\mathbf{6}} \equiv i\partial_\mu \psi_{\mathbf{6}} + g_{4W} A_\mu^{\bar{I}} (T_{SU(4)}^{\bar{I}}) \cdot \psi_{\mathbf{6}} + \psi_{\mathbf{6}} \cdot T_{SU(4)}^{\bar{I}, T} + g_{X_1} \mathcal{X}_1 X_{1\mu} \psi_{\mathbf{6}}. \quad (39)$$

The corresponding kinematic term for the rank-2 anti-symmetric field of $\psi_{\mathbf{6}}$ should be $\text{Tr}(\bar{\psi}_{\mathbf{6}} i \not{D} \psi_{\mathbf{6}})$. For the singlet of $\psi_{\mathbf{1}}$, the covariant derivative is simply given by

$$iD_\mu \psi_{\mathbf{1}} \equiv (i\partial_\mu + g_{X_1} \mathcal{X}_1 X_{1\mu}) \psi_{\mathbf{1}}, \quad (40)$$

The explicit form for the gauge fields of $g_{4W}A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}} + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu}$ can be expressed in terms of a 4×4 matrix below

$$\begin{aligned}
& g_{4W}A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}} + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu} \\
&= \frac{1}{2}g_{4W} \begin{pmatrix} 0 & A_\mu^1 - iA_\mu^2 & A_\mu^4 - iA_\mu^5 & A_\mu^9 - iA_\mu^{10} \\ A_\mu^1 + iA_\mu^2 & 0 & A_\mu^6 - iA_\mu^7 & A_\mu^{11} - iA_\mu^{12} \\ A_\mu^4 + iA_\mu^5 & A_\mu^6 + iA_\mu^7 & 0 & A_\mu^{13} - iA_\mu^{14} \\ A_\mu^9 + iA_\mu^{10} & A_\mu^{11} + iA_\mu^{12} & A_\mu^{13} + iA_\mu^{14} & 0 \end{pmatrix} \\
&+ \frac{g_{4W}}{2} \text{diag} \left(A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -\frac{2}{\sqrt{3}}A_\mu^8, 0 \right) \\
&+ \frac{g_{4W}}{2\sqrt{6}} \text{diag} \left([A_\mu^{15} + 12t_{\vartheta_G}\mathcal{X}_1X_{1\mu}] \mathbb{I}_{3 \times 3}, -3A_\mu^{15} + 12t_{\vartheta_G}\mathcal{X}_1X_{1\mu} \right), \tag{41}
\end{aligned}$$

where we have defined an $\text{SU}(4)_W$ mixing angle of

$$t_{\vartheta_G} \equiv \tan \vartheta_G = \frac{g_{X_1}}{\sqrt{6}g_{4W}}. \tag{42}$$

By using the relation of $[\hat{Q}_e(\mathbf{15}_W), A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}}] = Q_e A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}}$ with $\hat{Q}_e(\mathbf{15}_W)$ in Eq. (109d), we find the electrical charges of gauge bosons as follows

$$Q_e(A_\mu^1 \mp iA_\mu^2) = Q_e(A_\mu^4 \mp iA_\mu^5) = Q_e(A_\mu^9 \mp iA_\mu^{10}) = \pm 1, \tag{43}$$

while all other gauge bosons are electrically neutral. Correspondingly, the gauge fields of $g_{4W}A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}} + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu}$ can also be expressed as

$$\begin{aligned}
& g_{4W}A_\mu^{\bar{I}}T_{\text{SU}(4)}^{\bar{I}} + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu} = \\
& \frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ & W_\mu'^+ & W_\mu''^+ \\ W_\mu^- & & & \\ W_\mu'^- & & 0_{3 \times 3} & \\ W_\mu''^- & & & \end{pmatrix} + \frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & N_\mu & N_\mu' \\ 0 & \bar{N}_\mu & 0 & \bar{N}_\mu'' \\ 0 & \bar{N}_\mu' & \bar{N}_\mu'' & 0 \end{pmatrix} \\
&+ \frac{g_{4W}}{2} \text{diag} \left(A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -\frac{2}{\sqrt{3}}A_\mu^8, 0 \right) \\
&+ \frac{g_{4W}}{2\sqrt{6}} \text{diag} \left([A_\mu^{15} + 12t_{\vartheta_G}\mathcal{X}_1X_{1\mu}] \mathbb{I}_{3 \times 3}, -3A_\mu^{15} + 12t_{\vartheta_G}\mathcal{X}_1X_{1\mu} \right). \tag{44}
\end{aligned}$$

Here, all diagonal gauge bosons are neutral, while the off-diagonal components contain both the electrically charged gauge bosons of $(W_\mu^\pm, W_\mu'^\pm, W_\mu''^\pm)$ as well as the neutral gauge bosons of (N_μ, N_μ', N_μ'') .

This stage of symmetry breaking is achieved by Higgs VEVs in Eq. (18b), and the Higgs kinematic terms lead to the $(W_\mu''^\pm, N_\mu', N_\mu'')$ gauge boson masses of

$$\frac{1}{4}g_{4W}^2v_{341}^2 \cdot \left(W_\mu''^+W_\mu''^- + N_\mu'\bar{N}_\mu'' + N_\mu''\bar{N}_\mu' \right), \quad v_{341}^2 = (w_{\bar{4},V})^2 + (w_{\bar{4},I})^2 + (w_{\bar{4},VII})^2. \tag{45}$$

For the flavor-conserving neutral gauge bosons of $(A_\mu^{15}, X_{1\mu})$, the mass squared matrix reads

$$\frac{3}{16}g_{4W}^2v_{341}^2(A_\mu^{15}, X_{1\mu}) \cdot \begin{pmatrix} 1 & -t_{\vartheta_G} \\ -t_{\vartheta_G} & t_{\vartheta_G}^2 \end{pmatrix} \cdot \begin{pmatrix} A_\mu^{15} \\ X_{1\mu} \end{pmatrix}. \tag{46}$$

Obviously it contains a zero eigenvalue which corresponds to the massless gauge boson of $X_{2\mu}$ after the $SU(4)_W$ symmetry breaking. The gauge eigenstates of $(A_\mu^{15}, X_{1\mu})$ are diagonalized as follows

$$\begin{pmatrix} Z''_\mu \\ X_{2\mu} \end{pmatrix} = \begin{pmatrix} c_{\vartheta_G} & -s_{\vartheta_G} \\ s_{\vartheta_G} & c_{\vartheta_G} \end{pmatrix} \cdot \begin{pmatrix} A_\mu^{15} \\ X_{1\mu} \end{pmatrix}. \quad (47)$$

The $SU(4)_W \otimes U(1)_{X_1}$ gauge couplings of $(\alpha_{4W}, \alpha_{X_1})$ match with the $SU(3)_W \otimes U(1)_{X_2}$ gauge couplings as follows

$$\begin{aligned} \alpha_{3W}^{-1}(v_{341}) &= \alpha_{4W}^{-1}(v_{341}), \quad \alpha_{X_2}^{-1}(v_{341}) = \frac{1}{6}\alpha_{4W}^{-1}(v_{341}) + \alpha_{X_1}^{-1}(v_{341}), \\ \frac{1}{6}\alpha_{4W}^{-1} &= \alpha_{X_2}^{-1}s_{\vartheta_G}^2, \quad \alpha_{X_1}^{-1} = \alpha_{X_2}^{-1}c_{\vartheta_G}^2. \end{aligned} \quad (48)$$

According to the definitions of Eqs. (42) and (61), we find a relation of

$$\boxed{\sin \vartheta_G = \frac{1}{\sqrt{2}} \tan \vartheta_S}. \quad (49)$$

The tree-level masses for seven gauge bosons at this stage read

$$m_{W_\mu^{\prime\pm}}^2 = m_{N'_\mu, \bar{N}'_\mu}^2 = m_{N''_\mu, \bar{N}''_\mu}^2 = \frac{g_{X_2}^2}{4s_{\vartheta_G}^2} v_{341}^2, \quad m_{Z''_\mu}^2 = \frac{g_{X_2}^2}{16s_{\vartheta_G}^2 c_{\vartheta_G}^2} v_{341}^2. \quad (50)$$

The remaining massless gauge bosons are $(W_\mu^\pm, W_\mu^{\prime\pm}, N_\mu, \bar{N}_\mu)$ and $(A_\mu^3, A_\mu^8, X_{2\mu})$.

In terms of mass eigenstates, the vector bosons from the covariant derivative in Eq. (41) becomes

$$\begin{aligned} & g_{4W} A_\mu^{\bar{I}} T_{SU(4)}^{\bar{I}} + g_{X_1} \mathcal{X}_1 \mathbb{I}_4 X_{1\mu} \\ &= \frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ & W_\mu^{\prime+} & 0 \\ W_\mu^- & 0 & N_\mu & 0 \\ W_\mu^{\prime-} & \bar{N}_\mu & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & W_\mu^{\prime\prime+} \\ 0 & 0 & 0 & N'_\mu \\ 0 & 0 & 0 & N''_\mu \\ W_\mu^{\prime\prime-} & \bar{N}'_\mu & \bar{N}''_\mu & 0 \end{pmatrix} \\ &+ \frac{g_{4W}}{2} \text{diag}\left(A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -\frac{2}{\sqrt{3}}A_\mu^8, 0\right) \\ &+ g_{X_2} \text{diag}\left(\left(\frac{1}{12} + \mathcal{X}_1\right)\mathbb{I}_{3\times 3}, -\frac{1}{4} + \mathcal{X}_1\right) X_{2\mu} \\ &+ \frac{g_{X_2}}{s_{\vartheta_G} c_{\vartheta_G}} \text{diag}\left(\left[\frac{1}{12} - \left(\frac{1}{12} + \mathcal{X}_1\right)s_{\vartheta_G}^2\right]\mathbb{I}_{3\times 3}, -\frac{1}{4} + \left(\frac{1}{4} - \mathcal{X}_1\right)s_{\vartheta_G}^2\right) Z''_\mu. \end{aligned} \quad (51)$$

We split terms with massless vector bosons and massive bosons (marked with underlines), respectively. As a consistent check, the (44)-component in Eq. (51) is reduced to $-\frac{g_{X_2}}{2s_{\vartheta_G} c_{\vartheta_G}} Z''_\mu$ when setting $\mathcal{X}_1 = +\frac{1}{4}$ for the fundamental representation. Likewise, we find the explicit form the gauge fields of

$-g_{4W}A_\mu^{\bar{I}}(T_{\text{SU}(4)}^{\bar{I}})^T + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu}$ as follows

$$\begin{aligned}
& -g_{4W}A_\mu^{\bar{I}}(T_{\text{SU}(4)}^{\bar{I}})^T + g_{X_1}\mathcal{X}_1\mathbb{I}_4X_{1\mu} = \\
& -\frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^- & W_\mu'^- & 0 \\ W_\mu^+ & 0 & N_\mu & 0 \\ W_\mu'^+ & N_\mu & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} - \frac{g_{4W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & W_\mu''- \\ 0 & 0 & 0 & N_\mu' \\ 0 & 0 & 0 & N_\mu'' \\ W_\mu''+ & N_\mu' & N_\mu'' & 0 \end{pmatrix} \\
& - \frac{g_{4W}}{2} \text{diag} \left(A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -A_\mu^3 + \frac{1}{\sqrt{3}}A_\mu^8, -\frac{2}{\sqrt{3}}A_\mu^8, 0 \right) \\
& + g_{X_2} \text{diag} \left(\left(-\frac{1}{12} + \mathcal{X}_1\right)\mathbb{I}_{3 \times 3}, \frac{1}{4} + \mathcal{X}_1 \right) X_{2\mu} \\
& + \frac{g_{X_2}}{s_{\vartheta_G}c_{\vartheta_G}} \text{diag} \left(\left[-\frac{1}{12} + \left(\frac{1}{12} - \mathcal{X}_1\right)s_{\vartheta_G}^2\right]\mathbb{I}_{3 \times 3}, \frac{1}{4} - \left(\frac{1}{4} + \mathcal{X}_1\right)s_{\vartheta_G}^2 \right) Z_\mu''. \tag{52}
\end{aligned}$$

3.2.2 Gauge couplings of the \mathcal{G}_{341} fermions

For the anti-fundamental fermions of $\overline{\mathbf{8}}_{\mathbf{F}}^\Omega$, the covariant derivatives lead to the neutral currents of

$$i\bar{\psi}_{(\mathbf{3}, \mathbf{1}, -\frac{1}{3})^\Omega} \mathcal{D} \psi_{(\mathbf{3}, \mathbf{1}, -\frac{1}{3})^\Omega} \supset \frac{g_{X_2}}{s_{\vartheta_G}c_{\vartheta_G}} \left(+\frac{1}{3}s_{\vartheta_G}^2 \right) \overline{\mathcal{D}}_R^\Omega \gamma^\mu \mathcal{D}_R^\Omega \cdot Z_\mu'', \tag{53a}$$

$$\begin{aligned}
& i\bar{\psi}_{(\mathbf{1}, \overline{\mathbf{4}}, -\frac{1}{4})^\Omega} \mathcal{D} \psi_{(\mathbf{1}, \overline{\mathbf{4}}, -\frac{1}{4})^\Omega} \\
& \supset \frac{g_{X_2}}{s_{\vartheta_G}c_{\vartheta_G}} \left[\left(+\frac{1}{4} \right) \overline{\mathcal{N}}_L^{\Omega''} \gamma^\mu \check{\mathcal{N}}_L^{\Omega''} + \left(-\frac{1}{12} + \frac{1}{3}s_{\vartheta_G}^2 \right) \left(\overline{\mathcal{E}}_L^\Omega \gamma^\mu \mathcal{E}_L^\Omega + \overline{\mathcal{N}}_L^\Omega \gamma^\mu \mathcal{N}_L^\Omega + \overline{\check{\mathcal{N}}}_L^{\Omega'} \gamma^\mu \check{\mathcal{N}}_L^{\Omega'} \right) \right] \cdot Z_\mu''. \tag{53b}
\end{aligned}$$

For the rank-2 anti-symmetric fermionic representation of $(\mathbf{1}, \mathbf{6}, +\frac{1}{2})_{\mathbf{F}}$ from the $\mathbf{28}_{\mathbf{F}}$, we first transform to its conjugate of $(\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{F}} = (\mathbf{1}, \mathbf{3}, -\frac{2}{3})_{\mathbf{F}} \oplus (\mathbf{1}, \overline{\mathbf{3}}, -\frac{1}{3})_{\mathbf{F}}$, with $(\mathbf{1}, \mathbf{3}, -\frac{2}{3})_{\mathbf{F}} \equiv (\mathbf{n}'_R, -\mathbf{e}'_R, \tau_R)^T$ and $(\mathbf{1}, \overline{\mathbf{3}}, -\frac{1}{3})_{\mathbf{F}} \equiv (\mathbf{e}_R, \mathbf{n}_R, \check{\mathbf{n}}_R)^T$. Then, we denote the rank-2 anti-symmetric fermion of $(\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{F}}$ as follows

$$\psi_{(\mathbf{1}, \mathbf{6}, -\frac{1}{2})} \equiv \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & \check{\mathbf{n}}_R & -\mathbf{n}_R & \mathbf{n}'_R \\ -\check{\mathbf{n}}_R & 0 & \mathbf{e}_R & -\mathbf{e}'_R \\ \mathbf{n}_R & -\mathbf{e}_R & 0 & \tau_R \\ -\mathbf{n}'_R & \mathbf{e}'_R & -\tau_R & 0 \end{pmatrix}, \tag{54}$$

and its covariant derivative should be obtained according to Eq. (39). The covariant derivatives lead to the neutral currents of

$$i\bar{\psi}_{(\mathbf{3},\mathbf{1},-\frac{1}{3})}\not{D}\psi_{(\mathbf{3},\mathbf{1},-\frac{1}{3})} \supset \frac{gX_2}{s_{\vartheta_G}c_{\vartheta_G}}\left(+\frac{1}{3}s_{\vartheta_G}^2\right)\overline{\mathcal{D}}_L\gamma^\mu\mathcal{D}_L\cdot Z''_\mu, \quad (55a)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{1},+\frac{2}{3})}\not{D}\psi_{(\mathbf{3},\mathbf{1},+\frac{2}{3})} \supset \frac{gX_2}{s_{\vartheta_G}c_{\vartheta_G}}\left(-\frac{2}{3}s_{\vartheta_G}^2\right)\overline{t}_R\gamma^\mu t_R\cdot Z''_\mu, \quad (55b)$$

$$\begin{aligned} \text{Tr}\left[i\bar{\psi}_{(\mathbf{1},\mathbf{6},-\frac{1}{2})}\not{D}\psi_{(\mathbf{1},\mathbf{6},-\frac{1}{2})}\right] &\supset \frac{gX_2}{s_{\vartheta_G}c_{\vartheta_G}}\left[\left(\frac{1}{6}+\frac{1}{3}s_{\vartheta_G}^2\right)(\overline{\mathbf{n}}_R\gamma^\mu\check{\mathbf{n}}_R+\overline{\mathbf{n}}_R\gamma^\mu\mathbf{n}_R+\overline{\mathbf{e}}_R\gamma^\mu\mathbf{e}_R)\right. \\ &\left.+\left(-\frac{1}{6}+\frac{2}{3}s_{\vartheta_G}^2\right)(\overline{\mathbf{n}}'_R\gamma^\mu\mathbf{n}'_R+\overline{\mathbf{e}}'_R\gamma^\mu\mathbf{e}'_R+\overline{\tau}_R\gamma^\mu\tau_R)\right]\cdot Z''_\mu, \end{aligned} \quad (55c)$$

$$\begin{aligned} i\bar{\psi}_{(\mathbf{3},\mathbf{4},-\frac{1}{12})}\not{D}\psi_{(\mathbf{3},\mathbf{4},-\frac{1}{12})} &\supset \frac{gX_2}{s_{\vartheta_G}c_{\vartheta_G}}\left[\left(+\frac{1}{12}\right)(\overline{t}_L\gamma^\mu t_L+\overline{b}_L\gamma^\mu b_L+\overline{\mathcal{D}}'_L\gamma^\mu\mathcal{D}'_L)\right. \\ &\left.+\left(-\frac{1}{4}+\frac{1}{3}s_{\vartheta_G}^2\right)\overline{\mathcal{D}}''_L\gamma^\mu\mathcal{D}''_L\right]\cdot Z''_\mu, \end{aligned} \quad (55d)$$

$$\begin{aligned} &i\bar{\psi}_{(\mathbf{1},\overline{\mathbf{4}},-\frac{1}{4})}\not{D}\psi_{(\mathbf{1},\overline{\mathbf{4}},-\frac{1}{4})} \\ &\supset \frac{gX_2}{s_{\vartheta_G}c_{\vartheta_G}}\left[\left(-\frac{1}{12}+\frac{1}{3}s_{\vartheta_G}^2\right)(\overline{\mathbf{e}}''_R\gamma^\mu\mathbf{e}''_R+\overline{\mathbf{n}}''_R\gamma^\mu\mathbf{n}''_R+\overline{\check{\mathbf{n}}}'_R\gamma^\mu\check{\mathbf{n}}'_R)+\left(+\frac{1}{4}\right)\overline{\check{\mathbf{n}}}_R\gamma^\mu\check{\mathbf{n}}_R\right]\cdot Z''_\mu. \end{aligned} \quad (55e)$$

For the rank-3 anti-symmetric fermions of $\mathbf{56}_F$, we denote the rank-2 anti-symmetric fermion of $(\mathbf{3},\mathbf{6},+\frac{1}{6})_F$ and $(\mathbf{1},\mathbf{6},-\frac{1}{2})'_F$ as follows

$$\psi_{(\mathbf{3},\mathbf{6},+\frac{1}{6})} \equiv \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & \mathfrak{u}_L & \mathbf{u}_L & c_L \\ -\mathfrak{u}_L & 0 & \mathfrak{d}_L & s_L \\ -\mathbf{u}_L & -\mathfrak{d}_L & 0 & \mathfrak{D}'''_L \\ -c_L & -s_L & -\mathfrak{D}'''_L & 0 \end{pmatrix}, \quad (56a)$$

$$\psi_{(\mathbf{1},\mathbf{6},-\frac{1}{2})}' \equiv \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & \check{\mathbf{n}}'''_R & -\mathbf{n}''''_R & \mathbf{n}''''_R \\ -\check{\mathbf{n}}'''_R & 0 & \mathbf{e}''''_R & -\mathbf{e}''''_R \\ \mathbf{n}''''_R & -\mathbf{e}''''_R & 0 & e_R \\ -\mathbf{n}''''_R & \mathbf{e}''''_R & -e_R & 0 \end{pmatrix}. \quad (56b)$$

Their covariant derivatives should be obtained according to Eq. (39). The covariant derivatives from the $\mathbf{56}_F$ lead to the neutral currents of

$$i\bar{\psi}_{(1,4,-\frac{3}{4})}\not{D}\psi_{(1,4,-\frac{3}{4})} \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left[\left(-\frac{1}{4} + s_{\partial_G}^2\right) \overline{\mathbf{e}}_R \gamma^\mu \mathbf{e}_R \right. \\ \left. + \left(\frac{1}{12} + \frac{2}{3}s_{\partial_G}^2\right) (\overline{\mathbf{n}}_R''' \gamma^\mu \mathbf{n}_R''' + \overline{\mathbf{e}}_R''' \gamma^\mu \mathbf{e}_R''' + \overline{\mu}_R \gamma^\mu \mu_R) \right] \cdot Z_\mu'', \quad (57a)$$

$$i\bar{\psi}_{(3,1,+\frac{2}{3})'}\not{D}\psi_{(3,1,+\frac{2}{3})'} \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left(-\frac{2}{3}s_{\partial_G}^2\right) \overline{u}_R \gamma^\mu u_R \cdot Z_\mu'', \quad (57b)$$

$$i\bar{\psi}_{(1,1,-1)}\not{D}\psi_{(1,1,-1)} \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left(+s_{\partial_G}^2\right) \overline{\mathbf{e}}_L \gamma^\mu \mathbf{e}_L \cdot Z_\mu'', \quad (57c)$$

$$\text{Tr} \left[i\bar{\psi}_{(3,6,+\frac{1}{6})}\not{D}\psi_{(3,6,+\frac{1}{6})} \right] \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left[\left(\frac{1}{6} - \frac{1}{3}s_{\partial_G}^2\right) (\overline{\mathbf{u}}_L \gamma^\mu \mathbf{u}_L + \overline{\mathbf{u}}_L \gamma^\mu \mathbf{u}_L + \overline{\mathbf{d}}_L \gamma^\mu \mathbf{d}_L) \right. \\ \left. + \left(-\frac{1}{6}\right) (\overline{c}_L \gamma^\mu c_L + \overline{s}_L \gamma^\mu s_L + \overline{\mathfrak{D}}_L''' \gamma^\mu \mathfrak{D}_L''') \right] \cdot Z_\mu'', \quad (57d)$$

$$\text{Tr} \left[i\bar{\psi}_{(1,6,-\frac{1}{2})'}\not{D}\psi_{(1,6,-\frac{1}{2})'} \right] \\ \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left[\left(\frac{1}{6} + \frac{1}{3}s_{\partial_G}^2\right) (\overline{\mathbf{n}}_R'''' \gamma^\mu \mathbf{n}_R'''' + \overline{\mathbf{n}}_R'''' \gamma^\mu \mathbf{n}_R'''' + \overline{\mathbf{e}}_R'''' \gamma^\mu \mathbf{e}_R'''') \right. \\ \left. + \left(-\frac{1}{6} + \frac{2}{3}s_{\partial_G}^2\right) (\overline{\mathbf{n}}_R'''' \gamma^\mu \mathbf{n}_R'''' + \overline{\mathbf{e}}_R'''' \gamma^\mu \mathbf{e}_R'''' + \overline{e}_R \gamma^\mu e_R) \right] \cdot Z_\mu'', \quad (57e)$$

$$i\bar{\psi}_{(3,4,-\frac{1}{12})'}\not{D}\psi_{(3,4,-\frac{1}{12})'} \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left[\left(+\frac{1}{12}\right) (\overline{u}_L \gamma^\mu u_L + \overline{d}_L \gamma^\mu d_L + \overline{\mathfrak{D}}_L'''' \gamma^\mu \mathfrak{D}_L'''') \right. \\ \left. + \left(-\frac{1}{4} + \frac{1}{3}s_{\partial_G}^2\right) \overline{\mathfrak{D}}_L'''' \gamma^\mu \mathfrak{D}_L'''' \right] \cdot Z_\mu'', \quad (57f)$$

$$i\bar{\psi}_{(3,4,+\frac{5}{12})}\not{D}\psi_{(3,4,+\frac{5}{12})} \supset \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} \left[\left(-\frac{1}{12} - \frac{1}{3}s_{\partial_G}^2\right) (\overline{\mathbf{d}}_R \gamma^\mu \mathbf{d}_R + \overline{\mathbf{u}}_R \gamma^\mu \mathbf{u}_R + \overline{\mathbf{u}}_R \gamma^\mu \mathbf{u}_R) \right. \\ \left. + \left(\frac{1}{4} - \frac{2}{3}s_{\partial_G}^2\right) \overline{c}_R \gamma^\mu c_R \right] \cdot Z_\mu''. \quad (57g)$$

The neutral currents are expressed as follows in the $V - A$ basis

$$\mathcal{L}_{\text{SU}(4)_W}^{\text{NC,F}} = \frac{g_{X_2}}{s_{\partial_G}c_{\partial_G}} (g_f^V \bar{f} \gamma^\mu f + g_f^A \bar{f} \gamma^\mu \gamma_5 f) Z_\mu'', \quad (58)$$

and we tabulate the vectorial and axial couplings of (g_f^V, g_f^A) in Tab. 8. The flavor non-universality of the neutral currents mediated by Z_μ'' among three generational SM quarks and leptons are displayed explicitly. The heavy partner fermions of $(\mathfrak{D}, \mathbf{e}'', \mathbf{n}'', \mathbf{n}'', \mathbf{n}'')$ only exhibit vectorial couplings, since they have already obtained their masses at the first stage of symmetry breaking.

3.3 The $\text{SU}(3)_W \otimes \text{U}(1)_{X_2}$ gauge bosons

3.3.1 Covariant derivatives and gauge boson masses

We express the $\text{SU}(3)_W \otimes \text{U}(1)_{X_2}$ covariant derivatives for the $\text{SU}(3)_W$ fundamental and anti-fundamental representations as follows

$$iD_\mu \psi_{\mathbf{3}} \equiv i\partial_\mu \psi_{\mathbf{3}} + (g_{3W} A_\mu^{\bar{I}} T_{\text{SU}(3)}^{\bar{I}} + g_{X_2} \mathcal{X}_2 \mathbb{I}_3 X_{2\mu}) \cdot \psi_{\mathbf{3}}, \quad (59a)$$

$$iD_\mu \psi_{\bar{\mathbf{3}}} \equiv i\partial_\mu \psi_{\bar{\mathbf{3}}} + (-g_{3W} A_\mu^{\bar{I}} (T_{\text{SU}(3)}^{\bar{I}})^T + g_{X_2} \mathcal{X}_2 \mathbb{I}_3 X_{2\mu}) \cdot \psi_{\bar{\mathbf{3}}}. \quad (59b)$$

	u	c	t	u	\mathfrak{U}
$g_f^{V''}$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$
$g_f^{A''}$	$-\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$\frac{5}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{8}$	$-\frac{1}{8}$
	d	s	b	\mathfrak{D}	
$g_f^{V''}$	$\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{3}s_{\vartheta_G}^2$	
$g_f^{A''}$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{8}$	
	\mathfrak{D}	\mathfrak{D}''	\mathfrak{D}'	\mathfrak{D}''''	\mathfrak{D}'''
$g_f^{V''}$	$\frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{8} + \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{8} + \frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$
$g_f^{A''}$	0	$\frac{1}{8}$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{8}$	$\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$
	e	μ	τ	\mathfrak{E}	
$g_f^{V''}$	$-\frac{1}{8} + \frac{1}{2}s_{\vartheta_G}^2$	$\frac{1}{2}s_{\vartheta_G}^2$	$-\frac{1}{8} + \frac{1}{2}s_{\vartheta_G}^2$	$-\frac{1}{8} + s_{\vartheta_G}^2$	
$g_f^{A''}$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{8}$	
	$(\mathfrak{e}, \mathfrak{n})$	$(\mathfrak{e}', \mathfrak{n}')$	$(\mathfrak{e}'', \mathfrak{n}'')$	$(\mathfrak{e}''', \mathfrak{n}''')$	$(\mathfrak{e}'''' , \mathfrak{n}'''')$
$g_f^{V''}$	$\frac{1}{24} + \frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{8} + \frac{1}{2}s_{\vartheta_G}^2$	$-\frac{1}{12} + \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{2}s_{\vartheta_G}^2$	$\frac{1}{24} + \frac{1}{3}s_{\vartheta_G}^2$
$g_f^{A''}$	$\frac{1}{8}$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	0	$\frac{1}{12} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{8}$
	$\nu_{eL}/\nu_{\mu L}/\nu_{\tau L}$	$\tilde{\mathcal{N}}_L^{\Omega, \Omega', \Omega''}$	$\tilde{\mathfrak{n}}$	$\tilde{\mathfrak{n}}'$	$\tilde{\mathfrak{n}}''$
$g_f^{V''}$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$-\frac{1}{24} + \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{24} + \frac{1}{3}s_{\vartheta_G}^2$	$-\frac{1}{12} + \frac{1}{3}s_{\vartheta_G}^2$	$\frac{1}{4}$
$g_f^{A''}$	$\frac{1}{24} - \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{24} - \frac{1}{6}s_{\vartheta_G}^2$	$\frac{1}{8}$	0	0
					$\frac{1}{8}$

Table 8: The couplings of the flavor-conserving neutral currents mediated by Z'' in the $V - A$ basis.

The $SU(3)_W$ generators are normalized such that $\text{Tr}\left(T_{SU(3)}^{\tilde{I}} T_{SU(3)}^{\tilde{J}}\right) = \frac{1}{2}\delta^{\tilde{I}\tilde{J}}$. The gauge fields from Eq. (59a) can be expressed in terms of a 3×3 matrix

$$\begin{aligned}
& g_{3W} A_{\mu}^{\tilde{I}} T_{SU(3)}^{\tilde{I}} + g_{X_2} \mathcal{X}_2 \mathbb{I}_3 X_{2\mu} \\
&= \frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & W_{\mu}^{+} & W_{\mu}^{\prime+} \\ W_{\mu}^{-} & 0 & 0 \\ W_{\mu}^{\prime-} & 0 & 0 \end{pmatrix} + \frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & N_{\mu} \\ 0 & \bar{N}_{\mu} & 0 \end{pmatrix} \\
&+ \frac{g_{3W}}{2} \text{diag}\left(A_{\mu}^3 + \frac{1}{\sqrt{3}}A_{\mu}^8 + 2\sqrt{3}t_{\vartheta_S} \mathcal{X}_2 X_{2\mu}, \right. \\
&\quad \left. -A_{\mu}^3 + \frac{1}{\sqrt{3}}A_{\mu}^8 + 2\sqrt{3}t_{\vartheta_S} \mathcal{X}_2 X_{2\mu}, -\frac{2}{\sqrt{3}}A_{\mu}^8 + 2\sqrt{3}t_{\vartheta_S} \mathcal{X}_2 X_{2\mu}\right), \tag{60}
\end{aligned}$$

with an $SU(3)_W$ mixing angle defined by

$$t_{\vartheta_S} \equiv \tan \vartheta_S = \frac{g_{X_2}}{\sqrt{3}g_{3W}}. \tag{61}$$

One can identify that

$$\begin{aligned}
W_{\mu}^{\pm} &= \frac{1}{\sqrt{2}}(A_{\mu}^1 \mp iA_{\mu}^2), & W_{\mu}^{\prime\pm} &= \frac{1}{\sqrt{2}}(A_{\mu}^4 \mp iA_{\mu}^5), \\
N_{\mu} &= \frac{1}{\sqrt{2}}(A_{\mu}^6 - iA_{\mu}^7), & \bar{N}_{\mu} &= \frac{1}{\sqrt{2}}(A_{\mu}^6 + iA_{\mu}^7). \tag{62}
\end{aligned}$$

This stage of symmetry breaking is achieved by Higgs VEVs in Eq. (18c), and the corresponding gauge boson masses from the Higgs kinematic terms are

$$\begin{aligned} & \frac{1}{4}g_{3W}^2 v_{331}^2 \left[\left(W_\mu'^+ W'^{-\mu} + N_\mu \bar{N}^\mu \right) + \frac{2}{3} \left(A_\mu^8 - t_{\vartheta_S} X_{2\mu} \right)^2 \right], \\ v_{331}^2 &= (V_{\mathbf{3},3})^2 + (V_{\mathbf{3},\text{VI}})^2 + (V_{\mathbf{3},\text{IX}})^2 + (V_{\mathbf{3},\dot{2}})^2 + (V_{\mathbf{3},\text{VIII}}')^2. \end{aligned} \quad (63)$$

It is straightforward to obtain the mass eigenstates in terms of the mixing angle in Eq. (61) for this case

$$\begin{pmatrix} Z'_\mu \\ B_\mu \end{pmatrix} = \begin{pmatrix} c_{\vartheta_S} & -s_{\vartheta_S} \\ s_{\vartheta_S} & c_{\vartheta_S} \end{pmatrix} \cdot \begin{pmatrix} A_\mu^8 \\ X_{2\mu} \end{pmatrix}. \quad (64)$$

The $\text{SU}(3)_W \otimes \text{U}(1)_{X_2}$ gauge couplings of $(\alpha_{3W}, \alpha_{X_1})$ match with the EW gauge couplings as follows

$$\begin{aligned} \alpha_{2W}^{-1}(v_{331}) &= \alpha_{3W}^{-1}(v_{331}), \quad \alpha_Y^{-1}(v_{331}) = \frac{1}{3}\alpha_{3W}^{-1}(v_{331}) + \alpha_{X_2}^{-1}(v_{331}), \\ \frac{1}{3}\alpha_{3W}^{-1} &= \alpha_Y^{-1} s_{\vartheta_S}^2, \quad \alpha_{X_2}^{-1} = \alpha_Y^{-1} c_{\vartheta_S}^2. \end{aligned} \quad (65)$$

From the definitions of two mixing angles in Eqs. (61) and (76), we find a relation of

$$\boxed{\sin \vartheta_S = \frac{1}{\sqrt{3}} \tan \vartheta_W}, \quad (66)$$

from the gauge coupling matching in Eq. (65). The tree-level masses for five gauge bosons at this stage read

$$m_{W_\mu'^\pm}^2 = m_{N_\mu, \bar{N}_\mu}^2 = \frac{g_Y^2}{12s_{\vartheta_S}^2} v_{331}^2, \quad m_{Z'_\mu}^2 = \frac{g_{X_1}^2}{9s_{\vartheta_S}^2 c_{\vartheta_S}^2} v_{331}^2. \quad (67)$$

In terms of mass eigenstates, the gauge bosons from the covariant derivative in Eq. (60) become

$$\begin{aligned} & \frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ & 0 \\ W_\mu^- & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & W_\mu'^+ \\ 0 & 0 & N_\mu \\ W_\mu'^- & \bar{N}_\mu & 0 \end{pmatrix} \\ & + g_{3W} \text{diag} \left(\frac{1}{2}, -\frac{1}{2}, 0 \right) A_\mu^3 + g_Y \text{diag} \left(\left(\frac{1}{6} + \mathcal{X}_2 \right) \mathbb{I}_{2 \times 2}, -\frac{1}{3} + \mathcal{X}_2 \right) B_\mu \\ & + \frac{g_Y}{s_{\vartheta_S} c_{\vartheta_S}} \text{diag} \left(\left[\frac{1}{6} - \left(\frac{1}{6} + \mathcal{X}_2 \right) s_{\vartheta_S}^2 \right] \mathbb{I}_{2 \times 2}, -\frac{1}{3} + \left(\frac{1}{3} - \mathcal{X}_2 \right) s_{\vartheta_S}^2 \right) Z'_\mu. \end{aligned} \quad (68)$$

Likewise, the covariant derivative for the $\text{SU}(3)_W$ anti-fundamental representation gives

$$\begin{aligned} & -\frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^- & 0 \\ W_\mu^+ & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \frac{g_{3W}}{\sqrt{2}} \begin{pmatrix} 0 & 0 & W_\mu'^- \\ 0 & 0 & \bar{N}_\mu \\ W_\mu'^+ & N_\mu & 0 \end{pmatrix} \\ & - g_{3W} \text{diag} \left(\frac{1}{2}, -\frac{1}{2}, 0 \right) A_\mu^3 + g_Y \text{diag} \left(\left(-\frac{1}{6} + \mathcal{X}_1 \right) \mathbb{I}_{2 \times 2}, \frac{1}{3} + \mathcal{X}_1 \right) B_\mu \\ & + \frac{g_Y}{s_{\vartheta_S} c_{\vartheta_S}} \text{diag} \left(\left[-\frac{1}{6} + \left(\frac{1}{6} - \mathcal{X}_1 \right) s_{\vartheta_S}^2 \right] \mathbb{I}_{2 \times 2}, \frac{1}{3} - \left(\frac{1}{3} + \mathcal{X}_1 \right) s_{\vartheta_S}^2 \right) Z'_\mu. \end{aligned} \quad (69)$$

3.3.2 Gauge couplings of the \mathcal{G}_{331} fermions

From the anti-fundamental fermions of $\overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}$, we find the neutral currents of

$$i\overline{\psi}_{(\mathbf{3},\mathbf{1},-\frac{1}{3})^{\Omega}}\not{D}\psi_{(\mathbf{3},\mathbf{1},-\frac{1}{3})^{\Omega}} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left(+\frac{1}{3}s_{\vartheta_S}^2\right)\overline{\mathcal{D}}_R^{\Omega}\gamma^{\mu}\mathcal{D}_R^{\Omega}\cdot Z'_{\mu}, \quad (70a)$$

$$\begin{aligned} & i\overline{\psi}_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})^{\Omega}}\not{D}\psi_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})^{\Omega}} \\ \supset & \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left[\left(-\frac{1}{6}+\frac{1}{2}s_{\vartheta_S}^2\right)(\overline{\mathcal{E}}_L^{\Omega}\gamma^{\mu}\mathcal{E}_L^{\Omega}+\overline{\mathcal{N}}_L^{\Omega}\gamma^{\mu}\mathcal{N}_L^{\Omega})+\left(+\frac{1}{3}\right)\overline{\mathcal{N}}_L^{\Omega'}\gamma^{\mu}\mathcal{N}_L^{\Omega'}\right]\cdot Z'_{\mu}. \end{aligned} \quad (70b)$$

From the rank-2 anti-symmetric fermion of $\mathbf{28}_{\mathbf{F}}$, we find the neutral currents of

$$i\overline{\psi}_{(\mathbf{3},\mathbf{1},-\frac{1}{3})}\not{D}\psi_{(\mathbf{3},\mathbf{1},-\frac{1}{3})} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left(+\frac{1}{3}s_{\vartheta_S}^2\right)\overline{\mathcal{D}}_L\gamma^{\mu}\mathcal{D}_L\cdot Z'_{\mu}, \quad (71a)$$

$$i\overline{\psi}_{(\mathbf{3},\mathbf{1},+\frac{2}{3})}\not{D}\psi_{(\mathbf{3},\mathbf{1},+\frac{2}{3})} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left(-\frac{2}{3}s_{\vartheta_S}^2\right)\overline{t}_R\gamma^{\mu}t_R\cdot Z'_{\mu}, \quad (71b)$$

$$\begin{aligned} & i\overline{\psi}_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})}\not{D}\psi_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})} \\ \supset & \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left[\left(-\frac{1}{6}+\frac{1}{2}s_{\vartheta_S}^2\right)(\overline{\mathbf{e}}_R\gamma^{\mu}\mathbf{e}_R+\overline{\mathbf{n}}_R\gamma^{\mu}\mathbf{n}_R)+\left(+\frac{1}{3}\right)\overline{\mathbf{n}}_R\gamma^{\mu}\mathbf{n}'_R\right]\cdot Z'_{\mu}, \end{aligned} \quad (71c)$$

$$\begin{aligned} & i\overline{\psi}_{(\mathbf{1},\mathbf{3},-\frac{2}{3})}\not{D}\psi_{(\mathbf{1},\mathbf{3},-\frac{2}{3})} \\ \supset & \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left[\left(\frac{1}{6}+\frac{1}{2}s_{\vartheta_S}^2\right)(\overline{\mathbf{e}}'_R\gamma^{\mu}\mathbf{e}'_R+\overline{\mathbf{n}}'_R\gamma^{\mu}\mathbf{n}'_R)+\left(-\frac{1}{3}+s_{\vartheta_S}^2\right)\overline{\tau}_R\gamma^{\mu}\tau_R\right]\cdot Z'_{\mu}, \end{aligned} \quad (71d)$$

$$\begin{aligned} & i\overline{\psi}_{(\mathbf{3},\mathbf{3},0)}\not{D}\psi_{(\mathbf{3},\mathbf{3},0)} \\ \supset & \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left[\frac{1}{6}(1-s_{\vartheta_S}^2)(\overline{t}_L\gamma^{\mu}t_L+\overline{b}_L\gamma^{\mu}b_L)+\frac{1}{3}(-1+s_{\vartheta_S}^2)\overline{\mathcal{D}}_L'\gamma^{\mu}\mathcal{D}_L'\right]\cdot Z'_{\mu}, \end{aligned} \quad (71e)$$

$$i\overline{\psi}_{(\mathbf{3},\mathbf{1},-\frac{1}{3})''}\not{D}\psi_{(\mathbf{3},\mathbf{1},-\frac{1}{3})''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left(+\frac{1}{3}s_{\vartheta_S}^2\right)\overline{\mathcal{D}}_L''\gamma^{\mu}\mathcal{D}_L''\cdot Z'_{\mu}, \quad (71f)$$

$$\begin{aligned} & i\overline{\psi}_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})''}\not{D}\psi_{(\mathbf{1},\overline{\mathbf{3}},-\frac{1}{3})''} \\ \supset & \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}}\left[\left(-\frac{1}{6}+\frac{1}{2}s_{\vartheta_S}^2\right)(\overline{\mathbf{e}}_R''\gamma^{\mu}\mathbf{e}_R''+\overline{\mathbf{n}}_R''\gamma^{\mu}\mathbf{n}_R'')+ \left(+\frac{1}{3}\right)\overline{\mathbf{n}}_R''\gamma^{\mu}\mathbf{n}'_R''\right]\cdot Z'_{\mu}. \end{aligned} \quad (71g)$$

From the rank-3 anti-symmetric fermion of $\mathbf{56_F}$, we find the neutral currents of

$$i\bar{\psi}_{(\mathbf{1},\mathbf{3},-\frac{2}{3})'}\mathcal{D}\psi_{(\mathbf{1},\mathbf{3},-\frac{2}{3})'} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[\left(\frac{1}{6} + \frac{1}{2}s_{\vartheta_S}^2 \right) (\bar{\mathbf{e}}_R^{\prime\prime\prime}\gamma^\mu\mathbf{e}_R^{\prime\prime\prime} + \bar{\mathbf{n}}_R^{\prime\prime\prime}\gamma^\mu\mathbf{n}_R^{\prime\prime\prime}) + \left(-\frac{1}{3} + s_{\vartheta_S}^2 \right) \bar{\mu}_R\gamma^\mu\mu_R \right] \cdot Z'_\mu, \quad (72a)$$

$$i\bar{\psi}_{(\mathbf{1},\mathbf{1},-1)''}\mathcal{D}\psi_{(\mathbf{1},\mathbf{1},-1)''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} (+s_{\vartheta_S}^2) \bar{\mathbf{e}}_R\gamma^\mu\mathbf{e}_R \cdot Z'_\mu, \quad (72b)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{1},+\frac{2}{3})'}\mathcal{D}\psi_{(\mathbf{3},\mathbf{1},+\frac{2}{3})'} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left(-\frac{2}{3}s_{\vartheta_S}^2 \right) \bar{u}_R\gamma^\mu u_R \cdot Z'_\mu, \quad (72c)$$

$$i\bar{\psi}_{(\mathbf{1},\mathbf{1},-1)}\mathcal{D}\psi_{(\mathbf{1},\mathbf{1},-1)} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} (+s_{\vartheta_S}^2) \bar{\mathbf{e}}_L\gamma^\mu\mathbf{e}_L \cdot Z'_\mu, \quad (72d)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{3},0)'}\mathcal{D}\psi_{(\mathbf{3},\mathbf{3},0)'} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[\frac{1}{6}(1-s_{\vartheta_S}^2)(\bar{c}_L\gamma^\mu c_L + \bar{s}_L\gamma^\mu s_L) + \frac{1}{3}(-1+s_{\vartheta_S}^2)\bar{\mathfrak{D}}_L^{\prime\prime\prime}\gamma^\mu\mathfrak{D}_L^{\prime\prime\prime} \right] \cdot Z'_\mu, \quad (72e)$$

$$i\bar{\psi}_{(\mathbf{3},\bar{\mathbf{3}},+\frac{1}{3})}\mathcal{D}\psi_{(\mathbf{3},\bar{\mathbf{3}},+\frac{1}{3})} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[-\frac{1}{6}(1+s_{\vartheta_S}^2)(\bar{u}_L\gamma^\mu u_L + \bar{d}_L\gamma^\mu d_L) + \left(\frac{1}{3} - \frac{2}{3}s_{\vartheta_S}^2 \right) \bar{\mathfrak{U}}_L\gamma^\mu\mathfrak{U}_L \right] \cdot Z'_\mu, \quad (72f)$$

$$i\bar{\psi}_{(\mathbf{1},\bar{\mathbf{3}},-\frac{1}{3})'}\mathcal{D}\psi_{(\mathbf{1},\bar{\mathbf{3}},-\frac{1}{3})'} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[\left(-\frac{1}{6} + \frac{1}{2}s_{\vartheta_S}^2 \right) (\bar{\mathbf{e}}_R^{\prime\prime\prime\prime}\gamma^\mu\mathbf{e}_R^{\prime\prime\prime\prime} + \bar{\mathbf{n}}_R^{\prime\prime\prime\prime}\gamma^\mu\mathbf{n}_R^{\prime\prime\prime\prime}) + \left(+\frac{1}{3} \right) \bar{\mathfrak{n}}_R^{\prime\prime\prime}\gamma^\mu\mathfrak{n}_R^{\prime\prime\prime} \right] \cdot Z'_\mu, \quad (72g)$$

$$i\bar{\psi}_{(\mathbf{1},\mathbf{3},-\frac{2}{3})''}\mathcal{D}\psi_{(\mathbf{1},\mathbf{3},-\frac{2}{3})''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[\left(\frac{1}{6} + \frac{1}{2}s_{\vartheta_S}^2 \right) (\bar{\mathbf{e}}_R^{\prime\prime\prime\prime\prime}\gamma^\mu\mathbf{e}_R^{\prime\prime\prime\prime\prime} + \bar{\mathbf{n}}_R^{\prime\prime\prime\prime\prime}\gamma^\mu\mathbf{n}_R^{\prime\prime\prime\prime\prime}) + \left(-\frac{1}{3} + s_{\vartheta_S}^2 \right) \bar{e}_R\gamma^\mu e_R \right] \cdot Z'_\mu, \quad (72h)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{3},0)''}\mathcal{D}\psi_{(\mathbf{3},\mathbf{3},0)''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[\frac{1}{6}(1-s_{\vartheta_S}^2)(\bar{u}_L\gamma^\mu u_L + \bar{d}_L\gamma^\mu d_L) + \frac{1}{3}(-1+s_{\vartheta_S}^2)\bar{\mathfrak{D}}_L^{\prime\prime\prime\prime}\gamma^\mu\mathfrak{D}_L^{\prime\prime\prime\prime} \right] \cdot Z'_\mu, \quad (72i)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{1},-\frac{1}{3})''''}\mathcal{D}\psi_{(\mathbf{3},\mathbf{1},-\frac{1}{3})''''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left(+\frac{1}{3}s_{\vartheta_S}^2 \right) \bar{\mathfrak{D}}_L^{\prime\prime\prime\prime\prime}\gamma^\mu\mathfrak{D}_L^{\prime\prime\prime\prime\prime} \cdot Z'_\mu, \quad (72j)$$

$$i\bar{\psi}_{(\mathbf{3},\bar{\mathbf{3}},+\frac{1}{3})}\mathcal{D}\psi_{(\mathbf{3},\bar{\mathbf{3}},+\frac{1}{3})} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left[-\frac{1}{6}(1+s_{\vartheta_S}^2)(\bar{u}_R\gamma^\mu u_R + \bar{d}_R\gamma^\mu d_R) + \left(\frac{1}{3} - \frac{2}{3}s_{\vartheta_S}^2 \right) \bar{\mathfrak{U}}_R\gamma^\mu\mathfrak{U}_R \right] \cdot Z'_\mu, \quad (72k)$$

$$i\bar{\psi}_{(\mathbf{3},\mathbf{1},+\frac{2}{3})''''}\mathcal{D}\psi_{(\mathbf{3},\mathbf{1},+\frac{2}{3})''''} \supset \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} \left(-\frac{2}{3}s_{\vartheta_S}^2 \right) \bar{c}_R\gamma^\mu c_R \cdot Z'_\mu. \quad (72l)$$

The neutral currents are expressed as follows

$$\mathcal{L}_{\text{SU}(3)_W}^{\text{NC,F}} = \frac{g_Y}{s_{\vartheta_S}c_{\vartheta_S}} (g_f^V f' \bar{f} \gamma^\mu f + g_f^A f' \bar{f} \gamma^\mu \gamma_5 f) Z'_\mu, \quad (73)$$

in the $V - A$ basis, and we tabulate the vectorial and axial couplings in Tab. 9. Three generational SM fermions with the same electric charges display the universal vectorial and axial couplings to the Z'_μ

	$u/c/t$	\mathbf{u}	\mathfrak{U}	
$g_f^{V'}$	$\frac{1}{12} - \frac{5}{12}s_{\vartheta_S}^2$	$-\frac{1}{6} - \frac{1}{6}s_{\vartheta_S}^2$	$\frac{1}{3} - \frac{2}{3}s_{\vartheta_S}^2$	
$g_f^{A'}$	$-\frac{1}{12} - \frac{1}{4}s_{\vartheta_S}^2$	0	0	
	$d/s/b$	\mathfrak{D}	$\mathfrak{D}/\mathfrak{D}''/\mathfrak{D}''''$	$\mathfrak{D}'/\mathfrak{D}'''/\mathfrak{D}''''$
$g_f^{V'}$	$\frac{1}{12} + \frac{1}{12}s_{\vartheta_S}^2$	$-\frac{1}{6} - \frac{1}{6}s_{\vartheta_S}^2$	$\frac{1}{3}s_{\vartheta_S}^2$	$-\frac{1}{6} + \frac{1}{3}s_{\vartheta_S}^2$
$g_f^{A'}$	$-\frac{1}{12} + \frac{1}{4}s_{\vartheta_S}^2$	0	0	$\frac{1}{6}$
	$e/\mu/\tau$	\mathfrak{E}	$(\mathbf{e}, \mathbf{n})/(\mathbf{e}'', \mathbf{n}'')$	$(\mathbf{e}', \mathbf{n}')/(\mathbf{e}''', \mathbf{n}''')$
			$/(\mathbf{e}'''' , \mathbf{n}'''')$	$/(\mathbf{e}'''' , \mathbf{n}'''')$
$g_f^{V'}$	$-\frac{1}{4} + \frac{3}{4}s_{\vartheta_S}^2$	$s_{\vartheta_S}^2$	$-\frac{1}{6} + \frac{1}{2}s_{\vartheta_S}^2$	$\frac{1}{2}s_{\vartheta_S}^2$
$g_f^{A'}$	$-\frac{1}{12} + \frac{1}{4}s_{\vartheta_S}^2$	0	0	$\frac{1}{6}$
	$\nu_e/\nu_\mu/\nu_\tau$	$\check{\mathbf{n}}/\check{\mathbf{n}}'/\check{\mathbf{n}}''/\check{\mathbf{n}}'''$		
$g_f^{V'}$	$-\frac{1}{12} + \frac{1}{4}s_{\vartheta_S}^2$	$\frac{1}{3}$		
$g_f^{A'}$	$\frac{1}{12} - \frac{1}{4}s_{\vartheta_S}^2$	0		

Table 9: The couplings of the neutral currents mediated by Z'_μ in the $V - A$ basis.

gauge boson. Both of the SM fermion vectorial and axial couplings depend on the mixing angle of ϑ_S . Besides of vectorlike fermions of $(\mathfrak{D}, \mathbf{e}'', \mathbf{n}'', \check{\mathbf{n}}', \check{\mathbf{n}}'')$, the $(\mathbf{u}, \mathfrak{D}, \mathfrak{U}, \mathfrak{D}'', \mathfrak{D}'''' , \mathfrak{E}, \mathbf{e}, \mathbf{n}, \mathbf{e}'''' , \mathbf{n}'''')$ also exhibit vectorial couplings to the Z'_μ gauge boson, since they have obtained their masses at the second stage of symmetry breaking. For vectorlike fermions of $(\mathfrak{D}', \mathfrak{D}''', \mathfrak{D}'''' , \mathbf{n}', \mathbf{e}', \mathbf{n}''', \mathbf{e}'''' , \mathbf{n}'''' , \mathbf{e}'''')$ that obtain masses at the third stage of symmetry breaking, they exhibit both the vectorial and axial couplings of $(g_f^{V'}, g_f^{A'})$. Different from the SM fermions, only their vectorial couplings depend on the mixing angle of ϑ_S , while their axial couplings are independent of the mixing angle of ϑ_S .

3.4 The $SU(2)_W \otimes U(1)_Y$ gauge bosons

3.4.1 Covariant derivatives and gauge boson masses

We express the $SU(2)_W \otimes U(1)_Y$ covariant derivatives in the EW sector for the fundamental and anti-fundamental representations as follows

$$iD_\mu \psi_{\mathbf{2}} \equiv i\partial_\mu \psi_{\mathbf{2}} + (g_{2W} A_\mu^I T_{SU(2)}^I + g_Y \mathcal{Y} \mathbb{I}_2 B_\mu) \cdot \psi_{\mathbf{2}}, \quad (74a)$$

$$iD_\mu \psi_{\bar{\mathbf{2}}} \equiv i\partial_\mu \psi_{\bar{\mathbf{2}}} + (-g_{2W} A_\mu^I (T_{SU(2)}^I)^T + g_Y \mathcal{Y} \mathbb{I}_2 B_\mu) \cdot \psi_{\bar{\mathbf{2}}}, \quad (74b)$$

with $T_{SU(2)}^I \equiv \frac{1}{2} \sigma^I$. The gauge fields from Eqs. (74) can be expressed in terms of a 2×2 matrix

$$\begin{aligned} & g_{2W} A_\mu^I T_{SU(2)}^I + g_Y \mathcal{Y} \mathbb{I}_2 B_\mu \\ = & \frac{g_{2W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ \\ W_\mu^- & 0 \end{pmatrix} + \frac{g_{2W}}{2} \text{diag} \left(A_\mu^3 + 2t_{\vartheta_W} \mathcal{Y} B_\mu, -A_\mu^3 + 2t_{\vartheta_W} \mathcal{Y} B_\mu \right), \end{aligned} \quad (75a)$$

$$\begin{aligned} & -g_{2W} A_\mu^I (T_{SU(2)}^I)^T + g_Y \mathcal{Y} \mathbb{I}_2 B_\mu \\ = & -\frac{g_{2W}}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^- \\ W_\mu^+ & 0 \end{pmatrix} + \frac{g_{2W}}{2} \text{diag} \left(-A_\mu^3 + 2t_{\vartheta_W} \mathcal{Y} B_\mu, A_\mu^3 + 2t_{\vartheta_W} \mathcal{Y} B_\mu \right), \end{aligned} \quad (75b)$$

with the $SU(2)_W$ Weinberg angle defined by

$$t_{\vartheta_W} \equiv \tan \vartheta_W = \frac{g_Y}{g_{2W}}. \quad (76)$$

One can identify that

$$W_\mu^\pm = \frac{1}{\sqrt{2}}(A_\mu^1 \mp iA_\mu^2). \quad (77)$$

3.4.2 Fermion \mathcal{G}_{SM} gauge couplings

	$u/c/t$	\mathbf{u}	\mathfrak{U}
g_f^V	$\frac{1}{4} - \frac{2}{3}s_{\vartheta_W}^2$	$\frac{1}{2} - \frac{2}{3}s_{\vartheta_W}^2$	$-\frac{2}{3}s_{\vartheta_W}^2$
g_f^A	$-\frac{1}{4}$	0	0
	$d/s/b$	\mathfrak{d}	$\mathfrak{D}/\mathfrak{D}'/\mathfrak{D}''/\mathfrak{D}'''/\mathfrak{D}''''/\mathfrak{D}'''''$
g_f^V	$-\frac{1}{4} + \frac{1}{3}s_{\vartheta_W}^2$	$-\frac{1}{2} + \frac{1}{3}s_{\vartheta_W}^2$	$+\frac{1}{3}s_{\vartheta_W}^2$
g_f^A	$+\frac{1}{4}$	0	0
	$e/\mu/\tau$	\mathfrak{E}	$\nu_e/\nu_\mu/\nu_\tau$
g_f^V	$-\frac{1}{4} + s_{\vartheta_W}^2$	$+s_{\vartheta_W}^2$	$+\frac{1}{4}$
g_f^A	$+\frac{1}{4}$	0	$-\frac{1}{4}$

Table 10: The couplings of the flavor-conserving neutral currents mediated by Z_μ in the $V - A$ basis.

The flavor-conserving neutral currents are expressed as follows in the $V - A$ basis

$$\mathcal{L}_{\text{SU}(2)_W}^{\text{NC,F}} = \frac{e}{s_{\vartheta_W} c_{\vartheta_W}} (g_f^V \bar{f} \gamma^\mu f + g_f^A \bar{f} \gamma^\mu \gamma_5 f) Z_\mu, \quad (78)$$

and we recovered the SM vectorial and axial couplings in Tab. 10.

4 The gauge coupling unification in the SU(8) theory

The gauge coupling unification relies on the RGEs of the SU(8) theory. The two-loop RGE of a gauge coupling of α_Υ in the $\overline{\text{MS}}$ scheme is given by [30]

$$\frac{d\alpha_\Upsilon^{-1}(\mu)}{d \log \mu} = \frac{b_\Upsilon^{(1)}}{2\pi} + \sum_{\Upsilon'} \frac{b_{\Upsilon\Upsilon'}^{(2)}}{8\pi^2 \alpha_\Upsilon^{-1}(\mu)}, \quad (79)$$

where the one-loop and two-loop β coefficients for the non-Abelian groups are

$$b_\Upsilon^{(1)} = -\frac{11}{3}C_2(\mathcal{G}_\Upsilon) + \frac{2}{3} \sum_F T(\mathcal{R}_\Upsilon^F) + \frac{1}{3} \sum_S T(\mathcal{R}_\Upsilon^S), \quad (80a)$$

$$\begin{aligned} b_{\Upsilon\Upsilon'}^{(2)} = & -\frac{34}{3}C_2(\mathcal{G}_\Upsilon)^2 \alpha_\Upsilon - \sum_F \left[2 \sum_{\Upsilon'} C_2(\mathcal{R}_{\Upsilon'}^F) \alpha_{\Upsilon'} + \frac{10}{3}C_2(\mathcal{G}_\Upsilon) \alpha_\Upsilon \right] T(\mathcal{R}_\Upsilon^F) \\ & - \sum_S \left[4 \sum_{\Upsilon'} C_2(\mathcal{R}_{\Upsilon'}^S) \alpha_{\Upsilon'} + \frac{2}{3}C_2(\mathcal{G}_\Upsilon) \alpha_\Upsilon \right] T(\mathcal{R}_\Upsilon^S). \end{aligned} \quad (80b)$$

Here, $C_2(\mathcal{G}_\Upsilon)$ is the quadratic Casimir of the group \mathcal{G}_Υ , $T(\mathcal{R}_\Upsilon^F)$ and $T(\mathcal{R}_\Upsilon^S)$ are trace invariants of the chiral fermions in the irrep of \mathcal{R}_Υ^F and complex scalars in the irrep of \mathcal{R}_Υ^S . For the U(1) Abelian groups

with charges denoted as $\mathcal{X}^{F/S}$, the one-loop β coefficients become

$$b_{\Upsilon}^{(1)} = \frac{2}{3} \sum_F (\mathcal{X}_{\Upsilon}^F)^2 + \frac{1}{3} \sum_S (\mathcal{X}_{\Upsilon}^S)^2. \quad (81)$$

4.1 RGEs of the minimal setup

By following the survival hypothesis [5, 31–35], we first list the minimal set of massless Higgs fields and the massless fermions between different symmetry breaking stages according the minimal Higgs VEVs in Eqs. (18). The two-loop RGEs within the minimal SU(8) setup are derived by using the PyR@TE [36].

Between the $v_{441} \leq \mu \leq v_U$, almost all massless Higgs fields are in the first lines of Eqs. (15), (16), and (17), which are

$$\begin{aligned} (\bar{\mathbf{4}}, \mathbf{1}, +\frac{1}{4})_{\mathbf{H},\omega} \oplus (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\omega} &\subset \overline{\mathbf{8}}_{\mathbf{H},\omega}, \\ (\mathbf{6}, \mathbf{1}, +\frac{1}{2})_{\mathbf{H},\dot{\omega}} \oplus (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{\omega}} \oplus (\bar{\mathbf{4}}, \bar{\mathbf{4}}, 0)_{\mathbf{H},\dot{\omega}} &\subset \overline{\mathbf{28}}_{\mathbf{H},\dot{\omega}}, \\ (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_{\mathbf{H}} \oplus (\bar{\mathbf{4}}, \mathbf{4}, -\frac{1}{2})_{\mathbf{H}} \oplus (\mathbf{6}, \mathbf{6}, 0)_{\mathbf{H}} &\subset \mathbf{70}_{\mathbf{H}}. \end{aligned} \quad (82)$$

All SU(8) fermions in Eq. (3) remain massless after the decomposition into the \mathcal{G}_{441} irreps. Correspondingly, we have the \mathcal{G}_{441} β coefficients of

$$\begin{aligned} (b_{\text{SU}(4)_s}^{(1)}, b_{\text{SU}(4)_W}^{(1)}, b_{\text{U}(1)_{X_0}}^{(1)}) &= (+\frac{13}{3}, +\frac{13}{3}, +\frac{55}{3}), \\ b_{\mathcal{G}_{441}}^{(2)} &= \begin{pmatrix} 2299/6 & 405/2 & 12 \\ 405/2 & 2299/6 & 12 \\ 180 & 180 & 31 \end{pmatrix}. \end{aligned} \quad (83)$$

Between the $v_{341} \leq \mu \leq v_{441}$, the massless Higgs fields are

$$\begin{aligned} (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},3,\text{V},\text{VI}} &\subset \overline{\mathbf{8}}_{\mathbf{H},3,\text{V},\text{VI}}, \\ (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{1},\dot{2},\text{VIII}} &\subset (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{1},\dot{2},\text{VIII}} \subset \overline{\mathbf{28}}_{\mathbf{H},\dot{1},\dot{2},\text{VIII}}, \\ (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\dot{1},\dot{2},\text{VII},\text{IX}} &\subset (\bar{\mathbf{4}}, \bar{\mathbf{4}}, 0)_{\mathbf{H},\dot{1},\dot{2},\text{VII},\text{IX}} \subset \overline{\mathbf{28}}_{\mathbf{H},\dot{1},\dot{2},\text{VII},\text{IX}}, \\ (\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})'_{\mathbf{H}} &\subset (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_{\mathbf{H}} \subset \mathbf{70}_{\mathbf{H}}. \end{aligned} \quad (84)$$

The massless \mathcal{G}_{341} fermions are listed as follows

$$\begin{aligned} &\left[(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega} \right] \oplus (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{F}}^{\Omega} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}, \\ \Omega &= (\omega, \dot{\omega}), \quad \omega = (3, \text{V}, \text{VI}), \quad \dot{\omega} = (\dot{1}, \dot{2}, \text{VII}, \text{VIII}, \text{IX}), \\ &(\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IV}} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\text{IV}}, \\ &(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}} \oplus (\mathbf{1}, \mathbf{6}, +\frac{1}{2})_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{4}, -\frac{1}{12})_{\mathbf{F}} \subset \mathbf{28}_{\mathbf{F}}, \\ &(\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})_{\mathbf{F}} \oplus (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}} \oplus (\mathbf{1}, \mathbf{1}, -1)_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{6}, +\frac{1}{6})_{\mathbf{F}} \oplus (\mathbf{1}, \mathbf{6}, +\frac{1}{2})'_{\mathbf{F}} \\ \oplus &(\mathbf{3}, \mathbf{4}, -\frac{1}{12})'_{\mathbf{F}} \oplus (\bar{\mathbf{3}}, \mathbf{4}, -\frac{5}{12})_{\mathbf{F}} \subset \mathbf{56}_{\mathbf{F}}. \end{aligned} \quad (85)$$

Correspondingly, we have the \mathcal{G}_{341} β coefficients of

$$\begin{aligned} (b_{\text{SU}(3)_c}^{(1)}, b_{\text{SU}(4)_W}^{(1)}, b_{\text{U}(1)_{X_1}}^{(1)}) &= \left(-\frac{5}{3}, -3, +\frac{247}{18}\right), \\ b_{\mathcal{G}_{341}}^{(2)} &= \begin{pmatrix} 226/3 & 75/2 & 97/36 \\ 20 & 571/4 & 187/24 \\ 194/9 & 935/8 & 8639/432 \end{pmatrix}. \end{aligned} \quad (86)$$

Between the $v_{331} \leq \mu \leq v_{341}$, the massless Higgs fields are

$$\begin{aligned} (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},3,\text{VI}} &\subset (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},3,\text{VI}} \subset \overline{\mathbf{8}}_{\mathbf{H},3,\text{VI}}, \\ (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})'_{\mathbf{H},\dot{2},\text{VIII}} &\subset (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{2},\text{VIII}} \subset (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H},\dot{2},\text{VIII}} \subset \overline{\mathbf{28}}_{\mathbf{H},\dot{2},\text{VIII}}, \\ (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},\dot{2},\text{IX}} &\subset (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H},\dot{2},\text{IX}} \subset (\bar{\mathbf{4}}, \bar{\mathbf{4}}, 0)_{\mathbf{H},\dot{2},\text{IX}} \subset \overline{\mathbf{28}}_{\mathbf{H},\dot{2},\text{IX}}, \\ (\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})'''_{\mathbf{H}} &\subset (\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})'_{\mathbf{H}} \subset (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_{\mathbf{H}} \subset \mathbf{70}_{\mathbf{H}}. \end{aligned} \quad (87)$$

The massless \mathcal{G}_{331} fermions are listed as follows

$$\begin{aligned} & \left[(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega} \right] \oplus \left[(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{F}}^{\Omega} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega'} \right] \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}, \\ & \Omega = (\omega, \dot{\omega}), \quad \omega = (3, \text{VI}), \quad \dot{\omega} = (\dot{1}, \dot{2}, \text{VIII}, \text{IX}), \\ & (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IV}} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\text{IV}}, \quad (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{V},\text{V}''} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\text{V}}, \\ & (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{VII}} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{VII}''} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\text{VII}}, \\ & (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}} \oplus (\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{3}, 0)_{\mathbf{F}} \subset \mathbf{28}_{\mathbf{F}}, \\ & (\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})'_{\mathbf{F}} \oplus (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{3}, 0)'_{\mathbf{F}} \\ & \oplus (\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})''_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{3}, 0)''_{\mathbf{F}} \oplus (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})'''_{\mathbf{F}} \subset \mathbf{56}_{\mathbf{F}}. \end{aligned} \quad (88)$$

Correspondingly, we have the \mathcal{G}_{331} β coefficients of

$$\begin{aligned} (b_{3c}^{(1)}, b_{3W}^{(1)}, b_{X_2}^{(1)}) &= \left(-5, -\frac{23}{6}, +\frac{82}{9}\right), \\ b_{\mathcal{G}_{331}}^{(2)} &= \begin{pmatrix} 12 & 12 & 2 \\ 12 & 113/3 & 38/9 \\ 16 & 304/9 & 304/27 \end{pmatrix}. \end{aligned} \quad (89)$$

Between the $v_{\text{EW}} \leq \mu \leq v_{331}$, there is only one massless SM Higgs field of

$$(\mathbf{1}, \bar{\mathbf{2}}, +\frac{1}{2})'''_{\mathbf{H}} \subset (\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})'''_{\mathbf{H}} \subset (\mathbf{1}, \bar{\mathbf{4}}, +\frac{3}{4})'_{\mathbf{H}} \subset (\mathbf{4}, \bar{\mathbf{4}}, +\frac{1}{2})_{\mathbf{H}} \subset \mathbf{70}_{\mathbf{H}}. \quad (90)$$

The massless SM fermions (together with massless sterile neutrinos) are listed as follows

$$\begin{aligned}
& \left[(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}^{\Omega} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega} \right] \oplus \left[(\mathbf{1}, \bar{\mathbf{2}}, -\frac{1}{2})_{\mathbf{F}}^{\Omega} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega'} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\Omega''} \right] \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}, \quad \Omega = (\dot{1}, \dot{2}, 3), \\
& (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IV}} \oplus \dots \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IX}} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega}, \\
& (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{VI}'} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{VIII}'} \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IX}'} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega'}, \\
& (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{V}''} \oplus \dots \oplus (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{\text{IX}''} \subset \overline{\mathbf{8}}_{\mathbf{F}}^{\Omega''}, \\
& (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}} \oplus (\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}} \oplus (\mathbf{3}, \mathbf{2}, +\frac{1}{6})_{\mathbf{F}} \subset \mathbf{28}_{\mathbf{F}}, \\
& (\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}}' \oplus (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}}' \oplus (\mathbf{3}, \mathbf{2}, +\frac{1}{6})_{\mathbf{F}}' \\
& \oplus (\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}}''' \oplus (\mathbf{3}, \mathbf{2}, +\frac{1}{6})_{\mathbf{F}}''' \oplus (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\mathbf{F}}''' \subset \mathbf{56}_{\mathbf{F}}.
\end{aligned} \tag{91}$$

Correspondingly, we have the usual SM β coefficients of

$$\begin{aligned}
(b_{3c}^{(1)}, b_{2W}^{(1)}, b_Y^{(1)}) &= (-7, -\frac{19}{6}, +\frac{41}{6}), \\
b_{\mathcal{G}_{\text{SM}}}^{(2)} &= \begin{pmatrix} -26 & 9/2 & 11/6 \\ 12 & 35/6 & 3/2 \\ 44/3 & 9/2 & 199/18 \end{pmatrix}.
\end{aligned} \tag{92}$$




Figure 1: The RGEs of the minimal SU(8) setup. The dashed lines and the solid lines represent the one-loop and two-loop RGEs, respectively.

With the one- and two-loop β coefficients defined in Eqs. (83), (86), (89), and (92), we plot the RGEs of the minimal SU(8) setup in Fig. 1. Three intermediate symmetry breaking scales follow from the benchmark point in Eq. (2). The discontinuities of the U(1) gauge couplings at three intermediate scales follow from their definitions in Eqs. (28), (48), (65), respectively. The RG evolutions within $10^3 \text{ GeV} \lesssim \mu \lesssim 10^{13} \text{ GeV}$ are zoomed out in order to highlight the behaviors in three intermediate symmetry breaking scales. The current plot indicates $\alpha_{4S}^{-1}(\mu) \sim \alpha_{4W}^{-1}(\mu) \simeq 40$ and $\alpha_{X_0}^{-1}(\mu) \simeq 20$, for $v_{441} \lesssim \mu \lesssim 10^{18} \text{ GeV}$. Furthermore, the one-loop β coefficients of $b_{\text{SU}(4)_S}^{(1)} = b_{\text{SU}(4)_W}^{(1)} < b_{\text{U}(1)_{X_0}}^{(1)}$ from Eq. (83) guarantee that two non-Abelian SU(4)_{S,W} gauge couplings can never unify with the Abelian U(1)_{X₀} gauge coupling in the minimal setup.

The one-loop threshold effect was tentatively neglected in Fig. 1. This was usually considered as a major source to modify the RG behavior within the field theory [12, 37, 38], and it can be generically expressed as

$$\begin{aligned}
\alpha_{\text{UV}}^{-1}(\mu) &= \alpha_{\text{IR}}^{-1}(\mu) + \frac{\Delta_{\text{IR}}(\mu)}{12\pi}, \\
\Delta_{\text{IR}}(\mu) &= C_2(\mathcal{G}_{\text{UV}}) - C_2(\mathcal{G}_{\text{IR}}) - 21 \sum_{V_k \in \mathcal{G}_{\text{IR}}} \dim(V_k) \log \frac{M_{V_k}}{\mu} \\
&\quad + 2 \sum_{S_k \in \mathcal{G}_{\text{IR}}} \dim(S_k) \log \frac{M_{S_k}}{\mu} + 4 \sum_{\mathcal{F}_k \in \mathcal{G}_{\text{IR}}} \dim(\mathcal{F}_k) \log \frac{M_{\mathcal{F}_k}}{\mu}, \tag{93}
\end{aligned}$$

where V_k , S_k and \mathcal{F}_k represent the vector bosons, complex scalars and Weyl fermions that are integrated out at the threshold scale μ . Their dimensions are always given in terms of the broken subgroup of \mathcal{G}_{IR} . One may evaluate contributions due to the threshold effects in Eq. (93) to see if they can alter the gauge coupling evolutions significantly. Let us take the renormalization scale of $\mu = \overline{M}_{V_k}$ (with \overline{M}_{V_k} being the average of the gauge boson masses) in Eq. (93), and list the contributions from the massive scalars and fermions at each symmetry breaking stage. Given the RG behaviors depicted in Fig. 1, we will parametrize the differences between the Abelian and the non-Abelian gauge couplings. For simplicity, we take a common mass for both massive fermions and massive scalar fields at each stage, and we estimate the largest possible threshold effects by assuming these masses to be 10 times larger than the renormalization scale of $\mu = \overline{M}_{V_k}$. The massive vectorlike fermions at each symmetry breaking stage have been summarized in Tab. 6.

At the \mathcal{G}_{331} -breaking stage, we find

$$\begin{aligned}
\{\text{massive } S_k\} &= (\mathbf{1}, \overline{\mathbf{2}}, -\frac{1}{2})_{\mathbf{H}, 3, \text{VI}, \dot{2}, \text{IX}} \oplus (\mathbf{1}, \overline{\mathbf{2}}, -\frac{1}{2})'_{\mathbf{H}, \dot{2}, \text{VIII}}, \\
(\Delta_{3c}, \Delta_{2W}, \Delta_Y) &= (0, 1, 1) + (6, 0, 4) \log \frac{M_{(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}}}}{\overline{M}_{V_{331}}} + (6, 0, 4) \log \frac{M_{(\overline{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}}}{\overline{M}_{V_{331}}} \\
&+ (0, 6, 6) \log \frac{M_{(\mathbf{1}, \overline{\mathbf{2}}, +\frac{1}{2})_{\mathbf{F}}}}{\overline{M}_{V_{331}}} + (0, 6, 6) \log \frac{M_{(\mathbf{1}, \overline{\mathbf{2}}, -\frac{1}{2})_{\mathbf{F}}}}{\overline{M}_{V_{331}}} + (0, 6, 6) \log \frac{M_{(\mathbf{1}, \overline{\mathbf{2}}, -\frac{1}{2})_{\mathbf{H}}}}{\overline{M}_{V_{331}}} \\
&= (12 \log \frac{M_{\mathcal{F}}}{\overline{M}_{V_{331}}}, 1 + 12 \log \frac{M_{\mathcal{F}}}{\overline{M}_{V_{331}}} + 6 \log \frac{M_{\mathcal{S}}}{\overline{M}_{V_{331}}}, 1 + 20 \log \frac{M_{\mathcal{F}}}{\overline{M}_{V_{331}}} + 6 \log \frac{M_{\mathcal{S}}}{\overline{M}_{V_{331}}}), \\
\Rightarrow \Delta_Y - \frac{\Delta_{3c} + \Delta_{2W}}{2} &= \frac{1}{2} + 8 \log \frac{M_{\mathcal{F}}}{\overline{M}_{V_{331}}} + 3 \log \frac{M_{\mathcal{S}}}{\overline{M}_{V_{331}}} \lesssim 25. \tag{94}
\end{aligned}$$

At the \mathcal{G}_{341} -breaking stage, we find

$$\begin{aligned}
\{\text{massive } S_k\} &= (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},\mathbf{V}} \oplus (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H},\mathbf{i},\mathbf{V}\text{II}} \oplus (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})'_{\mathbf{H},\mathbf{i}} \oplus (\mathbf{1}, \mathbf{3}, -\frac{2}{3})_{\mathbf{H},\mathbf{i}}, \\
(\Delta_{3c}, \Delta_{3W}, \Delta_{X_2}) &= (0, 1, \frac{2}{3}) + (4, 0, \frac{8}{3}) \log \frac{M_{(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (4, 0, \frac{8}{3}) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} \\
&+ (0, 4, \frac{8}{3}) \log \frac{M_{(\mathbf{1}, \mathbf{3}, +\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (0, 4, \frac{8}{3}) \log \frac{M_{(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (6, 6, 4) \log \frac{M_{(\mathbf{3}, \bar{\mathbf{3}}, +\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (6, 6, 4) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{3}, -\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{341}}} \\
&+ (0, 0, 4) \log \frac{M_{(\mathbf{1}, \mathbf{1}, +1)_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (0, 0, 4) \log \frac{M_{(\mathbf{1}, \mathbf{1}, -1)_{\mathbf{F}}}}{\bar{M}_{V_{341}}} + (0, 4, \frac{8}{3}) \log \frac{M_{(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\mathbf{H}}}}{\bar{M}_{V_{341}}} + (0, 1, \frac{8}{3}) \log \frac{M_{(\mathbf{1}, \mathbf{3}, -\frac{2}{3})_{\mathbf{H}}}}{\bar{M}_{V_{341}}} \\
&= (20 \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{341}}}, 1 + 20 \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{341}}} + 5 \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{341}}}, \frac{2}{3} + \frac{80}{3} \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{341}}} + \frac{16}{3} \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{341}}}), \\
\Rightarrow \Delta_{X_2} - \frac{\Delta_{3c} + \Delta_{3W}}{2} &= \frac{1}{6} + \frac{20}{3} \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{341}}} + \frac{17}{6} \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{341}}} \lesssim 21. \tag{95}
\end{aligned}$$

At the \mathcal{G}_{441} -breaking stage, we find

$$\begin{aligned}
\{\text{massive } S_k\} &= (\bar{\mathbf{3}}, \mathbf{1}, -\frac{1}{3})_{\mathbf{H}, \Omega=(\omega, \dot{\omega})} \oplus (\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H}, \mathbf{IV}} \oplus (\mathbf{3}, \mathbf{1}, +\frac{2}{3})_{\mathbf{H}, \dot{\omega}} \\
&\oplus (\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H}, \mathbf{VII}, \mathbf{IX}} \oplus (\mathbf{1}, \mathbf{4}, -\frac{3}{4})_{\mathbf{H}} \\
&\oplus (\mathbf{1}, \mathbf{1}, -1)_{\mathbf{H}} \oplus (\mathbf{1}, \mathbf{1}, +1)_{\mathbf{H}} \oplus (\mathbf{3}, \bar{\mathbf{4}}, +\frac{5}{12})_{\mathbf{H}} \oplus (\bar{\mathbf{3}}, \mathbf{4}, -\frac{5}{12})_{\mathbf{H}} \oplus (\mathbf{3}, \mathbf{6}, +\frac{1}{6})_{\mathbf{H}} \oplus (\bar{\mathbf{3}}, \mathbf{6}, -\frac{1}{6})_{\mathbf{H}} \\
(\Delta_{3c}, \Delta_{4W}, \Delta_{X_1}) &= (1, 0, \frac{2}{3}) + (2, 0, \frac{4}{3}) \log \frac{M_{(\mathbf{3}, \mathbf{1}, -\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{441}}} + (2, 0, \frac{4}{3}) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{F}}}}{\bar{M}_{V_{441}}} \\
&+ (0, 2, 1) \log \frac{M_{(\mathbf{1}, \mathbf{4}, +\frac{1}{4})_{\mathbf{F}}}}{\bar{M}_{V_{441}}} + (0, 2, 1) \log \frac{M_{(\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{F}}}}{\bar{M}_{V_{441}}} \\
&+ (9, 0, 6) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (0, 2, 1) \log \frac{M_{(\mathbf{1}, \bar{\mathbf{4}}, -\frac{1}{4})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (5, 0, \frac{40}{3}) \log \frac{M_{(\mathbf{3}, \mathbf{1}, +\frac{2}{3})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} \\
&+ (0, 4, 6) \log \frac{M_{(\mathbf{1}, \mathbf{6}, -\frac{1}{2})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (20, 15, \frac{5}{6}) \log \frac{M_{(\bar{\mathbf{3}}, \bar{\mathbf{4}}, +\frac{1}{12})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} \\
&+ (4, 3, \frac{25}{6}) \log \frac{M_{(\mathbf{3}, \bar{\mathbf{4}}, +\frac{5}{12})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (4, 3, \frac{25}{6}) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{4}, -\frac{5}{12})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} \\
&+ (0, 0, 2) \log \frac{M_{(\mathbf{1}, \mathbf{1}, +1)_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (0, 0, 2) \log \frac{M_{(\mathbf{1}, \mathbf{1}, -1)_{\mathbf{H}}}}{\bar{M}_{V_{441}}} \\
&+ (6, 6, 1) \log \frac{M_{(\mathbf{3}, \mathbf{6}, +\frac{1}{6})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (6, 6, 1) \log \frac{M_{(\bar{\mathbf{3}}, \mathbf{6}, -\frac{1}{6})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} + (0, 1, \frac{9}{2}) \log \frac{M_{(\mathbf{1}, \mathbf{4}, -\frac{3}{4})_{\mathbf{H}}}}{\bar{M}_{V_{441}}} \\
&= (1 + 4 \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{441}}} + 54 \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{441}}}, 4 \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{441}}} + 40 \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{441}}}, \frac{2}{3} + \frac{14}{3} \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{441}}} + 46 \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{441}}}), \\
\Rightarrow \Delta_{X_1} - \frac{\Delta_{3c} + \Delta_{4W}}{2} &= \frac{1}{6} + \frac{2}{3} \log \frac{M_{\mathcal{F}}}{\bar{M}_{V_{441}}} - \log \frac{M_{\mathcal{S}}}{\bar{M}_{V_{441}}} \lesssim -0.6. \tag{96}
\end{aligned}$$

By collecting all threshold effects from Eqs. (94), (95), and (96) into Eq. (93), we find the overall threshold shift of $\Delta_{U(1)}/(12\pi)$ relative to the threshold effects of two non-Abelian couplings to be at most $\sim \mathcal{O}(1)$. Hence, the one-loop threshold effect cannot significantly modify the RG evolutions in Fig. 1 such that the large discrepancy of $|\alpha_{X_0}^{-1} - (\alpha_{4S}^{-1} + \alpha_{4W}^{-1})/2| \sim 20$ can be eliminated.

4.2 The gauge coupling unification with additional adjoint Higgs fields

The RG evolutions of the non-Abelian and Abelian gauge couplings with the minimal SU(8) setup as depicted in Fig. 1 do not achieve the unification, since the U(1) gauge couplings get enhanced much faster than two non-Abelian couplings with the current field contents. A tentative solution within the field theoretical framework is to further introduce adjoint Higgs fields of $\mathbf{63}_H$'s. If one assumes the $(\mathbf{15}, \mathbf{1}, 0)_H \oplus (\mathbf{1}, \mathbf{15}, 0)_H$ components in Eq. (9) remain massless after the GUT-scale symmetry breaking, they bring additional positive contributions to the β coefficients of two non-Abelian gauge couplings. Naturally, one expects the unification scale to locate below the reduced Planck scale. We display two examples of this possibility in Fig. 2, while they both require as many as 150 – 200 adjoint massless Higgs fields of $(\mathbf{15}, \mathbf{1}, 0)_H \oplus (\mathbf{1}, \mathbf{15}, 0)_H$ within the range of $v_{441} \lesssim \mu \lesssim v_U$ such that $v_U \simeq 5 \times 10^{17}$ GeV.




Figure 2: The RGEs of the SU(8) theory with 150 additional Higgs fields of $(\mathbf{15}, \mathbf{1}, 0)_H \oplus (\mathbf{1}, \mathbf{15}, 0)_H$ (left panel), and 200 additional Higgs fields of $(\mathbf{15}, \mathbf{1}, 0)_H \oplus (\mathbf{1}, \mathbf{15}, 0)_H$ (right panel). The dashed lines and the solid lines represent the one-loop and two-loop RGEs, respectively.

Given that the unification scale is precious close to the Planck scale, there were also consideration of the gravitational effects to the gauge coupling unification, such as in the context of the SU(5) and SO(10) GUTs [15, 16, 39, 40]. At the leading order, we consider the following $d = 5$ HSW operator in the SU(8) theory

$$\mathcal{O}_{\text{HSW}} \equiv -\frac{1}{2} \frac{c_{\text{HSW}}}{M_{\text{pl}}} \text{Tr}[\mathbf{63}_H U^{\mu\nu} U_{\mu\nu}], \quad (97)$$

with $U_{\mu\nu} = U_{\mu\nu}^{\mathbf{21}} T^{\mathbf{21}}$ being the SU(8) field strength. After the GUT-scale symmetry breaking with the VEV in Eq. (8), this HSW operator shifts two SU(4) $_{S/W}$ field strengths of $G_{\mu\nu} = G_{\mu\nu}^{\bar{A}} T^{\bar{A}}$ and $A_{\mu\nu} = A_{\mu\nu}^{\bar{I}} T^{\bar{I}}$ as follows

$$-\frac{1}{2} \left(1 - \frac{1}{4} c_{\text{HSW}} \zeta_0\right) \text{Tr}(G_{\mu\nu} G^{\mu\nu}) - \frac{1}{2} \left(1 + \frac{1}{4} c_{\text{HSW}} \zeta_0\right) \text{Tr}(A_{\mu\nu} A^{\mu\nu}), \quad (98)$$

with $\zeta_0 \equiv v_U/M_{\text{pl}}$. The gauge coupling unification is modified into

$$\left(1 - \frac{1}{4} c_{\text{HSW}} \zeta_0\right) \alpha_{4S}(v_U) = \left(1 + \frac{1}{4} c_{\text{HSW}} \zeta_0\right) \alpha_{4W}(v_U) = \alpha_{X_0}(v_U) = \alpha_U(v_U). \quad (99)$$

Accordingly, we find the following results

$$\begin{aligned} n_{\text{Adj}} = 150 & : c_{\text{HSW}} = 1.91, & v_U \simeq 4.9 \times 10^{17} \text{ GeV}, & \alpha_U^{-1}(v_U) \simeq 21.4, \\ n_{\text{Adj}} = 200 & : c_{\text{HSW}} = 2.54, & v_U \simeq 3.5 \times 10^{17} \text{ GeV}, & \alpha_U^{-1}(v_U) \simeq 22.5, \end{aligned} \quad (100)$$

in accordance to two plots in Fig. 2.

5 Summary and outlook

5.1 The main results

We study the RGEs of the gauge couplings in an SU(8) theory through its SWW symmetry breaking pattern in Eq. (1). Three intermediate symmetry breaking scales follow from the previous Ref. [10], which were obtained such that three generational SM quark/lepton masses and the CKM mixing pattern are consistently predicted. The gauge couplings at each symmetry breaking stage are defined in Sec. 3. Together with the symmetry breaking scales determined in Eq. (2), we obtain the RG evolution of three gauge couplings in Fig. 1 for the minimal setup. The gauge coupling unification cannot be achieved even when we include the one-loop threshold effects into account. The only possible solution within the field theory is to include addition 150 – 200 adjoint massless Higgs fields of $(\mathbf{15}, \mathbf{1}, 0)_{\mathbf{H}} \oplus (\mathbf{1}, \mathbf{15}, 0)_{\mathbf{H}}$ within the range of $v_{441} \lesssim \mu \lesssim v_U$.

We wish to mention that the examples shown in Fig. 2 are only possible accommodation when the current SWW symmetry breaking pattern is considered. Within the field theoretical framework, alternative symmetry breaking patterns other than the one in Eq. (1) are possible, such as

$$\text{WSW} \quad : \quad \text{SU}(8) \xrightarrow{v_U} \mathcal{G}_{441} \xrightarrow{v_{441}} \mathcal{G}_{431} \xrightarrow{v_{431}} \mathcal{G}_{331} \xrightarrow{v_{331}} \mathcal{G}_{\text{SM}} \xrightarrow{v_{\text{EW}}} \text{SU}(3)_c \otimes \text{U}(1)_{\text{EM}}, \quad (101a)$$

$$\text{WWS} \quad : \quad \text{SU}(8) \xrightarrow{v_U} \mathcal{G}_{441} \xrightarrow{v_{441}} \mathcal{G}_{431} \xrightarrow{v_{431}} \mathcal{G}_{421} \xrightarrow{v_{421}} \mathcal{G}_{\text{SM}} \xrightarrow{v_{\text{EW}}} \text{SU}(3)_c \otimes \text{U}(1)_{\text{EM}}, \quad (101b)$$

or even the one where the SU(8) was not maximally broken, such as in Ref. [14].

5.2 On the SUSY extension and the Kač-Moody extension

More broadly speaking, the unified theories were extended to an $\mathcal{N} = 1$ SUSY theory, such as the SUSY SU(5) theory [13]. The gauge coupling unification was achieved at $v_U \simeq 2 \times 10^{16}$ GeV, with the sparticle masses assumed to be $\sim \mathcal{O}(1)$ TeV therein. A SUSY SU(8) theory with non-trivially embedded SM fermions was not known before, and it is useful to sketch some underlying issues below for the future work.

- The current minimal SU(8) theory is anomalous, when one promotes the fields in Eqs. (3) and (7) into chiral superfields. At least two types of SUSY extensions to the minimal SU(8) theory are possible, with the additional chiral superfields of

$$\{H\}_{\text{I}} = \mathbf{8}_{\mathbf{H}}^{\omega} \oplus \mathbf{28}_{\mathbf{H}}^{\dot{\omega}}, \quad \omega = (3, \text{IV}, \text{V}, \text{VI}), \quad \dot{\omega} = (\dot{1}, \dot{2}, \text{VII}, \text{VIII}, \text{IX}), \quad (102a)$$

$$\{H\}_{\text{II}} = \mathbf{36}_{\mathbf{H}} \oplus \mathbf{36}_{\mathbf{H}}'. \quad (102b)$$

- With the superpotential of

$$W = \frac{1}{2} M_U \text{Tr} \Phi_{\text{Adj}}^2 + \frac{\lambda}{3} \text{Tr} \Phi_{\text{Adj}}^3, \quad \Phi_{\text{Adj}} \equiv \mathbf{24}_{\mathbf{H}}, \quad (103)$$

in the SUSY SU(5) theory, Witten has argued that two distinctive symmetry breaking patterns of SU(5) $\rightarrow \mathcal{G}_{\text{SM}}$ and SU(5) $\rightarrow \mathcal{G}_{41}$ are degenerate [41]. For the SU(8) theory with $\Phi_{\text{Adj}} \equiv \mathbf{63}_{\mathbf{H}}$, the symmetry breaking patterns of

$$\text{SU}(8) \rightarrow \mathcal{G}_{441}, \quad \text{SU}(8) \rightarrow \mathcal{G}_{351}/\mathcal{G}_{531}, \quad \text{SU}(8) \rightarrow \mathcal{G}_{621}, \quad \text{SU}(8) \rightarrow \mathcal{G}_{71}, \quad (104)$$

at the GUT scale all lead to degenerate minima. The subgroups of all four distinctive symmetry breaking patterns contain the \mathcal{G}_{SM} as their subgroup, hence they are phenomenologically reasonable. Correspondingly, the gauge couplings will be defined differently from the maximally breaking pattern of $\text{SU}(8) \rightarrow \mathcal{G}_{441}$.

- The other issue in the SUSY $\text{SU}(8)$ theory is the sparticle masses, which are determined by the SUSY breaking mechanisms. The modifications of the RGEs between the non-SUSY sector and the SUSY sector will be determined accordingly. Additional contributions to the one-loop threshold effect in Eq. (93) are necessary to account for the difference between the dimensional reduction scheme in the SUSY sector and the $\overline{\text{MS}}$ scheme in the non-SUSY sector [42].

Lastly, we briefly comment on a possible unification scenario within the string theoretical framework, which was previously studied in the $\text{SU}(5)/\text{SO}(10)$ theories and their SUSY extensions [43, 44]. In the heterotic string theory, Ginsparg has shown in Ref. [45] that the gauge and gravitational couplings must unify according to

$$k_i g_i^2 = g_{\text{string}}^2 = \frac{8\pi G_N}{\alpha'}, \quad (105)$$

with k_i being the Kač-Moody levels of each gauge couplings, and α' being the Regge slope. The string unification scale was previously given by [46]

$$M_{\text{string}} \approx g_{\text{string}} \times 5 \times 10^{17} \text{ GeV}, \quad (106)$$

which is quite close to the v_{441} in Eq. (2). It is notable that our benchmark scales given in Ref. [10] were estimated by fitting the SM flavor parameters within the $\text{SU}(8)$ framework, and are independent of the derivation of the string scale in Ref. [45, 46]. To achieve the string unification based on some extensions from the minimal setup of the $\text{SU}(8)$ theory, one has to fully study all possible symmetry breaking patterns under the SUSY extensions described above, as well as to look for the constraints of the unitarity and conformal invariance from the Kač-Moody Lie algebra.

Acknowledgements

We would like to thank Tianjun Li, Kaiwen Sun, Yuan Sun, Yinan Wang, Zhong-Zhi Xianyu, and Wenbin Yan for very enlightening discussions and communications. N.C. would like to thank Southeast University, and Nanjing University for hospitality when preparing this work. N.C. is partially supported by the National Natural Science Foundation of China (under Grants No. 12035008 and No. 12275140) and Nankai University. Y.N.M. is partially supported by the National Natural Science Foundation of China (under Grant No. 12205227), the Fundamental Research Funds for the Central Universities (WUT: 2022IVA052), and Wuhan University of Technology.

A Conventions, decomposition rules and charge quantizations in the $\text{SU}(8)$ theory

According to the symmetry breaking pattern in Eq. (1), we define the conventions of gauge group indices in Tab. 11. The fundamental and anti-fundamental representations will be denoted by superscripts and subscripts, respectively. We also define the decomposition rules in the $\text{SU}(8)$ theory. After

Indices	group	irrep	range
$\mathbf{a}, \mathbf{b}, \mathbf{c}$	$SU(8)$	fundamental anti-fundamental	$1, \dots, 8$
$\mathfrak{A}, \mathfrak{B}, \mathfrak{C}$	$SU(8)$	adjoint	$1, \dots, 63$
$\bar{a}, \bar{b}, \bar{c}$	$SU(4)_s$	fundamental anti-fundamental	$\clubsuit, \diamond, \heartsuit, \spadesuit$
$\bar{A}, \bar{B}, \bar{C}$	$SU(4)_s$	adjoint	$1, \dots, 15$
a, b, c	$SU(3)_c$	fundamental anti-fundamental	$\clubsuit, \diamond, \heartsuit$
A, B, C	$SU(3)_c$	adjoint	$1, \dots, 8$
$\bar{i}, \bar{j}, \bar{k}$	$SU(4)_W$	fundamental anti-fundamental	$1, 2, 3, 4$
$\bar{I}, \bar{J}, \bar{K}$	$SU(4)_W$	adjoint	$1, \dots, 15$
$\tilde{i}, \tilde{j}, \tilde{k}$	$SU(3)_W$	fundamental anti-fundamental	$1, 2, 3$
$\tilde{I}, \tilde{J}, \tilde{K}$	$SU(3)_W$	adjoint	$1, \dots, 8$
i, j	$SU(2)_W$	fundamental anti-fundamental	$1, 2$
I, J, K	$SU(2)_W$	adjoint	$1, 2, 3$

Table 11: Definition of indices for various gauge groups from the maximal $SU(8)$ symmetry breaking pattern. The fundamental and anti-fundamental indices will be distinguished by superscripts and subscripts, respectively.

the GUT-scale symmetry breaking, we define the $U(1)_{X_0}$ charges as follows

$$\hat{X}_0(\mathbf{8}) \equiv \text{diag}\left(\underbrace{-\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}}_{4_s}, \underbrace{+\frac{1}{4}, +\frac{1}{4}, +\frac{1}{4}, +\frac{1}{4}}_{4_w}\right). \quad (107)$$

Sequentially, the $U(1)_{X_1}$, $U(1)_{X_2}$, and $U(1)_Y$ charges are defined according to the $SU(4)_s$ and the $SU(4)_W$ fundamental representations as follows

$$\hat{X}_1(\mathbf{4}_s) \equiv \text{diag}\left(\underbrace{\left(-\frac{1}{12} + \mathcal{X}_0\right)\mathbb{I}_{3 \times 3}}_{\mathbf{3}_c}, \frac{1}{4} + \mathcal{X}_0\right), \quad (108a)$$

$$\hat{X}_2(\mathbf{4}_w) \equiv \text{diag}\left(\underbrace{\left(\frac{1}{12} + \mathcal{X}_1\right)\mathbb{I}_{3 \times 3}}_{\mathbf{3}_w}, -\frac{1}{4} + \mathcal{X}_1\right), \quad (108b)$$

$$\begin{aligned} \hat{Y}(\mathbf{4}_w) &\equiv \text{diag}\left(\left(\frac{1}{6} + \mathcal{X}_2\right)\mathbb{I}_{2 \times 2}, -\frac{1}{3} + \mathcal{X}_2, \mathcal{X}_2\right) \\ &= \text{diag}\left(\underbrace{\left(\frac{1}{4} + \mathcal{X}_1\right)\mathbb{I}_{2 \times 2}}_{\mathbf{2}_w}, \left(-\frac{1}{4} + \mathcal{X}_1\right)\mathbb{I}_{2 \times 2}\right), \end{aligned} \quad (108c)$$

$$\hat{Q}_e(\mathbf{4}_w) \equiv T_{SU(4)}^3 + \hat{Y}(\mathbf{4}_w) = \text{diag}\left(\frac{3}{4} + \mathcal{X}_1, \left(-\frac{1}{4} + \mathcal{X}_1\right)\mathbb{I}_{3 \times 3}\right). \quad (108d)$$

Based on the above definitions for the fundamental representations, the rules for other higher rank anti-symmetric representations can be derived by tensor productions. For adjoint representations of the $SU(4)_s$ and the $SU(4)_W$ groups, these charges are defined by setting all relevant $U(1)$ charges to zero in Eqs. (108)

$$\hat{X}_1(\mathbf{15}_s) \equiv \text{diag}\left(-\frac{1}{12}\mathbb{I}_{3 \times 3}, \frac{1}{4}\right), \quad (109a)$$

$$\hat{X}_2(\mathbf{15}_w) \equiv \text{diag}\left(\frac{1}{12}\mathbb{I}_{3 \times 3}, -\frac{1}{4}\right), \quad (109b)$$

$$\hat{Y}(\mathbf{15}_w) \equiv \text{diag}\left(\frac{1}{4}\mathbb{I}_{2 \times 2}, -\frac{1}{4}\mathbb{I}_{2 \times 2}\right), \quad (109c)$$

$$\hat{Q}_e(\mathbf{15}_w) \equiv \text{diag}\left(\frac{3}{4}, -\frac{1}{4}\mathbb{I}_{3 \times 3}\right). \quad (109d)$$

B Note added in proof of the $SU(8)$ maximal breaking pattern

Let us consider the maximally symmetry breaking pattern of an $SU(N)$ group of

$$SU(N) \rightarrow SU(n_S) \otimes SU(n_W) \otimes U(1), \quad (110)$$

with $n_S = n_W = N/2$ for the even- N , or $n_{S,W} = (N \pm 1)/2$ for the odd- N . The $U(1)$ charges of the $SU(N)$ fundamental representation after the symmetry breaking are given by

$$\text{diag}\left(\underbrace{-\frac{1}{n_S}, \dots, -\frac{1}{n_S}}_{\times n_S}, \underbrace{+\frac{1}{n_W}, \dots, +\frac{1}{n_W}}_{\times n_W}\right). \quad (111)$$

Meanwhile, the VEV of the adjoint Higgs for the symmetry-breaking pattern is given by

$$\langle \Phi_{\text{Adj}} \rangle = \frac{1}{\sqrt{2n_S n_W (n_S + n_W)}} \text{diag} \left(\underbrace{-n_W, \dots}_{\times n_S}, \underbrace{+n_S, \dots}_{\times n_W} \right) v_U, \quad (112)$$

with the normalization of $\text{Tr} \Phi_{\text{Adj}}^2 = \frac{1}{2} v_U^2$. If the U(1) coupling does not need to be normalized, one has the following relations

$$\begin{aligned} \frac{1}{n_S} &= \frac{n_W}{\sqrt{2n_S n_W (n_S + n_W)}} \Rightarrow n_S n_W = 2(n_S + n_W) = 2N, \\ \text{even} - N &: n_S = n_W = N/2 \Rightarrow N = 8, \\ \text{odd} - N &: n_S = (N + 1)/2, n_W = (N - 1)/2 \Rightarrow N^2 - 1 = 8N. \end{aligned} \quad (113)$$

Apparently, the condition for the odd- N case has no integer solution. Thus, the unique gauge group with its unique symmetry breaking pattern where the U(1) charge does not need additional normalization factor is the $\text{SU}(8) \rightarrow \mathcal{G}_{441}$.

References

- [1] H. Georgi and S. L. Glashow, “Unity of All Elementary Particle Forces,” *Phys. Rev. Lett.* **32** (1974) 438–441.
- [2] H. Fritzsch and P. Minkowski, “Unified Interactions of Leptons and Hadrons,” *Annals Phys.* **93** (1975) 193–266.
- [3] N. Cabibbo, “Unitary Symmetry and Leptonic Decays,” *Phys. Rev. Lett.* **10** (1963) 531–533.
- [4] M. Kobayashi and T. Maskawa, “CP Violation in the Renormalizable Theory of Weak Interaction,” *Prog. Theor. Phys.* **49** (1973) 652–657.
- [5] H. Georgi, “Towards a Grand Unified Theory of Flavor,” *Nucl. Phys. B* **156** (1979) 126–134.
- [6] CMS Collaboration, A. Tumasyan *et al.*, “A portrait of the Higgs boson by the CMS experiment ten years after the discovery,” *Nature* **607** no. 7917, (2022) 60–68, [arXiv:2207.00043 \[hep-ex\]](#).
- [7] ATLAS Collaboration, G. Aad *et al.*, “A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery,” *Nature* **607** no. 7917, (2022) 52–59, [arXiv:2207.00092 \[hep-ex\]](#). [Erratum: *Nature* 612, E24 (2022)].
- [8] S. M. Barr, “Doubly Lopsided Mass Matrices from Unitary Unification,” *Phys. Rev. D* **78** (2008) 075001, [arXiv:0804.1356 \[hep-ph\]](#).
- [9] N. Chen, Y.-n. Mao, and Z. Teng, “The global B – L symmetry in the flavor-unified SU(N) theories,” *JHEP* **04** (2024) 046, [arXiv:2307.07921 \[hep-ph\]](#).
- [10] N. Chen, Y.-n. Mao, and Z. Teng, “The Standard Model quark/lepton masses and the Cabibbo-Kobayashi-Maskawa mixing in an SU(8) theory,” [arXiv:2402.10471 \[hep-ph\]](#).
- [11] H. Georgi, H. R. Quinn, and S. Weinberg, “Hierarchy of Interactions in Unified Gauge Theories,” *Phys. Rev. Lett.* **33** (1974) 451–454.

- [12] L. J. Hall, “Grand Unification of Effective Gauge Theories,” *Nucl. Phys. B* **178** (1981) 75–124.
- [13] S. Dimopoulos and H. Georgi, “Softly Broken Supersymmetry and SU(5),” *Nucl. Phys. B* **193** (1981) 150–162.
- [14] Z.-q. Ma, T.-s. Tu, P.-y. Xue, and X.-j. Zhou, “An SU(8) Grand Unified Model Accomodating Three Generations With $V - A$ Coupling,” *Phys. Lett. B* **100** (1981) 399–402.
- [15] C. T. Hill, “Are There Significant Gravitational Corrections to the Unification Scale?,” *Phys. Lett. B* **135** (1984) 47–51.
- [16] Q. Shafi and C. Wetterich, “Modification of GUT Predictions in the Presence of Spontaneous Compactification,” *Phys. Rev. Lett.* **52** (1984) 875.
- [17] R. D. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys. Rev. Lett.* **38** (1977) 1440–1443.
- [18] L.-F. Li, “Group Theory of the Spontaneously Broken Gauge Symmetries,” *Phys. Rev. D* **9** (1974) 1723–1739.
- [19] X. Li and E. Ma, “Gauge Model of Generation Nonuniversality,” *Phys. Rev. Lett.* **47** (1981) 1788.
- [20] E. Ma, X. Li, and S. F. Tuan, “Gauge Model of Generation Nonuniversality Revisited,” *Phys. Rev. Lett.* **60** (1988) 495–498.
- [21] X.-y. Li and E. Ma, “Gauge model of generation nonuniversality reexamined,” *J. Phys. G* **19** (1993) 1265–1278, [arXiv:hep-ph/9208210](#).
- [22] N. Craig, D. Green, and A. Katz, “(De)Constructing a Natural and Flavorful Supersymmetric Standard Model,” *JHEP* **07** (2011) 045, [arXiv:1103.3708 \[hep-ph\]](#).
- [23] M. Bordone, C. Cornella, J. Fuentes-Martin, and G. Isidori, “A three-site gauge model for flavor hierarchies and flavor anomalies,” *Phys. Lett. B* **779** (2018) 317–323, [arXiv:1712.01368 \[hep-ph\]](#).
- [24] J. Fuentes-Martin, G. Isidori, J. Pagès, and B. A. Stefanek, “Flavor non-universal Pati-Salam unification and neutrino masses,” *Phys. Lett. B* **820** (2021) 136484, [arXiv:2012.10492 \[hep-ph\]](#).
- [25] J. Davighi and J. Tooby-Smith, “Electroweak flavour unification,” *JHEP* **09** (2022) 193, [arXiv:2201.07245 \[hep-ph\]](#).
- [26] J. Fuentes-Martin, G. Isidori, J. M. Lizana, N. Selimovic, and B. A. Stefanek, “Flavor hierarchies, flavor anomalies, and Higgs mass from a warped extra dimension,” *Phys. Lett. B* **834** (2022) 137382, [arXiv:2203.01952 \[hep-ph\]](#).
- [27] J. Davighi, G. Isidori, and M. Pesut, “Electroweak-flavour and quark-lepton unification: a family non-universal path,” *JHEP* **04** (2023) 030, [arXiv:2212.06163 \[hep-ph\]](#).
- [28] M. Fernández Navarro and S. F. King, “Tri-hypercharge: a separate gauged weak hypercharge for each fermion family as the origin of flavour,” *JHEP* **08** (2023) 020, [arXiv:2305.07690 \[hep-ph\]](#).

- [29] M. Fernández Navarro, S. F. King, and A. Vicente, “Tri-unification: a separate $SU(5)$ for each fermion family,” [arXiv:2311.05683 \[hep-ph\]](#).
- [30] M. E. Machacek and M. T. Vaughn, “Two Loop Renormalization Group Equations in a General Quantum Field Theory. 1. Wave Function Renormalization,” *Nucl. Phys. B* **222** (1983) 83–103.
- [31] S. L. Glashow, “The Future of Elementary Particle Physics,” *NATO Sci. Ser. B* **61** (1980) 687.
- [32] R. Barbieri, D. V. Nanopoulos, G. Morchio, and F. Strocchi, “Neutrino Masses in Grand Unified Theories,” *Phys. Lett. B* **90** (1980) 91–97.
- [33] R. Barbieri and D. V. Nanopoulos, “An Exceptional Model for Grand Unification,” *Phys. Lett. B* **91** (1980) 369–375.
- [34] R. Barbieri and D. V. Nanopoulos, “Hierarchical Fermion Masses From Grand Unification,” *Phys. Lett. B* **95** (1980) 43–46.
- [35] F. del Aguila and L. E. Ibanez, “Higgs Bosons in $SO(10)$ and Partial Unification,” *Nucl. Phys. B* **177** (1981) 60–86.
- [36] L. Sartore and I. Schienbein, “PyR@TE 3,” *Comput. Phys. Commun.* **261** (2021) 107819, [arXiv:2007.12700 \[hep-ph\]](#).
- [37] S. Weinberg, “Effective Gauge Theories,” *Phys. Lett. B* **91** (1980) 51–55.
- [38] A. Ernst, A. Ringwald, and C. Tamarit, “Axion Predictions in $SO(10) \times U(1)_{PQ}$ Models,” *JHEP* **02** (2018) 103, [arXiv:1801.04906 \[hep-ph\]](#).
- [39] M. K. Parida, P. K. Patra, and A. K. Mohanty, “Gravity Induced Large Grand Unification Mass in $SU(5)$ With Higher Dimensional Operators,” *Phys. Rev. D* **39** (1989) 316–322.
- [40] A. Datta, S. Pakvasa, and U. Sarkar, “Corrections to mass scale predictions in $SO(10)$ GUT with higher dimensional operators,” *Phys. Rev. D* **50** (1994) 2192–2198, [arXiv:hep-ph/9402356](#).
- [41] E. Witten, “Dynamical Breaking of Supersymmetry,” *Nucl. Phys. B* **188** (1981) 513.
- [42] P. Langacker and N. Polonsky, “Uncertainties in coupling constant unification,” *Phys. Rev. D* **47** (1993) 4028–4045, [arXiv:hep-ph/9210235](#).
- [43] A. Font, L. E. Ibanez, and F. Quevedo, “Higher Level Kac-Moody String Models and Their Phenomenological Implications,” *Nucl. Phys. B* **345** (1990) 389–430.
- [44] K. R. Dienes, “String theory and the path to unification: A Review of recent developments,” *Phys. Rept.* **287** (1997) 447–525, [arXiv:hep-th/9602045](#).
- [45] P. H. Ginsparg, “Gauge and Gravitational Couplings in Four-Dimensional String Theories,” *Phys. Lett. B* **197** (1987) 139–143.
- [46] V. S. Kaplunovsky, “One Loop Threshold Effects in String Unification,” *Nucl. Phys. B* **307** (1988) 145–156, [arXiv:hep-th/9205068](#). [Erratum: *Nucl.Phys.B* 382, 436–438 (1992)].