

COMPUTING THE CENTER OF A FUSION CATEGORY

FABIAN MÄURER AND ULRICH THIEL

ABSTRACT. We present a (Las Vegas) algorithm for explicitly computing the simple objects of the categorical (Drinfeld) center of a spherical fusion category. Our approach is based on decomposing the images of simple objects under the induction functor from the category to its center. We have implemented this algorithm in a general-purpose software framework `TENSORCATEGORIES.JL` for tensor categories that we develop within the open-source computer algebra system `OSCAR`. While the required computations are still too heavy to investigate standard examples whose center is not yet known up to some equivalence, our algorithm has the advantage of determining explicit half-braidings for the simple central objects, avoiding abstract equivalences and the like. Furthermore, it also works over not necessarily algebraically closed fields, and this yields new explicit examples of non-split modular categories.

1. INTRODUCTION

Let \mathcal{C} be a monoidal category with associators

$$(1.1) \quad a_{X,Y,Z}: (X \otimes Y) \otimes Z \xrightarrow{\cong} X \otimes (Y \otimes Z).$$

Throughout, we will use conventions as in the standard reference [9]. We are concerned with the *categorical (Drinfeld) center* $\mathcal{Z}(\mathcal{C})$ of \mathcal{C} . This is a categorification of the notion of the center of a monoid where, as usual, equalities are replaced by isomorphisms, and we keep track of these. Specifically, a *half-braiding* for an object $X \in \mathcal{C}$ is a natural isomorphism

$$(1.2) \quad \gamma_X = \{\gamma_X(Y): X \otimes Y \xrightarrow{\cong} Y \otimes X \mid Y \in \mathcal{C}\}$$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN–LANDAU, POSTFACH 3049, 67663 KAISERSLAUTERN, GERMANY

E-mail addresses: maeurer@mathematik.uni-kl.de, thiel@mathematik.uni-kl.de.

Date: June 18, 2024.

such that the diagram

$$(1.3) \quad \begin{array}{ccccc} & & (Y \otimes X) \otimes Z & \xrightarrow{a_{Y,X,Z}} & Y \otimes (X \otimes Z) & & \\ & \nearrow^{\gamma_X(Y) \otimes \text{id}_Z} & & & & \searrow^{\text{id}_Y \otimes \gamma_X(Z)} & \\ (X \otimes Y) \otimes Z & & & & & & Y \otimes (Z \otimes X) \\ & \searrow_{a_{X,Y,Z}} & & & & \nearrow_{a_{Y,Z,X}} & \\ & & X \otimes (Y \otimes Z) & \xrightarrow{\gamma_X(Y \otimes Z)} & (Y \otimes Z) \otimes X & & \end{array}$$

commutes for all $Y, Z \in \mathcal{C}$ and $\gamma_1 = \text{id}_X$. The center $\mathcal{Z}(\mathcal{C})$ is the category whose objects are pairs (X, γ_X) of an object $X \in \mathcal{C}$ and a half-braiding γ_X for X , and the morphisms $(X, \gamma_X) \rightarrow (Y, \gamma_Y)$ are morphisms $f: X \rightarrow Y$ in \mathcal{C} such that the diagram

$$(1.4) \quad \begin{array}{ccc} X \otimes Z & \xrightarrow{f \otimes \text{id}_Z} & Y \otimes Z \\ \downarrow \gamma_X(Z) & & \downarrow \gamma_Y(Z) \\ Z \otimes X & \xrightarrow{\text{id}_Z \otimes f} & Z \otimes Y \end{array}$$

commutes for all $Z \in \mathcal{C}$. The category $\mathcal{Z}(\mathcal{C})$ inherits a natural monoidal structure from \mathcal{C} , see [9, Section 7.13]. Moreover, the half-braidings induce the structure of a braiding (“commutativity constraint”) on $\mathcal{Z}(\mathcal{C})$, so that $\mathcal{Z}(\mathcal{C})$ is a braided monoidal category, see [9, Section 8.5]. Braided monoidal categories play an important role in knot theory and physics [1], which is one of the reasons why the center construction is important.

Among monoidal categories the fusion categories [10] play a major role since they behave somewhat like categorical analogues of groups and arise in several areas of mathematics and physics. The center of a fusion category is a braided fusion category, in fact a modular category [1], and thus of particular interest.

In this paper we will present an algorithm for computing the center of a spherical fusion category (Section 4). The algorithm is of Las Vegas type, meaning that if it returns a result, the result is mathematically correct, but it may happen that the algorithm does not succeed. This is due to the fact that we need to find a simple subobject of an object which amounts to finding primitive idempotents in its endomorphism algebra. No algorithm is known solving this problem over a general field but there are Las Vegas algorithms which perform very well in practice, namely the MeatAxe algorithm and its variants [22, 15, 17].

Our algorithm is theoretically straightforward. The main point is that we realized it in practice. To this end, we have begun developing a general-purpose software framework `TENSORCATEGORIES.JL` [12] for tensor categories and categorical representation theory. This framework is based on the open-source computer algebra system `OSCAR` [21, 5] which incorporates and extends powerful systems like `GAP` [13] and `Singular` [6]. `OSCAR`

in turn is based on the high-performance programming language Julia [3, 26] which, due its type system design and multiple dispatch paradigm, is nicely suited for working with categorical structures. Julia has a simple high-level syntax like Python but is much faster. OSCAR provides, e.g., fast algebra over number fields and over the algebraic closure of \mathbb{Q} , which is crucial for our purposes.

As with group representations, the theory of fusion categories shows its greatest development so far over an algebraically closed field of characteristic zero. An important aspect of our algorithm and implementation is that it works over any field. This leads to interesting new results, one of which we want to mention here.

Let \mathcal{C}' be the famous Ising fusion category. This is a fusion category over the complex numbers with three simple objects denoted by $\mathbb{1}$, χ , and X . The multiplication is given by $\chi \otimes \chi = \mathbb{1}$, $\chi \otimes X = X \otimes \chi = X$ and $X \otimes X = \mathbb{1} \oplus \chi$. The Ising category is a special case of Tambara–Yamagami fusion categories [25]. From general theory it is known [14, Proposition 4.1] that the center $\mathcal{Z}(\mathcal{C}')$ has 9 simple objects. The non-trivial associators of \mathcal{C}' are given by

$$\begin{aligned} a_{\chi, X, \chi} &= (-1) \text{id}_X \\ a_{X, \mathbb{1}, X} &= \text{id}_{\mathbb{1}} \oplus (-1) \text{id}_\chi \\ a_{X, \chi, X} &= (-1) \text{id}_{\mathbb{1}} \oplus \text{id}_\chi \\ a_{X, X, X} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \text{id}_{2X} , \end{aligned}$$

see [25]. It follows that \mathcal{C}' is actually defined over $\mathbb{Q}(\sqrt{2})$, i.e., it has a $\mathbb{Q}(\sqrt{2})$ -rational form \mathcal{C} . This is a fusion category over $\mathbb{Q}(\sqrt{2})$, and it is known from general theory that $\mathcal{Z}(\mathcal{C})$ is a semisimple tensor category—more precisely, it is what we call a weak pre-modular category, see Section 2. What is the structure of $\mathcal{Z}(\mathcal{C})$? As far as we are aware, this does not appear in the literature yet. Via explicit calculations with our software in Section 6 we will prove:

Theorem 1.1. *Let \mathcal{C} be the Ising fusion category considered over $\mathbb{Q}(\sqrt{2})$. Then $\mathcal{Z}(\mathcal{C})$ has five simple objects with dimensions 1, 1, 2, 2 and $4 \cdot \sqrt{2}$. The multiplication table is given by*

\otimes	$\mathbb{1}$	$\bar{\mathbb{1}}$	χ^2	$\mathbb{1} + \chi$	X^4
$\mathbb{1}$	$\mathbb{1}$	$\bar{\mathbb{1}}$	χ^2	$\mathbb{1} + \chi$	X^4
$\bar{\mathbb{1}}$	$\bar{\mathbb{1}}$	$\mathbb{1}$	χ^2	$\mathbb{1} + \chi$	X^4
χ^2	χ^2	χ^2	$2\mathbb{1} + 2\bar{\mathbb{1}}$	$2(\mathbb{1} + \chi)$	$2X^4$
$\mathbb{1} + \chi$	$\mathbb{1} + \chi$	$\mathbb{1} + \chi$	$2(\mathbb{1} + \chi)$	$\mathbb{1} + \bar{\mathbb{1}} + \chi^2$	$2X^4$
X^4	X^4	X^4	$2X^4$	$2X^4$	$4\mathbb{1} + 4\bar{\mathbb{1}} + 4\chi^2 + 8(\mathbb{1} + \chi)$

and the S -matrix is given by

$$S = \begin{pmatrix} 1 & 1 & 2 & 2 & 4 \cdot \sqrt{2} \\ 1 & 1 & 2 & 2 & -4 \cdot \sqrt{2} \\ 2 & 2 & 4 & -4 & 0 \\ 2 & 2 & -4 & 0 & 0 \\ 4 \cdot \sqrt{2} & -4 \cdot \sqrt{2} & 0 & 0 & 0 \end{pmatrix}$$

In particular, $\mathcal{Z}(\mathcal{C})$ is a non-split modular category. It splits over $\mathbb{Q}(\xi_{16})$ where ξ_{16} is a primitive 16-th root of unity.

We emphasize that our computational construction of the center is completely explicit so that we obtain the simple central objects as objects of \mathcal{C} together with half-braidings (omitted here to save space). We can also analyze the splitting of the central objects when passing to the algebraic closure of \mathbb{Q} , see Section 6.

As another illustration of the capabilities of our implementation, let us consider a fusion category coming from a Haagerup subfactor. There are three such fusion categories. Let \mathcal{H} be the one with six simple objects. Its multiplication table is given in Figure 1. The associators can be found in [29].

	$\mathbb{1}$	α	α^*	ρ	$\alpha\rho$	$\alpha^*\rho$
$\mathbb{1}$	$\mathbb{1}$	α	α^*	ρ	$\alpha\rho$	$\alpha^*\rho$
α	α	α^*	$\mathbb{1}$	$\alpha\rho$	$\alpha^*\rho$	ρ
α^*	α^*	$\mathbb{1}$	α	$\alpha^*\rho$	ρ	$\alpha\rho$
ρ	ρ	$\alpha^*\rho$	$\alpha\rho$	$\mathbb{1} \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\alpha \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$
$\alpha\rho$	$\alpha\rho$	$\alpha^*\rho$	ρ	$\alpha \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\mathbb{1} \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$
$\alpha^*\rho$	$\alpha^*\rho$	ρ	$\alpha\rho$	$\alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\alpha \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$	$\mathbb{1} \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$

FIGURE 1. Multiplication table of the Haagerup fusion category.

The modular structure of the center of \mathcal{H} is known, and it is known to have twelve simple objects [11]. The forget functor from the center is known as well [28], see Figure 1. We can recover these results explicitly with our software. If we define \mathcal{H} over the smallest field possible, which is a number field of degree 16, the center is computed within six

$$\begin{array}{l}
\mathbb{1} \\
\alpha \\
\alpha^* \\
\rho \\
\alpha\rho \\
\alpha^*\rho
\end{array}
\begin{pmatrix}
1 & 1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\
0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\
0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1
\end{pmatrix}$$

FIGURE 2. The forgetful functor $\mathcal{Z}(\mathcal{H}) \rightarrow \mathcal{H}$. The columns correspond to the simple objects of $\mathcal{Z}(\mathcal{H})$.

minutes. The center has two non-split objects. Splitting requires passing to a larger number field and takes another eight minutes. See Section 6 for further details.

In Section 2 we will specify our setting and make precise what we mean by “computing the center”. In Section 3 we discuss the computation of half-braidings to illustrate the complexity of the problem. In Section 4 we discuss our algorithm. In Section 5 we derive a theoretical result on the splitting of central objects under field extension. In Section 6 we present our software framework and illustrate it with the proof of Theorem 1.1.

Acknowledgments. This work is a contribution to the SFB-TRR 195 “Symbolic Tools in Mathematics and their Application” of the German Research Foundation (DFG), Project-ID 28623755. We thank Liam Rogel for testing our software and providing helpful feedback. We thank Gunter Malle for comments on a preliminary version of this paper.

CONTENTS

1. Introduction	1
2. Basic assumptions, notation, and facts	6
3. Computing half-braidings	11
4. An algorithm based on the induction functor	14
5. Splitting of simple central objects	20
6. Software framework and examples	23
References	29

2. BASIC ASSUMPTIONS, NOTATION, AND FACTS

We first want to emphasize that we do *not* require monoidal categories to be strict, i.e. the associators (1.1) may be non-identity morphisms. It is true that by Mac Lane's strictness theorem [9, Theorem 2.8.5], any monoidal category is monoidally equivalent to a strict monoidal category. But such a monoidal equivalence may be difficult to represent in the computer, which is against our algorithmic approach. Moreover, to represent a monoidal category in the computer one will usually choose a skeleton. Even though every category is equivalent to a skeletal category, a monoidal category may in general not be monoidally equivalent to a skeletal strict monoidal category, i.e. one cannot achieve strict and skeletal simultaneously, see [9, Remark 2.8.7]. We will thus incorporate associators everywhere.

We do assume, however, strictness for the unit object $\mathbb{1}$ of a monoidal category \mathcal{C} in the sense that we require the unit morphism $\iota: \mathbb{1} \otimes \mathbb{1} \rightarrow \mathbb{1}$ to be the identity, and for any $X \in \mathcal{C}$ we identify $\mathbb{1} \otimes X$ and $X \otimes \mathbb{1}$ with X . This is a basic assumption also in [9], see [9, Remark 2.2.9], and it is not problematic as was strictness because one can simply choose the skeleton appropriately, see [9, Exercise 2.8.8].

We let \mathbb{k} be a field. We do *not* require \mathbb{k} to be algebraically closed, and we do not make any assumptions on its characteristic. This level of generality is also addressed in [9, Section 4.16]. Even though we can work with an algebraic closure of \mathbb{Q} , say, in a computer algebra system like OSCAR, the arithmetic will be slower than in a number field. It will thus be of advantage to work over smaller fields. Moreover, this will lead us to interesting rationality questions. Since terminology in the theory of tensor categories over general fields is in parts not yet entirely consistent throughout the literature, we will be very precise about our definitions below. In general, our basis will be [9].

2.1. Fusion categories. We propose, and use, the following generalization of [9, Definition 4.1.1] to an arbitrary base field. By a *weak fusion category* over \mathbb{k} we mean a \mathbb{k} -linear abelian semisimple rigid monoidal category \mathcal{C} which has only finitely many non-isomorphic simple objects, the unit object is simple, the Hom-spaces are finite-dimensional over \mathbb{k} , and the bifunctor \otimes is \mathbb{k} -bilinear.¹ The Grothendieck ring $\text{Gr}(\mathcal{C})$ of such a category is a *weak fusion ring* as in [9, Section 3.8], see [8, Proposition 4.2], hence our terminology.

We call an object $X \in \mathcal{C}$ *scalar* if $\mathbb{k} \simeq \text{End}_{\mathcal{C}}(X)$ via the canonical map. We say that \mathcal{C} *splits* if all simple objects of \mathcal{C} are scalar, and then we call \mathcal{C} a *fusion category*. In case \mathbb{k} is algebraically closed, this definition coincides with [9, Definition 4.1.1]. Also, for a general field \mathbb{k} , it coincides with the definition in [4, Section 1.G], this paper being important

¹Such a category is finite in the sense of [9, Definition 1.8.6], the requirement of enough projectives following from semisimplicity. Moreover, the simplicity of the unit object implies that the category is indecomposable as required in [9, Definition 4.1.1].

for us since it generalizes many results and constructions from algebraically closed fields to arbitrary base fields (and rings). We want to incorporate the non-split case (with the notion of *weak*) since this will happen for the center, even if we start with a split category.

2.2. Coends. If \mathcal{C} is any category and $F: \mathcal{C} \times \mathcal{C}^{op} \rightarrow \mathcal{C}$ is a functor, then a *coend* of F is an object $C \in \mathcal{C}$ together with morphisms $c_X: F(X, X) \rightarrow C$ for each $X \in \mathcal{C}$ which are universal with the property that the diagram

$$(2.1) \quad \begin{array}{ccc} F(X, Y) & \xrightarrow{F(f, \text{id}_Y)} & F(Y, Y) \\ \downarrow F(\text{id}_X, f) & & \downarrow c_Y \\ F(X, X) & \xrightarrow{c_X} & C \end{array}$$

commutes for each morphism $f: X \rightarrow Y \in \mathcal{C}$. The coend object C , if it exists, is unique up to isomorphism. We refer to [18, Section IX.6] for more details on coends.

If \mathcal{C} is a rigid monoidal category, then if the functor $\mathcal{C} \times \mathcal{C}^{op} \rightarrow \mathcal{C}$ defined by $(X, Y) \mapsto X \otimes Y^*$ and $(f, g) \mapsto f \otimes g^*$, has a coend, it is called a (canonical) *coend* of \mathcal{C} . It follows from [24, Theorem 3.4] that any weak fusion category \mathcal{C} has a coend. If \mathcal{C} is a fusion category, a coend C is explicitly given by

$$(2.2) \quad C = \bigoplus_{i=1}^n X_i \otimes X_i^*,$$

where X_1, \dots, X_n is a complete set of representatives of the simple objects of \mathcal{C} , see [4, Section 3D].

2.3. Dimensions. If \mathcal{C} is a pivotal monoidal category, we have the notions of *left dimension* $\dim_l(X)$ and *right dimension* $\dim_r(X)$ for any object $X \in \mathcal{C}$, see [9, Section 4.7]. These dimensions are elements of $\text{End}_{\mathcal{C}}(\mathbb{1})$. Let us call an object X *spherical* if the left and right dimensions of X coincide. We then denote this common dimension by $\dim(X)$. If \mathcal{C} admits a coend C and if C is spherical, we set

$$(2.3) \quad \dim(\mathcal{C}) := \dim(C)$$

and call this the (categorical) *dimension* of \mathcal{C} . This definition is as in [4, Section 3D]. If \mathcal{C} is a spherical fusion category, it follows from (2.2) and the multiplicativity of dimension, [9, Proposition 4.7.12], that

$$(2.4) \quad \dim(\mathcal{C}) = \sum_{i=1}^n \dim(X_i)^2 \in \text{End}_{\mathcal{C}}(\mathbb{1}) \simeq \mathbb{k}.$$

Our definition of $\dim(\mathcal{C})$ thus agrees with [9, Definition 7.21.3].

2.4. Modular categories. By a (weak) *pre-modular category* we mean a braided spherical (weak) fusion category. To such a category \mathcal{C} we associate a matrix S defined as follows: if X_1, \dots, X_n are the simple objects of \mathcal{C} up to isomorphism, then S is the square matrix S of size n with entries

$$(2.5) \quad S_{ij} = \text{Tr}(c_{X_j X_i} c_{X_i X_j}) \in \mathbb{k} .$$

Here, $c_{X_i X_j}$ denotes the braiding $X_i \otimes X_j \rightarrow X_j \otimes X_i$ and Tr denotes the trace [9, Equation 8.40], which relies on the spherical structure. We call S the *S-matrix* of \mathcal{C} . We say that a pre-modular category is *modular* if its S -matrix is invertible. Our definitions of pre-modular, the S -matrix, and modular are the same as in [9] (Definitions 8.13.1, 8.13.2, and 8.14.1), but we note that in [9] all this is only considered in case where \mathbb{k} is algebraically closed of characteristic zero. We nonetheless think they make sense in general and they will be convenient for stating some facts. But we emphasize that our work does not depend on these definitions. In [4, Section 3E] the notion of a modular category over a general field (or ring) is defined which involves a non-degeneracy condition for a pairing defined using the coend. A modular category as defined here is modular as defined in [4] by [4, Remark 3.4].

2.5. Structure of the center. The key fact about the center we use here is the following.

Theorem 2.1 (Bruguières–Virelizier [4]). *Let \mathcal{C} be a spherical fusion category over a field \mathbb{k} with $\dim(\mathcal{C}) \neq 0$. Then $\mathcal{Z}(\mathcal{C})$ is a weak pre-modular category and*

$$(2.6) \quad \dim(\mathcal{Z}(\mathcal{C})) = \dim(\mathcal{C})^2 .$$

If $\mathcal{Z}(\mathcal{C})$ splits, it is a modular category.

Let us make some remarks on this fact. It is a deep conjecture [10, Conjecture 2.8] that any fusion category over an algebraically closed field of characteristic zero is pivotal. As far as we are aware, all known fusion categories are spherical. We note, however, that a given pivotal structure may not be spherical, see [27, Example 4.59]. If \mathbb{k} is of characteristic zero, the condition $\dim(\mathcal{C}) \neq 0$ is always true by [10, Theorem 2.3].² All the relevant structures on \mathcal{C} (direct sums, linearity, duals, pivotal, spherical) naturally transfer to $\mathcal{Z}(\mathcal{C})$, see [4, Section 1J] and [24, Theorem 3.6]. The only property left to establish that $\mathcal{Z}(\mathcal{C})$ is a weak fusion category is semisimplicity, and this is indeed not true in general, see [4, Example 2.5]. In fact, it is proven in [4, Corollary 2.3] that $\mathcal{Z}(\mathcal{C})$ is semisimple if and only if $\dim(\mathcal{C}) \neq 0$. So, if $\dim(\mathcal{C}) \neq 0$, it follows that $\mathcal{Z}(\mathcal{C})$ is a weak pre-modular category. The relation $\dim(\mathcal{Z}(\mathcal{C})) = \dim(\mathcal{C})^2$ is [4, Theorem 2.1]. Finally, if $\mathcal{Z}(\mathcal{C})$ splits, it follows from [4, Theorem 2.1] and our remarks in Section 2.4 that it is a modular category.

²In [10] the base field is assumed to be algebraically closed. But our definition of a fusion category requires the category to be split and then we can argue by scalar extension.

We note that in general it is not true that $\mathcal{Z}(\mathcal{C})$ splits: using our software we will show in Section 6 that the center of the Ising category over $\mathbb{Q}(\sqrt{2})$ is a non-split weak modular category. We note that the phenomenon of non-splitting of the center of a split fusion category is known, see e.g. [19]. We hope that our software will help to explore further examples of this exciting phenomenon. Finally, we note that the above theorem in case of an algebraically closed field was established much earlier by Müger [20]. The upshot of [4] was to establish it for any field.

2.6. Computing the center. We can now make precise what we mean by “computing the center”: given a spherical fusion category \mathcal{C} over a field \mathbb{k} with $\dim(\mathcal{C}) \neq 0$, we want to determine a complete set of representatives of the simple objects of $\mathcal{Z}(\mathcal{C})$. Since $\mathcal{Z}(\mathcal{C})$ is semisimple, this gives complete knowledge of $\mathcal{Z}(\mathcal{C})$.

In our computational perspective that we take on this problem we actually want to achieve more. First, to work explicitly with such a category \mathcal{C} in the computer we need to assume we have some sort of finite encoding (model) of \mathcal{C} . One, but not the only, possibility would be via the multiplication table of the Grothendieck ring $\text{Gr}(\mathcal{C})$ of \mathcal{C} and the 6j-symbols of \mathcal{C} , which encode the associators via a collection of matrices [9, Section 4.9]. Determining such a model may require hard work in specific examples, but we assume it exists and start from there. Our software provides a general framework to work with models of categories, the specifics being up to the user. We have already implemented models of several standard examples. We can now formulate our precise goal.

Goal. We want to compute a model of $\mathcal{Z}(\mathcal{C})$ in terms of the model of the original category \mathcal{C} , avoiding abstract equivalences and the like. In particular, we want to compute the simple objects of $\mathcal{Z}(\mathcal{C})$ explicitly as pairs of an object together with a half-braiding.

Example 2.2. Let us illustrate our goal in a simple example. Let G be a finite group and let $\mathbb{k}\text{-Vec}_G$ be the category of finite-dimensional G -graded vector spaces over \mathbb{k} , see [9, Example 2.3.6]. This is a spherical fusion category with simple objects $\{\delta_g\}_{g \in G}$, where δ_g is the 1-dimensional vector space concentrated in degree g . Here, the associators are trivial. There are versions $\mathbb{k}\text{-Vec}_G^\omega$ with non-trivial associator as well, see [9, Remark 2.6.2]. It follows from Equation (2.4) that $\dim(\mathbb{k}\text{-Vec}_G) = |G| \cdot 1_{\mathbb{k}}$. The center $\mathcal{Z}(\mathbb{k}\text{-Vec}_G)$ is well-known [9, Example 8.5.4]: the simple objects are parametrized by pairs (C, V) , where C is a conjugacy class in G and V is an irreducible finite-dimensional representation of the centralizer of an element $g \in C$.

But actual central objects are pairs (X, γ_X) of a G -graded vector space X and a half-braiding γ_X for X . How do the actual simple central objects look like? Answering this is

our goal, and precisely this is done by our algorithm and software implementation. We give a concrete example in Figure 3. Note that it is very easy to model $\mathbb{k}\text{-Vec}_G$ in the computer: we just need the group G and we describe morphisms as certain matrices over \mathbb{k} .

	(12)	(13)	(23)	(123)	(132)
()	1	1	1	1	1
()	-1	-1	-1	1	1
$2 \cdot ()$	$\begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix}$
$(23) \oplus (12) \oplus (13)$	$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$
$(23) \oplus (12) \oplus (13)$	$\begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 1 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}$
$(132) \oplus (123)$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	id	id
$(132) \oplus (123)$	$\begin{bmatrix} 0 & \xi_3 \\ \xi_3^2 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3^2 \\ \xi_3 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3^2 \\ \xi_3 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3 \\ \xi_3^2 & 0 \end{bmatrix}$
$(132) \oplus (123)$	$\begin{bmatrix} 0 & \xi_3^2 \\ \xi_3 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3 \\ \xi_3^2 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3 \\ \xi_3^2 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \xi_3^2 \\ \xi_3 & 0 \end{bmatrix}$

FIGURE 3. The actual center of $\mathbb{k}\text{-Vec}_G$ for G the symmetric group S_3 and $\mathbb{k} = \mathbb{Q}(\xi_3)$, where ξ_3 is a primitive third root of unity. The table lists the simple central objects (X, γ_X) . The first column lists the underlying object X of $\mathbb{k}\text{-Vec}_G$. The other columns are indexed by the simple objects Y of $\mathbb{k}\text{-Vec}_G$ and specify the half-braiding $\gamma_X(Y)$. The data in this table is readily provided by our software.

Remark 2.3. We note that our goal of computing the center makes sense whenever $\mathcal{Z}(\mathcal{C})$ is semisimple. For example, it is shown in [9, Theorem 9.3.2] that $\mathcal{Z}(\mathcal{C})$ is a fusion category, thus semisimple, whenever \mathcal{C} is a fusion category over an algebraically closed field of characteristic zero. So here, \mathcal{C} does not necessarily have to be spherical in contrast to our discussion above. For our algorithm, however, we need to assume the category is spherical, and we currently do not know how to compute the center without this assumption.

We also note that even without semisimplicity of the center $\mathcal{Z}(\mathcal{C})$ one could still ask how to compute it. But then knowledge of simple objects is not sufficient to really know the center itself. If $\mathcal{Z}(\mathcal{C})$ is a finite tensor category, meaning that objects are of finite length, then the Krull–Schmidt theorem holds [9, Theorem 1.5.7], so that the category is determined by its indecomposable objects. So, one needs to work with indecomposable objects instead of just simple objects. We are currently working on extending our software framework to support finite tensor categories, but we currently have no idea how to compute the center in this generality.

3. COMPUTING HALF-BRAIDINGS

Before we discuss our algorithm and software implementation, we first want to discuss the computation of half-braidings in order to illustrate the complexity of this problem.

Note that we have a forget functor

$$(3.1) \quad F: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}, \quad (X, \gamma_X) \mapsto X .$$

The image of the induced map $\text{Gr}(\mathcal{Z}(\mathcal{C})) \rightarrow \text{Gr}(\mathcal{C})$ between Grothendieck rings obviously lies in the center of $\text{Gr}(\mathcal{C})$. In the example in Figure 3 we can observe two crucial peculiarities of the center:

- (1) If (X, γ_X) is a simple object of $\mathcal{Z}(\mathcal{C})$, then X is not necessarily simple in \mathcal{C} .
- (2) If (X, γ_X) is a simple object of $\mathcal{Z}(\mathcal{C})$ there may exist another non-isomorphic half-braiding γ'_X for X , so that (X, γ'_X) is another simple object of $\mathcal{Z}(\mathcal{C})$. In other words, there may be several non-isomorphic simple objects of $\mathcal{Z}(\mathcal{C})$ lying above the same object of \mathcal{C} under the forget functor.

It seems a possible strategy to compute the center would be to inductively form direct sums Z of the simple objects in \mathcal{C} and determine the half-braidings on Z . The inductive procedure would ensure that this produces simple central objects and, over a splitting field, one could use the dimension formula (2.6) with the coend formula (2.2) to establish that one found all simple central objects. However, we will now argue that the determination of half-braidings and their isomorphism classes as objects of the center seems computationally infeasible.

First, semisimplicity of \mathcal{C} allows one to encode a half-braiding γ_Z by a finite amount of information, namely by the maps $\gamma_Z(X_i)$ for the simple objects X_i of \mathcal{C} . We used this already in the example in Figure 3. The precise statement is as follows.

Lemma 3.1 (Müger [20, Lemma 3.3]). *Let \mathcal{C} be a fusion category with simple objects $\{X_i\}$. Let $Z \in \mathcal{C}$. There is a bijection between half-braidings for Z and families of morphisms $\{\gamma_Z(X_i) \in \text{Hom}_{\mathcal{C}}(Z \otimes X_i, X_i \otimes Z)\}$ such that for all i, j, k and $t \in \text{Hom}_{\mathcal{C}}(X_k, X_i \otimes X_j)$ the*

diagram

$$(3.2) \quad \begin{array}{ccccc} Z \otimes X_k & \xrightarrow{\gamma_Z(X_k)} & X_k \otimes Z & \xrightarrow{t \otimes \text{id}_Z} & (X_i \otimes X_j) \otimes Z \\ \text{id}_Z \otimes t \downarrow & & & & \downarrow a_{X_i, X_j, Z} \\ Z \otimes (X_i \otimes X_j) & & & & X_i \otimes (X_j \otimes Z) \\ \downarrow a_{Z, X_i, X_j}^{-1} & & & & \uparrow \text{id}_{X_i} \otimes \gamma_Z(X_j) \\ (Z \otimes X_i) \otimes X_j & \xrightarrow{\gamma_Z(X_i) \otimes \text{id}_{X_j}} & (X_i \otimes Z) \otimes X_j & \xrightarrow{a_{X_i, Z, X_j}} & X_i \otimes (Z \otimes X_j) \end{array}$$

commutes and $\gamma_Z(\mathbb{1}) = \text{id}_Z$.

Remark 3.2. This statement is proven in [20] more generally for a semisimple monoidal category. In [20] the base field is assumed to be algebraically closed but this does not play a role in the proof. In [20] monoidal categories are assumed to be strict. We have given the non-strict version here. The statement and its proof are straightforward but there is one important detail which is less obvious: half-braidings need to be isomorphisms but we just consider morphisms $\gamma_Z(X_i)$ without requiring invertibility. The fact that we automatically get isomorphisms is due to the existence of dual objects in \mathcal{C} (rigidity) and [20, Lemma 2.2].

Lemma 3.1 allows us to set up a system of equations for half-braidings γ_Z for $Z \in \mathcal{C}$. Let X_i, X_j, X_k be simple objects of \mathcal{C} and let $t \in \text{Hom}_{\mathcal{C}}(X_k, X_i \otimes X_j)$. Then the lemma states an equation

$$(3.3) \quad \phi(\gamma_Z(X_k), t) = \psi(\gamma_Z(X_i), \gamma_Z(X_j), t),$$

where

$$(3.4) \quad \phi(\gamma_Z(X_k), t) = a_{X_i, X_j, X_k} \circ (t \otimes \text{id}_Z) \circ \gamma_Z(X_k)$$

and

$$(3.5) \quad \psi(\gamma_Z(X_i), \gamma_Z(X_j), t) = (\text{id}_{X_i} \otimes \gamma_Z(X_j)) \circ a_{X_i, Z, X_j} \circ (\gamma_Z X_i \otimes \text{id}_{X_j}) \circ a_{Z, X_i, X_j}^{-1} \circ (\text{id}_Z \otimes t).$$

Clearly ϕ and ψ are linear in t as well as in $\gamma_Z(X_i)$, $\gamma_Z(X_j)$, and $\gamma_Z(X_k)$ respectively. Thus, whenever the equations hold for a basis of $\text{Hom}_{\mathcal{C}}(X_k, X_i \otimes X_j)$, they hold for all t . After choosing a basis f_1, \dots, f_r for $\text{Hom}(Z \otimes X_k, X_i \otimes (Z \otimes X_j))$ and bases $g_1^l, \dots, g_{r_l}^l$ for $\text{Hom}_{\mathcal{C}}(Z \otimes X_l, X_l \otimes Z)$ we can replace $\gamma_Z(X_l)$ with

$$(3.6) \quad \gamma_Z(X_l) = a_1^l g_1^l + \dots + a_{r_l}^l g_{r_l}^l$$

and get a system of *quadratic* equations by comparing coefficients. Writing out the full details is technical, but we have implemented this system in our software framework and the exact details can be found in the source code. Using algebraic solvers like `msolve` [2]

this, in principle, yields an approach to computing the half-braidings for a fixed object. However, as we illustrate in Example 3.3 below, even in very small examples, the systems are extremely complicated and usually form a positive-dimensional ideal, so that there are infinitely many solutions and `msolve` cannot be used. If the base field is not a finite field or the field of rational numbers, the computational situation is even worse. Also, even if we have found (a parametric form of) the solutions, we still need to determine their isomorphism classes as objects of the center. Hence, without further simplifications from theory (which we do not know) this approach seems infeasible.

Example 3.3. Consider the category $\mathbb{k}\text{-Vec}_G$ for $G = S_3$ and $\mathbb{k} = \mathbb{Q}$. Consider the object $Z = (23) \oplus (12) \oplus (13)$. We can see in Figure 3, which we computed using our algorithm from the next section, that there are exactly two non-isomorphic half-braidings for Z . Note that they are indeed defined over \mathbb{Q} , we do not need the bigger field $\mathbb{Q}(\xi_3)$ as in Figure 3. The ideal defined by the equations for half-braidings on Z is given in Figure 4. It is generated by 78 polynomials in $18 = 6 \cdot 3$ indeterminates. Its dimension is 2. In this (simple) example, one could derive a parametric solution from the Groebner basis.

$$\begin{aligned}
& \langle x_1 - x_4^2, x_2 - x_5x_6, x_3 - x_5x_6, x_1 - x_7x_9, x_2 - x_8^2, x_3 - x_7x_9, x_1 - x_{10}x_{14}, x_2 - x_{11}x_{15}, x_3 - x_{12}x_{13}, x_1 - x_{12}x_{13}, \\
& x_2 - x_{10}x_{14}, x_3 - x_{11}x_{15}, x_1 - x_{16}x_{17}, x_2 - x_{16}x_{17}, x_3 - x_{18}^2, x_4 - x_7x_{12}, x_5 - x_8x_{11}, x_6 - x_9x_{10}, x_4 - x_{10}x_{17}, \\
& x_5 - x_{11}x_{18}, x_6 - x_{12}x_{16}, x_4 - x_9x_{13}, x_5 - x_7x_{14}, x_6 - x_8x_{15}, x_4 - x_{14}x_{16}, x_5 - x_{13}x_{17}, x_6 - x_{15}x_{18}, -x_4x_{13} + x_7, \\
& -x_5x_{15} + x_8, -x_6x_{14} + x_9, -x_5x_{10} + x_7, -x_6x_{11} + x_8, -x_4x_{12} + x_9, x_7 - x_{13}x_{18}, x_8 - x_{14}x_{16}, x_9 - x_{15}x_{17}, \\
& x_7 - x_{11}x_{16}, x_8 - x_{10}x_{17}, x_9 - x_{12}x_{18}, -x_4x_{16} + x_{10}, -x_5x_{18} + x_{11}, -x_6x_{17} + x_{12}, -x_6x_7 + x_{10}, -x_5x_8 + x_{11}, \\
& -x_4x_9 + x_{12}, x_{10} - x_{13}x_{15}, x_{11} - x_{13}x_{14}, x_{12} - x_{14}x_{15}, -x_8x_{16} + x_{10}, -x_7x_{17} + x_{11}, -x_9x_{18} + x_{12}, -x_4x_7 + x_{13}, \\
& -x_5x_9 + x_{14}, -x_6x_8 + x_{15}, -x_7x_{18} + x_{13}, -x_8x_{17} + x_{14}, -x_9x_{16} + x_{15}, -x_{10}x_{11} + x_{13}, -x_{11}x_{12} + x_{14}, -x_{10}x_{12} + x_{15}, \\
& -x_5x_{16} + x_{13}, -x_4x_{17} + x_{14}, -x_6x_{18} + x_{15}, -x_4x_{10} + x_{16}, -x_5x_{12} + x_{17}, -x_6x_{11} + x_{18}, -x_7x_{15} + x_{16}, -x_8x_{14} + x_{17}, \\
& -x_9x_{13} + x_{18}, -x_8x_{10} + x_{16}, -x_9x_{11} + x_{17}, -x_7x_{12} + x_{18}, -x_6x_{13} + x_{16}, -x_4x_{14} + x_{17}, -x_5x_{15} + x_{18}, x_1 - 1, x_2 - 1, \\
& x_3 - 1 \rangle \\
& = \langle x_{18}^2 - 1, x_{16}x_{17} - 1, x_{14} - x_{17}x_{18}, x_{13}x_{15} - x_{14}x_{16}^2, x_{12} - x_{14}x_{15}, x_{11} - x_{13}x_{14}, x_{10} - x_{13}x_{15}, x_9 - x_{12}x_{18}, x_8 - x_{10}x_{17}, \\
& x_7 - x_{11}x_{16}, x_6 - x_{15}x_{18}, x_5 - x_{13}x_{17}, x_4 - x_{14}x_{16}, x_3 - 1, x_2 - 1, x_1 - 1 \rangle
\end{aligned}$$

FIGURE 4. The ideal for half-braidings for the object $(23) \oplus (12) \oplus (13)$ in $\mathbb{Q}\text{-Vec}_{S_3}$. The second expression is a Groebner basis with respect to the lexicographic ordering.

4. AN ALGORITHM BASED ON THE INDUCTION FUNCTOR

We will now present our algorithm for computing the center. We note that the idea is rather simple in theory. The difficulties lie in making it constructive and effective—a lot of work was spent on the actual software implementation.

Assumption 4.1. Throughout, we assume that \mathcal{C} is a spherical fusion category over a field \mathbb{k} with $\dim(\mathcal{C}) \neq 0$. We denote by X_1, \dots, X_n a complete set of the isomorphism classes of simple objects of \mathcal{C} . We denote by $\psi: (-) \rightarrow (-)^{**}$ the (spherical) pivotal structure on \mathcal{C} . For an object X we denote by $\text{ev}_X: X^* \otimes X \rightarrow \mathbb{1}$ and $\text{coev}_X: \mathbb{1} \rightarrow X \otimes X^*$ the evaluation and coevaluation morphisms, respectively.

We note that by [9, Theorem 4.8.4] we have $\dim(X_i) \neq 0$. In the proof of this fact in [9] the base field is assumed to be algebraically closed, but the proof remains valid for a split simple object, which is the case in our definition of a fusion category.

4.1. The induction functor and the basic idea. Recall the forget functor $F: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ from Equation 3.1. It admits a two-sided adjoint

$$(4.1) \quad I: \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C}),$$

called the *induction functor* [9, Section 9.2]. Explicitly, for $V \in \mathcal{C}$ the underlying object of $I(V)$ is given by

$$(4.2) \quad FI(V) = \bigoplus_{i=1}^n (X_i \otimes V) \otimes X_i^*$$

and the half-braiding for this object is given by the component morphisms

$$(4.3) \quad \gamma(Z)_{ij} = \sum_{f \in B, g \in B'} \begin{array}{ccc} ((X_i \otimes V) \otimes X_i^*) \otimes W & \xrightarrow{a_{X_i \otimes V, X_i^*, W}} & (X_i \otimes V) \otimes (X_i^* \otimes W) \\ & & \downarrow (f \otimes \text{id}_V) \otimes g \\ & & ((W \otimes X_j) \otimes V) \otimes X_j^* \\ & & \downarrow a_{W, X_j, V} \otimes \text{id}_{X_j^*} \\ W \otimes ((X_j \otimes V) \otimes X_j^*) & \xleftarrow{a_{W, X_j \otimes V, X_j^*}} & (W \otimes (X_j \otimes V)) \otimes X_j^* \end{array},$$

where B and B' are a pair of dual bases of $\text{Hom}_{\mathcal{C}}(X_i, W \otimes X_j)$ and $\text{Hom}_{\mathcal{C}}(W \otimes X_i, X_j) \cong \text{Hom}_{\mathcal{C}}(X_i^* \otimes W, X_j^*)$. These details can be found [4, Section 2]. We have added the associators here.

We want to give the explicit adjunction isomorphisms between the forget functor and the induction functor. To this end, let us assume here that $X_1 = \mathbb{1}$. The left adjunction isomorphisms given by

$$(4.4) \quad \begin{aligned} \text{Hom}_{\mathcal{C}}(V, Y) &\cong \text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(V), (Y, \gamma)) \\ f &\mapsto \sum_i \dim X_i \cdot \phi_i(f) \circ p_i \end{aligned}$$

where $p_i: \bigoplus_i (X_i \otimes V) \otimes X_i^* \rightarrow (X_i \otimes V) \otimes X_i^*$ are the projections and $\phi_i(f)$ is given by

$$(4.5) \quad \begin{array}{ccc} (X_i \otimes V) \otimes X_i^* & \xrightarrow{\text{id}_{X_i} \otimes f \otimes \text{id}_{X_i^*}} & (X_i \otimes Y) \otimes X_i^* & \xrightarrow{\gamma(X_i)^{-1} \otimes \text{id}_{X_i^*}} & (Y \otimes X_i) \otimes X_i^* \\ \downarrow \text{:=}\phi_i(f) & & & & \downarrow a_{Y, X_i, X_i^*} \\ Y & \xleftarrow{\text{id}_Y \otimes \text{ev}_{X_i^*}} & Y \otimes (X_i^{**} \otimes X_i^*) & \xleftarrow{\text{id}_Y \otimes \psi_{X_i} \otimes \text{id}_{X_i^*}} & Y \otimes (X_i \otimes X_i^*) \end{array}$$

and an inverse $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(V), (Y, \gamma)) \rightarrow \text{Hom}_{\mathcal{C}}(V, Y)$ is given by $g \mapsto p_1 \circ g$. The proof is analogous to [16, Theorem 2.3] where a different pairing is used. The right adjunction isomorphisms are given by

$$(4.6) \quad \begin{aligned} \text{Hom}_{\mathcal{C}}(Y, V) &\cong \text{Hom}_{\mathcal{Z}(\mathcal{C})}((Y, \gamma), I(V)) \\ f &\mapsto \sum_i \iota_i \circ \sigma_i(f) \end{aligned}$$

where $\iota_i: X_i \otimes V \otimes X_i^* \rightarrow FI(V)$ are the inclusions and $\sigma_i(f)$ is defined by

$$(4.7) \quad \begin{array}{ccc} Y & \xrightarrow{\text{id}_Y \otimes \text{coev}_{X_i}} & Y \otimes (X_i \otimes X_i^*) & \xrightarrow{a_{Y, X_i, X_i^*}^{-1}} & (Y \otimes X_i) \otimes X_i^* \\ \downarrow \text{:=}\sigma_i(f) & & & & \downarrow \gamma(X_i) \otimes \text{id}_{X_i^*} \\ (X_i \otimes V) \otimes X_i^* & \xleftarrow{\text{id}_{X_i} \otimes f \otimes \text{id}_{X_i^*}} & (X_i \otimes Y) \otimes X_i^* & & \end{array}$$

and an inverse is given by $\text{Hom}_{\mathcal{C}}(Y, V) \rightarrow \text{Hom}_{\mathcal{Z}(\mathcal{C})}((Y, \gamma), I(V)): g \mapsto \iota_1 \circ g$. Again the proof follows the exact same arguments as the left adjoint case.

At the heart of our algorithm lies the following well-known property of the induction functor which is an immediate consequence of the adjunction.

Lemma 4.2. *Every simple object $(Z, \gamma_Z) \in \mathcal{Z}(\mathcal{C})$ arises as a subobject of $I(X_i)$ for some simple object $X_i \in \mathcal{C}$.*

Proof. Since $F((Z, \gamma_Z))$ is a non-zero object of \mathcal{C} , it has a simple quotient X_j and so there is a non-zero morphism $F((Z, \gamma_Z)) \rightarrow X_j$. Hence, using the adjunction, we conclude

$$(4.8) \quad 0 \neq \text{Hom}_{\mathcal{C}}(F((Z, \gamma_Z)), X_j) \simeq \text{Hom}_{\mathcal{Z}(\mathcal{C})}((Z, \gamma_Z), I(X_j)).$$

We conclude there is a non-zero morphism $f: (Z, \gamma_Z) \rightarrow I(X_j)$. Since (Z, γ_Z) is simple, the kernel of f must be zero, so f is a monomorphism and therefore (Z, γ_Z) is a subobject of $I(X_j)$. \square

In other words, the induction functor I is *surjective* as in [9, Definition 1.8.3], see also [9, Corollary 7.13.11]. The lemma yields an obvious idea how to compute a complete set $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ of the isomorphism classes of simple objects of $\mathcal{Z}(\mathcal{C})$. We formalize this in Algorithm 1.

Algorithm 1: Compute $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ – First version

```

1 Initialize  $\text{Irr}(\mathcal{Z}(\mathcal{C})) = \emptyset$ 
2 for  $i = 1$  to  $n$  do
3   Compute the object  $I(X_i)$ 
4   Compute a complete set  $\text{Irr}(I(X_i))$  of isomorphism classes of simple subobjects
   of  $I(X_i)$ 
5   for  $(Z, \gamma) \in \text{Irr}(I(X_i))$  do
6     if  $(Z, \gamma)$  is not isomorphic to any object in  $\text{Irr}(\mathcal{Z}(\mathcal{C}))$  then
7       Add  $(Z, \gamma)$  to  $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ 
8 return  $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ 

```

Using the explicit formulas in Equations (4.1) and (4.3), we were able to implement the induction functor in our software framework, so Step 3 in Algorithm 1 is constructive. The problems are Step 4 (the computation of simple subobjects) and Step 6 (the isomorphism check). Decomposing a semisimple object $X \in \mathcal{C}$ in an abelian category is equivalent to decomposing the endomorphism algebra $\text{End}_{\mathcal{C}}(X)$, i.e. if

$$X = \bigoplus_{i=1}^k n_i \cdot X_i$$

for simple objects X_i then

$$\text{End}_{\mathcal{C}}(X) = \bigoplus_{i=1}^k \text{Mat}_{n_i \times n_i} \text{End}_{\mathcal{C}}(X_i)$$

and $X_i \cong \text{Im}(f_i)$ for any $0 \neq f_i \in \text{End}_{\mathcal{C}}(X_i)$ considered as an element of $\text{End}_{\mathcal{C}}(X_i)$. The decomposition of algebras is in general a very hard task and there is no algorithm known. In our implementation we use the MeatAxe (Las Vegas) algorithm [22, 15, 17], enhanced by various further technical tweaks. Details can be found in the source code of our implementation. In practice, for small examples, this works reasonably well.

In order to decompose endomorphism algebras at all, we need to be able to determine these algebras explicitly. More generally, we need to be able to compute Hom-spaces in the center. We will discuss this in Section 4.3. First, we will discuss a simple optimization of Algorithm 1.

4.2. Center-generating simple objects. In most examples the computation of $I(X_i)$ for all simple X_i will result in redundant computations: 1) it might happen that $I(X_i) \cong I(X_j)$ for $X_i \not\cong X_j$; 2) there are examples where all simple objects are contained in just a few inductions. This leads us to the following concept.

Definition 4.3. A set of center-generating simples for $\mathcal{Z}(\mathcal{C})$ is a collection of simple objects $S_1, \dots, S_l \in \mathcal{C}$ such that every simple object $Z \in \mathcal{Z}(\mathcal{C})$ is a subquotient of at least one $I(S_i)$.

Now, the goal is to (algorithmically) find a small set of center-generating simples. Here is one approach. Recall that an object $X \in \mathcal{C}$ is called *invertible* if $X \otimes X^* \cong \mathbb{1}$. This immediately implies that X is simple. We will write $\text{Inv}(\mathcal{C}) \subseteq \text{Irr}(\mathcal{C})$ for the invertible objects in \mathcal{C} .

Lemma 4.4. Let $V, W \in \mathcal{C}$ such that $W \cong J \otimes V \otimes J^*$ for some $J \in \text{Inv}(\mathcal{C})$. Then $I(V) \cong I(W)$.

Proof. We can assume that \mathcal{C} is strict. Since J is invertible, the object $X_i \otimes J$ is simple and $- \otimes J$ is an auto-equivalence of \mathcal{C} . The half-braidings on the underlying object of $I(W) \cong I(J \otimes V \otimes J^*)$ are given by the components

$$\gamma(Z)_{ij} = \sum_{f \in B, g \in B'} f \otimes \text{id}_J \otimes \text{id}_V \otimes \text{id}_{J^*} \otimes g.$$

The $f \otimes \text{id}_J$ and $\text{id}_{J^*} \otimes g$ form again a pair of dual basis. Hence, $I(V)$ and $I(J \otimes V \otimes J^*)$ carry the same half-braiding up to reordering of the summands. \square

The action of $\text{Inv}(\mathcal{C})$ defines an equivalence relation on $\text{Irr}(\mathcal{C})$. So, by Lemma 4.4 the representatives for the equivalence classes form a center-generating set. This set does not need to be minimal but it is one that we can algorithmically compute.

Using the concept of center-generating simples we give in Algorithm 2 a refined version of Algorithm 1. We restricted the computations to a set of center-generating simples and replaced the condition for whether a simple object is accepted as new. The latter works since for every simple object $X_i \in \text{Irr}(\mathcal{C})$ and simple object $Z \in \text{Irr}(\mathcal{Z}(\mathcal{C}))$ we have

$$\text{Hom}_{\mathcal{C}}(X_i, Z) \cong \text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(X_i), Z)$$

and therefore whenever $\text{Hom}_{\mathcal{C}}(X_i, Z) \neq 0$ for any X_i processed before, then Z is already in the list of simple objects of $\mathcal{Z}(\mathcal{C})$.

Algorithm 2: Compute $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ – Refined version

```

1 Initialize  $\text{Irr}(\mathcal{Z}(\mathcal{C})) = \emptyset$ 
2 Compute a set  $S = \{S_1, \dots, S_l\} \subseteq \text{Irr}(\mathcal{C})$  of center-generating simples, e.g., via
   computing the conjugacy classes on  $\text{Irr}(\mathcal{C})$ 
3 for  $i \in \{1, \dots, l\}$  do
4   | Compute  $I(S_i)$ 
5   | Compute all non-isomorphic simple subobjects  $Z_1, \dots, Z_k$  of  $I(S_i)$ 
6   | for  $j \in \{1, \dots, k\}$  do
7   |   | if  $\text{Hom}_{\mathcal{C}}(Z_j, S_t) = 0$  for all  $t \in \{1, \dots, i-1\}$  then
8   |   |   | Add  $Z_j$  to  $\text{Irr}(\mathcal{C})$ 
9 return  $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ 

```

4.3. Computing Hom-spaces in $\mathcal{Z}(\mathcal{C})$. There are multiple ways to obtain morphisms in the center of a fusion category \mathcal{C} . The first and most straight forward approach is to solve for the condition 1.4. If $f \in \text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y) \subset \text{Hom}_{\mathcal{C}}(F(X), F(Y))$ we can write

$$f = a_1 g_1 + \dots + a_k g_k$$

where g_1, \dots, g_k is a basis of $\text{Hom}_{\mathcal{C}}(F(X), F(Y))$. Condition 1.4 yields now a set of equations

$$\gamma_Y(Z) \circ (f \otimes \text{id}_Z) = (\text{id}_Z \otimes f) \circ \gamma_X(Z)$$

for each simple object $Z \in \mathcal{C}$ which are linear in the a_i . By solving this system we obtain a basis of the space $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y)$. This approach scales badly since k grows quickly with the number of direct summands in $F(X)$ and $F(Y)$, and the setup of the system is expensive.

A much better approach is to take advantage of the adjunction isomorphisms 4.4 and 4.6. With those we immediately obtain endomorphism spaces of all objects of the form $I(X)$ since

$$(4.9) \quad \text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(X), I(X)) \cong \text{Hom}_{\mathcal{C}}(X, FI(X))$$

and with some effort also arbitrary Hom-spaces as follows. Let $X, Y \in \mathcal{Z}(\mathcal{C})$ and denote by Z_1, \dots, Z_r the simple objects of $\mathcal{Z}(\mathcal{C})$. Since $\mathcal{Z}(\mathcal{C})$ is semisimple we can express any morphism $f: X \rightarrow Y$ as a sum of morphisms $X \rightarrow Z_i \rightarrow Y$. If $\{S_1, \dots, S_l\}$ is a set

of center-generating simples, then for all Z_i there is some j such that there is a non-zero morphism $Z_i \hookrightarrow I(S_j)$. Then $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y)$ is spanned by morphisms of the form $X \rightarrow I(S_j) \rightarrow Y$. We summarize this approach in Algorithm 3.

Algorithm 3: Computing Hom-spaces in $\mathcal{Z}(\mathcal{C})$

Input: Objects $X, Y \in \mathcal{Z}(\mathcal{C})$. A set $\{S_1, \dots, S_l\}$ of center-generating simples

Output: A basis of $\text{Hom}_{\mathcal{Z}(\mathcal{C})}((X, \gamma_X), (Y, \gamma_Y))$

- 1 Compute a basis of the spaces $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(S_j), Y)$ and $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, I(S_j))$ for all $j = 1, \dots, l$ via the adjunction (4.9)
 - 2 Collect the set of maps $M_j = \{p \circ i \in \text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y) \mid p \in \text{Hom}_{\mathcal{Z}(\mathcal{C})}(I(S_j), Y), i \in \text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, I(S_j))\}$ for all $j = 1, \dots, l$
 - 3 Use linear algebra to find a basis in the union of the M_j
-

Still another way is to project onto $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y) \subseteq \text{Hom}_{\mathcal{C}}(F(X), F(Y))$ with the formula in the following lemma. In practice, however, algorithm 3 is the most efficient way to get to morphisms.

Lemma 4.5 ([16, Lemma 2.2.]). *For $(X, \gamma_X), (Y, \gamma_Y) \in \mathcal{Z}(\mathcal{C})$ the map $E_{X,Y} : \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(X, Y)$ given by*

$$E_{X,Y}(t) = \frac{1}{\dim \mathcal{C}} \sum_{i=1}^n \dim X_i \phi_i(t),$$

where $\phi_i(t)$ is given by

$$\begin{array}{ccccc}
 X & \xrightarrow{\text{id}_X \otimes \text{coev}(X_i)} & X \otimes (X_i \otimes X_i^*) & \xrightarrow{a_{X, X_i, X_i^*}^{-1}} & (X \otimes X_i) \otimes X_i^* \\
 \downarrow \phi_i(t) & & & & \downarrow \gamma_X(X_i) \otimes \text{id}_{X_i^*} \\
 & & & & (X_i \otimes X) \otimes X_i^* \\
 & & & & \downarrow \text{id}_{X_i} \otimes t \otimes \text{id}_{X_i^*} \\
 & & & & (X_i \otimes Y) \otimes X_i^* \\
 & & & & \downarrow a_{X_i, Y, X_i^*} \\
 & & & & X_i \otimes (Y \otimes X_i^*) \\
 & & & & \downarrow \psi_{X_i} \otimes \gamma_Y(X_i^*) \\
 Y & \xleftarrow{\text{ev}_{X_i^*} \otimes \text{id}_Y} & (X_i^{**} \otimes X_i^*) \otimes Y & \xleftarrow{a_{X_i^{**}, X_i^*, Y}^{-1}} & X_i^{**} \otimes (X_i^* \otimes Y)
 \end{array}$$

is a projection from $\text{Hom}_{\mathcal{C}}(X, Y)$ onto $\text{Hom}_{\mathcal{Z}(\mathcal{C})}((X, \gamma_X), (Y, \gamma_Y))$.

5. SPLITTING OF SIMPLE CENTRAL OBJECTS

In this section we want to take a closer look at some properties of the center when the ground field \mathbb{k} is not algebraically closed.

5.1. Scalar extension. For the following concepts we also refer to [8] and [19]. Let \mathcal{C} be a \mathbb{k} -linear semisimple abelian category and $\mathbb{k} \hookrightarrow \mathbb{K}$ be a field extension. We define the category $\mathbb{K} \otimes \mathcal{C}$ to have the same objects as \mathcal{C} and morphism spaces are given by

$$(5.1) \quad \mathrm{Hom}_{\mathbb{K} \otimes \mathcal{C}}(X, Y) = \mathbb{K} \otimes_{\mathbb{k}} \mathrm{Hom}_{\mathcal{C}}(X, Y) .$$

The category $\mathbb{K} \otimes \mathcal{C}$ is again \mathbb{k} -linear. However, it may not be abelian if \mathcal{C} does not split since the scalar extension of an endomorphism algebra might gain new idempotents which are not split (i.e. the object they would project to does not exist). Consider the example $\mathcal{C} = \mathrm{Rep}_{\mathbb{Q}}(C_3)$ of rational representations of the cyclic group of order 3. There is a 2-dimensional representation with endomorphism ring isomorphic to $\mathbb{K} = \mathbb{Q}(\xi_3)$, where ξ_3 is a third root of unity. Tensoring with $\mathbb{Q}(\xi_3)$ splits the algebra and hence produces projectors onto the completely irreducible subrepresentations. Formally, these objects are not yet contained in the category $\mathbb{K} \otimes \mathcal{C}$.

Hence, we want to consider the *Karoubian closure* $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$ of $\mathbb{K} \otimes \mathcal{C}$, which formally adds the corresponding objects for all idempotents. We refer to [7, Chapter 11] for basics on this construction. The category $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$ is a semisimple \mathbb{K} -linear abelian category. We call it the *scalar extension* of \mathcal{C} to \mathbb{K} . If \mathcal{C} has a monoidal structure, then so does $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$. If \mathcal{C} is a weak fusion category, then so is $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$. We say \mathbb{K} is a *splitting field* for \mathcal{C} if $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$ is split.

Notice that, by construction, every simple object of $\mathrm{Kar}(\mathbb{K} \otimes \mathcal{C})$ is a direct summand of a scalar extension $\mathbb{K} \otimes X$ of a simple object $X \in \mathcal{C}$. Moreover, if X and Y are non-isomorphic simple objects, then no direct summand of $\mathbb{K} \otimes X$ is isomorphic to a direct summand of $\mathbb{K} \otimes Y$ since otherwise there would be a non-zero morphism $\mathbb{K} \otimes X \rightarrow \mathbb{K} \otimes Y$, given by projection and inclusion, but this is not possible since

$$\mathrm{Hom}_{\mathbb{K} \otimes \mathcal{C}}(\mathbb{K} \otimes X, \mathbb{K} \otimes Y) = \mathbb{K} \otimes \mathrm{Hom}_{\mathcal{C}}(X, Y) = 0 .$$

5.2. Frobenius–Perron dimension. From now on, let \mathcal{C} be a weak fusion category. The Frobenius–Perron dimension of an object $X \in \mathcal{C}$ is defined to be the largest real eigenvalue of the matrix of the action of the isomorphism class of X on the Grothendieck ring $\mathrm{Gr}(\mathcal{C})$ of \mathcal{C} by multiplication, see [9, Section 4.5]. In case \mathbb{k} is algebraically closed, the Frobenius–Perron dimension of the category \mathcal{C} itself is defined in [9, Definition 6.1.7] as

$$(5.2) \quad \mathrm{FPdim}(\mathcal{C}) := \sum_{i=1}^n \mathrm{FPdim}(X_i)^2 .$$

In [23] this definition was extended to a general base field \mathbb{k} via

$$(5.3) \quad \text{FPdim}(\mathcal{C}) := \sum_{i=1}^n \frac{\text{FPdim}(X_i)^2}{\dim_{\text{End}_{\mathcal{C}}(\mathbb{1})} \text{End}_{\mathcal{C}}(X_i)} = \frac{\text{FPdim}(X_i)^2}{\dim_{\mathbb{k}} \text{End}_{\mathcal{C}}(X_i)},$$

where we use our assumption that $\mathbb{1}$ is split simple. In analogy with Equation (2.6) it is proven in [23, Theorem 4.1.8] that

$$(5.4) \quad \text{FPdim}(\mathcal{Z}(\mathcal{C})) = \text{FPdim}(\mathcal{C})^2.$$

This formula implies nice behavior of the simple central objects under extension to a splitting field.

Proposition 5.1. *Let \mathcal{C} be a fusion category. Let $\bar{\mathbb{k}}$ be an algebraic closure of \mathbb{k} and let $\overline{\mathcal{Z}(\mathcal{C})}$ be the Karoubian closure of $\bar{\mathbb{k}} \otimes \mathcal{Z}(\mathcal{C})$. If $Z \in \mathcal{Z}(\mathcal{C})$ is a simple object, then all direct summands of $\bar{\mathbb{k}} \otimes Z$ in $\overline{\mathcal{Z}(\mathcal{C})}$ occur with the same multiplicity and all have the same Frobenius–Perron dimension.*

Proof. Let Z_1, \dots, Z_r be the non-isomorphic simple objects of $\mathcal{Z}(\mathcal{C})$. Let

$$(5.5) \quad \bar{\mathbb{k}} \otimes Z_i = a_{i,1} Z_{i,1} \oplus \dots \oplus a_{i,n_i} Z_{i,n_i}$$

be the decomposition of $\bar{\mathbb{k}} \otimes Z_i$ in $\overline{\mathcal{Z}(\mathcal{C})}$ into pairwise non-isomorphic simple objects $Z_{i,j}$. By our remarks above, the $Z_{i,j}$ are pairwise non-isomorphic and yield all the simple objects of $\overline{\mathcal{Z}(\mathcal{C})}$. For any i the Cauchy–Schwarz inequality yields

$$(5.6) \quad \left(\sum_{j=1}^{n_i} a_{i,j} \cdot \text{FPdim}(Z_{i,j}) \right)^2 \leq \left(\sum_{j=1}^{n_i} \text{FPdim}(Z_{i,j})^2 \right) \left(\sum_{j=1}^{n_i} a_{i,j}^2 \right).$$

This is equivalent to

$$(5.7) \quad \frac{\left(\sum_{j=1}^{n_i} a_{i,j} \cdot \text{FPdim}(Z_{i,j}) \right)^2}{\sum_{j=1}^{n_i} a_{i,j}^2} \leq \sum_{j=1}^{n_i} \text{FPdim}(Z_{i,j})^2.$$

Notice that

$$(5.8) \quad \sum_{j=1}^{n_i} a_{i,j}^2 = \dim_{\bar{\mathbb{k}}} \text{End}_{\overline{\mathcal{Z}(\mathcal{C})}}(\bar{\mathbb{k}} \otimes Z_i) = \dim_{\mathbb{k}} \text{End}_{\mathcal{Z}(\mathcal{C})}(Z_i).$$

Moreover, since the Frobenius–Perron dimension is a ring morphism by [9, Proposition 3.3.6], we have

$$(5.9) \quad \text{FPdim}(\bar{\mathbb{k}} \otimes Z_i) = \sum_{j=1}^{n_i} a_{i,j} \text{FPdim}(Z_{i,j}).$$

Hence, $\text{FPdim}(\bar{\mathbb{k}} \otimes -)$ can be considered as a ring morphism on the Grothendieck ring of $\mathcal{Z}(\mathcal{C})$ taking positive values on the isomorphism classes of simple objects. Since this property uniquely characterizes the Frobenius–Perron dimension by [9, Proposition 3.3.6], it follows that

$$(5.10) \quad \text{FPdim}(Z_i) = \text{FPdim}(\bar{\mathbb{k}} \otimes Z_i) .$$

Now,

$$(5.11) \quad \text{FPdim}(\overline{\mathcal{Z}(\mathcal{C})}) = \sum_{i=1}^r \sum_{j=1}^{n_i} \text{FPdim}(Z_{i,j})^2 \geq \sum_{i=1}^r \frac{\left(\sum_{j=1}^{n_i} a_{i,j} \cdot \text{FPdim}(Z_{i,j}) \right)^2}{\sum_{j=1}^{n_i} a_{i,j}^2}$$

$$(5.12) \quad = \sum_{i=1}^r \frac{\text{FPdim}(Z_i)^2}{\dim_{\mathbb{k}} \text{End}_{\mathcal{Z}(\mathcal{C})}(Z_i)} = \text{FPdim}(\mathcal{Z}(\mathcal{C})) .$$

Since \mathcal{C} is split, it follows that $\bar{\mathbb{k}} \otimes \mathcal{C}$ is semisimple, hence equal to its Karoubian closure, so \mathcal{C} is a “rational form” of $\bar{\mathbb{k}} \otimes \mathcal{C}$ in the terminology of Theorem [19, Section 2]. It thus follows from [19, Lemma 5.1] that $\mathcal{Z}(\mathcal{C})$ is a rational form of $\mathcal{Z}(\bar{\mathbb{k}} \otimes \mathcal{C})$, i.e.,

$$(5.13) \quad \mathcal{Z}(\bar{\mathbb{k}} \otimes \mathcal{C}) = \overline{\mathcal{Z}(\mathcal{C})} .$$

Using Equation (5.4), we thus get

$$(5.14) \quad \text{FPdim}(\mathcal{Z}(\mathcal{C})) = \text{FPdim}(\mathcal{C})^2 = \text{FPdim}(\bar{\mathbb{k}} \otimes \mathcal{C})^2 = \text{FPdim}(\mathcal{Z}(\bar{\mathbb{k}} \otimes \mathcal{C})) = \text{FPdim}(\overline{\mathcal{Z}(\mathcal{C})}) .$$

We thus have equality in the Cauchy–Schwarz inequality (5.6). This implies that there is some $\lambda \in \mathbb{R}$ such that

$$(5.15) \quad \text{FPdim}(Z_{i,j}) = \lambda a_{i,j}$$

for all i, j . This holds if and only if

$$(5.16) \quad \frac{a_{i,j}}{\text{FPdim}(Z_{i,j})} = \lambda = \frac{a_{i',j'}}{\text{FPdim}(Z_{i',j'})}$$

for all i, j, i', j' .

Now, fix some i . Consider $A = \text{End}_{\mathcal{Z}(\mathcal{C})} Z_i$. By assumption, A is a simple division algebra. Consider the extension $Z(A) \otimes_{\mathbb{k}} A$ which is as $Z(A)$ -algebra isomorphic to $\bigoplus_{j=1}^{[Z(A):\mathbb{k}]} A$. Finally, A is central simple as $Z(A)$ -algebra and hence has a splitting field $\mathbb{K} \supseteq \mathbb{k}$ such that $A \cong \text{Mat}_{l \times l}(\mathbb{k})$. We conclude that $\mathbb{k} \otimes A \cong \bigoplus_{j=1}^{[Z(A):\mathbb{k}]} \text{Mat}_{l \times l}(\mathbb{k})$ as \mathbb{k} -algebras. This implies that $\mathbb{k} \otimes Z$ decomposes in $\text{Kar}(\mathbb{k} \otimes \mathcal{Z}(\mathcal{C}))$ into $[Z(A) : \mathbb{k}]$ non-isomorphic simple objects, each with multiplicity l . This forces $a_{i,j} = l$ for all j , and thus, by Equation (5.16), all $Z_{i,j}$ have the same Frobenius–Perron dimension. \square

Remark 5.2. If \mathbb{k} is a finite field the statement above is even stronger since all finite dimensional division algebras over a finite field are fields. Thus, a simple object Z decomposes into non-isomorphic simple objects with multiplicity one.

Example 5.3. The case that a simple object occurs with multiplicity greater than one from Lemma 5.1 can occur. Let \mathbf{Q} be the quaternion group and consider the category $\text{Vec}_{\mathbb{Q}}(\mathbf{Q})$ of finite-dimensional \mathbf{Q} -graded vector spaces over \mathbb{Q} . Then there is a central object lying over $4 \cdot \mathbf{1}_{\mathbf{Q}}$ corresponding to the four-dimensional irreducible \mathbb{Q} -representation of \mathbf{Q} . This object has (similar to the representation) an endomorphism ring isomorphic to the rational quaternions $\mathbb{H}_{\mathbb{Q}}$. Analogously to the representation it decomposes into two copies of the same simple object over $\mathbb{Q}(\sqrt{-1})$.

6. SOFTWARE FRAMEWORK AND EXAMPLES

Our software [12] is completely open source and comes with a documentation. Its general purpose is to serve as a framework for experiments with tensor categories and categorical representation theory. Instead of going into long discussions about its design and capabilities, we illustrate it by working with the two examples mentioned in the introduction: the Ising fusion category and a fusion category coming from the Haagerup subfactor. For users new to Julia we note Julia uses just-in-time compilation (JIT). This is, in parts, what makes Julia so fast but it means that the *first* execution of a function always takes a bit of time (since its code will be compiled)—afterwards it is instantaneous.

6.1. The Ising Category over $\mathbb{Q}(\sqrt{2})$. Users are completely free in how exactly they want to model a category. For encoding a fusion category via 6j-symbols we provide an own convenient structure called `SixJCategory`. The following code shows how to define the Ising fusion category over a general field K . An element $a \in K$ with $a^2 = 2$ needs to be provided to define the associators (see the introduction).

julia

```
function Ising(F::Ring, a::RingElem)
    C = SixJCategory(F, ["1", "χ", "X"])

    # Multiplication table of the Grothendieck ring
    M = zeros{Int,3,3,3}
    M[1,1,:] = [1,0,0]
    M[1,2,:] = [0,1,0]
    M[1,3,:] = [0,0,1]
    M[2,1,:] = [0,1,0]
```

```

M[2,2,:] = [1,0,0]
M[2,3,:] = [0,0,1]
M[3,1,:] = [0,0,1]
M[3,2,:] = [0,0,1]
M[3,3,:] = [1,1,0]

set_tensor_product!(C,M)

# The associators
set_associator!(C,2,3,2, matrices(-id(C[3])))
set_associator!(C,3,2,3, matrices((id(C[1])⊕(-id(C[2])))
z = zero(matrix_space(F,0,0))
set_associator!(C,3,3,3, [z, z, inv(a)*matrix(F,[1 1; 1 -1])])

set_one!(C,[1,0,0])

set_spherical!(C, [F(1) for s in simples(C)])

set_name!(C, "Ising fusion category")
return C
end

```

We can choose as K the algebraic closure of \mathbb{Q} , which is also available in OSCAR via `abelian_closure(QQ)`. But we can also work over $K = \mathbb{Q}(\sqrt{2})$, which is what we will do.

julia

```

julia> K,r2 = quadratic_field(2)
(Real quadratic field defined by x^2 - 2, sqrt(2))

julia> I = Ising(K,r2)
Ising fusion category

julia> a,b,c = simples(I)
3-element Vector{SixJObject}:
 1
 ̑
 ̒

```


Let us now come to the proof Theorem 1.1. We compute the simple objects of the center by invoking the algorithm in Section 4.

```

julia
julia> C = center(I)
Drinfeld center of Fusion Category with 3 simple objects

julia> simples(C)
5-element Vector{CenterObject}:
Central object:  $\mathbb{1}$ 
Central object:  $\mathbb{1}$ 
Central object:  $\mathbb{1} \oplus \chi$ 
Central object:  $2 \cdot \chi$ 
Central object:  $4 \cdot X$ 

```

We can see that there are five non-isomorphic simple objects. We show that two of them are not split over K and examine over which fields they will split. To do so we examine the endomorphism spaces. The object over $2 \cdot \chi$ will split if there is an endomorphism that is a zero-divisor, i.e. if there is a morphism with a non-trivial eigenvalue. Thus, we take a non-trivial endomorphism and consider the splitting field for its minimal polynomial.

```

julia
julia> H = End(C[4])
Vector space of dimension 2 over Real quadratic field defined by  $x^2 - 2$ .

julia> minpoly.(basis(H))
2-element Vector{AbstractAlgebra.Generic.Poly{nf_elem}}:
 $x^2 + 1$ 
 $x - 1$ 

```

So if we extend the field of definition to the splitting field of $x^2 + 1$, i.e. $\mathbb{Q}(\sqrt{2}, i)$, it will split. Similarly, one can show that the fifth simple object decomposes under this extension.

```

julia
julia> Kx,x = base_ring(I)[:x]
(Univariate polynomial ring in x over real quadratic field defined by  $x^2 - 2$ , x)

```

```

julia> L,i = number_field(x^2+1, "i")
(Relative number field of degree 2 over real quadratic field defined by x^2 - 2, i)

julia> C2 = C ⊗ L
Drinfeld center of Fusion Category with 3 simple objects

julia> simples(C2)
7-element Vector{CenterObject}:
Central object: 1
Central object: 1
Central object: 1 ⊕ χ
Central object: χ
Central object: χ
Central object: 2·X
Central object: 2·X

```

Repeat the splitting process one more time for the 6-th or 7-th simple.

```

julia
julia> _,f = minpoly.(basis(End(C2[6])))
2-element Vector{AbstractAlgebra.Generic.Poly{Hecke.NfRelElem{nf_elem}}}:
x - 1
x^2 + 1//4*sqrt(2)*i - 1//4*sqrt(2)

julia> M,a = number_field(f,"a")
(Relative number field of degree 2 over relative number field, a)

julia> simples(C2 ⊗ M)
9-element Vector{CenterObject}:
Central object: 1
Central object: 1
Central object: 1 ⊕ χ
Central object: χ
Central object: χ
Central object: X
Central object: X
Central object: X

```

Central object: X

```
julia> simplify(absolute_simple_field(M)[1])[1]
Number field with defining polynomial x^8 + 1
over rational field
```

Thus, we see that the center of the Ising category splits over the cyclotomic field $\mathbb{Q}(\xi_{16})$ with ξ_{16} a primitive 16-th root of unity. Next, we compute the multiplication table and S -matrix of the center over K .

julia

```
julia> print_multiplication_table(C)
5×5 Matrix{String}:
"X2" "X1" "X3"          "X4"          "X5"
"X1" "X2" "X3"          "X4"          "X5"
"X3" "X3" "X1 ⊕ X2 ⊕ X4" "2·X3"        "2·X5"
"X4" "X4" "2·X3"          "2·X1 ⊕ 2·X2" "2·X5"
"X5" "X5" "2·X5"          "2·X5"        "4·X1 ⊕ 4·X2 ⊕ 8·X3 ⊕ 4·X4"

julia> smatrix(C)
[ 1  1  2  2 -4*√2]
[ 1  1  2  2  4*√2]
[ 2  2  0 -4  0]
[ 2  2 -4  4  0]
[-4*√2 4*√2 0 0 0]
```

6.2. A fusion category from the Haagerup subfactor. Here, we consider the fusion category from the Haagerup subfactor as explained in the introduction. We define the category as a skeletal category with six simple objects over the smallest number field containing the $6j$ -symbols as found in [29] and compute the center accordingly.

julia

```
julia> H = HaagerupH3()
Fusion category from Haagerup H3 subfactor

julia> base_ring(H)
```

```

Number field with defining polynomial x^16 - 4*x^14 + 13*x^12 + 4*x^10 + 53*x^8 + 4*x^6
+ 13*x^4 - 4*x^2 + 1
over rational field

julia> ZH = center(H)
Drinfeld center of Fusion category from Haagerup H3 subfactor

julia> simples(ZH)
6-element Vector{CenterObject}:
Central object:  $2 \cdot \mathbb{1} \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$ 
Central object:  $\mathbb{1} \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$ 
Central object:  $\mathbb{1}$ 
Central object:  $2 \cdot \alpha \oplus 2 \cdot \alpha^* \oplus 2 \cdot \rho \oplus 2 \cdot \alpha\rho \oplus 2 \cdot \alpha^*\rho$ 
Central object:  $\alpha \oplus \alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$ 
Central object:  $6 \cdot \rho \oplus 6 \cdot \alpha\rho \oplus 6 \cdot \alpha^*\rho$ 

```

As with the Ising category the center does not split and we have to repeat the process from above. We can automatize this process with the function `split` from `TensorCategories.jl`.

```

julia
julia> dim.(End.(simples(ZH)))
6-element Vector{Int64}:
 1
 1
 1
 2
 1
 6

julia> split(ZH[4])
2-element Vector{CenterObject}:
Central object:  $\alpha \oplus \alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$ 
Central object:  $\alpha \oplus \alpha^* \oplus \rho \oplus \alpha\rho \oplus \alpha^*\rho$ 

julia> split(ZH[6])
6-element Vector{CenterObject}:
Central object:  $\rho \oplus \alpha\rho \oplus \alpha^*\rho$ 
Central object:  $\rho \oplus \alpha\rho \oplus \alpha^*\rho$ 

```

Central object: $\rho \oplus \alpha\rho \oplus \alpha*\rho$

Central object: $\rho \oplus \alpha\rho \oplus \alpha*\rho$

Central object: $\rho \oplus \alpha\rho \oplus \alpha*\rho$

Central object: $\rho \oplus \alpha\rho \oplus \alpha*\rho$

REFERENCES

- [1] Bojko Bakalov and Alexander Kirillov Jr. *Lectures on tensor categories and modular functors*. Vol. 21. University Lecture Series. American Mathematical Society, Providence, RI, 2001, pp. x+221. DOI: 10.1090/ulect/021.
- [2] Jérémy Berthomieu, Christian Eder, and Mohab Safey El Din. “msolve: A Library for Solving Polynomial Systems.” In: *2021 International Symposium on Symbolic and Algebraic Computation*. 46th International Symposium on Symbolic and Algebraic Computation. Saint Petersburg, Russia: ACM, July 2021, pp. 51–58. DOI: 10.1145/3452143.3465545.
- [3] Jeff Bezanson, Alan Edelman, Stefan Karpinski, and Viral B Shah. “Julia: A fresh approach to numerical computing.” In: *SIAM review* 59.1 (2017), pp. 65–98. DOI: 10.1137/141000671.
- [4] Alain Bruguières and Alexis Virelizier. “On the center of fusion categories.” In: *Pacific Journal of Mathematics* 264.1 (2013), pp. 1–30.
- [5] Wolfram Decker, Christian Eder, Claus Fieker, Max Horn, and Michael Joswig, eds. *The Computer Algebra System OSCAR: Algorithms and Examples*. 1st ed. Vol. 32. Algorithms and Computation in Mathematics. Springer, Aug. 2024. URL: <https://link.springer.com/book/9783031621260>.
- [6] Wolfram Decker, Gert-Martin Greuel, Gerhard Pfister, and Hans Schönemann. *SINGULAR 4-3-0 — A computer algebra system for polynomial computations*. <http://www.singular.uni-kl.de>. 2022.
- [7] Ben Elias, Shotaro Makisumi, Ulrich Thiel, and Geordie Williamson. *Introduction to Soergel bimodules*. Vol. 5. RSME Springer Series. Springer, Cham, 2020, pp. xxv+588. DOI: 10.1007/978-3-030-48826-0.
- [8] Pavel Etingof and Shlomo Gelaki. “Descent and forms of tensor categories.” In: *Int. Math. Res. Not. IMRN* 13 (2012), pp. 3040–3063. DOI: 10.1093/imrn/rnr119.
- [9] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*. Vol. 205. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015, pp. xvi+343. DOI: 10.1090/surv/205.
- [10] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik. “On fusion categories.” In: *Ann. of Math. (2)* 162.2 (2005), pp. 581–642. DOI: 10.4007/annals.2005.162.581.
- [11] David E. Evans and Paulo R. Pinto. “Modular invariants and the double of the Haagerup subfactor.” In: *Advances in operator algebras and mathematical physics*. Vol. 5. Theta Ser. Adv. Math. Theta, Bucharest, 2005, pp. 67–88.
- [12] Fabian Mäurer. *TensorCategories.jl*. Version 0.2.7. URL: <https://fabianmaurer.github.io/TensorCategories.jl/stable/>.
- [13] *GAP – Groups, Algorithms, and Programming, Version 4.13.0*. The GAP Group. 2024. URL: <https://www.gap-system.org>.
- [14] Shlomo Gelaki, Deepak Naidu, and Dmitri Nikshych. “Centers of graded fusion categories.” In: *Algebra Number Theory* 3.8 (2009), pp. 959–990. DOI: 10.2140/ant.2009.3.959.

- [15] Derek F. Holt and Sarah Rees. “Testing modules for irreducibility.” In: *J. Austral. Math. Soc. Ser. A* 57.1 (1994), pp. 1–16.
- [16] Alexander Kirillov Jr and Benjamin Balsam. “Turaev-Viro invariants as an extended TQFT.” In: *arXiv:1004.1533* (2010).
- [17] Klaus Lux and Herbert Pahlings. *Representations of groups*. Vol. 124. Cambridge Studies in Advanced Mathematics. A computational approach. Cambridge University Press, Cambridge, 2010, pp. x+460. DOI: 10.1017/CBO9780511750915.
- [18] Saunders MacLane. *Categories for the working mathematician*. Vol. 5. Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1971, pp. ix+262.
- [19] Scott Morrison and Noah Snyder. “Non-cyclotomic fusion categories.” In: *Trans. Amer. Math. Soc.* 364.9 (2012), pp. 4713–4733.
- [20] Michael Müger. “From subfactors to categories and topology II: The quantum double of tensor categories and subfactors.” In: *Journal of Pure and Applied Algebra* 180.1 (2003), pp. 159–219. DOI: [https://doi.org/10.1016/S0022-4049\(02\)00248-7](https://doi.org/10.1016/S0022-4049(02)00248-7).
- [21] *OSCAR – Open Source Computer Algebra Research system, Version 1.1.0-DEV*. The OSCAR Team, 2024. URL: <https://www.oscar-system.org>.
- [22] R. A. Parker. “The computer calculation of modular characters (the meat-axe).” In: *Computational group theory (Durham, 1982)*. Academic Press, London, 1984, pp. 267–274.
- [23] Sean C. Sanford. “Fusion Categories over Non-algebraically Closed Fields.” PhD thesis. 2022. URL: <https://www.proquest.com/dissertations-theses/fusion-categories-over-non-algebraically-closed/docview/2715420938/se-2>.
- [24] Kenichi Shimizu. “On unimodular finite tensor categories.” In: *Int. Math. Res. Not. IMRN* 1 (2017), pp. 277–322. DOI: 10.1093/imrn/rnv394.
- [25] Daisuke Tambara and Shigeru Yamagami. “Tensor Categories with Fusion Rules of Self-Duality for Finite Abelian Groups.” In: *Journal of Algebra* 209.2 (1998), pp. 692–707. DOI: <https://doi.org/10.1006/jabr.1998.7558>.
- [26] *The Julia Programming Language*. URL: <https://julialang.org>.
- [27] Daniel Tubbenhauer. “Quantum topology without topology.” Draft (v0.99 from May 23, 2022). 2022.
- [28] Robijn Vanhove, Laurens Lootens, Maarten Van Damme, Ramona Wolf, Tobias J. Osborne, Jutho Haegeman, and Frank Verstraete. “Critical Lattice Model for a Haagerup Conformal Field Theory.” In: *Phys. Rev. Lett.* 128 (23 June 2022), p. 231602. DOI: 10.1103/PhysRevLett.128.231602.
- [29] Ramona Wolf. “Microscopic models for fusion categories.” In: *arXiv:2101.04154* (2021).