

Application of Haldane's statistical correlation theory in classical systems

Projesh Kumar Roy^{1,*}

¹*Department of Chemistry, National Institute of Technology Tiruchirappalli,
Tanjore Main Road, NH83, Tiruchirappalli, Tamil Nadu-620015, India*

(Dated: June 25, 2024)

This letter investigates the application of Haldane's statistical correlation theory in classical systems. A modified statistical correlation theory has been proposed by including non-linearity in the form of an exponent into the original theory of Haldane. The dependence of the statistical correlation on indistinguishability is highlighted. Using this modified theory, a quasi-classical derivation of intermediate statistics is shown where indistinguishability can be introduced into distinguishable systems in the form of a statistical correlation. The final result is equivalent to the classical fractional exclusion statistics (CFES), which was derived earlier using a purely classical route. An extended non-linear correlation model based on power series expansion is also proposed, which can produce various intermediate statistical models.

INTRODUCTION

Since the inception of statistical mechanics, there were several attempts [1–5] to unify all fundamental statistical theories, i.e., Fermi-Dirac (FD) statistics, Bose-Einstein (BE) statistics, and Maxwell-Boltzmann (MB) statistics. The first comprehensive approach for such a unified theory was taken by Haldane [6] for quantum systems. In a nutshell, the postulate of Haldane states that the statistical correlation in quantum systems can be described as a reduction of the total available degenerate states (\tilde{g}_i) at an energy level ($\epsilon_i \in [0, \infty]$) due to the population (n_i) of the building blocks at the same energy level. The nature of the reduction was assumed to be linear, which resulted in the following equation,

$$\tilde{g}_i = g_i - \gamma n_i \quad (1)$$

where g_i is the number of ideal degenerate states at $n_i = 0$, and γ is the correlation factor independent of the system size. Applying this idea in Bosonic systems, Wu [7] was able to derive an equation for *intermediate* statistics which interpolates between Bose-Einstein ($\gamma = 0$) and Fermi-Dirac ($\gamma = 1$) statistics.

Application of Haldane-Wu (HW) statistics can be found in the field of 2D-anyons [8, 9], Laughlin liquids [10, 11], adsorption of polyatomic molecules [12–19], etc. A similar statistical theory was proposed by A. Polychronakos—namely the Polychronakos statistics (AP) [20–23]—which avoids the negative probability problem in HW statistics [20, 21, 24–27]. Various other forms of intermediate statistics also exist in the literature [28–32]. Such models can be termed quantum fractional exclusion statistics (QFES) models as they inherently assume the quantum nature of the systems.

Recently, a theory of intermediate statistics based on classical MB statistics—namely, the classical fractional exclusion statistics (CFES) [33]—was derived using the maximum entropy (MaxEnt) methods, i.e.,

$$n_i = g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - a \gamma \left(\frac{n_i}{g_i} \right)^{(a-1)} \right\} \quad (2)$$

Where a is an integer and $a > 0$. Following the MaxEnt method, three major constraints were used along with the corresponding Lagrange parameters during the derivation: (i) constant energy, (ii) constant particle number, and (iii) a special constrain, which can be written as,

$$C = \sum_i \left(\frac{n_i^a}{g_i^{a-1}} \right) \quad (3)$$

Interestingly, it was possible to recover FD and BE statistics at $a = 2$ using these constraints in a purely classical MB statistics framework. However, the origin of the special constraint is still not clear. A concept of *varying indistinguishability* was proposed in Ref. [33], which theorized that this factor can be regarded as a mathematical form of indistinguishability, and it is essentially a bridge between the MB and BE statistics.

This letter proposes a similar derivation of CFES starting from a quantum point of view. Using an *ab-initio* method, I show that a non-linear form of Haldane's theory can be used to highlight the effect of indistinguishability on statistical correlation, which clarifies the relationship between Eq. 3 and indistinguishability. Using this modified theory, a quasi-classical derivation was used to arrive at CFES-equivalent intermediate statistics. The letter is organized as follows—in section I, I describe the non-linear modifications to Eq. 1. In section II, I describe a self-correlating system following the modified statistical correlation theory. In section III, I describe extended non-linear modifications using power series expansion. Finally, I conclude in section IV with an outlook.

**CONCEPT OF INDISTINGUISHABILITY:
ZERO-MOMENT THEOREM**

In the subsequent letter, the word *particle* will be used to denote an entity whose energy distribution follows the MaxEnt principle. Let's consider that a single species of identical particles is present in our system. As described in the previous section, Haldane's theory of statistical correlations in quantum systems proposes that the reduction of the degeneracy factor (g_i) in a quantum energy level ϵ_i is proportional to the population (n_i) at the same level. In this section, I demonstrate an *ab initio* approach to understanding the concept of indistinguishability through the lens of statistical correlation theory. I consider that the reduction in g_i of energy level ϵ_i is dependent on not just n_i but also on the populations at all available energy levels ϵ_j . Moreover, I consider the possibility that the net contribution to statistical correlation from ϵ_j level to ϵ_i level may depend on the different population distribution functions of the particles ($p(n_{jk})$) at different degenerate states (g_{jk}) of energy level ϵ_j . The overall contribution of the statistical correlation to level ϵ_i is weighted by correlation factors γ_{jk} , which may differ for different energy levels and degenerate states. Lastly, I impose non-linearity to the net correlation contribution from g_{jk} , by raising an exponent m to the population factor n_{jk} . Then, the overall equation takes the shape,

$$\tilde{g}_i = g_i - \sum_j \sum_k \gamma_{jk} n_{jk}^m \quad (4)$$

In this model, all particles are distinguishable since the correlation contributions from each g_{jk} level can be different. In the next section, I will show that m and γ factors become dependent on each other for intermediate statistics.

Next, I decompose this model by imposing indistinguishability conditions. Firstly, I assume γ_{jk} to be a universal constant and independent of n_{jk} . Moreover, I assume that the correlation factor $\gamma_{jk} \equiv \gamma_j$; i.e., it does not depend on the properties of degenerate states. This enables us to assume that the correlation effects exerted by the particles belonging to different g_{jk} are similar. Secondly, I assume that for all practical purposes, the probability of finding a particle at any of the degenerate states k of energy level ϵ_j is equal and independent of k ; i.e., $p(n_{jk}) \equiv n_j/g_j$. Because of the uniform distribution of the particles at different degenerate states, one cannot distinguish between the individual particles present at the same energy level ϵ_j using their correlation contributions γ_{jk} and populations n_{jk} . As a result, they become indistinguishable from each other under these approximations.

Therefore, for such indistinguishable particles, one can state that, the standard deviation of the $p(n_{jk})$ function

concerning k degenerate states is zero. Going one step further, I propose a theorem as described below.

Zero-moment theorem: The m^{th} order moment ($\sigma_m(\epsilon_j) | m \geq 2$) of $p(n_{jk})$ function at different k degenerate states of an energy level ϵ_j is zero, i.e.,

$$\sigma_m(\epsilon_j) = \frac{\sum_k n_{jk}^m}{g_j} - \left(\frac{\sum_k n_{jk}}{g_j} \right)^m = 0 \quad (5)$$

which leads to,

$$\sum_k n_{jk}^m = \frac{n_j^m}{g_j^{m-1}} \quad (6)$$

A corollary of this theorem is that m has to be an integer to satisfy the condition in Eq. 6. We can now fully eliminate the k -summand in Eq. 4 by substituting Eq. 6 to Eq. 4 as,

$$\tilde{g}_i = g_i - \sum_j \gamma_j \left(\frac{n_j^m}{g_j^{m-1}} \right) \quad (7)$$

It is straightforward to show that if one disregards all contributions from the other energy levels except the i^{th} level; i.e., $\gamma_j \equiv \gamma \delta_i$ where δ_i is the Kronecker δ ; Eq. 1 can be recovered from Eq. 7 at the linear correlation case at $m = 1$. Note that, the linear correlation case exerts a maximum correlation effect on g_i .

Eq. 6 represents the mathematical description of the concept of indistinguishability, derived in the light of statistical correlation theory. It clearly describes the origin of the constraint C in Eq. 3 and how it is linked to indistinguishability. The dependency of statistical correlation on indistinguishability is an interesting aspect of this analysis. It provides an alternate path for introducing indistinguishability in a classical distinguishable system, rather than directly changing the microstate counting method. In the following section, I show that the application of Eq. 7 to classical MB statistics directly gives rise to the CFES-equivalent equation.

SELF-CORRELATING CLASSICAL SYSTEM

The non-linear correlation theory can certainly be used to modify the microstate counting formula of BE statistics (W_{BE}) and derive different forms of QFES, as described in reference 7. In this paper, I apply this theory to the classical systems and investigate the changes in classical MB statistics (W_{MB}) due to statistical correlations. Let's assume that the system is self-correlating, i.e., a single species of identical particles is present in the system which are themselves involved in correlation. Let's

denote their un-optimized population with a slightly different notation, ν_i . Using a quasi-classical method, I introduce indistinguishability in this classical system in the form of a statistical correlation. Note that, this derivation is different from reference 33, as the particles are assumed to have quantum nature *a priori*. Using Eq. 7, I can show that W_{MB} takes the form,

$$W_{\text{MB}} = W_{\text{MB}}^{\text{id}} \prod_i \left\{ 1 - \frac{1}{g_i} \sum_j \gamma_j \left(\frac{\nu_j^m}{g_j^{m-1}} \right) \right\}^{\nu_i} \quad (8)$$

where $W_{\text{MB}}^{\text{id}} = g_i^{\nu_i}/\nu_i!$ is the ideal number of the microstates for distinguishable classical systems. At a dilute condition; i.e., $\nu_i/g_i \sim 0$; one can derive the corresponding particle distribution in the following manner. Using MaxEnt method, one can maximize W_{MB} with respect to ν_i to get the optimized population, $\nu_{i,\text{max}} \equiv n_i$, as,

$$\begin{aligned} \frac{\partial \ln W_{\text{MB}}}{\partial \nu_i} &= \frac{\partial \ln W_{\text{MB}}^{\text{id}}}{\partial \nu_i} + \ln \left\{ 1 - \frac{1}{g_i} \sum_j \gamma_j \left(\frac{\nu_j^m}{g_j^{m-1}} \right) \right\} \\ &\quad - \frac{m\gamma_i \nu_i^m / g_i^{m-1}}{\{g_i - \sum_j \gamma_j (\nu_j^m / g_j^{m-1})\}} \\ &= 0 \end{aligned} \quad (9)$$

which leads to,

$$\begin{aligned} n_i &= g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - \frac{1}{g_i} \sum_j \gamma_j \left(\frac{n_j^m}{g_j^{m-1}} \right) \right\} \\ &\quad \exp \left\{ - \frac{m\gamma_i n_i^m / g_i^{m-1}}{g_i - \sum_j \gamma_j (n_j^m / g_j^{m-1})} \right\} \\ &\approx \tilde{g}_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - m\gamma_i \left(\frac{n_i^m}{\tilde{g}_i g_i^{m-1}} \right) \right\}; \text{ using } g_i \gg n_i \\ &= g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ \frac{\tilde{g}_i}{g_i} - m\gamma_i \left(\frac{n_i^m}{g_i^m} \right) \right\} \end{aligned} \quad (10)$$

where $\alpha = \beta\mu$ and $\beta = (1/k_{\text{B}}T)$ are the Lagrange parameters related to the chemical potential (μ) and temperature (T) of the system. One can show, if $\gamma_i \equiv \gamma\delta_i$, Eq. 10 reduces to a CFES-type equation [33] as,

$$n_i = g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - (m+1)\gamma \left(\frac{n_i^m}{g_i^m} \right) \right\} \quad (11)$$

where γ corresponds to the third Lagrange parameter and m is the exponent [34] described in Eq. 2 [33]. Consequently, one can derive the constraint C for the population distribution used in Eq. 2 in a retrospective way as,

$$C = \sum_i \left(\frac{n_i^{m+1}}{g_i^m} \right) \quad (12)$$

For $\gamma > 0$ case (abbr: γ^+), the relationship between γ and m satisfy the equation,

$$n_c = \left\{ \frac{1}{(m+1)\gamma^+} \right\}^{1/m} \quad (13)$$

where n_c is the maximum occupancy numbers at all available degenerate states. An interesting question comes up about the limits of m . In the previous section, I have already shown that m has to be an integer to satisfy the zero-moment theorem. If one only considers Eq. 2, $m = [-1, 0, \infty]$ will produce traditional MB statistics, as proved in reference [33]. However, if one considers Eq. 7, $m = 0$ system become unphysical as $\gamma \rightarrow \infty$ according to Eq. 13, which leads to $\tilde{g}_i \rightarrow -\infty$. Similarly, for $m = -1$, $\tilde{g}_i \rightarrow \infty$ at $n_i \rightarrow 0$. Therefore, if one considers the quasi-classical derivation of CFES under the modified statistical correlation theory, the limit of m is $\infty \geq m \geq 1$. Here, $m \rightarrow \infty$ and $\gamma \rightarrow 0$ system is the only allowed system where MB statistics can be recovered. Thus, Eq. 11 is truly unifying statistical theory that encompasses properties MB ($m = \infty, \gamma = 0$), FD ($m = 1, \gamma^+ = 1$), and BE ($m = 1, \gamma^- = -1$) statistics, as well as their intermediate statistics.

EXTENDED NON-LINEAR MODELS

It might be possible to use functions other than the simple exponent in Eq. 7 to introduce non-linearity in the statistical correlation. However, the elimination of the k -summand in Eq. 4 is necessary to introduce indistinguishability in any non-linear model. The exponent model was perhaps the simplest where such elimination was easy because of the zero-moment theorem. Another way to study other forms of non-linearity, which will also ensure the elimination of the k -summand, is to use a power series expansion of n_{jk} as,

$$\tilde{g}_j = g_j - \sum_j \sum_k \gamma_{jk} \left(\sum_{l=0}^{\infty} b_l n_{jk}^l \right) \quad (14)$$

where b_l 's are arbitrary constants chosen in a way that $\sum_l b_l n_{jk}^l$ is converging and finite valued function. Using zero-moment theorem and $\gamma_{jk} \equiv \gamma_j$, I get,

$$\tilde{g}_j = g_j - \sum_j \gamma_j \left(\sum_{l=0}^{\infty} b_l \frac{n_j^l}{g_j^{l-1}} \right) \quad (15)$$

which removes all k -summands in Eq. 14 and introduce indistinguishability in the model. Such elimination of k -summands is not possible for a similar power series expansion n_{jk} with $l < 0$, hence they are not allowed. Using the similar methods as described in the previous section, one can derive the optimal population distribution (for $\gamma_i \equiv \delta_i \gamma$) as,

$$n_i = g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - \gamma \left(\sum_{l=0}^{\infty} (l+1) b_l \frac{n_i^l}{g_i^l} \right) \right\} \quad (16)$$

The maximum occupancy numbers at each degenerate state will have an upper bound for $\gamma > 0$ as,

$$0 \leq \sum_{l=0}^{\infty} (l+1) b_l m_c^l \leq \frac{1}{\gamma^+} \quad (17)$$

Using different values for the coefficients of b_l , one can derive different statistical models from Eq 16. Clearly, choice of b_l has to be such that $b_l \rightarrow 0$ as $l \rightarrow \infty$ for $\gamma > 0$.

An interesting example is $b_l = 1/(l+1)!$, which leads to,

$$n_i = g_i e^{-(\alpha + \beta \epsilon_i)} \left\{ 1 - \gamma e^{(n_i/g_i)} \right\} \quad (18)$$

As for dilute systems $\exp(n_i/g_i) \sim (1 + n_i/g_i)$, Eq. 18 roughly converges to Eq. 11 with $m = 1$. In this particular case, one can show that, $\gamma^+(1 + n_i/g_i) < 1$, which leads to, $\gamma^+ < 1$.

CONCLUSION AND OUTLOOK

In this work, I have given clear-cut evidence that the special constraint used in the purely classical derivation of the CFES statistics is indeed linked to the indistinguishability of the system. Usually, the linear form of Haldane's theory is used in deriving QFES-type equations. The non-linear modification introduced in this letter helps us to understand the relationship between statistical correlation and indistinguishability using the zero-moment theorem. For a self-correlating system, a quasi-classical derivation using the non-linear statistical correlation theory directly produces CFES-equivalent intermediate statistics. Thus, I have shown that Haldane's statistical correlation theory lies at the core of CFES statistics.

I have shown that one can derive many intermediate statistical models using appropriate choices for the b_l coefficients, which will obey Eq. 17 and produce physically meaningful population distribution. An interesting question can be asked: Is self-correlation necessary

to show quantum effects? Currently, I am exploring a binary inter-correlating (not self-correlating) system, which can answer this question and produce interesting results. The application of CFES statistics in enhanced sampling methods in molecular dynamics simulations of classical systems is also being investigated.

ACKNOWLEDGEMENT

I am thankful to the National Institute of Technology, Tiruchirappalli, for providing the necessary funding for this research and to Prof. Sanjib Senapati for useful discussions.

* projesh@nitt.edu

- [1] G. Gentile j., itosservazioni sopra le statistiche intermedie, *Il Nuovo Cimento* 17 (1940) 493–497. doi:doi:10.1007/BF02960187.
- [2] C. Tsallis, Possible generalization of boltzmann-gibbs statistics, *Journal of Statistical Physics* 52 (1988) 479–487. doi:doi:10.1007/BF01016429.
- [3] N. H. March, N. Gidopoulos, A. K. Theophilou, M. J. Lea, W. Sung, The statistical distribution function for an anyon liquid, *Physics and Chemistry of Liquids* 26 (1993) 135–141. doi:doi:10.1080/00319109308030827.
- [4] N. H. March, Chemical collision model and statistical distribution function for an anyon liquid, *Physics and Chemistry of Liquids* 34 (1997) 61–64. doi:doi:10.1080/00319109708035914.
- [5] R. Acharya, P. N. Swamy, Statistical mechanics of anyons, *Journal of Physics A: Mathematical and General* 27 (1994) 7247–7263. doi:doi:10.1088/0305-4470/27/22/005.
- [6] F. D. M. Haldane, “fractional statistics” in arbitrary dimensions: A generalization of the pauli principle, *Phys. Rev. Lett.* 67 (1991) 937–940. doi:doi:10.1103/PhysRevLett.67.937.
- [7] Y.-S. Wu, Statistical distribution for generalized ideal gas of fractional-statistics particles, *Phys. Rev. Lett.* 73 (1994) 922–925. doi:doi:10.1103/PhysRevLett.73.922.
- [8] A. Dasnières de Veigy, S. Ouvry, One-dimensional statistical mechanics for identical particles: The calogero and anyon cases, *Modern Physics Letters A* 10 (1995) 1–13. doi:doi:10.1142/S0217732395000028.
- [9] A. D. de Veigy, S. Ouvry, One-dimensional statistical mechanics for identical particles: The calogero and anyon cases, *Modern Physics Letters B* 09 (1995) 271–283. doi:doi:10.1142/S0217984995000267.
- [10] F. E. Camino, W. Zhou, V. J. Goldman, Realization of a Laughlin quasiparticle interferometer: Observation of fractional statistics, *Phys. Rev. B* 72 (2005) 075342. doi:doi:10.1103/PhysRevB.72.075342.
- [11] D. Arovas, J. R. Schrieffer, F. Wilczek, Fractional statistics and the quantum hall effect, *Phys. Rev. Lett.* 53 (1984) 722–723. doi:doi:10.1103/PhysRevLett.53.722.
- [12] J. L. Riccardo, A. J. Ramirez-Pastor, F. Romá, Fractional statistical theory of adsorption of poly-

- atomics, *Phys. Rev. Lett.* 93 (2004) 186101. doi:doi:10.1103/PhysRevLett.93.186101.
- [13] J. Riccardo, F. Romá, A. Ramirez-Pastor, Generalized statistical description of adsorption of polyatomics, *Applied Surface Science* 252 (2005) 505–511. doi:doi:10.1016/j.apsusc.2005.02.067, fifth International Symposium, Effects on surface heterogeneity in adsorption and catalysis on solids.
- [14] G. F. Cerofolini, Fractional statistical analysis of adsorption, *Journal of Physics A: Mathematical and General* 39 (2006) 3195–3201. doi:doi:10.1088/0305-4470/39/13/001.
- [15] M. Dávila, J. Riccardo, A. Ramirez-Pastor, Fractional statistical theory and use of quasi-chemical approximation for adsorption of interacting k-mers, *Surface Science* 603 (2009) 683–689. doi:doi:10.1016/j.susc.2008.12.032.
- [16] M. Dávila, J. L. Riccardo, A. J. Ramirez-Pastor, Fractional statistics description applied to adsorption of alkane binary mixtures in zeolites, *The Journal of Chemical Physics* 130 (2009) 174715. doi:doi:10.1063/1.3124163.
- [17] D. A. Matoz-Fernandez, D. H. Linares, A. J. Ramirez-Pastor, New isotherm for multisite occupancy adsorption of long, straight rigid rods, *Langmuir* 27 (2011) 2456–2463. doi:doi:10.1021/la104122h.
- [18] D. Matoz-Fernandez, A. Ramirez-Pastor, Adsorption preference reversal phenomenon from multisite-occupancy theory for two-dimensional lattices, *Chemical Physics Letters* 610-611 (2014) 131–134. doi:doi:10.1016/j.cplett.2014.06.055.
- [19] J. J. Riccardo, J. L. Riccardo, A. J. Ramirez-Pastor, P. M. Pasinetti, Multiple exclusion statistics, *Phys. Rev. Lett.* 123 (2019) 020602. doi:doi:10.1103/PhysRevLett.123.020602.
- [20] A. P. Polychronakos, Probabilities and path-integral realization of exclusion statistics, *Physics Letters B* 365 (1996) 202–206. doi:doi:10.1016/0370-2693(95)01302-4.
- [21] A. P. Polychronakos, Solitons and fractional statistics, *Nuclear Physics B - Proceedings Supplements* 45 (1996) 81–87. doi:doi:10.1016/0920-5632(95)00617-6, proceedings of the Trieste Conference on Recent Developments in Statistical Mechanics and Quantum Field Theory.
- [22] W. S. Chung, H. Hassanabadi, Polychronakos statistics and α -deformed bose condensation of α -bosons, *Modern Physics Letters B* 32 (2018) 1850052. doi:doi:10.1142/S0217984918500525.
- [23] M. Hoyuelos, From creation and annihilation operators to statistics, *Physica A: Statistical Mechanics and its Applications* 490 (2018) 944–952. doi:doi:10.1016/j.physa.2017.09.006.
- [24] C. Nayak, F. Wilczek, Exclusion statistics: Low-temperature properties, fluctuations, duality, and applications, *Phys. Rev. Lett.* 73 (1994) 2740–2743. doi:doi:10.1103/PhysRevLett.73.2740.
- [25] S. Chaturvedi, V. Srinivasan, Interpolations between bose and fermi statistics, *Physica A: Statistical Mechanics and its Applications* 246 (1997) 576–586. doi:doi:10.1016/S0378-4371(97)00348-8.
- [26] S. Chaturvedi, V. Srinivasan, Microscopic interpretation of haldane’s semion statistics, *Phys. Rev. Lett.* 78 (1997) 4316–4319. doi:doi:10.1103/PhysRevLett.78.4316.
- [27] M. V. N. Murthy, R. Shankar, Exclusion statistics: A resolution of the problem of negative weights, *Phys. Rev. B* 60 (1999) 6517–6521. doi:doi:10.1103/PhysRevB.60.6517.
- [28] R. K. Niven, Combinatorial entropies and statistics, *The European Physical Journal B* 70 (2009) 49–63. doi:doi:10.1140/epjb/e2009-00168-5.
- [29] R. K. Niven, M. Grendar, Generalized classical, quantum and intermediate statistics and the p \tilde{A}^3 lya urn model, *Physics Letters A* 373 (2009) 621–626. doi:doi:10.1016/j.physleta.2008.12.025.
- [30] K. Ourabah, M. Tribeche, Fractional superstatistics from a kinetic approach, *Phys. Rev. E* 97 (2018) 032126. doi:doi:10.1103/PhysRevE.97.032126.
- [31] A. A. Abutaleb, Unified statistical distribution of quantum particles and symmetry, *International Journal of Theoretical Physics* 53 (2014) 3893–3900. doi:doi:10.1007/s10773-014-2140-7.
- [32] C.-T. Yan, Statistical model describing bose-einstein and fermi-dirac statistics, *Phys. Rev. E* 104 (2021) 064118. doi:doi:10.1103/PhysRevE.104.064118.
- [33] P. K. Roy, Derivation of a statistical model for classical systems obeying the fractional exclusion principle, *Phys. Rev. E* 106 (2022) 014141. doi:doi:10.1103/PhysRevE.106.014141.
- [34] Note that, the exponent m is shifted by 1 in this derivation method as compared to Eq. 2