

Excitation energies from state-specific ensemble density functionals with density-driven correlations

Tim Gould*

Qld Micro- and Nanotechnology Centre, Griffith University, Nathan, Qld 4111, Australia

Stephen G. Dale

*Qld Micro- and Nanotechnology Centre, Griffith University, Nathan, Qld 4111, Australia and
Institute of Functional Intelligent Materials, National University of Singapore, 4 Science Drive 2, Singapore 117544*

Leeor Kronik

*Department of Molecular Chemistry and Materials Science,
Weizmann Institute of Science, Rehovoth 7610000, Israel*

Stefano Pittalis

CNR-Istituto Nanoscienze, Via Campi 213A, I-41125 Modena, Italy

We present an effective ensemble density functional approximation, which we call GX24 as it is suitable for both ground (G) and excited state (X) energy prediction. This functional is derived from first principles based on a novel approximation for density-driven correlations. We show that GX24 exploits recent advances in ensemble density functional theory to resolve major paradigmatic failures (double excitations, charge transfer) of standard time-dependent density functional approximations. It thus represents a major step toward accurate modelling of neutral and charged excitations in a unified fashion.

Introduction: The simulation and understanding of electronic structure has been transformed by inexpensive and useful density functional approximations (DFAs) [1–4] built on exact density functional theory (DFT) expressions and constraints. Going beyond ground states usually requires time-dependent DFAs (TDDFAs) which, like their ground state counterparts, have transformed our understanding of excited states [5]. However, standard TDDFAs struggle to describe physically and technologically important processes, notably charge transfer [6] and double excitations. [7] Limitations may be overcome via specialised TDDFA-based approaches [8–10], but a generally applicable TDDFA is as yet unavailable. Thus, not only must interested users be familiar with an arsenal of different excited state approaches, they must also understand their applicability and limitations for specific problems.

Fortunately, a large class of excitations involve stationary states and do not call for a truly time-dependent approach, [11, 12] which is needed only for non-equilibrium physics. State-specific approaches which reuse regular DFT approximations (DFAs) are flourishing. [13–15] However, predictions can be extremely sensitive to functional choice [15] because of non-systematic errors.

An alternative idea that is attracting growing attention is the development of state-specific ensemble DFAs (EDFAs), built based on a first principles understanding of excited-states [16–18]. Modern EDFAs systematically and successfully address ‘typical’ excited state problems [18–23] and even multi-configurational problems like bond dissociation. [16] The key advantage of EDFAs is that they draw from excited state ensemble

DFT (EDFT), [24, 25] to embed understanding of the electronic structure of excited states [26–28] and thereby upgrade, rather than reuse, existing ground state DFAs for excited state predictions.

A significant difficulty that has been recognized only recently is that EDFT introduces an additional form of “density-driven” (dd) correlations [27, 29, 30] which must be accounted for in order to obtain accurate and general predictions. Originally, dd-correlations were brought forth by the realization that the individual auxiliary pure states forming the Kohn-Sham ensembles of EDFT do not necessarily need to reproduce the individual interacting densities [29]. Rather, what counts at the functional level is the ensemble-averaged density. The definition of dd-correlations was revisited and generalized to handle more direct calculations and confirm its importance. [30] Eventually, a formally exact expression was derived to inspire approximations involving, explicitly, pure-state densities and density-like overlaps of different states, an issue elaborated below [27].

This Letter derives, from first principles, a *so-far-elusive* practical approximation for dd-correlations. The derivation exploits understanding of the recently-derived weak and strongly correlated limits of EDFT [23] to interpolate the dd correlations and thus obtain a simple, closed-form approximate expression. For regular correlations, that are already captured by (E)DFAs, we build on modern long-range corrected hybrid DFT techniques. We thus produce a unified EDFA for ground (G) and excited (X) states, which we call GX24, that simultaneously and reliably addresses double and charge transfer excitations, as well as singlet-triplet splitting – all neutral

excitations of great relevance in photochemistry for which conventional TDDFAs struggle. We further demonstrate that GX24 is equally useful for charged excitations of importance in other settings, with different physics. In the following, we first provide a brief overview of state-specific contributions in EDFT, with an emphasis on the exact theory of dd-correlations. We then use these exact results to construct a useful approximation that leads to the GX24 functional. Finally, we demonstrate the usefulness of the new functional for selected problems.

EDFT in terms of state-specific contributions: To begin, let us briefly summarise how EDFT lets us address excited states. Seminal work by Gross, Oliveira and Kohn [24] (GOK) showed that ensembles of ground and excited states are subject to an extended variational (stationary) formalism. Specifically, $\mathcal{E}^w := \min_{\{\Psi\}} \text{Tr}[\hat{H}\hat{\Gamma}^w]$, where $\hat{\Gamma}^w = \sum_{\kappa} w_{\kappa} |\Psi_{\kappa}\rangle\langle\Psi_{\kappa}|$ is formed on orthonormal states, $\{|\Psi_{\kappa}\rangle\}$, averaged by weights w_{κ} . They also showed that \mathcal{E}^w can be obtained via an ensemble density functional, $\mathcal{E}^w[n^w]$, of the ensemble density $n = \text{Tr}[\hat{n}\hat{\Gamma}^w] = \sum_{\kappa} w_{\kappa} n_{\kappa}$ [25]. A recent reformulation of GOK theory, obtained by invoking the same weights for the interacting and the auxiliary symmetry-adapted KS states (denoted by the subscript ‘s’), yields [26, 27]

$$\mathcal{E}^w = \sum_{\kappa} w_{\kappa} \left[T_{s,\kappa} + \int n_{\kappa}(\mathbf{r}) v_{\text{ext}}(\mathbf{r}) d\mathbf{r} + E_{\text{Hxc},\kappa} \right], \quad (1)$$

where $\hat{T}_{s,\kappa} = \langle\kappa_s|\hat{T}|\kappa_s\rangle$ is the state-specific KS kinetic energy, v_{ext} is the external potential (typically the scalar potential of the electric field due to the nuclei of a molecule), and $E_{\text{Hxc},\kappa} = E_{\text{Hx},\kappa} + E_{c,\kappa}$, where $E_{\text{Hx},\kappa} = \langle\kappa_s|\hat{W}|\kappa_s\rangle$ is the state-specific KS Hartree-exchange energy and $E_{c,\kappa}$ is the state-specific correlation energy (more below). Notice also that the KS states, $|\kappa_s\rangle$, and their properties (e.g. densities) are implicitly weight-dependent and formally should be written as $|\kappa_s^w\rangle$. For notational simplicity we omit this weight dependence unless it is contextually essential.

KS states in EDFT are unavoidably non-interacting yet symmetry-adapted configuration-state functions (CSFs) which can combine a *finite* number of Slater determinants. As a result, the corresponding *conjoint* Hx energy does not lend itself to the usual determinant-based Hartree and Fock exchange separation.[26, 31]. With the goal of upgrading existing DFAs, we use the fluctuation dissipation theorem (FDT) [27] to split $E_{\text{Hx},\kappa}$ by recognising that the ground state Fock exchange energy obeys the relation, $E_x = \int \{\text{Tr}_{\omega}[\chi] - n\delta\} \frac{d\mathbf{r}d\mathbf{r}'}{2|\mathbf{r}-\mathbf{r}'|}$, with the KS (retarded density-density) response function, χ_s . Assuming the same relation for $\mathcal{E}_x^w = \sum_{\kappa} w_{\kappa} E_{x,\kappa}$ yields $E_{x,\kappa} = \int \{\text{Tr}_{\omega}[\chi_{s,\kappa}](\mathbf{r}, \mathbf{r}') - n_{s,\kappa}(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}')\} \frac{d\mathbf{r}d\mathbf{r}'}{2|\mathbf{r}-\mathbf{r}'|}$, where $\chi_{s,\kappa}$ is the response of $|\kappa_s\rangle$ and $\text{Tr}_{\omega}[\chi_{s,\kappa}](\mathbf{r}, \mathbf{r}') \equiv -\int_{0^-}^{\infty} \frac{d\omega}{\pi} \Im\chi_s^w[n](\mathbf{r}, \mathbf{r}'; \omega)$. We discuss the advantages of this definition below.




FIG. 1. Model adiabatic connection curves for the singly promoted triplet, 3S_1 , and singlet, 1S_1 , including exact limits. The teal dotted line and dashed purple line are $W_{3S_1}^\lambda$ and $W_{1S_1}^\lambda$, respectively. The red dotted line is parallel to the teal dotted line. The shaded areas are correlation energy components [see Eq. (6)] and are discussed in the text.

The ensemble Hartree energy, $\mathcal{E}_H^w = \sum_{\kappa} w_{\kappa} E_{H,\kappa}$, then captures the remaining Hx terms via

$$E_{H,\kappa} = U[n_{s,\kappa}] + 2 \sum_{\kappa' < \kappa} U[n_{s,\kappa\kappa'}]. \quad (2)$$

Here, $U[q] = U[q^*] = \int \frac{d\mathbf{r}d\mathbf{r}'}{2|\mathbf{r}-\mathbf{r}'|} q(\mathbf{r})q^*(\mathbf{r}')$ is a Coulomb-like integral adapted to admit complex inputs like transition densities, $n_{s,\kappa\kappa'} = \langle\kappa_s|\hat{n}|\kappa'_s\rangle$. The sum in the above equation is over states $|\kappa'\rangle$ that are lower in energy than $|\kappa\rangle$. Degenerate orbitals sometimes require a more sophisticated treatment [18]. Here we focus on non-degenerate singlets to avoid complications, unless otherwise stated.

Looking for suitable correlation companions, we employ the adiabatic coupling-constant integration and, again, the FDT. [27] The resulting ensemble correlation can be expressed as $\mathcal{E}_c^w[n] = \sum_{\kappa} w_{\kappa} E_{c,\kappa} = \sum_{\kappa} w_{\kappa} \int_{0^+}^1 (W_{\kappa}^\lambda - W_{\kappa}^{0^+}) d\lambda$. Here, $W_{\kappa}^\lambda \equiv \langle\kappa^\lambda|\hat{W}|\kappa^\lambda\rangle$, where $|\kappa^\lambda\rangle$ is an eigen-state of $\hat{T} + \hat{v}^\lambda + \lambda\hat{W}$ with $v^{0^+} = v_s$, $v^1 = v$ and v^λ otherwise chosen to preserve the (ensemble) density. The correlation energy is further resolved into state-driven (sd) correlations, $E_{c,\kappa}^{\text{sd}} = \int_{0^+}^1 d\lambda \int \text{Tr}_{\omega}[\Delta\chi_{\kappa}^\lambda](\mathbf{r}, \mathbf{r}') \frac{d\mathbf{r}d\mathbf{r}'}{2|\mathbf{r}-\mathbf{r}'|}$, where $\Delta\chi_{\kappa}^\lambda = \chi_{\kappa}^\lambda - \chi_{\kappa,s}$; and density-driven (dd) correlations,

$$E_{c,\kappa}^{\text{dd}} = \int_{0^+}^1 d\lambda (U[n_{\kappa}^\lambda] - U[n_{s,\kappa}]) + 2 \sum_{\kappa' < \kappa} \int_{0^+}^1 d\lambda (U[n_{\kappa\kappa'}^\lambda] - U[n_{s,\kappa\kappa'}]), \quad (3)$$

with $n_{\kappa\kappa'}^\lambda = \langle\kappa^\lambda|\hat{n}|\kappa'^\lambda\rangle$ and $n_{s,\kappa\kappa'} \equiv n_{\kappa\kappa'}^{0^+}$.

Both $E_{x,\kappa}$ and $E_{c,\kappa}^{\text{sd}}$ involve response functions analogous to the DFT case and are thus naturally paired. In fact, this analogy allows one to reuse ground state DFAs in order to model the same exchange and (sd) correlation terms in excited states. Ref. 18 further argued that *conventional* Fock exchange and DFA expressions can model unconventional exchange and sd correlations, by invoking

TABLE I. Ingredients for GX24. Here, ρ_{1S_0} and ρ_{3S_1} are the 1RDMs for the ground- and triplet- states. E'_{xc} and $E'_{ST} = 2(1 - \xi)[hl|lh]_0$ are defined in Eq. (10). 1D_2 is a double excitation ($h^2 \rightarrow l_1l_2$) to degenerate $l_{1,2}$.

State	$E_{\text{Hxc},\kappa}^{\text{GX24}} - U[n_\kappa]$
1S_0	$E'_{xc}[\rho_{1S_0}]$
3S_1	$E'_{xc}[\rho_{3S_1}]$
1S_1	$E'_{xc}[\rho_{3S_1}] + E'_{ST}[\phi_h, \phi_l]$
1S_2	$2E'_{xc}[\rho_{3S_1}] - E'_{xc}[\rho_{1S_0}] + E'_{ST}[\phi_h, \phi_l]$
1D_2	$E'_{xc}[\rho_{3S_1;l_1}] + E'_{xc}[\rho_{3S_1;l_2}] - E'_{xc}[\rho_{1S_0}]$ $+ \frac{1}{2}(E'_{ST}[\phi_h, \phi_{l_1}] + E'_{ST}[\phi_h, \phi_{l_2}]) + E'_{ST}[\phi_{l_1}, \phi_{l_2}]$

combination laws:

$$E_{x,\kappa}[n_\kappa] + E_{c,\kappa}^{\text{sd}}[n_\kappa] \approx \sum_P C_P^\kappa E_{xc}^{\text{DFA}}[n_P]. \quad (4)$$

Here, n_P are densities of Slater determinant states, $|\Phi_P\rangle$ that are lowest energy in their symmetry class, while C_P^κ are coefficients such that $\sum_P C_P^\kappa = 1$, $\sum_P C_P^\kappa n_P = n_\kappa$ and $\sum_P C_P^\kappa E_{x,P} = E_{x,\kappa}$ [32]. Combination laws have proved to be quite successful for excited state prediction [18, 23]. The outstanding challenge is thus with the density-driven correlations in Eq. (3).

Modeling density-driven correlations: The crux of the problem with dd correlations is that $E_{c,\kappa}^{\text{dd}}[n]$ does not resemble anything available from the glorious past of DFT. This is because for the regular non-degenerate ground state that is at the center of all popular DFA constructions, Eq. (3) yields exactly zero. All terms in Eq. (3) disappear because the KS state has the correct density at all λ (i.e. $n_s = n^\lambda = n$) and no eigenstate can be found that is lower in energy than the ground state.

For higher energy states we enter a largely unexplored territory. Let us first consider the situation in which we are able to select a specific state such that $n_{\kappa,s} = n_\kappa^\lambda = n_\kappa$ is constant along the adiabatic connection. Then, the first term in Eq. (3) vanishes. But, in general, $n_{s,\kappa\kappa'} \neq n_{\kappa\kappa'}^\lambda$ leaving us with,

$$E_{c,\kappa}^{\text{dd}}[n] = 2 \sum_{\kappa' < \kappa} \int_{0^+}^1 (U[n_{\kappa\kappa'}^\lambda] - U[n_{s,\kappa\kappa'}]) d\lambda. \quad (5)$$

Evaluating Eq. (5) involves the hitherto impervious task of modeling the variation of transition densities, $n_{\kappa\kappa'}^\lambda$, for all λ in the integration path. Fortunately, the high- and low-density limits, recently derived for excited electrons, offer an ideal means for interpolation over the whole range. Satisfying the known exact limits [28] allows us to write $W_\kappa^\lambda = E_{\text{Hxc},\kappa} + 2\lambda f^\lambda E_{c,\kappa}^{\text{GL2}} + \lambda^2 f^\lambda [E_{\text{Hxc}}^{\text{SCE}}[n_{s,\kappa}] - E_{\text{Hxc},\kappa}]$ where $f^\lambda \equiv f_\kappa[n](\lambda)$ is a monotone function obeying $f^0 = 1$ and $\lim_{\lambda \rightarrow \infty} \lambda^2 f^\lambda = 1$. Here, $E_{\text{Hxc},\kappa} \equiv W_\kappa^{0^+} = \langle \kappa_s | \hat{W} | \kappa_s \rangle$ and $E_{c,\kappa}^{\text{GL2}}$ (the *extension* of Görling-Levy second-order perturbation the-

ory, GL2, to ensemble DFT [33, 34]) belong to the high-density limit and are the leading order terms for $\lambda \rightarrow 0^+$. $E_{\text{Hxc}}^{\text{SCE}}[n] = \inf_{\Psi \rightarrow n} \langle \Psi | \hat{W} | \Psi \rangle$ is an energy functional for strictly correlated electrons (SCE) in the low-density and $\lambda \rightarrow \infty$ limit. In particular, we note that $E_{\text{Hxc}}^{\text{SCE}}$ only depends implicitly on κ , via the input density n_κ , and not explicitly like the other terms [28]. Integration yields

$$E_{c,\kappa} = E_{c,\kappa}^{\text{sd}} + E_{c,\kappa}^{\text{dd}} = \int_{0^+}^1 (W_\kappa^\lambda - W_\kappa^0) d\lambda \approx 2\tau E_{c,\kappa}^{\text{GL2}} + \xi (E_{\text{Hxc}}^{\text{SCE}}[n_\kappa] - E_{\text{Hxc},\kappa}). \quad (6)$$

If τ and ξ are taken as state-dependent density functionals, then Eq. (6) is exact. In the following, we obtain a useful approximation even by treating ξ as constant.

Eq. (6) yields a conjoint expression for the sd and dd correlation terms. To understand how to split them, we consider the case of a typical first excited singlet (1S_1) and triplet (3S_1) pair. These are obtained from a single promotion of an electron from the highest occupied molecular orbital (HOMO, h) to its lowest unoccupied counterpart (LUMO, l). 3S_1 has vanishing density-driven terms, as the only state lower in energy, 1S_0 , belongs to a different symmetry class and therefore $\langle {}^3S_1 | \hat{n} | {}^1S_0 \rangle = 0$. Furthermore, from Eq. (4) we can use a DFA for 3S_1 (representable as a Slater determinant) to model the exchange and state-driven correlations of 1S_1 (a CSF).

The situation is depicted in Figure 1. For the triplet, Eq. (6) only captures state-driven correlations: $E_{c,3S_1} = E_{c,3S_1}^{\text{sd}}$ and thus $E_{c,3S_1}^{\text{EDFA}} = E_{c,3S_1}^{\text{sd-EDFA}}$ (negative of the green shaded area). The excited singlet, 1S_1 , does have a density-driven term, $E_{c,1S_1}^{\text{dd}} = E_{c,1S_1} - E_{c,1S_1}^{\text{sd}}$, due to the transition densities with the singlet ground state, 1S_0 . Thus, $E_{c,1S_1}^{\text{DFA}} = E_{c,1S_1}^{\text{sd-EDFA}} + E_{c,1S_1}^{\text{dd-EDFA}}$ (negative of orange and violet, respectively) requires a density-driven correlation approximation, $E_{c,1S_1}^{\text{dd-EDFA}}$.

To proceed, we observe the following: i) we may express $E_{\text{Hxc}}^{\text{SCE-DFA}}[n_{s,\kappa}] \approx U[n_\kappa] + E_{xc}^{\text{SCE-GEA}}[n_\kappa]$, where $E_{xc}^{\text{SCE-GEA}}$ is a gradient expansion approximation which is known to be very good for SCE ground states [35–37]; ii) we use the same orbitals and thus same densities for 3S_1 and 1S_1 so that the corresponding Coulomb (U), DFA (for x and c), and SCE-GEA energies are identical [18, 27, 28]; iii) Combination rules [Eq. (4)] together with point (ii) imply that the DFA state-driven correlations of 1S_1 and 3S_1 are equal (in Fig. 1 green area = orange area and dotted lines are parallel). Then, $E_{c,1S_1}^{\text{dd-EDFA}} \approx E_{c,1S_1}^{\text{DFA}} - E_{c,3S_1}^{\text{sd-EDFA}}$ and, through geometric arguments, we see that $E_{c,1S_1}^{\text{dd-EDFA}} \propto -E_{ST}$ (negative of violet area). Matching terms to Eq. (6) yields, after some algebra:

$$E_{c,1S_1}^{\text{dd-EDFA}} \approx -2\xi U[n_{s,1S_0,1S_1}] \equiv -\xi E_{ST}. \quad (7)$$

Note that E_{ST} has the form of the singlet-triplet (ST) energy splitting of HF theory, but it is appropriately rescaled.

What about other excitations? Let us now consider the singlet, 1S_2 , obtained by a double excitation involving the promotion of both electrons from the HOMO to the (non-degenerate) LUMO. 1S_2 has no matching triplet to borrow SD physics from. However, the state-driven contributions can still be approximated successfully via ground-state functionals by invoking a pertinent combination rule [Eq. (4)], because $E_{x,^1S_2} = 2E_x[n_{^3S_1}] - E_x[n_{^1S_0}]$.

A first step towards a full approximation is to substitute $E_{\text{Hxc}}^{\text{SCE}} \approx U + E_{\text{xc}}^{\text{SCE-GEA}}$ into Eq. (6) to obtain $E_{c,\kappa} \approx 2\tau E_{c,\kappa}^{\text{GL2}} + \xi(E_{\text{xc}}^{\text{SCE-GEA}}[n_\kappa] - E_{x,\kappa}) + \xi(U[n_\kappa] - E_{\text{H},\kappa})$, as was done geometrically for the single promotion case. We next recognise that GL2, a state-driven term, should also obey the combination law. Then, we treat the SCE-GEA term using the same combination law, so that $E_{c,^1S_2}^{\text{sd-EDFA}} := 2\tau E_{c,^1S_2}^{\text{GL2}} + \xi(E_{\text{xc}}^{\text{SCE-GEA}}[n_{^1S_2}] - E_{x,^1S_2}) \approx 2E_c^{\text{DFA}}[n_{^3S_1}] - E_c^{\text{DFA}}[n_{^1S_0}]$. This covers all ‘x’ and ‘c’ terms in Eq. (6), meaning only the ‘H’ terms, $\xi(U[n_{^1S_2}] - E_{\text{H},^1S_2})$, remain. Thus,

$$E_{c,^1S_2}^{\text{dd-EDFA}} \approx -2\xi U[n_{s,^1S_1^1S_2}] \equiv -\xi E_{ST}, \quad (8)$$

follows from Eq. (2) and $n_{s,^1S_0^1S_2} = 0$. Extending this argument to higher states leads to our first major result:

$$E_{c,\kappa}^{\text{dd-EDFA}} \approx -2\xi \sum_{\kappa' < \kappa} U[n_{s,\kappa'\kappa}]. \quad (9)$$

Note, however, that some cases with orbital degeneracies require a more sophisticated treatment. The two-fold degenerate case reported in Table I (introduced in more detail below) can use Eq. (9) in unmodified form. General degeneracies must be treated using energy *levels*, not individual states, which can lead to contributions from the first term in Eq. (3).




FIG. 2. **a)** Excitation energies (y axis) as a function of reference values (x axis) for EX22. Colours indicate type of excitation. Dotted line indicates perfect agreement. **b)** GX24 (blue) and TDA CAM-B3LYP (green) energies versus reference energies for ‘easy’ excitation energies. **c)** Like **b)** but for ‘hard’ double excitations that TDDFT cannot tackle.

The GX24 approximation: We are now ready to combine the density-driven correlation model of Eq. (9) with a long-range corrected (LC) hybrid model that is known to be facilitate the description of charge transfer excitation [38]. Invoking Eqs (2), (4), and

(9) yields an EDFA of general form (specific examples are provided in Table I), $E_{\text{Hxc},\kappa}^{\text{EDFA}} = U[n_\kappa] + \sum_P C_P^\kappa E_{\text{xc}}^{\text{LC}}[\rho_P] + (1-2\xi) \sum_{\kappa' < \kappa} U[n_{s,\kappa'\kappa}]$. Here, $E_{\text{xc}}^{\text{LC}} := E_x^{\text{HF}} + (1-\alpha)(E_x^{\text{HJS},\gamma} - E_x^{\text{HF},\gamma}) + E_c^{\text{PBE}}$, where ‘PBE’ is the Perdew-Burke-Ernzerhof generalized-gradient approximation [39] and ‘HJS’ is the Henderson-Janesko-Scuseria PBE-like form [40] for short-range exchange. α denotes the fraction of short-range HF exchange, γ is a range-separation parameter that defines the range-partition of the Coulomb interaction, and ξ governs the weight of density-driven correlations.

The remaining task is to determine the numerical values for α , γ , and ξ . We choose $\alpha = \frac{3}{8}$ as a compromise between the usual PBE0 [41] ground state value, $\alpha = \frac{1}{4}$, and $\alpha = \frac{1}{2}$ that has been shown to offer some improvement for EDFT excited states. [18, 22, 23, 42] At this point, determining ξ and γ requires some empiricism and therefore we norm them over EX22, a benchmark set of 22 excitation energies (21 from QuestDB [43–45] plus the degenerate double excitation of diatomic BH [18]) that are selected to capture a diversity of physical and chemical behaviours. As revealed in **Supplementary Figure 1**, the global optimal value occurs for $\gamma = 0$, which is incompatible with our goal of capturing charge transfer with a LC hybrid. We therefore settle on a mid-ranged $\gamma = 0.2 \text{ Bohr}^{-1}$ with $\xi = 0.32$ to retain an appropriate treatment of charge transfer physics with a minimal impact (a 0.02 eV increase) on the overall optimal mean absolute error (MAE). Implementational details, including Refs 46–49, are in **Supplementary Material Section I (SM Sec I)**; details and discussion of EX22 are in **SM Sec II**.

Our second major result is the GX24 ground (G) and excited (X) state EDFA, defined by using

$$E'_{\text{xc}}[\rho] := E_x^{\text{HF}}[\rho] + E_c^{\text{PBE}}[n_\uparrow, n_\downarrow] + \frac{5}{8}(E_x^{\text{HJS},0.2}[n_\uparrow, n_\downarrow] - E_x^{\text{HF},0.2}[\rho]), \quad (10a)$$

$$E'_{ST}[\phi_h, \phi_l] := 2(1 - 0.32)[hl|lh] = 1.36[hl|lh], \quad (10b)$$

in the energy expressions in Table I. Eq. (10a) is defined in terms of spin-resolved one-body reduced density matrices (1RDMs), ρ_P , of Slater determinants; and thus accommodates 1S_0 ($\rho_{^1S_0,\uparrow} = \rho_{^1S_0,\downarrow} = \sum_{i \leq h} \rho_i$) and 3S_1 1RDMs ($\rho_{^3S_1,\uparrow} = \sum_{i \leq l} \rho_i$, $\rho_{^3S_1,\downarrow} = \sum_{i < h} \rho_i$) in the usual way – note, however, that in our calculations the single-particle orbitals are the same for the two spin channels. Eq (10b) involves electron repulsion integrals, $[ij|kl] = \int \phi_i(\mathbf{r})\phi_j^*(\mathbf{r})\phi_k(\mathbf{r}')\phi_l^*(\mathbf{r}') \frac{d\mathbf{r}d\mathbf{r}'}{|\mathbf{r}-\mathbf{r}'|}$.

Figure 2a reveals that Eq. (10) performs extremely well for $^1S_0 \rightarrow ^1S_1$ (SS) and $^3S_1 \rightarrow ^1S_1$ (ST) transitions in EX22, with impressive mean absolute errors (MAE) of 0.19 eV and 0.04 eV, respectively. It is worse, yet still respectable (in the absence of system-specific optimal tuning [9]) for difficult charge-transfer (CT, MAE 0.58 eV) and does extremely well for the notorious case of dou-




FIG. 3. Electron affinity of F (a), CN (b) and ozone (c) at fractional charges. Lime is without the dd-EDFA. Black dotted line is the theoretical linear behaviour. Black dots are experimental values. [51–53]

ble excitations (DX, 0.35 eV). Using only the sd-EDFA [Eq. (10a)] increases errors substantially for all four categories, overall by 83%, highlighting the importance of the dd-EDFA.

GX24 predictions: First, recognising that exact EDFT reduces to DFT for ground states, we evaluate the performance of GX24 on the P30-10 poison benchmark set [50], to test *worst-case* performance on ground states. Despite balancing the needs of ground and excited states, GX24 (MAE 0.65 eV) performs slightly better than the popular B3LYP (MAE 0.69 eV) on absolute errors and is the top performer on relative errors. Details of all predictions, including technical details and extended theory and data, are in [SM Sec III](#).

We next test GX24 on relatively ‘easy’ excitations that TDDFT can also address. Figure 2b shows the lowest-lying singlet-singlet transitions of four small atoms, with a focus on Rydberg-like excitations for which very high quality theoretical reference data is available [43]. GX24 performs exceptionally well with a MAE of 0.18 eV across all eight excitations. By contrast, CAM-B3LYP [54] (a popular LC hybrid) with the Tamm-Dancoff approximation (TDA) systematically underestimates gaps; and full TDDFT (not shown) is slightly worse. Figure 2c extends the tests to ‘hard’ problems with a substantial ‘double excitation’ character [55]. Again, GX24 performs exceptionally well considering the difficulty of these problems. TDDFT (not shown) can only detect three of these excitations, and even then does extremely poorly quantitatively (errors of 1.5, 2.8 and 3.2 eV).

Our final test steps outside the realm of neutral excitations to address *fractionally charged* excitation results. These excitations involve ground states, but reproducing non-integer electron number requires an ensemble treatment [56] and thus gives rise to dd correlations [57]. Figure 3 shows the electron affinity of F, CN, and ozone (O_3) at fractional charges, which in exact theory should be linear as a function of excess charge [56]. Applying only the sd-EDFA part of GX24 (i.e. Eq. (10a) adapted to known fractional charge combination rules [58–60]) to this problem leads to substantial non-linearities (lime curves in the figure). Using

$\mathcal{E}_c[n] \approx 2\tau\mathcal{E}_c^{\text{GL2}}[n] + \xi(E_{\text{Hxc}}^{\text{SCE}}[n] - \mathcal{E}_{\text{Hxc}}[n])$ and the combination rules leads to $\mathcal{E}_c^{\text{dd-EDFA}}(q) \approx -\xi\frac{q(1-q)}{2}[ll|ll]$ as the appropriate adaptation of Eq. (10b), where $\xi = 0.32$, q is the excess electron charge and ϕ_l is the frontier orbital. Including $\mathcal{E}_c^{\text{dd-EDFA}}(q)$ (blue curves) largely restores linearity in all cases, to yield excellent qualitative and quantitative results for all charges.

Conclusions: By transparent first principles modelling, we derived a simple yet effective approximation, Eq. (9), for density-driven correlations – a central quantity for dealing with excited states via ensemble density functionals. We then approximated the remaining (and better understood) state-driven correlations by a long-range corrected hybrid functional to produce a new EDFA, GX24 [Eq. (10)], that captures a variety of important ground- and excited state physics within a unified framework.

The performance of GX24 stands beyond the reach of state-of-the-art time-dependent DFAs for addressing neutral excitations of relevance to photochemistry, as it captures both charge transfer and double excitations. GX24’s performance on charged excitations is particularly astonishing, given that it was derived and normed on neutral excitations.

Moreover, some simplifying assumptions in the present work can be relaxed to offer further improvements: i) parameters, including ξ , are system and state-dependent, which can be accommodated in approximations; ii) other types of excitations, including challenging degenerate energy levels, can be accommodated within the framework; iii) the modeling of the correlation physics can be refined.

TG, SGD, and LK were supported by an Australian Research Council (ARC) Discovery Project (DP200100033). TG was supported by an ARC Future Fellowship (FT210100663). LK was supported by the Aryeh and Mintzi Katzman Professorial Chair and the Helen and Martin Kimmel Award for Innovative Investigation. SGD was supported by the Ministry of Education, Singapore, under its Research Centre of Excellence award to the Institute for Functional Intelligent Materials. Project No. EDUNC-33-18-279-V12. Computing resources were provided by the Australian National Computing Merit Application Scheme (NCMAS sp13).

* t.gould@griffith.edu.au

- [1] J. P. Perdew and K. Schmidt, Jacob’s ladder of density functional approximations for the exchange-correlation energy, AIP Conf. Proc. **577**, 1 (2001).
- [2] J. P. Perdew and S. Kurth, A primer in density functional theory (Springer, 2003) Chap. Chapter 1: Density functionals for non-relativistic Coulomb systems in the new century, pp. 1–55.
- [3] S. Kümmel and L. Kronik, Orbital-dependent density functionals: Theory and applications, Rev. Modern Phys.

- 80**, 3 (2008).
- [4] A. D. Becke, Perspective: Fifty years of density-functional theory in chemical physics, *J. Chem. Phys.* **140**, 18A301 (2014).
- [5] C. A. Ullrich, *Time-dependent density-functional theory*, reprinted, with corrections ed., Oxford graduate texts (Oxford University Press, Oxford, 2016) literaturverzeichnis: Seite 484-510.
- [6] N. T. Maitra, Charge transfer in time-dependent density functional theory, *J. Phys.: Cond. Matter* **29**, 423001 (2017).
- [7] P. Elliott, S. Goldson, C. Canahui, and N. T. Maitra, Perspectives on double-excitations in TDDFT, *Chem. Phys.* **391**, 110 (2011).
- [8] A. S. de Wijn, M. Lein, and S. Kümmel, Strong-field ionization in time-dependent density functional theory, *EPL (Europhysics Letters)* **84**, 43001 (2008).
- [9] T. Stein, L. Kronik, and R. Baer, Reliable prediction of charge transfer excitations in molecular complexes using time-dependent density functional theory, *J. Am. Chem. Soc.* **131**, 2818 (2009).
- [10] D. Casanova and A. I. Krylov, Spin-flip methods in quantum chemistry, *Phys. Chem. Chem. Phys.* **22**, 4326 (2020).
- [11] S. Matsika and A. I. Krylov, Introduction: Theoretical modeling of excited state processes, *Chem. Rev.* **118**, 6925 (2018).
- [12] T. Gould, Stationary conditions for excited states: the surprising impact of density-driven correlations, arXiv:2404.12593v1 [physics.chem-ph] (2024).
- [13] G. Levi, A. V. Ivanov, and H. Jónsson, Variational calculations of excited states via direct optimization of the orbitals in dft, *Faraday Discuss.* **224**, 448 (2020).
- [14] A. V. Ivanov, G. Levi, E. Ö. Jónsson, and H. Jónsson, Method for calculating excited electronic states using density functionals and direct orbital optimization with real space grid or plane-wave basis set, *J. Chem. Theory Comput.* **17**, 5034 (2021).
- [15] D. Hait and M. Head-Gordon, Orbital optimized density functional theory for electronic excited states, *J. Phys. Chem. Lett.* **12**, 4517 (2021).
- [16] M. Filatov, Spin-restricted ensemble-referenced Kohn-Sham method: Basic principles and application to strongly correlated ground and excited states of molecules, *WIREs Comput Mol Sci* **5**, 146 (2015).
- [17] F. Cernatic, B. Senjean, V. Robert, and E. Fromager, Ensemble density functional theory of neutral and charged excitations: Exact formulations, standard approximations, and open questions, *Top. Curr. Chem.* **380**, 10.1007/s41061-021-00359-1 (2021).
- [18] T. Gould, L. Kronik, and S. Pittalis, Double excitations in molecules from ensemble density functionals: Theory and approximations, *Phys. Rev. A* **104**, 022803 (2021).
- [19] M. Filatov and S. Shaik, A spin-restricted ensemble-referenced Kohn-Sham method and its application to diradicaloid situations, *Chem. Phys. Lett.* **304**, 429 (1999).
- [20] M. Filatov, M. Huix-Rotllant, and I. Burghardt, Ensemble density functional theory method correctly describes bond dissociation, excited state electron transfer, and double excitations, *J. Chem. Phys.* **142**, 184104 (2015).
- [21] Z.-h. Yang, A. Pribram-Jones, K. Burke, and C. A. Ullrich, Direct extraction of excitation energies from ensemble density-functional theory, *Phys. Rev. Lett.* **119**, 033003 (2017).
- [22] T. Gould, Approximately self-consistent ensemble density functional theory: Toward inclusion of all correlations, *J. Phys. Chem. Lett.* **11**, 9907 (2020).
- [23] T. Gould, Z. Hashimi, L. Kronik, and S. G. Dale, Single excitation energies obtained from the ensemble “HOMO–LUMO gap”: Exact results and approximations, *The Journal of Physical Chemistry Letters* **13**, 2452 (2022).
- [24] E. K. U. Gross, L. N. Oliveira, and W. Kohn, Rayleigh-Ritz variational principle for ensembles of fractionally occupied states, *Phys. Rev. A* **37**, 2805 (1988).
- [25] E. K. U. Gross, L. N. Oliveira, and W. Kohn, Density-functional theory for ensembles of fractionally occupied states. i. basic formalism, *Phys. Rev. A* **37**, 2809 (1988).
- [26] T. Gould and S. Pittalis, Hartree and exchange in ensemble density functional theory: Avoiding the nonuniqueness disaster, *Phys. Rev. Lett.* **119**, 243001 (2017).
- [27] T. Gould, G. Stefanucci, and S. Pittalis, Ensemble density functional theory: Insight from the fluctuation-dissipation theorem, *Phys. Rev. Lett.* **125**, 233001 (2020).
- [28] T. Gould, D. P. Kooi, P. Gori-Giorgi, and S. Pittalis, Electronic excited states in extreme limits via ensemble density functionals, *Phys. Rev. Lett.* **130**, 106401 (2023).
- [29] T. Gould and S. Pittalis, Density-driven correlations in many-electron ensembles: Theory and application for excited states, *Phys. Rev. Lett.* **123**, 016401 (2019).
- [30] E. Fromager, Individual correlations in ensemble density-functional theory: State-driven/density-driven decompositions without additional Kohn-Sham systems, *Phys. Rev. Lett.* **124**, 243001 (2020).
- [31] A. Pribram-Jones, Z.-h. Yang, J. R. Trail, K. Burke, R. J. Needs, and C. A. Ullrich, Excitations and benchmark ensemble density functional theory for two electrons, *J. Chem. Phys.* **140** (2014).
- [32] It follows from the expressions derived in Ref. 18 that the exchange energy depends linearly on orbital occupation factors, as do densities, for fixed orbitals and orbital energies. Thus, for any target density expressible via a combinations rules of densities, it follows that the exchange energy obeys the same combination rules.
- [33] A. Görling and M. Levy, Correlation-energy functional and its high-density limit obtained from a coupling-constant perturbation expansion, *Phys. Rev. B* **47**, 13105 (1993).
- [34] Z. Yang, Second-order perturbative correlation energy functional in the ensemble density-functional theory, *Phys. Rev. A* **104**, 052806 (2021).
- [35] M. Seidl, J. P. Perdew, and M. Levy, Strictly correlated electrons in density-functional theory, *Phys. Rev. A* **59**, 51 (1999).
- [36] M. Seidl, J. P. Perdew, and S. Kurth, Density functionals for the strong-interaction limit, *Phys. Rev. A* **62**, 012502 (2000).
- [37] S. Šmiga and L. A. Constantin, Modified interaction-strength interpolation method as an important step toward self-consistent calculations, *J. Chem. Theory Comput.* **16**, 4983 (2020).
- [38] H. Iikura, T. Tsuneda, T. Yanai, and K. Hirao, A long-range correction scheme for generalized-gradient-approximation exchange functionals, *J. Chem. Phys.* **115**, 3540 (2001).

- [39] J. P. Perdew, K. Burke, and M. Ernzerhof, Generalized gradient approximation made simple, *Phys. Rev. Lett.* **77**, 3865 (1996).
- [40] T. M. Henderson, B. G. Janesko, and G. E. Scuseria, Generalized gradient approximation model exchange holes for range-separated hybrids, *J. Chem. Phys.* **128**, 194105 (2008).
- [41] C. Adamo and V. Barone, Toward reliable density functional methods without adjustable parameters: The PBE0 model, *J. Chem. Phys.* **110**, 6158 (1999).
- [42] T. Gould and L. Kronik, Ensemble generalized Kohn-Sham theory: The good, the bad, and the ugly, *J. Chem. Phys.* **154**, 094125 (2021).
- [43] P.-F. Loos, A. Scemama, A. Blondel, Y. Garniron, M. Caffarel, and D. Jacquemin, A mountaineering strategy to excited states: Highly accurate reference energies and benchmarks, *J. Chem. Theory Comput.* **14**, 4360 (2018).
- [44] P.-F. Loos, N. Galland, and D. Jacquemin, Theoretical 0-0 energies with chemical accuracy, *J. Phys. Chem. Lett.* **9**, 4646 (2018).
- [45] P.-F. Loos, M. Boggio-Pasqua, A. Scemama, M. Caffarel, and D. Jacquemin, Reference energies for double excitations, *J. Chem. Theory Comput.* **15**, 1939 (2019).
- [46] T. Gould, *Broadway - an EDFT library for python* (2024).
- [47] R. M. Parrish, L. A. Burns, D. G. A. Smith, A. C. Simmonett, A. E. DePrince, E. G. Hohenstein, U. Bozkaya, A. Y. Sokolov, R. D. Remigio, R. M. Richard, J. F. Gonthier, A. M. James, H. R. McAlexander, A. Kumar, M. Saitow, X. Wang, B. P. Pritchard, P. Verma, H. F. Schaefer, K. Patkowski, R. A. King, E. F. Valeev, F. A. Evangelista, J. M. Turney, T. D. Crawford, and C. D. Sherrill, Psi4 1.1: An open-source electronic structure program emphasizing automation, advanced libraries, and interoperability, *J. Chem. Theory Comput.* **13**, 3185 (2017).
- [48] D. G. A. Smith, L. A. Burns, D. A. Sirianni, D. R. Nascimento, A. Kumar, A. M. James, J. B. Schriber, T. Zhang, B. Zhang, A. S. Abbott, E. J. Berquist, M. H. Lechner, L. A. Cunha, A. G. Heide, J. M. Waldrop, T. Y. Takeshita, A. Alenaizan, D. Neuhauser, R. A. King, A. C. Simmonett, J. M. Turney, H. F. Schaefer, F. A. Evangelista, A. E. DePrince, T. D. Crawford, K. Patkowski, and C. D. Sherrill, Psi4numpy: An interactive quantum chemistry programming environment for reference implementations and rapid development, *J. Chem. Theory Comput.* **14**, 3504 (2018).
- [49] Q. Sun, X. Zhang, S. Banerjee, P. Bao, M. Barbry, N. S. Blunt, N. A. Bogdanov, G. H. Booth, J. Chen, Z.-H. Cui, J. J. Eriksen, Y. Gao, S. Guo, J. Hermann, M. R. Hermes, K. Koh, P. Koval, S. Lehtola, Z. Li, J. Liu, N. Mardirossian, J. D. McClain, M. Motta, B. Mussard, H. Q. Pham, A. Pulkin, W. Purwanto, P. J. Robinson, E. Ronca, E. R. Sayfutyarova, M. Scheurer, H. F. Schurkus, J. E. T. Smith, C. Sun, S.-N. Sun, S. Upadhyay, L. K. Wagner, X. Wang, A. White, J. D. Whitfield, M. J. Williamson, S. Wouters, J. Yang, J. M. Yu, T. Zhu, T. C. Berkelbach, S. Sharma, A. Y. Sokolov, and G. K.-L. Chan, Recent developments in the pyscf program package, *J. Chem. Phys.* **153**, 10.1063/5.0006074 (2020).
- [50] T. Gould and S. G. Dale, Poisoning density functional theory with benchmark sets of difficult systems, *Phys. Chem. Chem. Phys.* **24**, 6398 (2022).
- [51] C. Blondel, P. Cacciani, C. Delsart, and R. Trainham, High-resolution determination of the electron affinity of fluorine and bromine using crossed ion and laser beams, *Phys. Rev. A* **40**, 3698 (1989).
- [52] S. E. Bradforth, E. H. Kim, D. W. Arnold, and D. M. Neumark, Photoelectron spectroscopy of CN^- , NCO^- , and NCS^- , *J. Chem. Phys.* **98**, 800 (1993).
- [53] S. E. Novick, P. C. Engelking, P. L. Jones, J. H. Futrell, and W. C. Lineberger, Laser photoelectron, photodetachment, and photodestruction spectra of O_3 , *J. Chem. Phys.* **70**, 2652 (1979).
- [54] T. Yanai, D. Tew, and N. Handy, A new hybrid exchange-correlation functional using the coulomb-attenuating method (CAM-B3LYP), *Chem. Phys. Lett.* **393**, 51 (2004).
- [55] F. Kossoski, M. Boggio-Pasqua, P.-F. Loos, and D. Jacquemin, Reference energies for double excitations: Improvement and extension, *J. Chem. Theory Comput.* 10.1021/acs.jctc.4c00410 (2024).
- [56] J. P. Perdew, R. G. Parr, M. Levy, and J. L. Balduz, Density-functional theory for fractional particle number: Derivative discontinuities of the energy, *Phys. Rev. Lett.* **49**, 1691 (1982).
- [57] Note, in the ensemble case there are no transition densities to worry about. Rather, the $\lambda = 0$ limit yields a weighted average of electrostatic energies, $(1-q)U[n_N] + qU[n_{N+1}]$ of N and $N + 1$ electron systems, whereas the $\lambda = \infty$ limit yields the electrostatic energy of the average, $U[(1-q)n_N + qn_{N+1}]$. The dd correlation energy is a fraction of the difference.
- [58] T. Gould and J. F. Dobson, The flexible nature of exchange, correlation, and Hartree physics: Resolving “delocalization” errors in a “correlation free” density functional, *J. Chem. Phys.* **138**, 014103 (2013).
- [59] E. Kraisler and L. Kronik, Piecewise linearity of approximate density functionals revisited: Implications for frontier orbital energies, *Phys. Rev. Lett.* **110**, 126403 (2013).
- [60] E. Kraisler and L. Kronik, Fundamental gaps with approximate density functionals: The derivative discontinuity revealed from ensemble considerations, *J. Chem. Phys.* **140**, 18A540 (2014).