

Evaluating Model Performance Under Worst-case Subpopulations

Mike Li¹ Hongseok Namkoong² Shangzhou Xia³

Decision, Risk, and Operations Division, Columbia Business School

{mli24, namkoong, sxia24}@gsb.columbia.edu

Abstract

The performance of ML models degrades when the training population is different from that seen under operation. Towards assessing distributional robustness, we study the worst-case performance of a model over *all* subpopulations of a given size, defined with respect to core attributes Z . This notion of robustness can consider arbitrary (continuous) attributes Z , and automatically accounts for complex intersectionality in disadvantaged groups. We develop a scalable yet principled two-stage estimation procedure that can evaluate the robustness of state-of-the-art models. We prove that our procedure enjoys several finite-sample convergence guarantees, including *dimension-free* convergence. Instead of overly conservative notions based on Rademacher complexities, our evaluation error depends on the dimension of Z only through the out-of-sample error in estimating the performance conditional on Z . On real datasets, we demonstrate that our method certifies the robustness of a model and prevents deployment of unreliable models.

1 Introduction

The training population typically does not accurately represent what the model will encounter under operation. Model performance has been observed to substantially degrade under distribution shift [20, 36, 96, 109, 68] in speech recognition [67], automated essay scoring [4], and wildlife conservation [11]. Similar trends persist for state-of-the-art (SOTA) NLP and computer vision models [107, 100], even on new data constructed under a near-identical process [78, 90]. Heavily engineered commercial models are no exception [24], performing poorly on rare entities in named entity linking and examples that require abstraction and distillation in summarization tasks [51].

A particularly problematic form of distribution shift comes from embedded power structures in data collection. Data forms the infrastructure on which we build prediction models [39], and they inherit socioeconomic and political inequities against marginalized communities. For example, out of 10,000+ cancer clinical trials the National Cancer Institute funds, less than 5% of participants were non-white [26]. Typical models replicate and perpetuate such bias, and their performance drops significantly on underrepresented groups. Speech recognition systems work poorly for Blacks [67] and those with minority accents [3]. More generally, model performance degrades across demographic attributes such as race, gender, or age, in facial recognition, video captioning, language identification, and academic recommender systems [54, 60, 21, 98, 108, 24].

Model training typically relies on varied engineering practices. It is crucial to *rigorously certify* model robustness prior to deployment for these heuristic approaches to bear fruit and transform consequential applications. Ensuring that models perform uniformly well across subpopulations is simultaneously critical for reliability, fairness, satisfactory user experience, and long-term business goals. While a natural approach is to evaluate performance across a small set of groups, disadvantaged subpopulations are hard to define a priori because of *intersectionality*. The most adversely affected are often determined by a complex combination of variables such as race, income,

and gender [24]. For example, performance on summarization tasks varies across demographic characteristics and document specific traits such as abstractiveness, distillation, and location and dispersion of information [51].

Motivated by these challenges, we study the worst-case subpopulation performance across *all* subpopulations of a given size. This conservative notion of performance evaluates robustness to unanticipated distribution shifts in Z , and automatically accounts for complex intersectionality by virtue of being agnostic to demographic groupings. Formally, let Z be a set of core attributes that we wish to guarantee uniform performance over. It may include protected demographic variables such as race, gender, income, age, or task-specific information such as length of the prompt or metadata on the input; notably, it can contain any continuous or discrete variables. We let $X \in \mathcal{X}$ be the input / covariate, and $Y \in \mathcal{Y}$ be the label. In NLP and vision applications, X is high-dimensional and typically $\dim(Z) \ll \dim(X)$.

We use $\theta(X)$ to denote a fixed prediction model and consider flexible and abstract losses $\ell(\theta(x); y)$. Our goal is to ensure that the model θ performs well over all subpopulations defined over Z . We evaluate model losses on a mixture component, which we call a subpopulation. Postulating a lower bound $\alpha \in (0, 1]$ on the demographic proportion (mixture weight), we consider the set of subpopulations of the data-generating distribution P_Z

$$\mathcal{Q}_\alpha := \{Q_Z \mid P_Z = aQ_Z + (1 - a)Q'_Z \text{ for some } a \geq \alpha, \text{ and subpopulation } Q'_Z\}. \quad (1.1)$$

The demographic proportion (mixture weight) a represents how underrepresented the subpopulation is under the data-generating distribution P_Z .

Before deploying the model θ , we wish to evaluate the worst-case subpopulation performance

$$W_\alpha^* := \sup_{Q_Z \in \mathcal{Q}_\alpha} \mathbb{E}_{Z \sim Q_Z} [\mathbb{E}[\ell(\theta(X), Y) \mid Z]]. \quad (1.2)$$

The worst-case subpopulation performance (1.2) guarantees uniform performance over subpopulations (1.1) and has a clear interpretation that can be communicated to diverse stakeholders. The minority proportion α can often be chosen from first principles, e.g., we wish to guarantee uniformly good performance over subpopulations comprising at least $\alpha = 20\%$ of the collected data. Alternatively, it is often informative to study the threshold level of α^* when $\alpha \mapsto W_\alpha^*$ crosses the *maximum level of acceptable loss*. The threshold α^* provides a *certificate of robustness* on the model $\theta(\cdot)$, guaranteeing that all subpopulations larger than α^* enjoy good performance.

We develop a principled and scalable procedure for estimating the worst-case subpopulation performance (1.2) and the certificate of robustness α^* . A key technical challenge is that for each data point, we observe the loss $\ell(\theta(X); Y)$ but never observe the conditional risk evaluated at the attribute Z

$$\mu^*(Z) := \mathbb{E}[\ell(\theta(X); Y) \mid Z]. \quad (1.3)$$

In Section 2, we propose a two-stage estimation approach where we compute an estimate $\hat{\mu}(\cdot) \in \mathcal{H}$ of the conditional risk $\mu^*(\cdot)$. Then, we compute a *debiased* estimate of the worst-case subpopulation performance under $\hat{\mu}(\cdot)$ using a dual reformulation of the worst-case problem (1.2). We show several theoretical guarantees for our estimator of the worst-case subpopulation performance (1.2). In particular, our first finite-sample result (Section 4) shows convergence at the rate $O_p(\sqrt{\mathbf{Comp}_n(\mathcal{H})/n})$, where \mathbf{Comp}_n denotes a notion of complexity for the model class estimating the conditional risk (1.3).

In some applications, it may be natural to define Z using images or natural languages describing the input and use deep networks to predict the conditional risk (1.3). As the complexity term

$\text{Comp}_n(\mathcal{H})$ becomes prohibitively large in this case [10, 121], our second result (Section 4.3) shows data-dependent *dimension-free* concentration of our two-stage estimator: our bound only depends on the complexity of the model class \mathcal{H} through the out-of-sample error for estimating the conditional risk (1.3). This error can be made small using overparameterized deep networks, allowing us to estimate the conditional risk (1.3) using even the largest deep networks and still obtain a theoretically principled upper confidence bound on the worst-case subpopulation performance. Leveraging these guarantees, we develop principled procedures for estimating the certificates of robustness α^* in Section E.

In Section 5, we demonstrate the effectiveness of our procedure on real data. By evaluating model robustness under subpopulation shifts, our methods allow the selection of robust models before deployment as we illustrate using the recently proposed CLIP model [86]. Finally, we generalize the notion of worst-case subpopulation performance we study in Section 6. We note that these measures in fact form an equivalence with coherent risk measures and distributionally robust losses. At a high level, our result uncovers a deeper connection between classical ideas in risk measures and the more recent ML fairness literature.

Related work. Our notion of worst-case subpopulation performance is also related to the by now vast literature on fairness in ML. We give a necessarily abridged discussion and refer readers to Barocas et al. [8] and Corbett-Davies and Goel [35] for a comprehensive treatment. A large body of work studies *equalizing* a notion of performance over fixed, pre-defined demographic groups for *classification tasks* [32, 45, 7, 57, 66, 117]. Kearns et al. [63, 64], Hébert-Johnson et al. [59] consider finite subgroups defined by a structured class of functions over Z , and study methods of equalizing performance across them. By contrast, our approach instantiates Rawls’ theory of distributive justice [88, 89], where we consider the allocation of the loss $\ell(\cdot; \cdot)$ as a resource. Rawls’ difference principle maximizes the welfare of the worst-off group and provides incentives for groups to maintain the status quo [88]. Similarly, Hashimoto et al. [58] studied negative feedback loops generated by user retention—they use a more conservative notion of worst-case loss than ours—as poor performance on a currently underrepresented user group can have long-term consequences.

The long line of works on distributionally robust optimization (DRO) aims to *train models* to perform well under distribution shifts. Previous approaches considered finite-dimensional worst-case regions such as constraint sets [37, 52, 5] and those based on notions of distances for probability measures such as f -divergences and likelihood ratios [13, 115, 14, 75, 74, 81, 72, 42, 41, 73], Levy-Prokhorov [43], Wasserstein distances [44, 99, 16, 15, 17, 76, 48, 16, 111, 19, 28, 49, 47, 18, 50, 114], and integral probability metrics based on reproducing kernels [105, 122, 113, 120]. The distribution shifts considered in these approaches are modeled after mathematical convenience and are often difficult to interpret. As a result, optimizing models under worst-case performance often results in overly conservative models and these approaches do not currently scale to modern large-scale NLP or vision applications, as those models could have hundreds of thousands of parameters, posing great computational challenges when DRO methods are applied. In this work, we approach distributional robustness from a different angle: instead of robust training, we study the problem of *evaluation* – given a model, can we certify its robustness properties in any way?

In particular, our work is most closely related to Duchi et al. [40], who proposed algorithms for *training* models with respect to the worst-case subpopulation performance (1.2), a more ambitious goal than our narrower viewpoint of *evaluating* model performance pre-deployment. Their (full-batch) training procedure requires solving a convex program with n^2 variables per gradient step, which

is often prohibitively expensive. Furthermore, training with respect to the worst-case conditional risk $\mathbb{E}[\ell(\theta(X); Y) | Z]$ does not scale to deep networks that can overfit to the training data [97]. By contrast, our evaluation perspective aims to take advantage of the rapid progress in deep learning. We build scalable evaluation methods that apply to arbitrary models, which allows leveraging state-of-the-art engineered approaches for training $\theta(\cdot)$. Our narrower focus on evaluation allows us to provide convergence rates that scale advantageously with the dimension of Z , compared to the nonparametric $O_p(n^{-1/d})$ rates for training [40]. Recently, Jeong and Namkoong [62] studied a similar notion of worst-case subpopulation performance in causal inference.

As we note later, our worst-case subpopulation performance gives the usual conditional value-at-risk for $\mathbb{E}[\ell(\theta(X); Y) | Z]$, a classical tail risk measure. Tail-risk estimation has attracted great interest: Wozabal and Wozabal [119], Pflug and Wozabal [84] derive asymptotic properties of plug-in estimates of coherent, law-invariant risk functionals, and Belomestny and Krätschmer [12], Guigues et al. [55] derive central limit results. A distinguishing aspect of our work is the unobservability of the conditional risk $\mathbb{E}[\ell(\theta(X), Y) | Z]$, which necessitates a shift to a *semiparametric estimation* paradigm. To tackle this challenge, we derive a debiased approach to tail-risk estimation and provide both asymptotic and finite-sample convergence guarantees. Concurrent to an earlier conference version of this work, Subbaswamy et al. [106] study a similar problem and propose another estimator different from ours. They claim their estimator is debiased and cite recent work [62] as inspiration; but to our knowledge, their estimator is not debiased as they incorrectly apply Jeong and Namkoong [62]’s main insights and we note important errors in their proof (Section 3 for a detailed discussion). In addition to this difference, all finite-sample convergence guarantees and connections to coherent risk measures are new in this work.

Our work significantly expands on the earlier conference version. We develop a debiased estimator instead of a plug-in estimator, a standard approach in the semiparametric statistics literature that corrects for the first-order error in estimating the nuisance parameter $z \mapsto \mathbb{E}[\ell(\theta(X); Y) | Z = z]$. We derive a central limit result for our debiased estimator in Section 4.1, showing that it is possible to achieve standard \sqrt{n} -rates of convergence even when the fitted $z \mapsto \hat{\mu}(z)$ converge at a slower $n^{-1/3}$ rate. We also extend our prior finite-sample concentration guarantees over uniformly bounded losses to heavy-tailed losses in Section 4.4. Finally, we study a natural extension of our worst-case subpopulation performance where we allow the subpopulation proportion α to be stochastic. We connect this generalized worst-case subpopulation performance to the vast literature on distributional robustness and coherent risk measures in Section 6.

2 Methodology

We begin by contrasting our approach to standard alternatives that consider pre-defined, fixed demographic groups [80]. Identifying disadvantaged subgroups a priori is often challenging as they are determined by *intersections* of multiple demographic variables. To illustrate such complex intersectionality, consider a drug dosage prediction problem for Warfarin [34], a common anti-coagulant (blood thinner). Taking the best prediction model for the optimal dosage on this dataset based on genetic, demographic and clinical factors [34], we present the squared error on the root dosage. In Figure 1, when age and race are considered *simultaneously* instead of *separately*, subpopulation performances vary significantly across intersectional groups.

The worst-case subpopulation performance (1.2) automatically accounts for latent intersectionality. It is agnostic to demographic groupings and allows considering infinitely many subpopulations




Figure 1. Conditional risk $\mu^*(Z) = \mathbb{E}[(Y - \theta(X))^2 | Z]$. Here $Z = \text{age}$ on the left panel, $Z = \text{race}$ in the center, and $Z = (\text{age}, \text{race})$ on the right. A = Asian, B = Black, U = Unknown, W = White.

that represent at least α -fraction of the training population P . By allowing the modeler to select arbitrary protected attributes Z , we are able to consider potentially complex subpopulations. For example, Z can even be defined with respect to a natural language description of the input X . The choice of Z —and subsequent worst-case subpopulation performance (1.2) of the conditional risk $\mu^*(Z) = \mathbb{E}[\ell(\theta(X); Y) | Z]$ —interpolates between the most conservative notion of subpopulations (when $Z = (X, Y)$) and simple counterparts defined over a single variable.

The choice of the subpopulation size α should be informed by domain knowledge—desired robustness of the system—and the dataset size relative to the complexity of Z . Often, proxy groups can be used for selecting α . If we wish to ensure good performance over patients of all races aged 50 years or older, we can choose α to be the proportion of the least represented (*race, age* ≥ 50) group—this leads to $\alpha = 5\%$ in the Warfarin data. The corresponding worst-case subpopulation performance (1.2) guarantees good performance over all groups of similar size.

When it is challenging to commit to a specific subpopulation size, it may be natural to postulate a *maximum level of acceptable loss* $\bar{\ell}$. To measure the robustness of a model, we define the smallest subpopulation size α^* for which the worst-case subpopulation performance is acceptable

$$\alpha^* := \inf\{\alpha : W_\alpha^* \leq \bar{\ell}\}. \quad (2.1)$$

This provides a *certificate of robustness*: if α^* is large, then θ is brittle against even majority subpopulations; if it is sufficiently small, then the model $\theta(X)$ performs well on underrepresented subpopulations.

We now derive estimators for the worst-case subpopulation performance (1.2) and the certificate of robustness (2.1), based on i.i.d. observations $(X_i, Y_i, Z_i)_{i=1}^n \sim P$. We assume our observations are independent from the data used to train the model $\theta(\cdot)$.

Dual reformulation The worst-case subpopulation performance (1.2) is unwieldy as it involves an infinite dimensional optimization problem over probabilities. Instead, we use its dual reformulation for tractable estimation.

We denote $(\cdot)_+ = \max(\cdot, 0)$, and denote by $W_\alpha(\mu)$ the worst-case subpopulation performance for a function $\mu(Z)$ (so that $W_\alpha^* = W_\alpha(\mu^*)$). Let $P_{1-\alpha}^{-1}(\mu)$ denote the $(1 - \alpha)$ -quantile of $\mu(Z)$.

Lemma 1 (Shapiro et al. [103, Theorem 6.2] and Rockafellar and Uryasev [92]). *If $\mathbb{E}[\mu(Z)_+] < \infty$, then for $\alpha \in (0, 1)$,*

$$W_\alpha(\mu) := \sup_{Q_Z \in \mathcal{Q}_\alpha} \mathbb{E}_{Z \sim Q_Z} [\mu(Z)] = \inf_{\eta \in \mathbb{R}} \left\{ \frac{1}{\alpha} \mathbb{E}_P (\mu(Z) - \eta)_+ + \eta \right\} = \frac{1}{\alpha} \int_0^\alpha P_{1-t}^{-1}(\mu) dt. \quad (2.2)$$

The infimum is attained at $\eta = P_{1-\alpha}^{-1}(\mu)$. Moreover, if $\mu(Z)$ has no probability mass at $P_{1-\alpha}^{-1}(\mu)$, then $W_\alpha(\mu) = \mathbb{E}[\mu(Z) \mid \mu(Z) \geq P_{1-\alpha}^{-1}(\mu)]$.

The dual (2.2) shows W_α^* is a tail-average of $\mu^*(Z)$, a popular risk measure known as the conditional value-at-risk (CVaR) in portfolio optimization [92]. The dual optimum is attained at the $(1 - \alpha)$ -quantile of the $\mu^*(Z)$ [93, Theorem 10], giving the worst-case subpopulation

$$\tau^*(Z) = \frac{1}{\alpha} \mathbf{1} \{ \mu^*(Z) \geq P_{1-\alpha}^{-1}(\mu^*) \}. \quad (2.3)$$

3 Estimation

A key challenge in estimating W_α^* is that we can only observe losses $\ell(\theta(X_i); Y_i)$ and never observe the conditional risk $\mu^*(\cdot)$ (1.3). To estimate $\mu^*(\cdot)$, we can solve an empirical approximation to the loss minimization problem

$$\underset{\mu \in \mathcal{H}}{\text{minimize}} \quad \mathbb{E} \left[(\ell(\theta(X); Y) - \mu(Z))^2 \right] \quad (3.1)$$

for some model class \mathcal{H} (class of mappings $\mathcal{Z} \rightarrow \mathbb{R}$). The loss minimization formulation (3.1) allows the use of any machine learning estimator, as well as standard tools for model selection (e.g. cross validation). Denoting by $\bar{\mu}$ a minimizer of (3.1), we may have $\bar{\mu}(\cdot) \neq \mu^*(\cdot) = \mathbb{E}[\ell(\theta(X); Y) \mid \cdot]$ if the model class \mathcal{H} is not sufficiently rich. In the following section, we provide guarantees that scale with the misspecification error $\bar{\mu} - \mu^*$.

As $\mu^*(\cdot)$ must be estimated, this yields a *semiparametric* tail-risk estimation problem: $\mu^*(\cdot)$ can be a complex nonparametric function, but ultimately we are interested in evaluating a one-dimensional statistical functional $W_\alpha(\mu^*)$. A natural plug-in approach is to split the data into auxiliary and main samples, where we first estimate $\hat{\mu}$ using a sample average approximation of the problem (3.1) on the auxiliary sample S_1 . On the main sample S_2 , we can estimate the worst-case subpopulation performance using the dual

$$\widehat{W}_\alpha(\mu) := \inf_{\eta} \left\{ \frac{1}{\alpha |S_2|} \sum_{i \in S_2} (\mu(Z) - \eta)_+ + \eta \right\}. \quad (3.2)$$

The final plug-in estimator is given by $\widehat{W}_\alpha(\hat{\mu})$.

The plug-in estimator is suboptimal since it does not take into account the potential error incurred by using the approximation $\hat{\mu}$ in the final step. To build intuition on this semiparametric error, we provide a heuristic analysis of $W_\alpha(\hat{\mu}) - W_\alpha(\mu^*)$ using a first-order Taylor expansion, ignoring the statistical error incurred in the second stage for a moment. Abusing notation, the functional $P \mapsto W_\alpha(P) := W_\alpha(\mathbb{E}_P[\ell(\theta(X); Y) \mid Z])$ can be seen to be suitably differentiable so that there is a continuous linear map \dot{W}_α on the space of square integrable functions such that

$$\frac{d}{dr} (W_\alpha(P + r(\bar{P} - P)) - W_\alpha(P)) = \dot{W}_\alpha(\bar{P} - P).$$

By the Riesz Representation Theorem, there is a function $\nabla W_\alpha(X, Y, Z; P)$ such that

$$\dot{W}_\alpha(\bar{P} - P) = \int \nabla W_\alpha(X, Y, Z; P) d(\bar{P} - P),$$

where we assume ∇W_α has mean zero without loss of generality; the random variable ∇W_α is often referred to as the pathwise derivative or the efficient influence function of the functional W_α [82].

Under appropriate regularity conditions, we can apply the chain rule for influence function calculus [82, 65] and use Danskin's theorem [22, Theorem 4.13] to calculate functional gradients. Noting that the dual optimum is attained at the $(1 - \alpha)$ -quantile $P_{1-\alpha}^{-1}(\mu_P)$ by first order conditions, we have

$$\begin{aligned} \nabla W_\alpha(X, Y, Z; P) &= \frac{1}{\alpha} (\mu_P(Z) - P_{1-\alpha}^{-1}(\mu_P))_+ - \frac{1}{\alpha} \mathbb{E}_P[(\mu_P(Z) - P_{1-\alpha}^{-1}(\mu_P))_+] \\ &\quad + \tau^*(Z)(\ell(\theta(X); Y) - \mu_P(Z)) \end{aligned}$$

where we use

$$\mu_P(Z) := \mathbb{E}_P[\ell(\theta(X); Y) \mid Z] \quad \text{and} \quad \tau_P(Z) := \alpha^{-1} \mathbf{1} \{ \mu_P(Z) \geq P_{1-\alpha}^{-1}(\mu_P) \}.$$

Taking a first-order Taylor expansion around the learned parameter $\hat{\mu}$, we arrive at

$$W_\alpha(\hat{\mu}) - W_\alpha(\mu^*) = -\mathbb{E}_P[\hat{\tau}(Z)(\ell(\theta(X); Y) - \hat{\mu}(Z))] + \text{Rem}_2 \quad (3.3)$$

where $\hat{\tau}_k(\cdot)$ is an estimator of the worst-case subpopulation (2.3) and Rem_2 is a second-order remainder term.

The Taylor expansion (3.3) provides a natural approach to correcting the first-order error of the plug-in estimator—a standard approach called *debiasing* in the semiparametric statistics literature [83, 82, 31]. Instead of the plug-in (3.2), the *debiased* estimator on the main sample S_2 is given by

$$\inf_{\eta} \left\{ \frac{1}{\alpha |S_2|} \sum_{i \in S_2} (\hat{\mu}(Z_i) - \eta)_+ + \eta \right\} + \frac{1}{|S_2|} \sum_{i \in S_2} \hat{\tau}(Z_i)(\ell(\theta(X_i); Y_i) - \hat{\mu}(Z_i)).$$

The the debiased estimator automatically achieves a second-order error and as we show in Section 4.1, it achieves parametric rates of convergence even when $\hat{\mu}$ converges more slowly.

Cross-fitting procedure To utilize the entire sample, we take a cross-fitting approach [31] where we partition the data into K folds. We assign a single fold as the main data and use the rest as auxiliary data, switching the role of the main fold to get K separate estimators; the final estimator is simply the average of the K versions as outlined in Algorithm 1. We take $\hat{\tau}_k(z) := \frac{1}{\alpha} \mathbf{1} \{ \hat{\mu}_k(z) \geq \hat{q} \}$, where \hat{q} is an estimator of $P_{1-\alpha}^{-1}(\hat{\mu}_k)$ based on the auxiliary data. The quantile estimator \hat{q} can be computed using *unsupervised* observations Z , which is typically cheap to collect. To estimate the threshold subpopulation size α^* , we simply take the cross-fitted version of the plug-in estimator

$$\hat{\alpha}_k := \inf \{ \alpha : \hat{W}_{\alpha,k}(\hat{\mu}_k) \leq \bar{\ell} \}. \quad (3.4)$$

Since $\alpha \mapsto \hat{W}_{\alpha,k}(\hat{\mu}_k)$ is decreasing, the threshold can be efficiently found by a simple bisection search.

In Section 4.1, we show that our cross-fitted augmented estimator $\hat{\omega}_\alpha$ has asymptotic variance

$$\sigma_\alpha^2 := \frac{1}{\alpha^2} \text{Var} \left((\mu^*(X) - P_{1-\alpha}^{-1}(\mu^*))_+ \right) + \text{Var}(\kappa(D; \mu^*, \tau^*)). \quad (3.5)$$

In particular, letting z_δ be the $(1 - \delta/2)$ -quantile of a standard normal distribution, our result gives confidence intervals with asymptotically exact coverage $\mathbb{P}(\mathbf{W}_\alpha^* \in [\hat{\omega}_\alpha \pm z_\delta \hat{\sigma}_\alpha / \sqrt{n}]) \rightarrow 1 - \delta$.

Algorithm 1 Cross-fitting procedure for estimating worst-case subpopulation performance (1.2)

- 1: **INPUT:** Subpopulation size α , model class \mathcal{H} , K -fold partition $\cup_{k=1}^K I_k = [n]$ of $\{(X_i, Y_i, Z_i)\}_{i=1}^n$
s.t. $|I_k| = \frac{n}{K}$
- 2: **For** $k \in [K]$
- 3: **Estimate nuisance parameters** Using the data $\{(X_i, Y_i, Z_i)\}_{i \in I_k^c}$, fit estimators
- 4: 1. Solve $\hat{\mu}_k \in \operatorname{argmin}_{\mu \in \mathcal{H}} \sum_{i \in I_k^c} (\ell(\theta(X_i); Y_i) - \mu(Z_i))^2$.
- 5: 2. $\hat{\tau}_k(z) := \frac{1}{\alpha} \mathbf{1} \{\hat{\mu}_k(z) \geq \hat{q}_k\}$, where \hat{q}_k is an estimator of $P_{1-\alpha}^{-1}(\hat{\mu}_k)$ on I_k^c
- 6: **Compute augmented estimator** Using the data $\{(X_i, Y_i, Z_i)\}_{i \in I_k}$, compute

$$\begin{aligned} \hat{\omega}_{\alpha,k} &:= \inf_{\eta} \left\{ \frac{1}{\alpha} \mathbb{E}_{Z \sim \hat{P}_k} (\hat{\mu}_k(Z) - \eta)_+ + \eta \right\} + \mathbb{E}_{(X,Y,Z) \sim \hat{P}_k} [\hat{\tau}_k(Z) (\ell(\theta(X); Y) - \hat{\mu}_k(Z))] \\ \hat{\sigma}_{\alpha,k}^2 &:= \frac{1}{\alpha^2} \operatorname{Var}_{Z \sim \hat{P}_k} (\hat{\mu}_k(Z) - \hat{q}_k)_+ + \operatorname{Var}_{(X,Y,Z) \sim \hat{P}_k} (\hat{\tau}_k(Z) (\ell(\theta(X); Y) - \hat{\mu}_k(Z))) \end{aligned}$$

- 7: **Return** Estimator $\hat{\omega}_\alpha = \frac{1}{K} \sum_{k \in [K]} \hat{\omega}_{\alpha,k}$, and variance estimate $\hat{\sigma}_\alpha^2 = \frac{1}{K} \sum_{k \in [K]} \hat{\sigma}_{\alpha,k}^2$
-

So far, we considered an estimand that involves an optimization problem over the dual variable η . We derived a debiased estimation approach accounting for errors in estimating μ_P , which gave us a formula involving (μ_P, τ_P) . Later, we will show that our final estimator is indeed debiased with respect to the nuisance parameters (μ_P, τ_P) . Concurrent to an earlier conference version of this work, Subbaswamy et al. [106] study a similar setting and propose another estimator different from ours. In contrast to our approach, they treat (μ_P, η_P) as nuisance parameters and define the estimand as

$$T(P; \mu, \eta) = \frac{1}{\alpha} \mathbb{E}_P (\mu(Z) - \eta)_+ + \eta.$$

As is evident here, this approach requires debiasing with respect to *both* μ and η .

Given black-box estimates $(\hat{\mu}, \hat{\eta})$, Subbaswamy et al. [106] estimate the following expression on a separate fold

$$\frac{1}{\alpha} \mathbb{E}_{Z \sim P} (\hat{\mu}(Z) - \hat{\eta})_+ + \hat{\eta} + \frac{1}{\alpha} \mathbb{E}_{Z \sim P} [\mathbf{1} \{\hat{\mu}(Z) \geq \hat{\eta}\} (\ell(\theta(X); Y) - \hat{\mu}(Z))].$$

They claim this approach is debiased and cite Jeong and Namkoong [62]’s related approach as the justification. However, Jeong and Namkoong [62]—which we also draw inspiration from—follow an exactly analogous approach as ours involving just (μ_P, τ_P) and do not consider η as a separate nuisance parameter. This difference results in an erroneous asymptotic result in Subbaswamy et al. [106, Theorem 1]; as far as we can tell, their proof of Neyman orthogonality is incorrect and their proposed estimator does not appear debiased. For example, they cite Danskin’s theorem despite their functional $T(P; \mu, \eta)$ not involving an optimization functional.

4 Convergence guarantees

To *rigorously* verify the robustness of a model prior to deployment, we present convergence guarantees for our estimator (Algorithm 1). We begin by proving a central limit result for our estimator,

showing that debiasing allows us to achieve standard \sqrt{n} -rates of convergence even when the fitted $\hat{\mu}_k$ converge at a slower rate. Since asymptotic guarantees ignore the dimensionality of Z , we then turn to finite sample concentration guarantees. Finally, we provide convergence guarantees for our estimator (3.4) for the certificate of robustness (2.1) in the appendix (Section E).

4.1 Asymptotics

In contrast to the literature on debiased estimation, our estimand is nonlinear in P , which requires a different proof approach than what is standard (e.g., [31]). Our asymptotic result proof approach is inspired by Jeong and Namkoong [62], using standard tools in empirical process theory.

Assumption A. *Bounded residuals* $\mathbb{E}[\ell(\theta(X); Y)^2] + \|\mathbb{E}[(\ell(\theta(X); Y) - \mu^*(Z))^2 \mid Z]\|_{L^\infty(\mathcal{X})} < \infty$

Assumption B. *Let $\|\hat{\mu}_k - \mu^*\|_{L^1(\mathcal{X})} \xrightarrow{a.s.} 0$, and let there exist an envelope function $\tilde{\mu} : \mathcal{Z} \rightarrow \mathbb{R}$ satisfying $\mathbb{E}[\tilde{\mu}(Z)^2] < \infty$ and $\max(|\hat{\mu}_{0,k}|, |\hat{\mu}_{1,k}|) \leq \tilde{\mu}$. There exists $\delta_n, \Delta_n \downarrow 0$, and $M > 0$ such that with probability at least $1 - \Delta_n$, for all $k \in [K]$, $|\hat{\tau}_k| \leq M$, $\|\hat{\mu}_k - \mu^*\|_{L^\infty(\mathcal{X})} \leq \delta_n n^{-1/3}$, and $|\hat{q}_k - P_{1-\alpha}^{-1}(\hat{\mu}_k)| \leq \delta_n n^{-1/3}$.*

To estimate the tail-average (2.2) (recall Lemma 1), we need to estimate the quantile $P_{1-\alpha}^{-1}(\mu^*)$. We assume that for functions around μ^* , their positive density exists at the $(1 - \alpha)$ -quantile [110, Chapter 3.7]. Let \mathcal{U} be a set of (measurable) functions $\mu : \mathcal{Z} \rightarrow \mathbb{R}$ such that

$F_{r,\mu}$, the cumulative distribution of $(\mu^* + r(\mu - \mu^*))(Z)$, is uniformly differentiable in $r \in [0, 1]$ at $P_{1-\alpha}^{-1}(\mu^* + r(\mu - \mu^*))$, with a positive and uniformly bounded density. Formally, if we let $q_{r,\mu} := P_{1-\alpha}^{-1}(\mu^* + r(\mu - \mu^*))$, then for each $r \in [0, 1]$, there is a positive density $f_{r,\mu}(q_{r,\mu}) > 0$ such that

$$\lim_{t \rightarrow 0} \sup_{r \in [0,1]} \left| \frac{1}{t} (F_{r,\mu}(q_{r,\mu} + t) - F_{r,\mu}(q_{r,\mu})) - f_{r,\mu}(q_{r,\mu}) \right| = 0. \quad (4.1)$$

We require this holds for our estimators $\mu = \hat{\mu}_k$ with high probability.

Assumption C. $\exists \Delta'_n \downarrow 0$ s.t. with probability at least $1 - \Delta'_n$, $\hat{\mu}_k \in \mathcal{U}$ for all $k \in [K]$.

Our debiased estimator $\hat{\omega}_\alpha$ enjoys central limit rates with the influence function

$$\psi(X, Y, Z) := \left[\frac{1}{\alpha} (\mu^*(Z) - P_{1-\alpha}^{-1}(\mu^*))_+ + P_{1-\alpha}^{-1}(\mu^*) \right] - W_\alpha^* + \tau^*(Z)(\ell(\theta(X); Y) - \mu^*(Z)). \quad (4.2)$$

See Section A for the proof of the following central limit result.

Theorem 1. *Under Assumptions A-C, $\sqrt{n}(\hat{\omega}_\alpha - W_\alpha^*) \overset{d}{\rightsquigarrow} N(0, \text{Var}(\psi(X, Y, Z)))$.*

4.2 Concentration using the localized Rademacher complexity

We give finite-sample convergence at the rate $O_p(\sqrt{\mathfrak{Comp}_n(\mathcal{H})/n})$, where $\mathfrak{Comp}_n(\mathcal{H})$ is the localized Rademacher complexity [9] of the model class \mathcal{H} for estimating the conditional risk $\mu^*(Z)$. We restrict attention to nonnegative and uniformly bounded losses, as is conventional in the literature.

Assumption D. *There is a B such that $\ell(\theta(X); Y) \in [0, B]$, and $\mu(Z) \in [0, B]$ a.s. for all $\mu \in \mathcal{H}$.*

Throughout this section, we do not stipulate well-specification, meaning that we allow the conditional risk $\mu^*(Z) = \mathbb{E}[\ell(\theta(X); Y) | Z]$ not to be in the model class \mathcal{H} .

To characterize the finite-sample convergence behavior of our estimator $\widehat{\omega}_\alpha$, we begin by decomposing the error of the augmented estimator $\widehat{\omega}_{\alpha,k}$ on each fold I_k into two terms relating to the two stages in Algorithm 1. Recalling the notation in Eq. (2.2) (so that $W_\alpha^* = W_\alpha(\mu^*)$), we have

$$W_\alpha^* - \widehat{\omega}_{\alpha,k} = \underbrace{T(P; \mu^*, \tau^*) - T(P; \widehat{\mu}_k, \widehat{\tau}_k)}_{(a): \text{ first stage}} + \underbrace{T(P; \widehat{\mu}_k, \widehat{\tau}_k) - T(\widehat{P}_k; \widehat{\mu}_k, \widehat{\tau}_k)}_{(b): \text{ second stage}},$$

because $W_\alpha^* = T(P; \mu^*, \tau^*)$ and $\widehat{\omega}_{\alpha,k} = T(P; \widehat{\mu}_k, \widehat{\tau}_k)$. To bound term (b), we prove concentration guarantees for estimators of the dual (2.2) (see Proposition 8 in Appendix B.1). To bound term (a), we use a localized notion of the Rademacher complexity.

Formally, for $\xi_1, \dots, \xi_n \in \Xi$ and i.i.d. random signs $\varepsilon_i \in \{-1, 1\}$ (independent of ξ_i), recall the standard notion of (empirical) Rademacher complexity of $\mathcal{G} \subseteq \{g : \Xi \rightarrow \mathbb{R}\}$

$$\mathfrak{R}_n(\mathcal{G}) := \mathbb{E}_\varepsilon \left[\sup_{g \in \mathcal{G}} \frac{1}{n} \sum_{i=1}^n \varepsilon_i g(\xi_i) \right].$$

We say that a function $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is *sub-root* [9] if it is nonnegative, nondecreasing, and $r \mapsto \psi(r)/\sqrt{r}$ is nonincreasing for $r > 0$. Any (non-constant) sub-root function is continuous, and has a unique positive fixed point. Let $\psi_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a sub-root upper bound on the localized Rademacher complexity $\psi_n(r) \geq \mathbb{E}[\mathfrak{R}_n \{g \in \mathcal{G} : \mathbb{E}[g^2] \leq r\}]$. (The localized Rademacher complexity itself is sub-root.) The fixed point of ψ_n characterizes generalization guarantees [9, 69].

Let $\bar{\mu}$ be the best model in the model class \mathcal{H}

$$\bar{\mu} := \operatorname{argmin}_{\mu \in \mathcal{H}} \mathbb{E}[(\ell(\theta; X, Y) - \mu(Z))^2].$$

Let $\psi_{|I_k^c|}(r)$ be a subroot upper bound on the localized Rademacher complexity around $\bar{\mu}$

$$\psi_{|I_k^c|}(r) \geq 2\mathbb{E} \left[\mathfrak{R}_{|I_k^c|} \{ \mu \in \mathcal{H} : \mathbb{E}[(\mu(Z) - \bar{\mu}(Z))^2] \leq rB^2/4 \} \right]. \quad (4.3)$$

We define $r_{|I_k^c|}^*$ as the fixed point of $\psi_{|I_k^c|}(r)$.

As we show shortly, we bound the estimation error of our procedure using the *square root* of the excess risk in the first-stage problem (3.1)

$$\mathbb{E} \left[(\ell(\theta; X, Y) - \widehat{\mu}_k(Z))^2 | I_k^c \right] - \mathbb{E} \left[(\ell(\theta; X, Y) - \bar{\mu}(Z))^2 \right] \quad (4.4)$$

By using a refined analysis offered by localized Rademacher complexities, we are able to use a fast rate of convergence of $O_p(\mathfrak{Comp}_n(\mathcal{H})/n)$ on the preceding excess risk. In turn, this provides the following $O_p(\sqrt{\mathfrak{Comp}_n(\mathcal{H})/n})$ bound on the estimation error as we prove in Appendix B.2. In the bound, we have made explicit the approximation error term $\|\bar{\mu} - \mu^*\|_{L^2}$. As the model class \mathcal{H} grows richer, there is tension as the approximation error term will shrink, yet the localized Rademacher complexity of \mathcal{H} will grow.

Theorem 2. *Let Assumption D hold. For some constant $C > 0$, for each fold $k \in [K]$, with probability at least $1 - 3\delta$,*

$$|W_\alpha^* - \widehat{\omega}_{\alpha,k}| \leq \frac{CB}{\alpha} \left(\sqrt{r_{(1-K^{-1})n}^*} + \sqrt{\frac{K \log(2/\delta)}{n}} \right) + \frac{2}{\alpha} \|\bar{\mu} - \mu^*\|_{L^2}.$$

By controlling the fixed point r_n^* of the localized Rademacher complexity, we are able to provide convergence of our estimator (6). For example, when \mathcal{H} is a bounded VC-class [110], it is known that its fixed point satisfy [9, Corollary 3.7]

$$r_n^* \asymp \log(n/\text{VC}(\mathcal{H})) \cdot \text{VC}(\mathcal{H})/n,$$

where $\text{VC}(\cdot)$ is the VC-dimension.

4.3 Data-dependent dimension-free concentration

In some situations, it may be appropriate to define subpopulations (Z) over features of an image, or natural language descriptions. For such high-dimensional variables Z and complex model classes \mathcal{H} such as deep networks, the complexity measure \mathbf{Comp}_n is often prohibitively conservative and renders the resulting concentration guarantee meaningless. We provide an alternative concentration result that depends on the size of model class \mathcal{H} only through the out-of-sample error in the first-stage problem (3.1). This finite-sample, data-dependent convergence result depends only on the out-of-sample generalization error for estimating $\mu^*(\cdot)$. In particular, the out-of-sample error can grow smaller as \mathcal{H} gets richer, and as a result of hyperparameter tuning and model selection, it is often very small for overparameterized models such as deep networks. This allows us to construct valid finite-sample upper confidence bounds for the worst-case subpopulation performance (1.2) even when Z is defined over high-dimensional features and \mathcal{H} represent deep networks.

For simplicity, denote

$$\Delta_S(\mu) := \frac{1}{|S|} \sum_{i \in S} (\ell(\theta(X_i); Y_i) - \mu(Z_i))^2. \quad (4.5)$$

for any function $\mu : \mathcal{Z} \rightarrow \mathbb{R}$ on any data set S . We prove the following result in Appendix B.3.

Theorem 3. *Let Assumption D hold. For some constant $C > 0$, for each fold $k \in [K]$, with probability at least $1 - 3\delta$,*

$$|\mathbf{W}_\alpha^* - \widehat{\omega}_{\alpha,k}| \leq \frac{2}{\alpha} \left(\sqrt{[\Delta_{I_k}(\widehat{\mu}_k) - \Delta_{I_k}(\bar{\mu})]_+} + \|\bar{\mu} - \mu^*\|_{L^2} + CB \left(\frac{2K \log(2/\delta)}{n} \right)^{1/4} \right).$$

Moreover, if the model class \mathcal{H} is convex, then $\|\bar{\mu} - \mu^*\|_{L^2}$ can be replaced with $\|\bar{\mu} - \mu^*\|_{L^1}$.

Following convention in learning theory, we refer to our data-dependent concentration guarantee *dimension-free*. For overparameterized model classes \mathcal{H} such as deep networks, the localized Rademacher complexity in Theorem 2 becomes prohibitively large [10, 121]. In contrast, the current result can still provide meaningful finite-sample bounds: model selection and hyperparameter tuning provides low out-of-sample performance in practice, and the difference $\Delta_{I_k}(\widehat{\mu}_k) - \Delta_{I_k}(\bar{\mu})$ can be often made very small. Concretely, it is possible to calculate an upper bound on this term as $\Delta_{I_k}(\bar{\mu})$ is lower bounded by $\min_{\mu \in \mathcal{H}} \Delta_{I_k}(\mu)$.

4.4 Extensions for heavy-tailed loss functions

While it is standard to study uniformly bounded losses when considering finite-sample convergence guarantees, it is nevertheless a rather restrictive assumption. We now show concentration guarantees for broader classes of losses such as sub-Gaussian or sub-exponential ones. Our analysis builds on Mendelson [77]’s framework for heavier-tailed losses based on one-sided concentration inequalities.

Denote as \mathcal{D}_{μ^*} the $L_2(\mathbb{P})$ unit ball centered at μ^* , the residuals $\zeta_i := \ell(\theta(X_i); Y_i) - \mu^*(Z_i)$ and $\varepsilon_i \in \{-1, 1\}$ iid random signs, we define

$$\begin{aligned}\phi_{I_k^c}(s) &:= \sup_{\mu \in \mathcal{H} \cap s\mathcal{D}_{\mu^*}} \left| \frac{1}{\sqrt{|I_k^c|}} \sum_{i \in I_k^c} \varepsilon_i \zeta_i (\mu(Z_i) - \mu^*(Z_i)) \right|, \\ \alpha_{I_k^c}(\gamma, \delta) &:= \inf \left\{ s > 0 : \mathbb{P} \left(\phi_{I_k^c}(s) \leq \gamma s^2 \sqrt{|I_k^c|} \right) \geq 1 - \delta \right\}, \\ \beta_{I_k^c}(\gamma) &:= \inf \left\{ r > 0 : \mathbb{E} \sup_{\mu \in \mathcal{H} \cap s\mathcal{D}_{\mu^*}} \left| \frac{1}{\sqrt{|I_k^c|}} \sum_{i \in I_k^c} \varepsilon_i (\mu(Z_i) - \mu^*(Z_i)) \right| \leq \gamma r \sqrt{|I_k^c|} \right\}, \\ Q_{\mathcal{H}-\mathcal{H}}(u) &:= \inf_{\mu_1, \mu_2 \in \mathcal{H}} \mathbb{P}(|\mu_1 - \mu_2| \geq u \|\mu_1 - \mu_2\|_{L^2}).\end{aligned}$$

Assumption E. *The conditional risk is in the model class \mathcal{H} : $\mu^*(Z) = \mathbb{E}[\ell(\theta(X); Y) \mid Z] \in \mathcal{H}$.*

Assumption F. *The model class \mathcal{H} is closed and convex.*

Assumption G. *The loss function ℓ is sub-Gaussian with parameter B^2 .*

Theorem 4 (Mendelson [77], Theorem 3.1). *Let Assumptions E, F, G hold. Fix $\tau > 0$ for which $Q_{\mathcal{H}-\mathcal{H}}(2\tau) > 0$ and set $\gamma < \tau^2 Q_{\mathcal{H}-\mathcal{H}}(2\tau)/16$. There is a numerical constant $C > 0$ such that for every $\delta \in (0, 1)$ and $k \in [K]$, with probability at least $1 - \delta - \exp(-(1 - K^{-1})nQ_{\mathcal{H}-\mathcal{H}}(2\tau)^2/2)$,*

$$|\mathbb{W}_\alpha(\mu^*) - \widehat{\omega}_{\alpha, k}| \leq \frac{C}{\alpha} \left(\alpha_{I_k^c} \left(\gamma, \frac{\delta}{4} \right) + \beta_{I_k^c} \left(\frac{\tau Q_{\mathcal{H}-\mathcal{H}}(2\tau)}{16} \right) + B \sqrt{\frac{K \log(2/\delta)}{n}} \right) \quad (4.6)$$

For an example of where this result provides a tighter bound than that of Theorem 2, we look at the persistence framework. Let $Z \in \mathbb{R}^m$ be a random vector with independent mean-zero, variance-1 random coordinates. Consider the linear model class $\mathcal{H} = \{ \langle t, \cdot \rangle : \mathbb{E} \|t\|_1 \leq R \}$, and suppose the conditional risk is $\mu^* = \langle t^*, \cdot \rangle + \varsigma$, where $\mathbb{E} \|t^*\|_1 \leq R$ and ς is an independent mean-zero random variable with variance at most B^2 .

Lemma 2 (Mendelson [77], Theorem 4.6). *Let Assumptions D, E, F hold. Fix $B > 1$. There exist constants c_1, c_2 and c_3 that depend only on B for which with probability at least $1 - 2 \exp(-c_3 |I_k^c| v_2 \min\{B^{-2}, R^{-1}\})$,*

$$\left| \mathbb{W}_\alpha(\mu^*) - \widehat{\mathbb{W}}_{\alpha, k}(\widehat{\mu}_k) \right| \leq \frac{C}{\alpha} \left(\sqrt{\max\{v_1, v_2\}} + B \sqrt{\frac{\log(2/\delta)}{|I_k^c|}} \right), \quad (4.7)$$

where we define

$$v_1 = \frac{R^2}{|I_k^c|} \log \left(\frac{2c_1 m}{|I_k^c|} \right) \mathbf{1}\{|I_k^c| \leq c_1 m\} \quad (4.8)$$

$$v_2 = \begin{cases} \frac{RB}{\sqrt{|I_k^c|}} \sqrt{\log \left(\frac{2c_2 m B}{\sqrt{|I_k^c|} R} \right)} & \text{if } |I_k^c| \leq c_2 m^2 B^2 / R^2, \\ \frac{B^2 m}{|I_k^c|} & \text{otherwise.} \end{cases} \quad (4.9)$$




Figure 2: $\hat{W}_{\alpha,k}(\hat{\mu}_1)$ and $W_{\alpha}(\theta)$ from simulation experiments with $\alpha = 0.3$

5 Experiments

We begin by verifying the asymptotic convergence of our proposed two-stage estimator with a simulation experiment on a classification task. We then demonstrate the practical applicability of our work through two model selection problems: a long-standing precision medicine problem affecting millions of patients, and a land-use image classification problem involving large-scale neural network models that is of particular interest to the machine learning community. We show that, by allowing certifying models’ performance over worst-case subpopulations, our procedure is able to instill insights into model selection by predicting model performance on unseen data, and, if necessary, raising awareness on brittle models without the knowledge of out-of-distribution samples.

5.1 Simulation

We illustrate the asymptotic convergence of our two-stage estimator $\hat{W}_{\alpha,k}(\hat{\mu}_1)$ of the worst-case subpopulation performance $W_{\alpha}(\theta)$. We conduct a binary classification experiment where we randomly generate and fix two vectors $\theta, \theta_0^* \in \mathbb{R}^d$ on the unit sphere. The data-generating distribution is given by $X \stackrel{iid}{\sim} \mathbf{N}(\gamma, \Sigma)$ and

$$Y \mid X = \begin{cases} \text{sgn}(X^\top \theta_0^*) & X^1 \leq z_{0.95} = 1.645 \\ -\text{sgn}(X^\top \theta_0^*) & \text{otherwise.} \end{cases}$$

In this data-generating distribution, there is a drastic difference between subpopulations generated by $X^1 \leq z_{0.95}$ and $X^1 > z_{0.95}$; typical prediction models will perform poorly on the latter rare group. The loss function is taken to be the hinge loss $\ell(\theta; x, y) = [1 - y \cdot \theta^\top x]_+$, where $y \in \{\pm 1\}$. We take the first covariate X^1 as our protected attribute Z . Let $d = 5$, $\Sigma = \mathbf{I}_5$, $\gamma = 0$.

We fix $\alpha = 0.3$. To analyze the asymptotic convergence of our two-stage estimator, for sample size ranging in 1,000 to 256,000 doubling each time, we run 40 repeated experiments of the estimation procedure on simulated data. We split each sample evenly into S_1 and S_2 and using gradient boosted trees in the package XGBoost [27] to estimate the conditional risk. On a log-scale, we report the mean estimate across random runs in Figure 2 alongside error bars. To compute the true worst-case subpopulation performance $W_{\alpha}(\mu^*)$ of the conditional risk $\mu^*(X^1)$, we first run a Monte Carlo simulation for 150,000 copies of $X^1 \sim \mathbf{N}(0, 1)$. For each sampled X^1 , we generate 100,000 copies of $(X^2, X^3, X^4, X^5) \sim \mathbf{N}(\mathbf{0}, \mathbf{I}_4)$ independent of X^1 and compute the mean loss among them to approximate the conditional risk $\mu^*(X^1)$. Finally, we approximate $W_{\alpha}(\mu^*)$ using the empirical

distribution of $\mu^*(\cdot)$, obtaining 6.47×10^{-1} . We observe convergence toward the true value as sample size n grows, verifying the consistency of our two-stage estimator $\widehat{W}_{\alpha,k}(\widehat{\mu}_1)$.

5.2 Warfarin Optimal Dosage

Precision Medicine—an emerging approach that treats diseases at a personal level incorporating individual variability in genes—has attracted great attention in recent years. One important area precision medicine intends to tackle is optimal dosage: given the patients’ individual characteristics such as demographic, genetic, and symptomatic information, is it possible to design an automated algorithm to predict the optimal dosage for the patients? Unfortunately, this task is often presented with much difficulty. In the case of Warfarin — one of the most widely used anticoagulant agents — its optimal dosage can differ substantially across genetics, demographics, and existing conditions of the patients by a factor as much as 10 [34]. Traditionally, physicians often determine the dosage through trial and error, but this large variation makes the appropriate dosage hard to establish, and an incorrect dosage can lead to highly undesirable side effects. It is therefore important to develop a more reliable method to help determine the optimal dosage for the patients. Furthermore, to ensure fair treatment to all patients, it is imperative that the model performs uniformly well over all subpopulations.

We use the Warfarin optimal dosage prediction problem to illustrate how our metric can be used as a robustness certificate of the model that informs model selection. We consider the Pharmacogenetics and Pharmacogenomics Knowledge Base dataset where the Warfarin optimal dosage is found through trial and error by clinicians. The dataset consists of 4,788 patients (after excluding missing data) with features representing *demographics, genetic markers, medication history, pre-existing conditions, and reason for treatment*. It has been observed empirically in Consortium [34] that a linear model outperforms a number of more complicated modeling approaches (including kernel methods, neural networks, splines, boosting) for predicting the optimal dosage, at least based on average prediction accuracy on the out-of-sample test set.

Comparable average case performance does not guarantee similar worst case performance; in dosage prediction problems fair treatment to all groups, including the underrepresented groups, is essential [25, 87, 53, 2]. With far fewer model parameters than other more expressive models, are linear models truly on par with other models in ensuring uniform good performance overall all subgroups?

To answer this question, we evaluate and compare the worst-case subpopulation performance of different models over $Z = X$. We take the entire feature vector including all available demographic and genetic information as core attributes defining the subpopulations. By taking such a core attribute vector we are being extra conservative, but this decision is motivated by the nature of the optimal dosage prediction problem in that one shall make decisions based on worst case performance guarantees for all patients, irrespective of their demographics, genetic markers, pre-existing conditions, etc.

More specifically and following the approach in Consortium [34], we take the root-dosage as our outcome Y , and consider minimizing the squared loss function

$$\ell(\theta(X); Y) = (Y - \theta(X))^2.$$

We consider four popular models common used in practice: *Linear Regression, Ridge Regression, XGBoost, Random Forests*. Past literature has shown that Linear Regression model does not underperform other more expressive models. If we can certify that Linear Regression model is at

least as robust as other models, then we provide a certificate to the Linear Regression model and one would naturally choose the linear models over others thanks to their simplicity and interpretability.




Figure 3. Worst-case subpopulation performance $W_\alpha(\theta)$, where $W_{1.0}(\theta) = \mathbb{E}[\ell(\theta(X); Y)]$. Results are averaged over 50 random seeds with error bars corresponding to 95% confidence interval over the random runs.

Figure 3 plots our metric against different choice of subpopulation size α , for the four models considered. We observe that the performance of linear model closely matches that of other more expressive models, and the trend holds over a range of different subpopulation sizes, even for small $\alpha = 5\%$. Our finding thus instills confidence in the linear regression model: our diagnostic is able to certify its advantageous performance even on tail subpopulations despite it is the simplest among the four models.

At the same time, our diagnostic raises concerns about poor tail subpopulation performance: all models suffer from significant performance deterioration on small subpopulation sizes (e.g. $\alpha = 5\%$), and the prediction loss is as much as six times worse than the average-case performance. This observation shows that achieving uniformly good performance across subgroups is a challenging task in the Warfarin example, and more attention is needed to address this significant deterioration of performance on the worst-case subpopulation.

5.3 Functional Map of the World (FMoW)

Training a robust model has attracted great interest in both operations research and machine learning communities. However, existing direct robustness treatment method is generally not scalable to modern day machine learning or deep learning experiments, where models such as deep neural networks have hundreds to thousands parameters. By only focusing on evaluation, our metric is able to evaluate and certify the robustness of these large-scale models that are otherwise hard to robustly train on. In the second experiment we look at an image classification application of our method and illustrate how our procedure is useful in evaluating robustness properties of SOTA, large-scale models such as neural networks.

5.3.1 Background

We study a satellite image classification problem [33]. Satellite images can impact economic and environmental policies globally by allowing large-scale measurements on poverty [1], population changes, deforestation, and economic growth [56]. It is therefore important to implement automated approaches that allow scientists and sociologists to provide continuous monitoring of land usage and analyze data from remote regions at a relatively low cost with models that perform reliably across time and space.

We consider the Functional Map of the World (FMoW) dataset [33] comprising of satellite images, where the goal is to predict building / land usage categories (62 classes). We take a recently published variant of this dataset in Huang et al. [61], Koh et al. [68], FMoW-WILDS, that is designed specifically for evaluating model performance under *temporal* and *spatial* distribution shifts. Due to the scale of the dataset (>10K images), traditional robust training approaches do not scale; we will show that our evaluation metric is fully scalable, and our metric provides insights on the performances of SOTA deep learning neural network models on future unseen data.

We take the SOTA models reported in Koh et al. [68] of FMoW as our benchmark. These models are deep neural networks that benefit from *transfer learning*: unlike traditional models that are trained from scratch to solve the problem on-hand, these benchmark models are built on *pre-trained models* — existing models whose parameters have been pre-trained on other, usually massive datasets such as *ImageNet* — and then adapted to the present problem through parameter fine-tuning. It is observed that transfer models exhibit stellar performance across numerous datasets and matching, if not surpassing, that of SOTA models trained from scratch for the specific problem. However, although these SOTA benchmark models on FMoW achieve satisfactory out-of-sample ID accuracy on the past data, all suffer from significant performance drop on the future OOD data, suggesting that these benchmark models are not robust to temporal shifts.

Recently, Radford et al. [86] published a self-supervising model, *Contrastive Language-Image Pre-training (CLIP)*, that has been observed to exhibit many exciting robustness properties. *CLIP* models are pre-trained on 400M image-text pairs using natural language supervision and contrastive losses. The pre-training data for *CLIP* is 400 times bigger than *ImageNet*, the standard pre-training dataset used for FMoW benchmark models, and Radford et al. [86] have observed that *CLIP* exhibits substantial *relative robustness gains* over other methods on natural distribution shifts of *ImageNet*. Previous work in Wortsman et al. [118] has shown significant performance gains using *CLIP*-based models on FMoW.

Motivated by the challenges and opportunities, we take the perspective of a practitioner, presented with different but comparable models and with historical data, wishing to select the “best” model that perform well across all subgroups into the future. More concretely, we will illustrate the use of our procedure on models that achieve similar average accuracy and loss and are indistinguishable in the view of traditional model diagnostic tools. We will demonstrate that our procedure is able to select the most robust models that perform best on future data across different subpopulations.

5.3.2 Problem Specifics

The input X_i is an RGB satellite image, each pixel represented by a vector in \mathbb{R}^3 , and the label $Y_i \in \{0, 1, 2, \dots, 61\}$ represents one of the 62 land use categories. We consider the following three non-overlapping subsets of the data based on image taken times:

- Training: select images in 2003 – 2013 (76,863 images)

- Validation: select images in 2003 – 2013 (11,483 images)
- Test: images in 2016 – 2018 (22,108 images)

We consider training and validation images (images from pre-2013) as in-distribution (ID) data, and test images in taken between 2016 and 2018 as out-of-distribution (OOD) data. Each image also comes with meta information, including the (longitude, latitude) location of which the photo was taken, the continent/region information, and the weather information (cloudiness on a scale from 1 to 10) [61, 68]. We define subpopulations based on meta information.

Similarly to the Warfarin example, we observe that model performances remain similar either temporally or spatially when each dimension is considered *separately*, but there is substantial variability across intersections of region and year. For a standard benchmark deep neural network model in Huang et al. [61], Koh et al. [68] that achieves near-SOTA performance, we present these trends in Figure 4(a). Furthermore, in Figure 4(b), we observe substantial variability in error rates across different labels, indicating there is a varying level of difficulty in classifying different classes. (We observed similar patterns for other models.)




Figure 4. For *DenseNet* ERM, spatiotemporal intersectionality (left) and performance by class (right)

Because of significant variation in difficulty in learning different classes, we consider the core attributes Z on which the subpopulation is defined to be the full meta information vector concatenated with the true class label y . As we impose no assumption on Z , we could also use the semantic meaning of each class labels in place of categorical variables. We will go back to this point in Section 5.4.

We take three benchmark SOTA transfer learning models (all pretrained on *ImageNet*). Two of the three models come from Huang et al. [61], Koh et al. [68]. Both models are *DenseNet*-121 based models, but trained with different objectives:

- *DenseNet*-121 ERM: the model is trained on FMoW to minimize the usual average training loss (mean squared loss);
- *DenseNet*-121 IRM: the model is trained on FMoW to minimize the invariant risk (the loss adds an extra penalty term that penalizes feature distributions that have different optimal linear classifiers for each domain), proposed in Arjovsky et al. [6].

We consider a third model, the *ImageNet*-pretrained Dual Path Network model (*DPN*-68 model) [29]. This model has a different architecture from that of *DenseNet* models. All these models achieve an out-of-sample ID accuracy of 60%.

However, these SOTA benchmark models suffer from significant performance drop on the OOD data: all models suffer from a performance drop of 7% in average accuracy, and this performance drop increases to a stunning 30% for images coming from Africa (the region where all model perform the achieve the lowest accuracy, which we will call the worst-case region). These observations reveal that there is natural distribution shift from past data to future unseen data. It also raises the flag that despite achieving great ID performance, these SOTA benchmark models are not robust against these shifts.

On the other hand, *CLIP* models have exhibited promising robustness properties [86]. To adapt CLIP-based models to the satellite image classification problem, we adopt a weight-space ensembling method (*CLIP WiSE-FT*) in Wortsman et al. [118]. This method has been observed to exhibit large Pareto improvements in the sense that it leads to a suite of models with improved performance with respect to both ID and OOD accuracy. Motivated by the observed robustness gains, we consider the *CLIP WiSE-FT* model that achieves comparable performance on the FMoW ID validation set to the benchmark *ImageNet* pre-trained models. Appendix C provides additional details on the experimental settings and training specifications.

Goal: Presented with both benchmark *ImageNet*-based models and *CLIP WiSE-FT*, our goal as the practitioner is to choose the one that generalizes best in the future unseen OOD test data uniformly across all subpopulations.

Challenge: In the view of traditional model diagnostic tools, these models are indistinguishable as they achieve comparable average ID accuracy and ID loss.

Nevertheless, our proposed method is able to select models that perform well “in the future” without requiring OOD data, and at the same time our method raises awareness of difficulty in domain generalization. We report the experiment results in the next section.

5.3.3 Results

We compute estimators of W_α (Algorithm 1) on the ID validation data using the standard cross entropy loss.




Figure 5. Left: $Z = (\text{all metadata})$; Right: $Z = (\text{all metadata}, Y)$. Results are averaged over 50 random seeds with error bars corresponding to a 95% confidence interval over the random runs.

Models	ID, 2002–2013		OOD, 2016–2018			
	Accuracy	Loss	Accuracy	Loss	Africa Accuracy	Africa Loss
<i>CLIP WiSE-FT</i>	0.61	2.78	0.56	2.84	0.38	3.08
<i>DenseNet</i> ERM	0.61	2.78	0.53	3.50	0.33	5.41
<i>DenseNet</i> IRM	0.59	2.44	0.51	2.94	0.31	4.46
<i>DPN-68</i>	0.61	2.75	0.53	3.55	0.31	5.61

Table 1. Model performance on ID validation and OOD test sets. All models suffer a performance drop on the OOD test set in both accuracy and average loss. The performance degradation is particularly significant on images from Africa. On the OOD data, *CLIP WiSE-FT* outperforms other models both in average accuracy/loss and worst-region accuracy/loss.

In Figure 5, we summarize the estimated worst-case subpopulation performances defined over the entire meta data, across different subpopulation sizes α . First, we note that all models have comparable ID accuracy of $\sim 60\%$ and *DenseNet* IRM has the best average-case cross entropy loss. However, the worst-case subpopulation performance of the *ImageNet* pre-trained models is substantially worse compared to that of *CLIP WiSE-FT*. This gap grows larger as the subpopulation size α becomes increasingly small. Evaluations on worst-case subpopulations suggest that *CLIP WiSE-FT* exhibits robustness against subpopulation shifts; in contrast, average-case evaluations will select *DenseNet* IRM. We observe a drastic performance deterioration on tail subpopulations. The inclusion of label information in Z significantly deteriorates worst-case performance, raising concerns about the distributional robustness of all models including changes in the label distribution.

In Table 1, we present model performances on the OOD data collected during 2016–2018. All models suffer a significant performance drop under temporal distribution shift, particularly on images collected in Africa where predictive accuracy drops by up to 20 percentage points. *CLIP WiSE-FT* exhibits the most robustness under spatiotemporal shift than any other model, as presaged by evaluations of worst-case subpopulation performance in Figure 5.

5.4 Flexibility in the choice of Z

A key advantage of our method is the flexibility in the choice of Z : the modeler can define granular or coarse subpopulations based on this choice. As defining subpopulations over all metadata can be conservative, we present additional results under $Z=(region, year)$ and $Z=(region, year, label Y)$ in Appendix C.

Moreover, the flexibility in the choice of Z allows us to exploit the *semantic meaning of class labels*. Instead of incorporating labels as a categorical variable, we can generate natural language description of the labels by concatenating each class label with engineered prompts, and pass it to the CLIP text encoder [86]. This process generates a feature representation for the label that is semantically meaningful, and we can use these feature vectors in our core attributes to define subpopulations. In Appendix C.3, we present evaluation results where we take the feature vector in place of the label Y when defining Z .

6 Connections to coherence and distributional robustness

The worst-case subpopulation performance (1.2) only considers subpopulations that comprise α -fraction of the data. In this section, we propose a generalized measure of model robustness that considers subpopulation sizes on various scales. We show that our generalized notion of worst-case subpopulation performance is closely related to coherent risk measures and distributional robustness. Concretely, instead of a single subpopulation size, we take the average over $\alpha \in (0, 1]$ using a probability measure λ

$$\int_{(0,1]} W_\alpha(\mu^*) d\lambda(\alpha). \quad (6.1)$$

The multi-scale average (6.1) introduces substantial modeling flexibility by incorporating prior beliefs on which subpopulation sizes are of higher concern. It can consider arbitrarily small subpopulations (see Proposition 6 to come), and by letting $\lambda = t\delta\{\alpha\} + (1-t)\delta\{1\}$ for $t \in [0, 1]$, it interpolates between average and the worst subpopulation performance $W_\alpha(\mu^*)$.

Equipped with the multi-scale average (6.1), we define the *generalized worst-case subpopulation performance* over a (nonempty) class Λ of probability measures on the half-open interval $(0, 1]$

$$W_\Lambda(\mu) := \sup_{\lambda \in \Lambda} \int_{(0,1]} W_\alpha(\mu) d\lambda(\alpha). \quad (6.2)$$

We show an equivalence between generalized worst-case subpopulation performances (6.2) and *coherent risk measures*, an axiomatic definition of risk-aversion. By utilizing a well-known duality between coherence and distributional robustness [38, 46, 30, 95], we can further show that $W_\Lambda(\mu)$ is in fact flexible enough to represent any worst-case performance over distribution shifts

$$\mu \mapsto \sup_{Q \in \mathcal{Q}} \mathbb{E}_Q[\mu], \quad (6.3)$$

where \mathcal{Q} is a collection of probability measures dominated by P . The worst-case subpopulation performance (1.2) is a particular distributionally robust formulation with \mathcal{Q} given by the set (1.1); our equivalence results to come show the converse by utilizing the generalization (6.2).

Formally, let (Ω, \mathcal{F}, P) be the underlying probability space over which all random variables are defined. We restrict attention to random variables with finite moments: for $k \in [1, \infty)$, define

$$\mathcal{L}^k := \left\{ \mu : (\Omega, \mathcal{F}, P) \rightarrow \mathbb{R} \text{ s.t. } \mathbb{E}_P[\mu^k] < \infty \right\}.$$

A risk measure $\rho : \mathcal{L}^k \rightarrow \mathbb{R} \cup \{\infty\}$ maps a random variable connoting a notion of loss, e.g., $\mu(Z) = \mathbb{E}[\ell(\theta(X); Y) \mid Z]$, to a single number representing the modeler's disutility. We say that a risk measure is *law-invariant* if $\rho(\mu) = \rho(\mu')$ whenever $\mu \stackrel{d}{=} \mu'$. A *coherent* risk measure incorporates sensible notions of risk-aversion through the following axioms.

Definition 1 (Shapiro et al. [103, Definition 6.4]). $\rho : \mathcal{L}^k \rightarrow \mathbb{R} \cup \{\infty\}$ is **coherent** if it satisfies

1. *Convexity:* $\rho(t\mu + (1-t)\mu') \leq t\rho(\mu) + (1-t)\rho(\mu')$ for all $t \in [0, 1]$ and $\mu, \mu' \in \mathcal{L}^k$
2. *Monotonicity:* $\rho(\mu) \leq \rho(\mu')$ if $\mu, \mu' \in \mathcal{L}^k$ and $\mu \leq \mu'$ P -almost surely
3. *Translation equivariance:* $\rho(\mu + c) = \rho(\mu) + c$ for all $c \in \mathbb{R}$ and $\mu \in \mathcal{L}^k$
4. *Positive homogeneity:* $\rho(c\mu) = c\rho(\mu)$ for all $c > 0$ and $\mu \in \mathcal{L}^k$

While the above axioms have originally been proposed with economic and financial applications in mind (e.g., see the tutorial Rockafellar [91]), they can also be interpreted from the perspective of predictive systems [116]. Convexity models diminishing marginal utility (for loss/disutility); the modeler incurs higher marginal disutility when the underlying prediction model $\theta(X)$ is already poor and incurring high prediction errors. Monotonicity is a natural property to enforce. Translation equivariance says that any addition of certain prediction error translates to a proportional change in the modeler’s disutility. Similarly, positive homogeneity says a tenfold increase in prediction error leads to a proportional increase in the modeler’s disutility.

Standard duality results give a one-to-one correspondence between coherent risk measures and particular distributionally robust formulations. For any space \mathcal{L}^k , consider its dual space \mathcal{L}^{k*} where $1/k + 1/k_* = 1$ and $k_* \in (1, \infty]$. Recall that for a risk measure $\rho : \mathcal{L}^k \rightarrow \mathbb{R} \cup \{\infty\}$, we say $\rho(\cdot)$ is proper if its domain is nonempty and define its *Fenchel conjugate* $\rho^* : \mathcal{L}^{k*} \rightarrow \mathbb{R} \cup \{\infty\}$ as $\rho^*(L) := \sup_{\mu \in \mathcal{L}^k} \{\mathbb{E}_P[L\mu] - \rho(\mu)\}$. For a (sufficiently regular) convex function $\rho : \mathcal{L}^k \rightarrow \mathbb{R} \cup \{\infty\}$, Fenchel-Moreau duality gives the biconjugacy relation

$$\rho(\mu) = \sup_{L \in \mathcal{L}^{k*}} \{\mathbb{E}_P[L\mu] - \rho^*(L)\}. \quad (6.4)$$

When $\rho(\cdot)$ is coherent, proper, and lower semi-continuous, its *dual set* (a.k.a. domain of ρ^*) can be characterized as the following

$$\text{dom } \rho^* = \left\{ L \in \mathcal{L}^{k*} : L \geq 0, \mathbb{E}_P[L] = 1, \text{ and } \mathbb{E}_P[L\mu] \leq \rho(\mu) \ \forall \mu \in \mathcal{L}^k \right\}, \quad (6.5)$$

where $\rho^*(L) = 0$ whenever $L \in \text{dom } \rho^*$. The dual set is weak* closed and is a set of probability density functions, so $\mathbb{E}_P[L\mu]$ can be viewed as the expectation $\mathbb{E}_Q[\mu]$ in under the probability measure defined by $\frac{dQ}{dP} = L$. Collecting these observations, the biconjugacy relation (6.4) gives the following result.

Lemma 3 (Shapiro et al. [103, Theorem 6.42]). *The set of coherent, proper, lower semi-continuous, and law-invariant risk measures is identical to the set of mappings given by $\mu \mapsto \sup_{Q \in \mathcal{Q}} \mathbb{E}_Q[\mu]$ for some nonempty class of probability measures \mathcal{Q} over (Ω, \mathcal{F}, P) satisfying $\frac{dQ}{dP} \in \mathcal{L}^{k*}$ for all $Q \in \mathcal{Q}$.*

In this equivalence, we do not require the class \mathcal{Q} to be convex since the dual set of $\mu \mapsto \sup_{Q \in \mathcal{Q}} \mathbb{E}_Q[\mu]$ is the convex hull of \mathcal{Q} .

The main result of this section gives an equivalence between generalized worst-case subpopulation performances (6.2) and distributionally robust losses (6.3), and in turn, coherent risk measures. Recall that a probability space (Ω, \mathcal{F}, P) is *nonatomic* if any $S \in \mathcal{F}$ with $\mathbb{P}(S) > 0$ contains a subset $S' \in \mathcal{F}$ such that $\mathbb{P}(S) > \mathbb{P}(S') > 0$.

Theorem 5. *A generalized worst-case subpopulation performance measure (6.2) is coherent, proper, lower semi-continuous, and law-invariant. If P is nonatomic, then the converse also holds.*

Nonatomicity is a mild assumption since most machine learning applications do not depend on the underlying probability space: we can simply take the underlying probability space to be the standard uniform space $\Omega = [0, 1]$ equipped with the Borel sigma algebra and the Lebesgue/uniform measure.

Theorem 5 is essentially a consequence of a well-known reformulation of coherent risk measures known as the Kusuoka representation [71, 101, 103]. Our proof is not novel, but we give it in Appendix D.1 for completeness; it is constructive so that given a coherent risk measure $\rho(\cdot)$, we

define the exact set of probability measures Λ_ρ such that $\rho(\cdot)$ is equal to the generalized worst-case subpopulation performance (6.2) defined with Λ_ρ . Our construction makes concrete how the axioms of coherence translate to multiple preferences over subpopulation sizes.

As an example, we consider the *higher-order conditional value-at-risk* [70], a more conservative risk measure than the worst-case subpopulation performance (1.2) we considered in prior sections: for $\alpha \in (0, 1]$,

$$\rho_k(\mu) := \inf_{\eta \in \mathbb{R}} \left\{ \frac{1}{\alpha} \left(\mathbb{E}_P (\mu - \eta)_+^k \right)^{1/k} + \eta \right\}.$$

When $k = 1$, we recover the worst-case subpopulation performance (1.2). The following result makes explicit the equivalence relation given in Theorem 5 for the risk measure family $\rho_k(\cdot)$.

Proposition 6. *For $k \in (1, \infty)$, let $\Lambda_k := \left\{ \lambda \in \Delta((0, 1]) : \int_0^1 \left(\int_u^1 a^{-1} d\lambda(a) \right)^{k_*} du \leq \alpha^{-k_*} \right\}$. If P is nonatomic,*

$$\rho_k(\mu) = \sup_{Q \ll P} \left\{ \mathbb{E}_Q[\mu] : \mathbb{E}_P \left(\frac{dQ}{dP} \right)^{k_*} \leq \alpha^{-k_*} \right\} = W_{\Lambda_k}(\mu).$$

In the above result, we can see that the set Λ_k allows arbitrarily small subpopulations by using the $L^{k_*}(P)$ -norm. See Appendix D.2 for its proof.

7 Discussion

To ensure models perform reliably under operation, we need to *rigorously* certify their performance under distribution shift prior to deployment. We study the *worst-case subpopulation performance* of a model, a natural notion of model robustness that is easy to communicate with users, regulators, and business leaders. Our approach allows flexible modeling of subpopulations over an arbitrary variable Z and automatically accounts for complex intersectionality. We develop scalable estimation procedures for the worst-case subpopulation performance (1.2) and the certificate of robustness (2.1) of a model. Our convergence guarantees apply even when we use high-dimensional inputs (e.g. natural language) to define Z . Our diagnostic may further inform data collection and model improvement by suggesting data collection efforts and model fixes on regions of \mathcal{Z} with high conditional risk (1.3).

The worst-case performance (1.2) over mixture components as subpopulations (1.1) provides a strong guarantee over arbitrary subpopulations, but it may be overly conservative in cases when there is a natural geometry in $Z \in \mathcal{Z}$. Incorporating such problem-specific structures in defining a tailored notion of subpopulation is a promising research direction towards operationalizing the concepts put forth in this work. As an example, Srivastava et al. [104] recently studied similar notions of worst-case performance defined over human annotations.

We focus on the narrow question of evaluating model robustness under distribution shift; our evaluation perspective is thus inherently limited. Data collection systems inherit socioeconomic inequities, and reinforce existing political power structures. This affects *all* aspects of the ML development pipeline, and our diagnostic is no panacea. A notable limitation of our approach is that we do not explicitly consider the power differential that often exists between those who deploy the prediction system and those for whom it gets used on. Systems must be deployed with considered analysis of its adverse impacts, and we advocate for a holistic approach towards addressing its varied implications.

References

- [1] B. Abelson, R. Kush, and J. Sun. Targeting direct cash transfers to the extremely poor. *Proceedings of the ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 2014.
- [2] American Medical Association. AMA passes first policy recommendations on augmented intelligence., 2018. URL www.ama-assn.org/ama-passes-first-policy-recommendations-augmented-intelligence.
- [3] D. Amodei, S. Ananthanarayanan, R. Anubhai, J. Bai, E. Battenberg, C. Case, J. Casper, B. Catanzaro, Q. Cheng, and G. Chen. Deep speech 2: end-to-end speech recognition in English and Mandarin. In *Proceedings of the 33rd International Conference on Machine Learning*, pages 173–182, 2016.
- [4] E. Amorim, M. Cançado, and A. Veloso. Automated essay scoring in the presence of biased ratings. In *Association for Computational Linguistics (ACL)*, pages 229–237, 2018.
- [5] Anonymous. Distributionally robust neural networks. In *Submitted to International Conference on Learning Representations*, 2020. URL <https://openreview.net/forum?id=ryxGuJrFvS>. under review.
- [6] M. Arjovsky, L. Bottou, I. Gulrajani, and D. Lopez-Paz. Invariant risk minimization. In *Proceedings of the Eighth International Conference on Learning Representations*, 2020.
- [7] S. Barocas and A. D. Selbst. Big data’s disparate impact. *104 California Law Review*, 3: 671–732, 2016.
- [8] S. Barocas, M. Hardt, and A. Narayanan. *Fairness and Machine Learning*. fairmlbook.org, 2019.
- [9] P. L. Bartlett, O. Bousquet, and S. Mendelson. Local Rademacher complexities. *Annals of Statistics*, 33(4):1497–1537, 2005.
- [10] P. L. Bartlett, D. J. Foster, and M. J. Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, pages 6241–6250, 2017.
- [11] S. Beery, E. Cole, and A. Gjoka. The iwildcam 2020 competition dataset. *arXiv:2004.10340 [cs.CV]*, 2020.
- [12] D. Belomestny and V. Krätschmer. Central limit theorems for law-invariant coherent risk measures. *Journal of Applied Probability*, 49(1):1–21, 2012.
- [13] A. Ben-Tal, D. den Hertog, A. D. Waegenaere, B. Melenberg, and G. Rennen. Robust solutions of optimization problems affected by uncertain probabilities. *Management Science*, 59(2): 341–357, 2013.
- [14] D. Bertsimas, V. Gupta, and N. Kallus. Data-driven robust optimization. *Mathematical Programming, Series A*, 167(2):235–292, 2018.
- [15] J. Blanchet and K. Murthy. Quantifying distributional model risk via optimal transport.

- Mathematics of Operations Research*, 44(2):565–600, 2019.
- [16] J. Blanchet, Y. Kang, F. Zhang, and K. Murthy. Data-driven optimal transport cost selection for distributionally robust optimization. *arXiv:1705.07152 [stat.ML]*, 2017.
- [17] J. Blanchet, Y. Kang, and K. Murthy. Robust Wasserstein profile inference and applications to machine learning. *Journal of Applied Probability*, 56(3):830–857, 2019.
- [18] J. Blanchet, F. He, and K. Murthy. On distributionally robust extreme value analysis. *Extremes*, pages 1–31, 2020.
- [19] J. Blanchet, K. Murthy, and N. Si. Confidence regions in Wasserstein distributionally robust estimation. *Biometrika*, 109(2):295–315, 2022.
- [20] J. Blitzer, R. McDonald, and F. Pereira. Domain adaptation with structural correspondence learning. In *Proceedings of the 2006 conference on empirical methods in natural language processing*, pages 120–128. Association for Computational Linguistics, 2006.
- [21] S. L. Blodgett, L. Green, and B. O’Connor. Demographic dialectal variation in social media: A case study of African-American English. In *Proceedings of Empirical Methods for Natural Language Processing*, pages 1119–1130, 2016.
- [22] J. F. Bonnans and A. Shapiro. *Perturbation Analysis of Optimization Problems*. Springer, 2000.
- [23] D. B. Brown. Large deviations bounds for estimating conditional value-at-risk. *Operations Research Letters*, 35(6):722–730, 2007.
- [24] J. Buolamwini and T. Gebru. Gender shades: Intersectional accuracy disparities in commercial gender classification. In *Conference on Fairness, Accountability and Transparency*, pages 77–91, 2018.
- [25] D. S. Char, N. H. Shah, and D. Magnus. Implementing machine learning in health care—addressing ethical challenges. *New England Journal of Medicine*, 378(11):981, 2018.
- [26] M. S. Chen, P. N. Lara, J. H. Dang, D. A. Paterniti, and K. Kelly. Twenty years post-NIH revitalization act: enhancing minority participation in clinical trials (EMPaCT): laying the groundwork for improving minority clinical trial accrual: renewing the case for enhancing minority participation in cancer clinical trials. *Cancer*, 120:1091–1096, 2014.
- [27] T. Chen and C. Guestrin. Xgboost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 785–794, 2016.
- [28] X. Chen, Q. Lin, and G. Xu. Distributionally robust optimization with confidence bands for probability density functions. *INFORMS Journal on Optimization*, 4(1):65–89, 2022.
- [29] Y. Chen, J. Li, H. Xiao, X. Jin, S. Yan, and J. Feng. Dual path networks. *arXiv:1707.01629 [cs.CV]*, 2017.
- [30] P. Cheridito, F. Delbaen, and M. Kupper. Coherent and convex monetary risk measures for bounded càdlàg processes. *Stochastic Processes and their Applications*, 112(1):1–22, 2004.

- [31] V. Chernozhukov, D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins. Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68, 2018.
- [32] A. Chouldechova. A study of bias in recidivism prediction instruments. *Big Data*, pages 153–163, 2017.
- [33] G. Christie, N. Fendley, J. Wilson, and R. Mukherjee. Functional map of the world. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pages 6172–6180, 2018.
- [34] I. W. P. Consortium. Estimation of the warfarin dose with clinical and pharmacogenetic data. *New England Journal of Medicine*, 360(8):753–764, 2009.
- [35] S. Corbett-Davies and S. Goel. The measure and mismeasure of fairness: A critical review of fair machine learning. *arXiv:1808.00023 [cs.CY]*, 2018.
- [36] H. Daume III and D. Marcu. Domain adaptation for statistical classifiers. *Journal of artificial Intelligence research*, 26:101–126, 2006.
- [37] E. Delage and Y. Ye. Distributionally robust optimization under moment uncertainty with application to data-driven problems. *Operations Research*, 58(3):595–612, 2010.
- [38] F. Delbaen. Coherent risk measures on general probability spaces. In *Advances in finance and stochastics*, pages 1–37. Springer, 2002.
- [39] E. Denton, A. Hanna, R. Amironesei, A. Smart, H. Nicole, and M. K. Scheuerman. Bringing the people back in: Contesting benchmark machine learning datasets. *arXiv:2007.07399 [cs.CY]*, 2020.
- [40] J. Duchi, T. Hashimoto, and H. Namkoong. Distributionally robust losses for latent covariate mixtures. *arXiv:2007.13982 [stat.ML]*, 2020.
- [41] J. C. Duchi and H. Namkoong. Learning models with uniform performance via distributionally robust optimization. *Annals of Statistics*, 49(3):1378–1406, 2021.
- [42] J. C. Duchi, P. W. Glynn, and H. Namkoong. Statistics of robust optimization: A generalized empirical likelihood approach. *Mathematics of Operations Research*, 46:946–969, 2021.
- [43] E. Erdoğan and G. Iyengar. Ambiguous chance constrained problems and robust optimization. *Mathematical Programming*, 107(1-2):37–61, 2006.
- [44] P. M. Esfahani and D. Kuhn. Data-driven distributionally robust optimization using the wasserstein metric: Performance guarantees and tractable reformulations. *Mathematical Programming, Series A*, 171(1-2):115–166, 2018.
- [45] M. Feldman, S. A. Friedler, J. Moeller, C. Scheidegger, and S. Venkatasubramanian. Certifying and removing disparate impact. In *proceedings of the 21th ACM SIGKDD international conference on knowledge discovery and data mining*, pages 259–268, 2015.
- [46] H. Föllmer and A. Schied. Convex measures of risk and trading constraints. *Finance and stochastics*, 6(4):429–447, 2002.

- [47] R. Gao. Finite-sample guarantees for Wasserstein distributionally robust optimization: Breaking the curse of dimensionality. *arXiv:2009.04382 [cs.LG]*, 2020.
- [48] R. Gao and A. J. Kleywegt. Distributionally robust stochastic optimization with Wasserstein distance. *Mathematics of Operations Research*, 2022.
- [49] R. Gao, X. Chen, and A. Kleywegt. Wasserstein distributional robustness and regularization in statistical learning. *arXiv:1712.06050 [cs.LG]*, 2017.
- [50] R. Gao, X. Chen, and A. J. Kleywegt. Wasserstein distributionally robust optimization and variation regularization. *Operations Research*, 72(3):1177–1191, 2024.
- [51] K. Goel, N. Rajani, J. Vig, S. Tan, J. Wu, S. Zheng, C. Xiong, M. Bansal, and C. Ré. Robustness gym: Unifying the nlp evaluation landscape. *arXiv:2101.04840 [cs.CL]*, 2021.
- [52] J. Goh and M. Sim. Distributionally robust optimization and its tractable approximations. *Operations research*, 58(4-part-1):902–917, 2010.
- [53] S. N. Goodman, S. Goel, and M. R. Cullen. Machine learning, health disparities, and causal reasoning. *Annals of Internal Medicine*, 2018.
- [54] P. J. Grother, G. W. Quinn, and P. J. Phillips. Report on the evaluation of 2d still-image face recognition algorithms. *NIST Interagency/Internal Reports (NISTIR)*, 7709, 2010.
- [55] V. Guigues, V. Kratschmer, and A. Shapiro. A central limit theorem and hypotheses testing for risk-averse stochastic programs. *SIAM Journal on Optimization*, 28(2):1337–1366, 2018.
- [56] S. Han, D. Ahn, S. Park, J. Yang, S. Lee, J. Kim, H. Yang, S. Park, and M. Cha. Learning to score economic development from satellite imagery. In *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, page 2970–2979, 2020.
- [57] M. Hardt, E. Price, and N. Srebro. Equality of opportunity in supervised learning. In *Advances in Neural Information Processing Systems 29*, 2016.
- [58] T. Hashimoto, M. Srivastava, H. Namkoong, and P. Liang. Fairness without demographics in repeated loss minimization. In *Proceedings of the 35th International Conference on Machine Learning*, 2018.
- [59] Ú. Hébert-Johnson, M. P. Kim, O. Reingold, and G. N. Rothblum. Calibration for the (computationally-identifiable) masses. *arXiv:1711.08513 [cs.LG]*, 2017.
- [60] D. Hovy and A. Søgaard. Tagging performance correlates with author age. In *Proceedings of the 53rd Annual Meeting of the Association for Computational Linguistics (Short Papers)*, volume 2, pages 483–488, 2015.
- [61] G. Huang, Z. Liu, L. Van Der Maaten, and K. Q. Weinberger. Densely connected convolutional networks. In *Proceedings of the International Conference on Computer Vision*, pages 4700–4708, 2017.
- [62] S. Jeong and H. Namkoong. Robust causal inference under covariate shift via worst-case subpopulation treatment effect. In *Proceedings of the Thirty Third Annual Conference on Computational Learning Theory*, 2020.

- [63] M. Kearns, S. Neel, A. Roth, and Z. S. Wu. Preventing fairness gerrymandering: Auditing and learning for subgroup fairness. *arXiv:1711.05144 [cs.LG]*, 2018.
- [64] M. Kearns, S. Neel, A. Roth, and Z. S. Wu. An empirical study of rich subgroup fairness for machine learning. In *Proceedings of the Conference on Fairness, Accountability, and Transparency*, pages 100–109. ACM, 2019.
- [65] E. H. Kennedy. Semiparametric doubly robust targeted double machine learning: a review. *arXiv:2203.06469 [stat.ME]*, 2022.
- [66] J. Kleinberg, S. Mullainathan, and M. Raghavan. Inherent trade-offs in the fair determination of risk scores. In *Proceedings of the 8th Conference on Innovations in Theoretical Computer Science (ITCS)*, 2016.
- [67] A. Koenecke, A. Nam, E. Lake, J. Nudell, M. Quartey, Z. Mengesha, C. Toups, J. R. Rickford, D. Jurafsky, and S. Goel. Racial disparities in automated speech recognition. *Proceedings of the National Academy of Sciences*, 117(14):7684–7689, 2020.
- [68] P. W. Koh, S. Sagawa, H. Marklund, S. M. Xie, M. Zhang, A. Balsubramani, W. Hu, M. Yasunaga, R. L. Phillips, S. Beery, et al. Wilds: A benchmark of in-the-wild distribution shifts. *arXiv:2012.07421 [cs.LG]*, 2020.
- [69] V. Koltchinskii. Local Rademacher complexities and oracle inequalities in risk minimization. *Annals of Statistics*, 34(6):2593–2656, 2006.
- [70] P. A. Krokmal. Higher moment coherent risk measures. *Quantitative Finance*, 7(4):373–387, 2007.
- [71] S. Kusuoka. On law invariant coherent risk measures. In *Advances in Mathematical Economics*, pages 83–95. Springer, 2001.
- [72] Y. Laguel, J. Malick, and Z. Harchaoui. First-order optimization for superquantile-based supervised learning. In *2020 IEEE 30th International Workshop on Machine Learning for Signal Processing (MLSP)*, pages 1–6. IEEE, 2020.
- [73] Y. Laguel, K. Pillutla, J. Malick, and Z. Harchaoui. Superquantiles at work: Machine learning applications and efficient subgradient computation. *Set-Valued and Variational Analysis*, 29(4):967–996, 2021.
- [74] H. Lam. Recovering best statistical guarantees via the empirical divergence-based distributionally robust optimization. *Operations Research*, 67(4):1090–1105, 2019.
- [75] H. Lam and E. Zhou. Quantifying input uncertainty in stochastic optimization. In *Proceedings of the 2015 Winter Simulation Conference*. IEEE, 2015.
- [76] J. Li, S. Huang, and A. M.-C. So. A first-order algorithmic framework for distributionally robust logistic regression. *Advances in Neural Information Processing Systems*, 32, 2019.
- [77] S. Mendelson. Learning without concentration. In *Proceedings of the Twenty Seventh Annual Conference on Computational Learning Theory*, 2014.
- [78] J. Miller, K. Krauth, B. Recht, and L. Schmidt. The effect of natural distribution shift

- on question answering models. In *International Conference on Machine Learning*, pages 6905–6916. PMLR, 2020.
- [79] J. Miller, R. Taori, A. Raghunathan, S. Sagawa, P. W. Koh, V. Shankar, P. Liang, Y. Carmon, and L. Schmidt. Accuracy on the line: on the strong correlation between out-of-distribution and in-distribution generalization. In *Proceedings of the 38th International Conference on Machine Learning*, 2021.
- [80] M. Mitchell, S. Wu, A. Zaldivar, P. Barnes, L. Vasserman, B. Hutchinson, E. Spitzer, I. D. Raji, and T. Gebru. Model cards for model reporting. In *Proceedings of the Conference on Fairness, Accountability, and Transparency*, pages 220–229, 2019.
- [81] T. Miyato, S.-i. Maeda, M. Koyama, K. Nakae, and S. Ishii. Distributional smoothing with virtual adversarial training. *arXiv:1507.00677 [stat.ML]*, 2015.
- [82] W. K. Newey. The asymptotic variance of semiparametric estimators. *Econometrica*, pages 1349–1382, 1994.
- [83] J. Neyman. Optimal asymptotic tests of composite statistical hypotheses. *Probability and Statistics*, 416(44), 1959.
- [84] G. Pflug and N. Wozabal. Asymptotic distribution of law-invariant risk functionals. *Finance and Stochastics*, 14(3):397–418, 2010.
- [85] L. Prashanth, K. Jagannathan, and R. K. Kolla. Concentration bounds for cvar estimation: The cases of light-tailed and heavy-tailed distributions. In *Proceedings of the 37th International Conference on Machine Learning*, 2020.
- [86] A. Radford, J. W. Kim, C. Hallacy, A. Ramesh, G. Goh, S. Agarwal, G. Sastry, A. Askell, P. Mishkin, J. Clark, et al. Learning transferable visual models from natural language supervision. In *Proceedings of the 38th International Conference on Machine Learning*, 2021.
- [87] A. Rajkomar, M. Hardt, M. D. Howell, G. Corrado, and M. H. Chin. Ensuring fairness in machine learning to advance health equity. *Annals of Internal Medicine*, 2018.
- [88] J. Rawls. *Justice as fairness: A restatement*. Harvard University Press, 2001.
- [89] J. Rawls. *A theory of justice*. Harvard university press, 2009.
- [90] B. Recht, R. Roelofs, L. Schmidt, and V. Shankar. Do ImageNet classifiers generalize to ImageNet? In *Proceedings of the 36th International Conference on Machine Learning*, 2019.
- [91] R. T. Rockafellar. Coherent approaches to risk in optimization under uncertainty. *Tutorials in Operations Research*, 3:38–61, 2007.
- [92] R. T. Rockafellar and S. Uryasev. Optimization of conditional value-at-risk. *Journal of Risk*, 2:21–42, 2000.
- [93] R. T. Rockafellar and S. Uryasev. Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance*, 26(7):1443–1471, 2002.
- [94] R. T. Rockafellar and R. J. B. Wets. *Variational Analysis*. Springer, New York, 1998.

- [95] A. Ruszczyński and A. Shapiro. Optimization of convex risk functions. *Mathematics of Operations Research*, 31(3):433–452, 2006.
- [96] K. Saenko, B. Kulis, M. Fritz, and T. Darrell. Adapting visual category models to new domains. In *Proceedings of the European Conference on Computer Vision*, pages 213–226. Springer, 2010.
- [97] S. Sagawa, P. W. Koh, T. B. Hashimoto, and P. Liang. Distributionally robust neural networks for group shifts: On the importance of regularization for worst-case generalization. In *Proceedings of the Seventh International Conference on Learning Representations*, 2019.
- [98] P. Sapiezynski, V. Kassarnig, and C. Wilson. Academic performance prediction in a gender-imbalanced environment. In *Proceedings of the Eleventh ACM Conference on Recommender Systems*, volume 1, pages 48–51, 2017.
- [99] S. Shafieezadeh-Abadeh, P. M. Esfahani, and D. Kuhn. Distributionally robust logistic regression. In *Advances in Neural Information Processing Systems 28*, pages 1576–1584, 2015.
- [100] V. Shankar, A. Dave, R. Roelofs, D. Ramanan, B. Recht, and L. Schmidt. Do image classifiers generalize across time? *arXiv:1906.02168 [cs.LG]*, 2019.
- [101] A. Shapiro. On Kusuoka representation of law invariant risk measures. *Mathematics of Operations Research*, 38(1):142–152, 2013.
- [102] A. Shapiro, D. Dentcheva, and A. Ruszczyński. *Lectures on Stochastic Programming: Modeling and Theory*. SIAM and Mathematical Programming Society, 2009.
- [103] A. Shapiro, D. Dentcheva, and A. Ruszczyński. *Lectures on stochastic programming: modeling and theory*. SIAM, second edition, 2014.
- [104] M. Srivastava, T. Hashimoto, and P. Liang. Robustness to spurious correlations via human annotations. In *Proceedings of the 37th International Conference on Machine Learning*, pages 9109–9119. PMLR, 2020.
- [105] M. Staib and S. Jegelka. Distributionally robust optimization and generalization in kernel methods. In *Advances in Neural Information Processing Systems*, pages 9131–9141, 2019.
- [106] A. Subbaswamy, R. Adams, and S. Saria. Evaluating model robustness and stability to dataset shift. In *Proceedings of the 24 International Conference on Artificial Intelligence and Statistics*, pages 2611–2619, 13–15 Apr 2021.
- [107] R. Taori, A. Dave, V. Shankar, N. Carlini, B. Recht, and L. Schmidt. Measuring robustness to natural distribution shifts in image classification. In *Advances in Neural Information Processing Systems 20*, 2020.
- [108] R. Tatman. Gender and dialect bias in YouTube’s automatic captions. In *First Workshop on Ethics in Natural Language Processing*, volume 1, pages 53–59, 2017.
- [109] A. Torralba and A. A. Efros. Unbiased look at dataset bias. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pages 1521–1528. IEEE, 2011.
- [110] A. W. van der Vaart and J. A. Wellner. *Weak Convergence and Empirical Processes: With*

Applications to Statistics. Springer, New York, 1996.

- [111] R. Volpi, H. Namkoong, O. Sener, J. Duchi, V. Murino, and S. Savarese. Generalizing to unseen domains via adversarial data augmentation. In *Advances in Neural Information Processing Systems 31*, 2018.
- [112] M. J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019.
- [113] J. Wang, R. Gao, and Y. Xie. Sinkhorn distributionally robust optimization. *arXiv:2109.11926 [math.OA]*, 2021.
- [114] S. Wang, N. Si, J. Blanchet, and Z. Zhou. On the foundation of distributionally robust reinforcement learning. *arXiv:2311.09018 [cs.LG]*, 2023.
- [115] Z. Wang, P. Glynn, and Y. Ye. Likelihood robust optimization for data-driven problems. *Computational Management Science*, pages 1–21, 2015.
- [116] R. Williamson and A. Menon. Fairness risk measures. In *Proceedings of the 36th International Conference on Machine Learning*, pages 6786–6797. PMLR, 2019.
- [117] B. Woodworth, S. Gunasekar, M. I. Ohanessian, and N. Srebro. Learning non-discriminatory predictors. In *Conference on Learning Theory*, pages 1920–1953. PMLR, 2017.
- [118] M. Wortsman, G. Ilharco, M. Li, J. W. Kim, H. Hajishirzi, A. Farhadi, H. Namkoong, and L. Schmidt. Robust fine-tuning of zero-shot models. *arXiv:2109.01903 [cs.CV]*, 2021.
- [119] D. Wozabal and N. Wozabal. Asymptotic consistency of risk functionals. *Journal of Nonparametric Statistics*, 21(8):977–990, 2009.
- [120] C. Xu, J. Lee, X. Cheng, and Y. Xie. Flow-based distributionally robust optimization. *IEEE Journal on Selected Areas in Information Theory*, 2024.
- [121] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals. Understanding deep learning requires rethinking generalization. In *Proceedings of the Fifth International Conference on Learning Representations*, 2017.
- [122] J.-J. Zhu, W. Jitkrittum, M. Diehl, and B. Schölkopf. Kernel distributionally robust optimization: Generalized duality theorem and stochastic approximation. In *International Conference on Artificial Intelligence and Statistics*, pages 280–288. PMLR, 2021.

A Proof of Theorem 1

Let $D := (X, Y, Z)$, $I_k^{c, \infty}$ be the set of indices *not* in I_k (as $n \rightarrow \infty$) and define T to be the debiased functional

$$T(P; \mu, \tau) := \inf_{\eta} \left\{ \frac{1}{\alpha} \mathbb{E}_P (\mu(Z) - \eta)_+ + \eta \right\} + \mathbb{E}_P [\tau(Z) (\ell(\theta(X); Y) - \mu(Z))]. \quad (\text{A.1})$$

The cross-fitted estimator is $\hat{\omega}_\alpha = \frac{1}{K} \sum_{k=1}^K T(\hat{P}_k; \hat{\mu}_k, \hat{\tau}_k)$, where \hat{P}_k is the empirical distribution on the k -th fold. Our goal is to show

$$\begin{aligned} & \sqrt{|I_k|} \left(T(\hat{P}_k; \hat{\mu}_k, \hat{\tau}_k) - T(P; \mu^*, \tau^*) \right) \\ &= \sqrt{|I_k|} \left(T(\hat{P}_k; \hat{\mu}_k, \hat{\tau}_k) - T(P; \hat{\mu}_k, \hat{\tau}_k) \right) + \sqrt{|I_k|} \left(T(P; \hat{\mu}_k, \hat{\tau}_k) - T(P; \mu^*, \tau^*) \right) \\ &= \frac{1}{\sqrt{|I_k|}} \sum_{i \in I_k} \psi(D_i) + o_p(1). \end{aligned}$$

We begin by establishing

$$\sqrt{|I_k|} \left(T(\hat{P}_k; \hat{\mu}_k, \hat{\tau}_k) - T(P; \hat{\mu}_k, \hat{\tau}_k) \right) = \frac{1}{\sqrt{|I_k|}} \sum_{i \in I_k} \psi(D_i) + o_p(1). \quad (\text{A.2})$$

Using basic epiconvergence results [94, Theorem 7.17] and their implications for convergence of optimal solution sets of convex optimization problems [94, Theorem 7.31], restrict the dual (2.2) to a compact set $S_\alpha := [P_{1-\alpha}^{-1}(\mu^*) \pm 1]$

$$\inf_{\eta \in S_\alpha} \left\{ \frac{1}{\alpha} \mathbb{E}_{Z \sim Q} (\hat{\mu}_k(Z) - \eta)_+ + \eta \right\} = \inf_{\eta} \left\{ \frac{1}{\alpha} \mathbb{E}_{Z \sim Q} (\hat{\mu}_k(Z) - \eta)_+ + \eta \right\} \quad \text{eventually} \quad (\text{A.3})$$

for $Q = \hat{P}_k, P$. Letting

$$T_{S_\alpha}(Q; \mu, \tau) := \inf_{\eta \in S_\alpha} \left\{ \frac{1}{\alpha} \mathbb{E}_{D \sim Q} (\mu(Z) - \eta)_+ + \eta \right\} + \mathbb{E}_{D \sim Q} [\tau(Z) (\ell(\theta(X); Y) - \mu(Z))], \quad (\text{A.4})$$

we show the convergence (A.2) with T replaced by T_{S_α} , conditional on the event where $\hat{\mu}_k \in \mathcal{U}$

$$\mathcal{E}_{n,k} := \{ \hat{\mu}_k \in \mathcal{U}, \text{ and conditions of Assumption B holds for } k \}. \quad (\text{A.5})$$

This implies the unconditional result (A.2) since $\mathbb{P}(\mathcal{E}_{n,k}) \rightarrow 1$ by Assumptions B, C.

Standard empirical process theory guarantees that the empirical measure \hat{P}_k satisfies the uniform CLT over random variables

$$\begin{aligned} f_{n,\eta}(D) &:= \frac{1}{\alpha} (\hat{\mu}_k(Z) - \eta)_+ + \eta \text{ for } \eta \in S_\alpha, \\ f_{n, P_{1-\alpha}^{-1}(\mu^*)+2}(D) &:= \hat{\tau}_k(Z) (\ell(\theta(X); Y) - \hat{\mu}_k(Z)) \end{aligned}$$

indexed by $\Lambda := S_\alpha \cup \{P_{1-\alpha}^{-1}(\mu^*) + 2\}$. Formally, let $\ell^\infty(\Lambda)$ be the usual space of uniformly bounded functions on Λ endowed with the sup norm. We treat measures as bounded functionals $\hat{P}_k : \eta \mapsto \mathbb{E}_{D \sim \hat{P}_k} f_{n,\eta}(D)$ and $P_n : \eta \mapsto \mathbb{E}_{D \sim P} f_{n,\eta}(D)$.

Proposition 7 (Jeong and Namkoong [62, Prop. 4]). *Conditional on $\mathcal{E}_{n,k}$,*

$$\sqrt{n} \left(\mathbb{E}_{D \sim \hat{P}_k} f_{n,\eta}(D) - \mathbb{E}_{D \sim P} f_{n,\eta}(D) \right) \xrightarrow{d} \mathbb{G} \text{ in } \ell^\infty(\Lambda),$$

where \mathbb{G} is a Gaussian process on $\Lambda = S_\alpha \cup \{P_{1-\alpha}^{-1}(\mu^*) + 2\}$ with covariance $\Sigma(\eta, \eta')$

$$\begin{aligned} & \frac{1}{\alpha^2} \mathbb{E} \left[(\mu^*(Z) - \eta)_+ (\mu^*(Z) - \eta')_+ \right] + \frac{\eta'}{\alpha} \mathbb{E} \left[(\mu^*(Z) - \eta)_+ \right] + \frac{\eta}{\alpha} \mathbb{E} \left[(\mu^*(Z) - \eta')_+ \right] \text{ if } \eta, \eta' \in S_\alpha \\ & \frac{1}{\alpha^2} \mathbb{E} \left[(\mu^*(Z) - \eta)_+ (\ell(\theta(X); Y) - \mu^*(Z)) \right] \text{ if } \eta \in S_\alpha, \eta' = P_{1-\alpha}^{-1}(\mu^*) + 2 \\ & \mathbb{E} \left[\tau^*(Z)^2 (\ell(\theta(X); Y) - \mu^*(Z))^2 \right] \text{ if } \eta = \eta' = P_{1-\alpha}^{-1}(\mu^*) + 2. \end{aligned}$$

Recall that Danskin's theorem [102, Theorem 7.21] says the optimization functional

$$\lambda_{\text{opt}}(H) := \inf_{\eta \in S_\alpha} H(\eta)$$

is uniformly Hadamard differentiable. This allows us to use the functional delta method [110, Delta method, Theorem 3.9.5] to the map $Q \mapsto T_{S_\alpha}(Q; \hat{\mu}_k, \hat{\tau}_k)$, which gives the convergence (A.2).

Next, we show the term $\sqrt{|I_k|} (T(P; \hat{\mu}_k, \hat{\tau}_k) - T(P; \mu^*, \tau^*))$ vanishes. Let $\mathfrak{R}_k : [0, 1] \rightarrow \mathbb{R}$

$$\mathfrak{R}_k(r) := T(P; (1-r)(\mu^*, \tau^*) + r(\hat{\mu}_k, \hat{\tau}_k)) - T(P; \mu^*, \tau^*), \quad (\text{A.7})$$

so that $\mathfrak{R}_k(0) = 0$, and $\mathfrak{R}_k(1)$ is equal to $T(P; \hat{\mu}_k, \hat{\tau}_k) - T(P; \mu^*, \tau^*)$. On the event $\mathcal{E}_{n,k}$, $\mathfrak{R}_k(r)$ is differentiable under Assumptions A, B

$$\mathfrak{R}'_k(r) = \mathbb{E}_{Z \sim P} \left[(\hat{\mu}_k - \mu^*)(Z) \left(\frac{1}{\alpha} \mathbf{1} \{ \hat{\mu}_{k,r}(Z) \geq P_{1-\alpha}^{-1}(\hat{\mu}_{k,r}) \} - \tau^*(Z) \right) \right] \quad (\text{A.8})$$

where $(\hat{\mu}_{k,r}, \hat{\tau}_{k,r}) := (\mu^*, \tau^*) + r((\hat{\mu}_k, \hat{\tau}_k) - (\mu^*, \tau^*))$. (It is easy to check this using Danskin's theorem.)

The mean value theorem then gives $\mathfrak{R}_k(1) = \mathfrak{R}_k(0) + \mathfrak{R}'_k(r) \cdot (1 - 0) = \mathfrak{R}'_k(r)$ for some $r \in [0, 1]$. Debiasing nominally guarantees $\mathfrak{R}'_k(0) = 0$, but going further we will now show $\sup_{r \in [0, 1]} |\mathfrak{R}'_k(r)| = o_p(n^{-1/2})$. Since $\hat{\tau}_k \in [-M, M]$ on $\mathcal{E}_{n,k}$, elementary calculations and repeated applications of Holder's inequality yield

$$\sup_{r \in [0, 1]} |\mathfrak{R}'_k(r)| \leq \|\hat{\mu}_k - \mu^*\|_{L^\infty(\mathcal{X})} \sup_{r \in [0, 1]} \left\| \frac{1}{\alpha} \mathbf{1} \{ \hat{\mu}_{k,r}(Z) \geq P_{1-\alpha}^{-1}(\hat{\mu}_{k,r}) \} - \tau^*(Z) \right\|_{L^1(\mathcal{X})} \quad (\text{A.9})$$

where C is a positive constant that only depends on c , and M . The last term in the bound is bounded by $\delta_n n^{-1/2}$ by the definition of $\mathcal{E}_{n,k}$.

Our uniform differentiability assumption (4.1) for $F_{\hat{\mu}_{k,r}}$ guarantees the following notions of smoothness. We omit its derivations as the calculations are elementary but tedious [62].

Lemma 4. *On the event $\mathcal{E}_{n,k}$, we have $\sup_{r \in [0, 1]} |P_{1-\alpha}^{-1}(\hat{\mu}_{k,r}) - P_{1-\alpha}^{-1}(\mu^*)| = O(\|\hat{\mu}_k - \mu^*\|_{L^\infty(\mathcal{X})}) = O(\delta_n n^{-1/3})$ and*

$$\begin{aligned} & \sup_{r \in [0, 1]} \left\| \frac{1}{\alpha} \mathbf{1} \{ \hat{\mu}_{k,r}(Z) \geq P_{1-\alpha}^{-1}(\hat{\mu}_{k,r}) \} - \tau^*(Z) \right\|_{L^1(\mathcal{X})} \\ & \lesssim n^{1/6} \left(\|\hat{\mu}_k - \mu^*\|_{L^1(\mathcal{X})} + |\hat{q}_k - P_{1-\alpha}^{-1}(\hat{\mu}_k)| + \sup_{r \in [0, 1]} |P_{1-\alpha}^{-1}(\hat{\mu}_{k,r}) - P_{1-\alpha}^{-1}(\mu^*)| \right) \\ & \quad + n^{-1/6} \delta_n + \|\hat{\mu}_k - \mu^*\|_{L^\infty(\mathcal{X})}. \end{aligned}$$

We conclude that the bound (A.9) is $O(\delta_n n^{-1/2})$ on the event $\mathcal{E}_{n,k}$, meaning $\sup_{r \in [0, 1]} |\mathfrak{R}'_k(r)| = o(n^{-1/2})$ on the event $\mathcal{E}_{n,k}$. Since $\mathbb{P}(\mathcal{E}_{n,k}) \rightarrow 1$ from Assumptions B, C, we have the final result.

B Proof of finite-sample concentration results

Our results are based on a general concentration guarantee for estimating the dual reformulation (2.2) for any given $\mu(Z)$. We give this result in Appendix B.1, and build on it in subsequent proofs of key results. In the following, we use \lesssim to denote inequality up to a numerical constant that may change line by line.

B.1 Concentration bounds for worst-case subpopulation performance

Since $\ell(\hat{y}; y) \geq 0$ for losses used in most machine learning problems, we assume that \mathcal{H} consists of nonnegative functions. To show exponential concentration guarantees, we consider sub-Gaussian conditional risk models $\mu(Z)$. Note the concentration results here are more general than needed for the purpose of proving the main results, because any random variable bounded in $[0, B]$ is inherently sub-Gaussian with parameter $B^2/4$.

Definition 2. A function $\mu : \mathcal{Z} \rightarrow \mathbb{R}$ with $\mathbb{E}|\mu(Z)| < \infty$ is sub-Gaussian with parameter σ^2 if

$$\mathbb{E}[\exp(\lambda(\mu(Z) - \mathbb{E}[\mu(Z)]))] \leq \exp\left(\frac{\sigma^2 \lambda^2}{2}\right) \quad \text{for all } \lambda \in \mathbb{R}.$$

The sub-Gaussian assumption can be relaxed to sub-exponential random variables, with minor and standard modifications to subsequent results. We omit these results for brevity.

Define a dual plug-in estimator for the worst-case subpopulation performance of $\mu(Z)$ on I_k

$$\widehat{W}_{\alpha,k}(\mu) = \inf_{\eta} \left\{ \frac{1}{\alpha |I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ + \eta \right\}. \quad (\text{B.1})$$

The following result shows that for any sub-Gaussian μ that is bounded from below, the plug-in estimator (6) converges at the rate $O_p(|I_k|^{-1/2})$.

Proposition 8. There is a universal constant $C > 0$ such that for all $h \geq 0$ that is sub-Gaussian with parameter σ^2 ,

$$|\widehat{W}_{\alpha,k}(\mu) - W_{\alpha}(\mu)| \leq \frac{C\sigma}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}} \quad \text{with probability at least } 1 - \delta.$$

We prove the proposition in the rest of the subsection. By a judicious application of the empirical process theory, our bounds—which apply to nonnegative random variables—are simpler than existing concentration guarantees for conditional value-at-risk [23, 85].

Our starting point is the following claim, which bounds $|\widehat{W}_{\alpha,k}(\mu) - W_{\alpha}(\mu)|$ in terms of the suprema of empirical process on $\{z \mapsto (\mu(z) - \eta)_+ : \eta \geq 0\}$.

Claim 5.

$$\left| \widehat{W}_{\alpha,k}(\mu) - W_{\alpha}(\mu) \right| \leq \frac{1}{\alpha} \sup_{\eta \geq 0} \left| \frac{1}{|I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ - \mathbb{E}(\mu(Z) - \eta)_+ \right| \quad (\text{B.2})$$

The crux of this claim is that η does not range over \mathbb{R} , but rather has a lower bound; the value 0 can be replaced with any almost sure lower bound on $\mu(Z)$. Deferring the proof of Claim 5 to

the end of the subsection, we proceed by bounding the suprema of the empirical process in the preceding display.

We begin by introducing requisite concepts in empirical process theory, which we use in the rest of the proof; we refer readers to van der Vaart and Wellner [110] for a comprehensive treatment. Recall the definition of Orlicz norms, which allows controlling the tail behavior of random variables.

Definition 3 (Orlicz norms). *Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ be a non-decreasing, convex function with $\psi(0) = 0$. For any random variable W , its Orlicz norm $\|W\|_\psi$ is*

$$\|W\|_\psi := \inf \left\{ t > 0 : \mathbb{E} \left[\psi \left(\frac{|W|}{t} \right) \right] \leq 1 \right\}.$$

Remark 1: From Markov's inequality, we have

$$\mathbb{P}(|W| > t) \leq \mathbb{P} \left(\psi \left(\frac{|W|}{\|W\|_\psi} \right) \geq \psi \left(\frac{t}{\|W\|_\psi} \right) \right) \leq \psi \left(\frac{t}{\|W\|_\psi} \right)^{-1}.$$

For $\psi_p(s) = e^{s^p} - 1$, a similar argument yields

$$\mathbb{P}(|W| > t) \leq 2 \exp \left(-t^p / \|W\|_{\psi_p}^p \right). \quad (\text{B.3})$$

◇

A sub-Gaussian random variable $\mu(Z)$ with parameter σ^2 has bounded Orlicz norm $\|\mu(Z)\|_{\psi_2} \leq 2\sigma$ (see, for example, Wainwright [112, Section 2.4] and van der Vaart and Wellner [110, Lemma 2.2.1]).

Remark 2: The converse also holds: for W such that $\mathbb{P}(|W| > t) \leq c_1 \exp(-c_2 t^p)$ for all t , and constants $c_1, c_2 > 0$ and $p \geq 1$, Fubini gives

$$\mathbb{E} \left[\exp \left(\frac{|W|^p}{t^p} \right) - 1 \right] = \mathbb{E} \left[\int_0^{|W|^p} t^{-1/p} \exp(t^{-1/p} s) ds \right] = \int_0^\infty \mathbb{P}(|W|^p > s) t^{-1/p} \exp \left(t^{-1/p} s \right) ds.$$

Using the tail probability bound, the preceding display is bounded by

$$c_1 \int_0^\infty \exp(-c_2 s) t^{-1/p} \exp(t^{-1/p} s) ds = \frac{c_1 t^{-1/p}}{c_2 - t^{-1/p}}.$$

So the Orlicz norm $\|W\|_{\psi_p}$ is bounded by $\left(\frac{1+c_1}{c_2} \right)^{1/p}$. ◇

In the following, we let W be the right hand side of the bound (B.2), and control its Orlicz norm $\|W\|_{\psi_2}$ using Dudley's entropy integral [110]. We use the standard notion of the covering number. For a vector space \mathcal{V} , let $V \subset \mathcal{V}$ be a collection of vectors. Letting $\|\cdot\|$ be a norm on \mathcal{V} , a collection $\{v_1, \dots, v_N\} \subset \mathcal{V}$ is an ϵ -cover of \mathcal{V} if for each $v \in \mathcal{V}$, there is a v_i satisfying $\|v - v_i\| \leq \epsilon$. The *covering number* of V with respect to $\|\cdot\|$ is

$$N(\epsilon, V, \|\cdot\|) := \inf \{ N \in \mathbb{N} : \text{there is an } \epsilon\text{-cover of } V \text{ with respect to } \|\cdot\| \}.$$

For a collection \mathcal{H} of functions $f : \mathcal{Z} \rightarrow \mathbb{R}$, let F be its envelope function such that $|f(z)| \leq F(z)$ for all $z \in \mathcal{Z}$. The following result controls the suprema of empirical processes using the (uniform) metric entropy. The result is based on involved chaining arguments [110, Section 2.14].

Lemma 6 (van der Vaart and Wellner [110, Theorem 2.14.1 and 2.14.5]).

$$\begin{aligned} & \sqrt{|I_k|} \left\| \sup_{f \in \mathcal{H}} \left| \frac{1}{|I_k|} \sum_{i \in I_k} f(Z_i) - \mathbb{E}f(Z) \right| \right\|_{\psi_2} \\ & \lesssim \|F\|_{\psi_2} + \|F\|_{L^2(P)} \sup_Q \int_0^1 \sqrt{1 + \log N(\epsilon \|F\|_{L^2(Q)}, \mathcal{H}, L^2(Q))} d\epsilon, \end{aligned}$$

where the supremum is over all discrete probability measures Q such that $\|F\|_{L^2(Q)} > 0$.

Evidently, $F(z) = (\mu(z))_+ = \mu(z)$ is an envelope function for the following class of functions

$$\mathcal{H} = \{z \mapsto (\mu(z) - \eta)_+ : \eta \geq 0\}.$$

Using the tail probability bound (B.3), we conclude

$$\begin{aligned} & \sup_{\eta \geq 0} \left| \frac{1}{|I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ - \mathbb{E}(\mu(Z) - \eta)_+ \right| \\ & \lesssim \sqrt{\frac{\log(2/\delta)}{|I_k|}} \left(\|F\|_{\psi_2} + \|F\|_2 \sup_Q \int_0^1 \sqrt{1 + \log N(\epsilon \|F\|_{L^2(Q)}, \mathcal{H}, L^2(Q))} d\epsilon \right), \end{aligned}$$

with probability at least $1 - \delta$.

Since we have $\|F\|_{L^2(P)} \leq \|F\|_{\psi_2} \lesssim \sigma$, it now suffices to show that the above uniform metric entropy is bounded by a universal constant. We use the standard notion of VC-dimension [110, Chapter 2.6, page 135].

Lemma 7 (van der Vaart and Wellner [110, Theorem 2.6.7]). *Let $\text{VC}(\mathcal{H})$ be the VC-dimension of the collection of subsets $\{(z, t) : t < f(x)\}$ for $f \in \mathcal{H}$. For any probability measure Q such that $\|F\|_{L^2(Q)} > 0$ and $0 < \epsilon < 1$, we have*

$$N(\epsilon \|F\|_{L^2(Q)}, \mathcal{H}, L^2(Q)) \lesssim \text{VC}(\mathcal{H})(16e)^{\text{VC}(\mathcal{H})} \left(\frac{1}{\epsilon}\right)^{2(\text{VC}(\mathcal{H})-1)}.$$

Translations of a monotone function on \mathbb{R} has VC-dimension 2.

Lemma 8 (van der Vaart and Wellner [110, Theorem 2.6.16]). *The class of functions $\mathcal{H}' = \{z \mapsto (\mu(z) - \eta)_+ : \eta \in \mathbb{R}\}$ has VC-dimension $\text{VC}(\mathcal{H}') = 2$.*

From Lemmas 7 and 8, we conclude that for the function class $\mathcal{H} = \{z \mapsto (\mu(z) - \eta)_+ : \eta \geq 0\}$, the uniform metric entropy

$$\sup_Q \int_0^1 \sqrt{1 + \log N(\epsilon \|F\|_{L^2(Q)}, \mathcal{H}, L^2(Q))} d\epsilon$$

is bounded by a universal constant. This gives our desired result.

Proof of Claim 5 To show the bound (B.2), we use the dual reformulation for both $W_\alpha(\mu)$ and its empirical approximation $\widehat{W}_{\alpha,k}(\mu)$ on I_k . For any probability measure P , recall two different definitions of the quantile of $\mu(Z)$

$$P_{1-\alpha}^{-1}(\mu(Z)) := \inf\{t : \mathbb{P}_Z(\mu(Z) \leq t) \geq 1 - \alpha\}$$

$$P_{1-\alpha,+}^{-1}(\mu(Z)) := \inf\{t : \mathbb{P}_Z(\mu(Z) \leq t) > 1 - \alpha\}.$$

We call $P_{1-\alpha,+}^{-1}(\mu(Z))$ the upper $(1 - \alpha)$ -quantile. The two values characterize the optimal solution set of the dual problem (2.2); they are identical when $\mu(Z)$ has a positive density at $P_{1-\alpha}^{-1}(\mu(Z))$.

Lemma 9 (Rockafellar and Uryasev [93, Theorem 10]). *For any probability measure P such that $\mu(Z) \geq 0$ P -a.s. and $\mathbb{E}_P[\mu(Z)_+] < \infty$, we have*

$$[P_{1-\alpha}^{-1}(\mu(Z)), P_{1-\alpha,+}^{-1}(\mu(Z))] = \operatorname{argmin}_{\eta \in \mathbb{R}} \left\{ \frac{1}{\alpha} \mathbb{E}_P(\mu(Z) - \eta)_+ + \eta \right\}.$$

Since P was an arbitrary measure in Lemmas 1 and 9, identical results follow for the empirical distribution on I_k . Hence, we have

$$\begin{aligned} \left| \widehat{W}_{\alpha,k}(\mu) - W_{\alpha}(\mu) \right| &= \left| \inf_{\eta \in \mathbb{R}} \left\{ \frac{1}{\alpha |I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ + \eta \right\} - \inf_{\eta \in \mathbb{R}} \left\{ \frac{1}{\alpha} \mathbb{E}(\mu(Z) - \eta)_+ + \eta \right\} \right| \\ &= \left| \inf_{\eta \geq 0} \left\{ \frac{1}{\alpha |I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ + \eta \right\} - \inf_{\eta \geq 0} \left\{ \frac{1}{\alpha} \mathbb{E}(\mu(Z) - \eta)_+ + \eta \right\} \right| \end{aligned}$$

where we used Lemma 9 to restrict the feasible region in the last equality. The preceding display is then bounded by

$$\sup_{\eta \geq 0} \left| \frac{1}{\alpha |I_k|} \sum_{i \in I_k} (\mu(Z_i) - \eta)_+ + \eta - \frac{1}{\alpha} \mathbb{E}(\mu(Z) - \eta)_+ - \eta \right|.$$

B.2 Proof of Theorem 2

We abuse notation and use C for a numerical constant that may change line to line. From the decomposition (4.4), it suffices to bound term (a) and term (b) separately.

Term (b) can be bounded with the help of Proposition 8 because $\widehat{\mu}_k(\cdot)$ is trained on a sample I_k^c independent from I_k used to estimate the worst-case subpopulation performance (Eq. (B.1)). More precisely, recalling that any bounded random variable random variable taking values in $[0, B]$ is sub-Gaussian with parameter $B^2/4$, Proposition 8 implies

$$\left| \widehat{W}_{\alpha,k}(\widehat{\mu}_k) - W_{\alpha}(\widehat{\mu}_k) \right| \leq \frac{CB}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}} \quad \text{with probability at least } 1 - \delta.$$

For the debiasing term, Hoeffding's inequality implies with probability at least $1 - \delta$ conditional on I_k^c ,

$$\left| \mathbb{E}_{\widehat{P}_k}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c] - \mathbb{E}_P[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c] \right| \lesssim \frac{B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}},$$

because $\widehat{\tau}_k \in \{0, 1/\alpha\}$ and $\ell(\cdot, \cdot), \mu(\cdot) \in [0, B]$. Then the same bound holds with total probability at least $1 - \delta$. Hence, with probability at least $1 - 2\delta$, term (b) is bounded by

$$(b) \equiv T(P; \widehat{\mu}_k, \widehat{\tau}_k) - T(\widehat{P}_k; \widehat{\mu}_k, \widehat{\tau}_k) \lesssim \frac{B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}}.$$

To bound term (a) in the decomposition (4.4), we first note

$$\begin{aligned} |\mathbf{W}_\alpha(\widehat{\mu}_k) - \mathbf{W}_\alpha(\mu^*)| &\leq \frac{1}{\alpha} \sup_{\eta} \left| \mathbb{E} \left[(\widehat{\mu}_k(Z) - \eta)_+ \mid I_k^{\mathcal{C}} \right] - \mathbb{E} (\mu^*(Z) - \eta)_+ \right| \\ &\leq \frac{1}{\alpha} \mathbb{E} \left[|\widehat{\mu}_k(Z) - \mu^*(Z)| \mid I_k^{\mathcal{C}} \right], \end{aligned}$$

where the first inequality follows from the dual (2.2), and the second inequality follows from the non-expansiveness of the function $(\cdot)_+$. Similarly for the debiasing term,

$$\begin{aligned} |\mathbb{E}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^{\mathcal{C}}]| &= |\mathbb{E}[\mathbb{E}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid Z, I_k^{\mathcal{C}}] \mid I_k^{\mathcal{C}}]| \\ &= |\mathbb{E}[\widehat{\tau}_k(Z)(\mu^*(Z) - \widehat{\mu}_k(Z)) \mid I_k^{\mathcal{C}}]| \\ &\leq \frac{1}{\alpha} \mathbb{E} \left[|\widehat{\mu}_k(Z) - \mu^*(Z)| \mid I_k^{\mathcal{C}} \right], \end{aligned}$$

where the first equality follows from the law of total probability, the second by definition of μ^* , and the inequality because $\widehat{\tau}_k(\cdot) \in \{0, 1/\alpha\}$. Hence,

$$\begin{aligned} |(a)| &\leq |\mathbf{W}_\alpha(\mu^*) - \mathbf{W}_\alpha(\widehat{\mu}_k)| + |\mathbb{E}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^{\mathcal{C}}]| \\ &\leq \frac{2}{\alpha} \mathbb{E} \left[|\widehat{\mu}_k(Z) - \mu^*(Z)| \mid I_k^{\mathcal{C}} \right], \\ &\leq \frac{2}{\alpha} \sqrt{\mathbb{E} \left[(\widehat{\mu}_k(Z) - \mu^*(Z))^2 \mid I_k^{\mathcal{C}} \right]} = \frac{2}{\alpha} \sqrt{\text{err}(\mathcal{H}, I_k^{\mathcal{C}})}, \end{aligned}$$

where the first inequality follows from the definition of (a), the second inequality follows from the bounds above, the last inequality uses Holder inequality, and we define the generalization error for the first-stage estimation problem (3.1) based on $I_k^{\mathcal{C}}$,

$$\begin{aligned} \text{err}(\mathcal{H}, I_k^{\mathcal{C}}) &:= \mathbb{E} \left[(\mu^*(Z) - \widehat{\mu}_k(Z))^2 \mid I_k^{\mathcal{C}} \right] \\ &= \mathbb{E} \left[(\ell(\theta(X); Y) - \widehat{\mu}_k(Z))^2 \mid I_k^{\mathcal{C}} \right] - \mathbb{E}(\ell(\theta(X); Y) - \mu^*(Z))^2 \\ &= \mathbb{E} \left[(\ell(\theta(X); Y) - \widehat{\mu}_k(Z))^2 \mid I_k^{\mathcal{C}} \right] - \mathbb{E}(\ell(\theta(X); Y) - \bar{\mu}(Z))^2 + \mathbb{E}(\mu^*(Z) - \bar{\mu}(Z))^2 \end{aligned}$$

We use the following concentration result based on the localized Rademacher complexity [9].

Lemma 10 (Bartlett et al. [9, Corollary 5.3]). *Let Assumption D hold. Then, with probability at least $1 - \delta$,*

$$\mathbb{E} \left[(\ell(\theta(X); Y) - \widehat{\mu}_k(Z))^2 \mid I_k^{\mathcal{C}} \right] - \mathbb{E}(\ell(\theta(X); Y) - \bar{\mu}(Z))^2 \leq CB^2 \left(r_{|I_k^{\mathcal{C}}|}^* + \frac{\log(1/\delta)}{|I_k^{\mathcal{C}}|} \right).$$

Using $\sqrt{a+b+c} \leq \sqrt{a} + \sqrt{b} + \sqrt{c}$ for $a, b, c \geq 0$, we have the desired result.

B.3 Proof of Theorem 3

Instead of the decomposition (4.4) we use for Theorem 2, we use an alternative form

$$\widehat{\omega}_{\alpha,k} - \mathbf{W}_\alpha(\mu^*) = \underbrace{\widehat{\mathbf{W}}_{\alpha,k}(\widehat{\mu}_k) - \widehat{\mathbf{W}}_{\alpha,k}(\mu^*)}_{(a): \text{ first stage}} + \underbrace{\widehat{\mathbf{W}}_{\alpha,k}(\mu^*) - \mathbf{W}_\alpha(\mu^*)}_{(b): \text{ second stage}}$$

$$+ \underbrace{\mathbb{E}_{\widehat{P}_k} [\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c]}_{(c): \text{ debiasing term}} \quad (\text{B.4})$$

Term (b) can be bounded using Proposition 8 as before. Without assuming $\mu^* \in \mathcal{H}$, recall that any bounded random variable taking values in $[0, B]$ is sub-Gaussian with parameter $B^2/4$, so Proposition 8 yields

$$|(b)| = \left| \widehat{W}_{\alpha, k}(\mu^*) - W_\alpha(\mu^*) \right| \leq C \frac{B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}} \quad \text{with probability at least } 1 - \delta.$$

We first notice the following bound on term (a).

$$\begin{aligned} (a) &= \left| \widehat{W}_{\alpha, k}(\widehat{\mu}_k) - \widehat{W}_{\alpha, k}(\mu^*) \right| \leq \frac{1}{\alpha} \sup_{\eta} \left| \mathbb{E}_{\widehat{P}_k} [(\widehat{\mu}_k(Z_i) - \eta)_+ - (\mu^*(Z_i) - \eta)_+ \mid I_k^c] \right| \\ &\leq \frac{1}{\alpha} \mathbb{E}_{\widehat{P}_k} [|\widehat{\mu}_k(Z) - \mu^*(Z)| \mid I_k^c] \equiv \frac{1}{\alpha} \|\widehat{\mu}_k(Z) - \mu^*(Z)\|_{L^1(\widehat{P}_k|I_k^c)}. \end{aligned} \quad (\text{B.5})$$

Next we will bound the debiasing term (c) with the same quantity. We start by observing Hoeffding's inequality implies

$$\left| \mathbb{E}_{\widehat{P}_k} [\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c] - \mathbb{E}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c] \right| \leq \frac{B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}}$$

with probability at least $1 - \delta$ because $\widehat{\tau}_k \in \{0, 1/\alpha\}$ and $\ell, \widehat{\mu}_k \in [0, B]$ almost surely. Then notice by definition of $\mu^* \equiv \mathbb{E}[\ell(\theta(X); Y) \mid Z]$ we know $\mathbb{E}[\widehat{\tau}_k(Z)(\ell(\theta(X); Y) - \widehat{\mu}_k(Z)) \mid I_k^c] = \mathbb{E}[\widehat{\tau}_k(Z)(\mu^*(Z) - \widehat{\mu}_k(Z)) \mid I_k^c]$ by conditioning on Z . Then we again invoke Hoeffding's inequality to argue with probability at least $1 - \delta$,

$$\left| \mathbb{E}[\widehat{\tau}_k(Z)(\mu^*(Z) - \widehat{\mu}_k(Z)) \mid I_k^c] - \mathbb{E}_{\widehat{P}_k} [\widehat{\tau}_k(Z)(\mu^*(Z) - \widehat{\mu}_k(Z)) \mid I_k^c] \right| \leq \frac{B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}}.$$

Lastly we notice

$$\begin{aligned} \left| \mathbb{E}_{\widehat{P}_k} [\widehat{\tau}_k(Z)(\mu^*(Z) - \widehat{\mu}_k(Z)) \mid I_k^c] \right| &\leq \frac{1}{\alpha} \mathbb{E}_{\widehat{P}_k} [|\mu^*(Z) - \widehat{\mu}_k(Z)| \mid I_k^c] \\ &\equiv \frac{1}{\alpha} \|\mu^*(Z) - \widehat{\mu}_k(Z)\|_{L^1(\widehat{P}_k|I_k^c)} \end{aligned}$$

because $\widehat{\tau}_k \in \{0, 1/\alpha\}$. Hence, we conclude with probability at least $1 - 2\delta$,

$$|(c)| \leq \frac{1}{\alpha} \|\mu^*(Z) - \widehat{\mu}_k(Z)\|_{L^1(\widehat{P}_k|I_k^c)} + \frac{2B}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}}.$$

Thus we have shown with probability at least $1 - 3\delta$,

$$|\widehat{W}_{\alpha, k} - W_\alpha^*| \leq \frac{2}{\alpha} \|\mu^*(Z) - \widehat{\mu}_k(Z)\|_{L^1(\widehat{P}_k|I_k^c)} + \frac{CB}{\alpha} \sqrt{\frac{\log(2/\delta)}{|I_k|}}.$$

We now present two approaches to bounding the empirical L^1 -norm of the error $\mu^*(Z) - \widehat{\mu}_k(Z)$ for whether the model class \mathcal{H} is convex. Before we move on, notice the following identity that is useful for both cases, which can be interpreted geometrically as the cosine theorem in the $L^2(\widehat{P}_k)$ space. For any two functions $\mu, \tilde{\mu} : \mathcal{Z} \rightarrow [0, B]$,

$$\|\tilde{\mu}(Z) - \mu(Z)\|_{L^2(\widehat{P}_k)} = \Delta_{I_k}(\tilde{\mu}) - \Delta_{I_k}(\mu) + 2\mathbb{E}_{\widehat{P}_k} [(\ell(\theta(X); Y) - \mu(Z))(\tilde{\mu}(Z) - \mu(Z))]. \quad (\text{B.6})$$

B.3.1 Continuing proof of Theorem 3 with a non-convex model class

First note Hölder's inequality implies

$$\|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^1(\hat{P}_k|I_k^c)} \leq \|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)},$$

so it suffices to bound the L^2 -norm. In order to use the identity B.6 with $\mu = \mu^*$ and $\tilde{\mu} = \hat{\mu}_k$, we notice by definition of $\mu^* \equiv \mathbb{E}[\ell(\theta(X); Y) | Z]$ that $\mathbb{E}[(\ell(\theta(X); Y) - \mu^*(Z))(\hat{\mu}_k(Z) - \mu^*(Z))] = 0$ for all $\tilde{\mu}$. Since $(\ell(\theta(X); Y) - \mu^*(Z))(\hat{\mu}_k(Z) - \mu^*(Z))$ is almost surely bounded in $[-B^2, B^2]$ and i.i.d., Hoeffding inequality [112, Ch. 2] yields

$$\left| \mathbb{E}_{\hat{P}_k}[(\ell(\theta(X); Y) - \mu^*(Z))(\hat{\mu}_k(Z) - \mu^*(Z))] \right| \leq B^2 \sqrt{\frac{2 \log(2/\delta)}{|I_k|}} \quad \text{with probability at least } 1 - \delta.$$

Hence, the identity B.6 implies with probability at least $1 - \delta$,

$$\|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)}^2 \leq [\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})] + [\Delta_{I_k}(\bar{\mu}) - \Delta_{I_k}(\mu^*)] + B^2 \sqrt{\frac{2 \log(2/\delta)}{|I_k|}}$$

Similarly, Hoeffding inequality implies with probability at least $1 - \delta$,

$$\begin{aligned} \Delta_{I_k}(\bar{\mu}) - \Delta_{I_k}(\mu^*) &\leq \mathbb{E}(\ell(\theta(X); Y) - \bar{\mu}(Z))^2 - \mathbb{E}(\ell(\theta(X); Y) - \mu^*(Z))^2 + B^2 \sqrt{\frac{2 \log(1/\delta)}{|I_k|}} \\ &= \|\bar{\mu} - \mu^*\|_{L^2}^2 + B^2 \sqrt{\frac{2 \log(1/\delta)}{|I_k|}}, \end{aligned}$$

where the equality follows by the definition of the conditional risk $\mu^*(Z) \equiv \mathbb{E}[\ell(\theta(X); Y) | Z]$. Hence, with probability at least $1 - 2\delta$,

$$\begin{aligned} \|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)} &\leq \sqrt{\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu}) + \|\bar{\mu} - \mu^*\|_{L^2}^2 + 2B^2 \sqrt{\frac{2 \log(2/\delta)}{|I_k|}}} \\ &\leq \sqrt{[\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})]_+} + \|\bar{\mu} - \mu^*\|_{L^2} + \sqrt{2}B \left(\frac{2 \log(2/\delta)}{|I_k|} \right)^{1/4}. \end{aligned}$$

Therefore, we conclude that with probability at least $1 - 5\delta$,

$$|\mathbf{W}_\alpha^* - \hat{\omega}_{\alpha,k}| \leq \frac{2}{\alpha} \left(\sqrt{[\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})]_+} + \|\bar{\mu} - \mu^*\|_{L^2} + CB \left(\frac{2 \log(2/\delta)}{|I_k|} \right)^{1/4} \right).$$

B.3.2 Continuing proof of Theorem 3 with a convex model class

First notice with probability at least $1 - \delta$,

$$\begin{aligned} \|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^1(\hat{P}_k|I_k^c)} &\leq \|\bar{\mu}(Z) - \hat{\mu}_k(Z)\|_{L^1(\hat{P}_k|I_k^c)} + \|\mu^*(Z) - \bar{\mu}(Z)\|_{L^1(\hat{P}_k)} \\ &\leq \|\bar{\mu}(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)} + \|\mu^*(Z) - \bar{\mu}(Z)\|_{L^1(\hat{P}_k)} \\ &\leq \|\bar{\mu}(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)} + \|\mu^*(Z) - \bar{\mu}(Z)\|_{L^1(P)} + \frac{B}{2} \sqrt{\frac{\log(1/\delta)}{|I_k|}}, \end{aligned}$$

where the first inequality follows by the triangle inequality, the second by Hölder’s inequality, and the last by Hoeffding inequality because $\mu^*, \hat{\mu}_k \in [0, B]$. The identity B.6 implies

$$\|\bar{\mu}(Z) - \hat{\mu}_k(Z)\|_{L^2(\hat{P}_k|I_k^c)}^2 = \Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu}) + 2\mathbb{E}_{\hat{P}_k}[(\ell(\theta(X_i); Y_i) - \bar{\mu}(Z_i))(\hat{\mu}_k(Z_i) - \bar{\mu}(Z_i))].$$

Since we assume the model class \mathcal{H} is convex and $\hat{\mu}_k \in \mathcal{H}$, the first-order condition of $\bar{\mu} \in \arg \min_{\mu \in \mathcal{H}} \mathbb{E}(\ell(\theta(X); Y) - \mu(Z))^2$ gives

$$\mathbb{E}[(\ell(\theta(X); Y) - \bar{\mu}(Z))(\hat{\mu}_k(Z) - \bar{\mu}(Z)) \mid Z, I_k^c] \leq 0,$$

so Hoeffding inequality implies with probability at least $1 - \delta$,

$$\mathbb{E}_{\hat{P}_k}[(\ell(\theta(X); Y) - \bar{\mu}(Z))(\hat{\mu}_k(Z) - \bar{\mu}(Z))] \leq B^2 \sqrt{\frac{2 \log(1/\delta)}{|I_k|}}. \quad (\text{B.7})$$

Hence, with probability at least $1 - 2\delta$,

$$\begin{aligned} \|\mu^*(Z) - \hat{\mu}_k(Z)\|_{L^1(\hat{P}_k|I_k^c)} &\leq \|\bar{\mu} - \mu^*\|_{L^1} + \sqrt{\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})} + 2B^2 \sqrt{\frac{2 \log(1/\delta)}{|I_k|}} + \frac{B}{2} \sqrt{\frac{\log(1/\delta)}{|I_k|}} \\ &\leq \sqrt{[\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})]_+} + \|\bar{\mu} - \mu^*\|_{L^1} + CB \left(\frac{2 \log(2/\delta)}{|I_k|} \right)^{1/4}. \end{aligned}$$

Therefore, we conclude that with probability at least $1 - 5\delta$,

$$|\mathbf{W}_\alpha^* - \hat{\omega}_{\alpha,k}| \leq \frac{2}{\alpha} \left(\sqrt{[\Delta_{I_k}(\hat{\mu}_k) - \Delta_{I_k}(\bar{\mu})]_+} + \|\bar{\mu} - \mu^*\|_{L^1} + CB \left(\frac{2 \log(2/\delta)}{|I_k|} \right)^{1/4} \right).$$

C Additional experiment details

In this section, we present additional experiments for the Functional Map of the World (FMoW) dataset. Due to the ever-changing nature of aerial images and the uneven availability of data from different regions, it is imperative that ML models maintain good performance under temporal (learn from the past and generalize to future) and spatial distribution shifts (learn from one region and generalize to another). Without having access to the out-of-distribution samples, our diagnostic raises awareness on brittleness of model performance against subpopulation shifts.

C.1 Dataset Description

The original Functional Map of the World (FMoW) dataset by [33] consists of over 1 million images from over 200 countries. We use a variant, FMoW-WILDS, proposed by Koh et al. [68], which temporally groups observations to simulate distribution shift across time. Each data point includes an RGB satellite image x , and a corresponding label y on the land / building use of the image (there are 62 different classes). FMoW-WILDS splits data into non-overlapping time periods: we train and validate models $\theta(\cdot)$ on data collected from years 2002-2013, and simulate distribution shift by looking at data collected during 2013-2018. Data collected during 2002-2013 (“in-distribution”) is split into training ($n=76,863$), validation ($n=19,915$), and test ($n=11,327$). Data collected during 2013-2018 (“out-of-distribution”) is split into two sets: one consisting of observations from years 2013-2016 ($n=19,915$), and another consisting of observations from years 2016-2018 ($n=22,108$).

All data splits contain images from a diverse array of geographic regions. We evaluate the worst-case subpopulation performance on in-distribution validation data, and study model performance under distribution shift on data after 2016.

C.2 Models Evaluated

We consider *DenseNet* models as reported by Koh et al. [68], including the vanilla empirical risk minimization (ERM) model and models trained with robustness interventions (IRM [6] method; Koh et al. [68] notes that ERM’s performance closely match or outperform “robust” counterparts even under distribution shift. We also evaluate *ImageNet* pre-trained *DPN-68* model from Miller et al. [79]. As separate experiments, we also consider *ResNet-18* and *VGG-11* from Miller et al. [79], and the results are reported in C.6.

CLIP (Contrastive Language-Image Pre-training) is a newly proposed model pre-trained on 400M image-text pairs, and has been shown to exhibit strong zero-shot performance on out-of-distribution samples [86]. Although not specifically designed for classification tasks, CLIP can be used for classification by predicting the class whose encoded text is the closest to the encoded image. We consider the weight-space ensembled *CLIP WiSE* models proposed in [118] as it is observed that these models exhibit robust behavior on FMoW. *CLIP WiSE* models are constructed by linearly combining the model weights of *CLIP ViT-B16 Zeroshot model* and *CLIP ViT-B16 FMoW end-to-end finetuned* model.

To illustrate the usage of our method, we choose the *CLIP WiSE* model that has similar ID validation accuracy as the *DenseNet* Models. This turns out to be putting 60% weight on *CLIP ViT-B16 Zeroshot model* and 40% weight on *CLIP ViT-B16 FMoW end-to-end finetuned*. *DenseNet* Models have average ID validation loss 2.4 – 2.8, but *CLIP WiSE* has average ID validation loss 1.6. To ensure fair comparison, we calibrate the temperature parameter such that the average loss of *CLIP WiSE* matches the worst average loss of the models considered. We deliberately make *CLIP WiSE* no better than any *DenseNet* Models, in the hope that our metric will recover its robustness property.

C.3 Flexibility of our metric

We implement Algorithm 1 by partitioning the ID validation data into two; we estimate $\mu^*(Z)$ using XGBoost on one sample, and estimate $W_\alpha(\cdot)$ at varying subpopulation size α on the other. By switching the role of each split, our final estimator averages two versions of $\widehat{W}_{\alpha,k}(\widehat{\mu})$.

C.3.1 A less conservative Z

In Section 5, we report results when Z is defined over all metadata consisting of (longitude, latitude, cloud cover, region, year), as well as the label Y . Defining subpopulations over such a wide range of variables may be overly conservative in some scenarios, and to illustrate the flexibility of our approach, we now showcase a more tailored definition of subpopulations. Since FMoW-WILDS is specifically designed for spatiotemporal shifts, a natural choice of Z is to condition on (region, year). Motivated by our observation that some classes are harder to predict than others (Figure 4(b)), we also consider $Z = (\text{region, year, label } Y)$. We plot our findings in Figure 6. If we simply define $Z = (\text{year, region})$, the corresponding worst-case subpopulation performance is less pessimistic. However, when we add labels to Z , we again see a drastic decrease in the worst-case subpopulation

#	Text Prompt
1	“CLASSNAME”
2	“a picture of a CLASSNAME.”
3	“a photo of a CLASSNAME.”
4	“an image of an CLASSNAME”
5	“an image of a CLASSNAME in asia.”
6	“an image of a CLASSNAME in africa.”
7	“an image of a CLASSNAME in the americas.”
8	“an image of a CLASSNAME in europe.”
9	“an image of a CLASSNAME in oceania.”
10	“satellite photo of a CLASSNAME”
11	“satellite photo of an CLASSNAME”
12	“satellite photo of a CLASSNAME in asia.”
13	“satellite photo of a CLASSNAME in africa.”
14	“satellite photo of a CLASSNAME in the americas.”
15	“satellite photo of a CLASSNAME in europe.”
16	“satellite photo of a CLASSNAME in oceania.”
17	“an image of a CLASSNAME”

Table 2: Text prompts for CLIP text encoders

performance, and that *CLIP WiSE-FT* outperforms all other models by a significant amount. This is consistent with our motivation in defining subpopulations over labels; our procedure automatically takes into account the interplay between class labels and spatiotemporal information.




Figure 6. In the left panel $Z = (\text{year}, \text{region})$; in the right panel $Z = (\text{year}, \text{region}, \text{label } Y)$. Here we take Z to contain only spatial and temporal information, a less conservative counterpart to the experiment reported in the main text. We again see that introduction of labels in Z drastically increase our metric, showing varying difficulty in learning different labels.

C.3.2 Using semantics of the labels

Alternatively, we may wish to define subpopulations over rich natural language descriptions on the input X . To illustrate the flexibility of our procedure in such scenarios, we consider subpopulations defined over the semantic meaning of the class names: CLIP-encoded class names using the 17

prompts reported in Table 2. For comparison, we report the (estimated) worst-case subpopulation performance (1.2) when we take $Z = (\text{all metadata, encoded labels})$ and $(\text{all metadata, label } Y, \text{ encoded labels})$ in Figure 7. We observe that in this case, the semantics of the class names do not contribute to further deterioration in robustness, and the relative ordering across models remains unchanged.




Figure 7. In the left panel, $Z = (\text{all meta, encoded labels})$; in the right panel, $Z = (\text{all meta, label } Y, \text{ encoded labels})$. We see that in this case no significant difference is introduced when semantics of the class names are included.

C.4 Analysis of spatiotemporal distribution shift

The significant performance drop in the Africa region on data collected from 2016-2018 was also observed in [68, 118]. In Figures 8-9, we plot the number of samples collected from Africa over data splits. In particular, we observe a large number of single-unit and multi-unit residential instances emerge in the OOD data. Data collection systems are often biased against the African continent—often as a result of remnants of colonialism—and addressing such bias is an important topic of future research.

C.5 Estimator of model loss

One potential limitation of our approach is \hat{h} does not always estimate the tail losses accurately, and this is important because our approach precisely is designed to counter ML models that perform poorly on tail subpopulations. Figure 10 plots a histogram of model losses and the estimated conditional risk \hat{h} for *DenseNet* ERM and *CLIP WiSE*, where the y-axis is plotted on a log-scale. It is clear that *DenseNet* ERM has more extreme losses compared to the *CLIP WiSE* model, suggesting that at least part of the reason why *DenseNet* ERM suffers poor loss on subpopulations: it is overly confident when it’s incorrect. While a direct comparison is not appropriate since the conditional risk $\mu(Z)$ represent *smoothed* losses, we observe that naive estimators of $\mu(\cdot)$ may consistently underestimate. In this particular instance, since the extent of underestimation is more severe for *ImageNet* pre-trained models, our experiments are fortuitously providing an even more conservative comparison between the two model classes, instilling confidence in the relative robustness of the *CLIP WiSE* model.

Alternatively, we can directly define the worst-case subpopulation performance (1.2) using the 0-1 loss. The discrete nature of the 0-1 loss pose some challenges in estimating $\mu(\cdot)$. While we chose




Figure 8: Instances by class, ID 2002-2013, Africa




Figure 9: Instances by class, test 2016-2018, Africa




Figure 10. Histograms of model losses and fitted losses \hat{h} . Y-axis count is plotted in **log-scale**. For *DenseNet* ERM model, fitted \hat{h} underestimates the extreme losses (right-tail).

to focus on the cross entropy loss that aligns with model training, we leave a thorough study of 0-1 loss to future work.

C.6 Additional comparisons

We use the ensembled *CLIP WiSE* model constructed by averaging the network weights of *CLIP zero-shot* and *CLIP finetuned* models. So far, we used proportion $\lambda = 0.4$ to match the ID validation accuracy of *CLIP WiSE* to that of *DenseNet* models and *DPN-68*. In this subsection, we provide alternative choices:

1. $\lambda = 0.24$ to match ID accuracy of *ResNet-18* of 47%
2. $\lambda = 0.27$ to match ID accuracy of *VGG-11* of 51%.

Similar to *DPN-68*, *ResNet-18* and *VGG-11* are *ImageNet* pretrained models fine-tuned on FMoW as evaluated by Miller et al. [79]. We refer to the two *CLIP WiSE* models as *CLIP WiSE-24* and *CLIP WiSE-27* respectively, and report all model performances below. Again, we observe that our approach successfully picks out the more robust *CLIP WiSE* models, in contrast to the non-robust models chosen by ID accuracy or ID loss.

D Proof of equivalence results

In this section, we discuss in detail how our notion of generalized worst-case subpopulation performance is closely connected to distributional robustness and coherent risk measures. The reader is recommended to refer to Section 6 of the lectures notes by Shapiro et al. [103] for more detail.

Model	ID, 2002-2013			OOD, 2016-2018	
	Accuracy	Loss	$W_{0.10}$	Accuracy	Loss
<i>CLIP WiSE-24</i>	0.47	2.84	3.47	0.45	2.85
<i>ResNet-18</i>	0.48	2.84	5.05	0.40	3.36
<i>CLIP WiSE-27</i>	0.51	3.07	3.54	0.48	3.08
<i>VGG-11</i>	0.51	3.06	6.07	0.45	3.68

Table 3. Additional experiments showcasing our approach successfully identifies more robust models.

D.1 Proof of Theorem 5

To see the first claim, fix a nonempty class Λ of probability measures on $(0, 1]$. Notice $W_\Lambda(\cdot)$ is proper because Λ is nonempty and $W_\Lambda(0) = 0 < \infty$. It is lower semi-continuous because it is a pointwise supremum of $\mu \mapsto \int_{(0,1]} W_\alpha(\mu) d\lambda(\alpha)$, which are lower semi-continuous. Coherence can be shown by verifying the definition. Clearly, W_Λ is law-invariant because it is defined using only $W_\alpha(\cdot)$ which is law-invariant.

For the converse relation, recall the discussion preceding Lemma 3. The biconjugacy relation (6.4) gives the variational representation

$$\rho(\mu) = \sup_{L \in \text{dom } \rho^*} \mathbb{E}_P[L\mu].$$

Since ρ is law-invariant, we have the tautological reformulation

$$\rho(\mu) = \sup_{\mu' \in \mathcal{L}^k} \left\{ \rho(\mu') : \mu' \stackrel{d}{=} \mu \right\} = \sup_{L \in \text{dom } \rho^*, \mu' \in \mathcal{L}^k} \left\{ \mathbb{E}_P[L\mu'] : \mu' \stackrel{d}{=} \mu \right\}. \quad (\text{D.1})$$

Next, we use a generalization of the Hardy–Littlewood inequality. Let $P_{1-\alpha}^{-1}(\mu)$ denote the $(1 - \alpha)$ -quantile of the random variable $\mu \in \mathcal{L}^k$.

Lemma 11 (Shapiro et al. [102, Lemma 6.25]). *Suppose P is nonatomic. For $\mu \in \mathcal{L}^k$ and $L \in \mathcal{L}^{k*}$,*

$$\sup_{\mu' \in \mathcal{L}^k} \left\{ \mathbb{E}_P[L\mu'] : \mu' \stackrel{d}{=} \mu \right\} = \int_0^1 P_{1-t}^{-1}(L) P_{1-t}^{-1}(\mu) dt.$$

In particular, $\mathbb{E}_P[\mu] = \int_0^1 P_{1-t}^{-1}(\mu) dt$.

We use the following elementary identity to rewrite $P_{1-t}^{-1}(L)$ in the preceding display. We defer its proof to the end of the subsection.

Lemma 12. *For any random variable $L \in \mathcal{L}^{k*}(P)$,*

$$P_{1-t}^{-1}(L) = \int_t^1 \alpha^{-1} d\lambda_L(\alpha) \quad \text{for } t \in (0, 1),$$

where $\lambda_L(\cdot)$ is a probability distribution on $(0, 1]$, if and only if $L \geq 0$ P -a.s., $\mathbb{E}_P[L] = 1$, $\lambda_L(\alpha) = \mathbb{E}_P(L - P_{1-\alpha}^{-1}(L))_+$ for $\alpha \in (0, 1)$, and $\lambda_L(1) = 1$.

For any fixed $L \in \mathcal{L}^{k*}$, conclude

$$\sup_{\mu' \in \mathcal{L}^k} \left\{ \mathbb{E}_P[L\mu'] : \mu' \stackrel{d}{=} \mu \right\} = \int_0^1 P_{1-t}^{-1}(L) P_{1-t}^{-1}(\mu) dt = \int_0^1 \int_{(t,1]} \alpha^{-1} d\lambda_L(\alpha) P_{1-t}^{-1}(\mu) dt.$$

Applying Fubini-Tonelli to the RHS of the preceding display,

$$\sup_{\mu' \in \mathcal{L}^k} \left\{ \mathbb{E}_P[L\mu'] : \mu' \stackrel{d}{=} \mu \right\} = \int_{(0,1]} \frac{1}{\alpha} \int_0^\alpha P_{1-t}^{-1}(\mu) dt d\lambda_L(\alpha) = \int_{(0,1]} W_\alpha(\mu) d\lambda_L(\alpha),$$

where the final equality follows from the change-of-variables reformulation (2.2). To obtain the representation (D.1), we take the supremum over $L \in \text{dom } \rho^*$

$$\rho(\mu) = \sup_{L \in \text{dom } \rho^*} \int_{(0,1]} W_\alpha(\mu) d\lambda_L(\alpha).$$

D.1.1 Proof of Lemma 12

Before proving the equivalence, we first show the following identity for any L and λ_L .

$$\int_0^\alpha [h(t) - h(\alpha)] dt = \mathbb{E}_P(L - P_{1-\alpha}^{-1}(L))_+ - \lambda_L(\alpha), \quad (\text{D.2})$$

where we define $h(t) := P_{1-t}^{-1}(L) - \int_{(t,1]} a^{-1} d\lambda_L(a)$ for convenience. We separately consider the two differences in the integrand

$$\int_0^\alpha [h(t) - h(\alpha)] dt = \int_0^\alpha P_{1-t}^{-1}(L) - P_{1-\alpha}^{-1}(L) - \left(\int_t^1 a^{-1} d\lambda_L(a) - \int_\alpha^1 a^{-1} d\lambda_L(a) \right) dt.$$

For $t \in (0, \alpha)$

$$\begin{aligned} \int_0^\alpha [P_{1-t}^{-1}(L) - P_{1-\alpha}^{-1}(L)] dt &= \int_0^1 [P_{1-t}^{-1}(L) - P_{1-\alpha}^{-1}(L)]_+ dt \\ &= \int_0^1 P_{1-t}^{-1} \left([L - P_{1-\alpha}^{-1}(L)]_+ \right) dt = \mathbb{E}_P [L - P_{1-\alpha}^{-1}(L)]_+, \end{aligned}$$

and

$$\int_0^\alpha \left[\int_{(t,1]} a^{-1} d\lambda_L(a) - \int_{(\alpha,1]} a^{-1} d\lambda_L(a) \right] dt = \int_{(0,\alpha]} \left(\int_0^a dt \right) a^{-1} d\lambda_L(a) = \lambda_L(\alpha).$$

Now we show the “if” part. It is clear that λ_L is nondecreasing, as $P_{1-\alpha}^{-1}(\cdot)$ is nonincreasing in α , and $\lambda_L(\alpha) \leq \mathbb{E}_P[L] = 1 = \lambda_L(1)$ because $L \geq 0$ P -a.s. We know λ_L is right-continuous because $P_{1-\alpha}^{-1}(\cdot)$ is left continuous in α . Then we conclude λ_L is a probability distribution on $(0, 1]$ by noticing $\lim_{\alpha \downarrow 0} \lambda_L(\alpha) = 0$ because

$$\begin{aligned} \lambda_L(\alpha) &= \left\| (L - P_{1-\alpha}^{-1}(L))_+ \right\|_1 \leq \left\| (L - P_{1-\alpha}^{-1}(L))_+ \right\|_{k_*} \\ &\leq \|L\|_{k_*} \mathbb{P}\{L > P_{1-\alpha}^{-1}(L)\}^{1/k_*} \\ &\leq \|L\|_{k_*} \alpha^{1/k_*}, \end{aligned}$$

where the first inequality follows by Hölder’s inequality, the second because $(L - P_{1-\alpha}^{-1}(L))_+ = (L - P_{1-\alpha}^{-1}(L)) \mathbf{1}\{L > P_{1-\alpha}^{-1}(L)\}$ and $L \geq 0$ P -a.s., and the last by definition of $P_{1-\alpha}^{-1}(L)$.

By the definition of λ_L , the RHS of Identity (D.2) is zero, so $\alpha h(\alpha) = \int_0^\alpha h(t) dt$. In particular, $h(\cdot)$ is differentiable in $(0, 1)$ since $\alpha \mapsto \int_0^\alpha h(t) dt$ is differentiable. Taking derivatives on both sides of the preceding display, we have $\alpha h'(\alpha) + h(\alpha) = h(\alpha)$ and in particular, $h'(\alpha) = 0$ for $\alpha \in (0, 1]$.

To show that $h(t) \equiv h$ is uniformly zero, notice

$$h = \int_0^1 h(t) dt = \int_0^1 \left[P_{1-t}^{-1}(L) - \int_t^1 a^{-1} d\lambda_L(a) \right] dt = \mathbb{E}_P L - 1 = 0,$$

where we used Fubini-Tonelli and $\lambda_L(1) = 1$ in the third equality.

Next, we show the “only if” part. Clearly the LHS of Identity (D.2) is zero, so $\lambda_L(\alpha) = \mathbb{E}_P (L - P_{1-\alpha}^{-1}(L))_+$. We know $L \geq 0$ P -a.s. because $\lambda_L \geq 0$. Lastly, $\mathbb{E}_P[L] = \int_0^1 P_{1-t}^{-1}(L) dt = \int_0^1 \int_t^1 \alpha^{-1} d\lambda_L(\alpha) dt = 1$ by Fubini-Tonelli.

D.2 Proof of Proposition 6

First, to see the duality result, notice the minimax theorem and Hölder’s inequality imply

$$\begin{aligned} \rho_k(\mu) &= \inf_{\eta \in \mathbb{R}} \{ \alpha^{-1} \|(\mu - \eta)_+\|_k + \eta \} \\ &= \inf_{\eta \in \mathbb{R}} \sup \{ \mathbb{E}_P[L(\mu - \eta)] + \eta : L \geq 0, \|L\|_{k_*} \leq \alpha^{-1} \} \\ &= \sup \left\{ \inf_{\eta \in \mathbb{R}} \{ \mathbb{E}_P[L(\mu - \eta)] + \eta \} : L \geq 0, \|L\|_{k_*} \leq \alpha^{-1} \right\} \\ &= \sup_L \{ \mathbb{E}_P[L\mu] : L \geq 0, \mathbb{E}_P[L] = 1, \|L\|_{k_*} \leq \alpha^{-1} \}. \end{aligned}$$

This means $\text{dom } \rho_k^* = \{L \in \mathcal{L}^k : L \geq 0, \mathbb{E}_P[L] = 1, \|L\|_{k_*} \leq \alpha^{-1}\}$. Theorem 5 implies

$$\rho_k(\mu) = \sup \left\{ \int_{(0,1]} W_\alpha(\mu) d\lambda_L(\alpha) : \lambda_L(\alpha) = \mathbb{E}_P (L - P_{1-\alpha}^{-1}(L))_+, L \geq 0, \mathbb{E}_P[L] = 1, \|L\|_{k_*} \leq \alpha^{-1} \right\}.$$

Lemma 12 further implies $\{L, \lambda_L : \lambda_L(\alpha) = \mathbb{E}_P (L - P_{1-\alpha}^{-1}(L))_+, L \geq 0, \mathbb{E}_P[L] = 1\} = \{L, \lambda_L \in \Delta((0, 1]) : P_{1-t}^{-1}(L) = \int_t^1 \alpha^{-1} d\lambda_L(\alpha)\}$. Notice $P_{1-t}^{-1}(L) = \int_t^1 \alpha^{-1} d\lambda_L(\alpha)$ implies

$$\|L\|_{k_*}^{k_*} = \int_\Omega L^{k_*} d\mathbb{P} = \int_0^1 (P_{1-t}^{-1}(L))^{k_*} dt = \int_0^1 \left(\int_t^1 \alpha^{-1} d\lambda_L(\alpha) \right)^{k_*} dt,$$

so

$$\rho_k(\mu) = \sup \left\{ \int_{(0,1]} W_\alpha(\mu) d\lambda_L(\alpha) : \lambda_L \in \Lambda_k, P_{1-t}^{-1}(L) = \int_t^1 \alpha^{-1} d\lambda_L(\alpha) \right\}.$$

Lastly, since L can always be defined based on any $\lambda_L \in \Delta((0, 1])$, we drop the last equality and conclude

$$\rho_k(\mu) = \sup \left\{ \int_{(0,1]} W_\alpha(\mu) d\lambda_L(\alpha) : \lambda_L \in \Lambda_k \right\} = W_{\Lambda_k}(\mu).$$

E Certificate of robustness

Instead of estimating the worst-case subpopulation performance for a fixed subpopulation size α , it may be natural to posit a level of acceptable performance (upper bound $\bar{\ell}$ on the loss) and study α^* , the smallest subpopulation size (2.1) over which the model $\theta(\cdot)$ can guarantee acceptable performance. Our plug-in estimator $\hat{\alpha}$ given in Eq. (3.4) enjoys similar concentration guarantees as

given above. The following theorem—whose proof we give in Appendix E.1—states that the true α^* is either close to our estimator $\hat{\alpha}$ or it is sufficiently small, certifying the robustness of the model against subpopulation shifts.

Theorem 9. *Let Assumption D hold, let $U(\delta) > 0$ be such that for any fixed $\alpha \in (0, 1]$, $|\widehat{W}_{\alpha,k}(\hat{\mu}) - W_\alpha(\mu^*)| \leq U(\delta)/\alpha$ with probability at least $1 - \delta$. Then given any $\underline{\alpha} \in (0, 1]$, either $\alpha^* < \underline{\alpha}$, or*

$$\left| \frac{\alpha^*}{\hat{\alpha}} - 1 \right| \leq \frac{U(\delta)}{\widehat{\mathbb{E}} \left[\widehat{\mu}(Z) - \widehat{P}_{1-\underline{\alpha}\wedge\hat{\alpha}}^{-1}(\widehat{\mu}(Z)) \right]_+}$$

with probability at least $1 - \delta$, where $\widehat{\mathbb{E}}$ and $\widehat{P}_{1-\alpha}^{-1}$ denote the expectation and the $(1 - \alpha)$ -quantile under the empirical probability measure induced by I_k .

Our approach simultaneously provides localized Rademacher complexity bounds and dimension-free guarantees. Our bound becomes large as $\underline{\alpha} \rightarrow 0$ and we conjecture this to be a fundamental difficulty as the worst-case subpopulation performance (1.2) focuses on α -fraction of the data.

E.1 Proof of Theorem 9

For ease of notation, we suppress any dependence on the prediction model $\theta(X)$ under evaluation. Consider any $\alpha_1, \alpha_2 \in (0, 1]$ with $\alpha_1 < \alpha_2$. Denote by $\widehat{\mathbb{P}}$ the empirical probability measure induced by $(Z_i : i \in I_k)$. For convenience denote $\xi_1 := \widehat{P}_{1-\alpha_1}^{-1}(\widehat{\mu}(Z))$ and $\xi_2 := \widehat{P}_{1-\alpha_2}^{-1}(\widehat{\mu}(Z))$, so $\xi_1 \geq \xi_2$ and $\widehat{W}_{\alpha_1}(\widehat{\mu}) \geq \widehat{W}_{\alpha_2}(\widehat{\mu})$. Notice that

$$\begin{aligned} \widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_2]_+ - \widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_1]_+ &= \widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_2; \widehat{\mu}(Z) > \xi_2] - \widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_1; \widehat{\mu}(Z) \geq \xi_1] \\ &= \underbrace{\widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_1; \xi_2 < \widehat{\mu}(Z) < \xi_1]}_{\leq 0} + (\xi_1 - \xi_2) \underbrace{\widehat{\mathbb{P}}[\widehat{\mu}(Z) > \xi_2]}_{\leq \alpha_2} \\ &\leq (\xi_1 - \xi_2)\alpha_2. \end{aligned}$$

Hence, by Lemma 9,

$$\widehat{W}_{\alpha_1}(\widehat{\mu}) - \widehat{W}_{\alpha_2}(\widehat{\mu}) = \left(\frac{\widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_1]_+}{\alpha_1} + \xi_1 \right) - \left(\frac{\widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_2]_+}{\alpha_2} + \xi_2 \right) \geq \frac{\widehat{\mathbb{E}}[\widehat{\mu}(Z) - \xi_1]_+}{\alpha_1\alpha_2}(\alpha_2 - \alpha_1),$$

meaning

$$|\alpha_1 - \alpha_2| \leq \frac{\alpha_1\alpha_2|\widehat{W}_{\alpha_1}(\widehat{\mu}) - \widehat{W}_{\alpha_2}(\widehat{\mu})|}{\widehat{\mathbb{E}}[\widehat{\mu}(Z) - \widehat{P}_{1-\alpha_1}^{-1}(\widehat{\mu}(Z))]_+}.$$

Now suppose $\alpha^* \geq \underline{\alpha}$. Notice W_α and $\widehat{W}_{\alpha,k}$ are continuous and nonincreasing in α , so the definitions (2.1) and (3.4) imply $W_{\alpha^*}(\mu^*) = \bar{\ell} = \widehat{W}_{\hat{\alpha}}(\widehat{\mu})$. Plugging $\hat{\alpha}$ and α^* into the inequality above, we know with probability at least $1 - \delta$,

$$|\alpha^* - \hat{\alpha}| \leq \frac{\hat{\alpha}\alpha^*|\widehat{W}_{\alpha^*}(\widehat{\mu}) - W_{\alpha^*}(\mu^*)|}{\widehat{\mathbb{E}} \left[\widehat{\mu}(Z) - \widehat{P}_{1-\alpha^*\wedge\hat{\alpha}}^{-1}(\widehat{\mu}(Z)) \right]_+} \leq \frac{\hat{\alpha}U(\delta)}{\widehat{\mathbb{E}} \left[\widehat{\mu}(Z) - \widehat{P}_{1-\underline{\alpha}\wedge\hat{\alpha}}^{-1}(\widehat{\mu}(Z)) \right]_+}.$$