

Solving the prisoner's dilemma trap in Hamilton's model of temporarily formed random groups

José F. Fontanari^{a,*}, Mauro Santos^b

^a*Instituto de Física de São Carlos, Universidade de São Paulo, 13560-970 São Carlos, São Paulo, Brazil*

^b*Departament de Genètica i de Microbiologia, Grup de Genòmica, Bioinformàtica i Biologia Evolutiva (GBBE), Universitat Autònoma de Barcelona, Spain*
cE3c - Centre for Ecology, Evolution and Environmental Changes & CHANGE - Global Change and Sustainability Institute, Lisboa, Portugal

Abstract

Explaining the evolution of cooperation in the strong altruism scenario, where a cooperator does not benefit from her contribution to the public goods, is a challenging problem that requires positive assortment among cooperators (i.e., cooperators must tend to associate with other cooperators) or punishment of defectors. The need for these drastic measures stems from the analysis of a group selection model of temporarily formed random groups introduced by Hamilton nearly fifty years ago to describe the fate of altruistic behavior in a population. Challenging conventional wisdom, we show analytically here that strong altruism evolves in Hamilton's original model in the case of biparental sexual reproduction. Moreover, when the cost of cooperation is small and the amplified contribution shared by group members is large, cooperation is the only stable strategy in equilibrium. Thus, our results provide a solution to the 'problem of origination' of strong altruism, i.e. how cooperation can take off from an initial low frequency of cooperators. We discuss a possible reassessment of cooperation in cases of viral co-infection, as cooperation may even be favored in situations where the prisoner's dilemma applies.

1. Introduction

As noted by Darwin in his discussion of courage and self-sacrifice, traits associated with altruistic behavior are counter-selected within a group, but increase the group's productivity and the group's chances of survival in the case of intergroup fighting [1]. Recognizing that the possible outcomes of this conflict between different levels of selection could only be determined by mathematical analysis, two slightly different models of group selection were proposed nearly

*Corresponding author

Email addresses: fontanari@ifsc.usp.br (José F. Fontanari), mauro.santos@uab.es (Mauro Santos)

fifty years ago to study the conditions for the evolution of altruistic behavior [2, 3]. Both models assume temporary random grouping of individuals, but differ in the way in which within-group competition between selfish and altruistic individuals is implemented. Hamilton's model follows the Darwinian view that altruists are always at disadvantage in local competition with selfish individuals, but groups with more altruists contribute more offspring to a pool of dispersers that will eventually form the new groups [2]. Wilson's trait group model ignores within-group competition altogether: although it assumes that individuals acquire their fitness locally, the competition between altruistic and selfish individuals takes place globally, i.e., in the population at large [3, 4]. Recently, we have pointed out the equivalence between Wilson's model and the evolutionary game approach to studying the emergence of cooperation in n -person public goods games, which is based on the replicator equation framework [5]. We are aware of the distinction between cooperation, which involves mutual benefit, and altruism, which implies 'others-only' benefit [6, 7]. However, we will use both terms, cooperation and altruism, more or less interchangeably here.

The outcome of these group selection models depends on whether an altruist benefits from her altruistic behavior or not. When the return from altruistic behavior exceeds the cost of performing it, we have the scenario of weak altruism [4, 8]. This is still altruism in the sense that selfish individuals benefit the most from this behavior. The traditional n -person prisoner's dilemma [9, 10, 11] and the n -person snowdrift game [5, 12, 13, 14] fall into this category. Both Hamilton's and Wilson's models of group selection with randomly formed groups predict that altruism will evolve in this scenario, a clearly impossible outcome for homogeneous or unstructured populations, which explains the notoriety of these models. When an altruist has no return from her altruistic behavior, as in the others-only variant of the n -person prisoner's dilemma [2, 15, 16], we have the more challenging scenario of strong altruism. Hamilton's seminal paper addressed this scenario and concluded that altruism cannot progress in randomly formed groups: cooperation is impossible unless there is some form of positive assortment among cooperators [2]. Wilson came to the same conclusion when groups are formed by binomially sampling individuals from the total population, as in Hamilton's model, but mistakenly suggested that environmental heterogeneity could produce a sufficiently large excess of variance over the binomial expectation that strong altruism might be favored [3]. This error was quickly corrected [17], so that both models now point to the impossibility of strong altruism evolving in the absence of explicit positive assortment among altruists, i.e., altruists must have a tendency to group with other altruists. Nonetheless, we show here, using both analytical and simulation results, that strong altruism can evolve in Hamilton's model without assuming positive assortment. This is achieved by not taking the phenotypic gambit in full [18].

The phenotypic gambit, often used in evolutionary game theory, assumes, among other things, the simplest genetic system (e.g., haploid individuals). Although, this is usually wrong, we can generally ignore genetics to study the evolutionary fate of behavioral strategies, except in the case of diploidy, when the heterozygote is the fittest genotype [18] (see also [19]). Let us accept the

gambit and assume that organisms are haploid and that each chromosome consists of L loci, one of which controls one of the only two pure strategies, altruistic (cooperator) and selfish (defector). Now consider three possible modes of reproduction that are commonly used in theoretical models: asexual reproduction, sexual reproduction (production of haploid gametes that combine in pairs to produce a recombinant offspring chromosome) that allows self-mating, and sexual reproduction that requires two different individuals, what Williams calls euphrasic [20]. Thus, mating pairs are formed by randomly drawing individuals (with or without replacement) according to their relative fitness, and each pair produces a single offspring. This is still a simplified picture of a more complex reality, where organisms can be diploid and the behavioral strategy is likely to be controlled by many loci with possible epistatic interactions. The question here is whether the phenotypic gambit, which in this case would also ignore the complications introduced by the mode of reproduction, is still valid. If so, we would conclude that strong altruism cannot evolve in Hamilton’s model regardless of how individuals reproduce.

Here, we show that the way we model reproduction can have a qualitative impact on the evolution of altruism in ways that the phenotypic approach cannot predict or explain. In particular, we show that strong altruism can evolve under biparental sexual reproduction in Hamilton’s model [2].

The rest of the paper is organized as follows. In Section 2 we describe Hamilton’s model of temporarily formed random groups, assuming G groups of size n , using the terminology of the n -person prisoner’s dilemma. In Section 3 we develop an analytical treatment to solve Hamilton’s model in the deterministic limit $G \rightarrow \infty$ and show that in the case of asexual reproduction, cooperation is doomed to extinction in the strong altruism scenario, but not in the weak altruism scenario. In addition, we show that sexual reproduction, allowing self-mating, leads to the same conclusion as in the asexual case. We then consider biparental sexual reproduction and, challenging conventional wisdom, derive the conditions that guarantee the emergence of cooperation in the strong altruism scenario. In all cases, we follow the fate of the altruistic or selfish strategy, which for sexual reproduction would be analogous to following the allelic changes in the recombinant offspring chromosomes hosting the behavioral gene. In Section 4 we use simulations to prove the robustness of the conclusions of the deterministic analysis to the effect of demographic noise that arises when G is finite. Finally, in Section 5 we summarize our results and also suggest a possible reassessment of cooperation in cases of viral co-infection, as cooperation may even be favored in situations where the prisoner’s dilemma applies. Returning to the phenotypic gambit, we cannot avoid ending this section by recalling a quote from Albert Einstein (possibly misattributed; see [21]): “Make things as simple as possible, but not simpler.”

2. Model

Here we recast Hamilton’s model of temporarily formed random groups [2] using the terminology of public goods games [9]. Consider a population of Gn

individuals, divided into G groups, each of which contains n individuals. Within each group, individuals play a n -person prisoner's dilemma game to determine their fitness: each individual may (or may not) contribute an amount $c > 0$ to the public goods, which is then multiplied by a factor $r > 1$, and the resulting amount rc is divided among the $n - 1$ other players. This is the others-only variant of the n -person prisoner's dilemma, where contributors do not benefit from their own contributions [15]. We refer to individuals that contribute to the public goods as cooperators and those that do not as defectors.

The fitness (or payoff) of a cooperator in a group with $i - 1$ other cooperators is

$$f_c(i) = 1 - c + (i - 1) \frac{rc}{n - 1}, \quad (1)$$

whereas the fitness of a defector in a group with i cooperators is

$$f_d(i) = 1 + i \frac{rc}{n - 1}. \quad (2)$$

These are the payoffs of the popular n -person prisoner's dilemma [9, 15, 16] with a constant baseline fitness value of 1. They are identical to the fitness attributed to altruistic and selfish individuals in Hamilton's model of randomly formed groups: in Hamilton's terminology, an altruist gives up $k = c$ units of her own fitness in order to add $K = rc$ units to the joint fitness of her $n - 1$ companions [2]. We will assume $c \in [0, 1]$ to ensure that the fitness values are not negative, so $c = 1$ is the worst scenario for cooperation. The parallel between the n -person prisoner's dilemma and Hamilton's model ends here, however. In fact, in Hamilton's model the fitness of a group with i cooperators,

$$\pi(i) = n + ic(r - 1), \quad (3)$$

determines the number of offspring m produced by the group. In particular, we assume that m is a random variable distributed by a Poisson distribution with mean $\pi(i)$. Group fitness plays no role in the n -person prisoner's dilemma, where individuals decide whether or not to keep their strategies by comparing their payoffs to those of randomly selected individuals in the population at large [22, 23], as in Wilson's trait group formulation [3, 4], so there is effectively no within-group competition.

In Hamilton's model, within-group competition takes place during the reproduction stage. As mentioned before, we will consider three modes of reproduction. In the asexual reproduction mode, which is the one considered in Hamilton's seminal paper, we select one parent within each group with probability equal to her relative fitness, and the single offspring inherits the strategy of the parent. For each group, exactly m parents are selected in this way, and the resulting m offspring are added to the offspring pool. Hamilton refers to the offspring pool as the migrant pool, which is formed by young individuals that reach maturity [2]. In sexual reproduction with selfing allowed, we randomly select two not necessarily different parents, and the single offspring inherits the strategy of either parent with equal probability. This selection process is repeated m times resulting in the addition of m offspring to the offspring pool.

This mode is clearly equivalent to asexual reproduction, as we will show in the following section. Finally, in the biparental sexual reproduction mode, the two randomly chosen parents must be different and, as before, the single offspring inherits the strategy of either parent with equal probability. In this mode, it is guaranteed that in a group with $i = n - 1$ cooperators, one cooperator will be chosen as a parent. Note that the groups contribute different numbers of offspring to the offspring pool, not only because m is a Poisson distributed random variable, but also because the parameter of the Poisson, i.e. $\pi(i)$, depends on the number of cooperators within the group, which varies from group to group. Finally, each group of the next generation is formed by sampling n individuals with replacement from the offspring pool, and the whole process is repeated until one of the strategies disappears and evolution stops.

In the others-only variant of the n -person prisoner's dilemma, altruistic behavior (i.e., contribution to public goods) reduces the fitness of the cooperator, corresponding to the strong altruism scenario. As noted above, the conventional wisdom in this case is that cooperation cannot evolve in randomly formed groups that last less than one generation [24]. The more traditional variant of the n -person prisoner's dilemma allows the amplified amount rc to be redistributed to all players, so that the cooperator gets the return cr/n [10]. If $cr/n - c > 0$ or $r > n$, then altruistic behavior confers a net benefit to the cooperator, corresponding to the weak altruism scenario. A similar situation occurs in the n -person snowdrift game when the payoff for completing a task is greater than the effort expended [5]. In this case, the cooperator may have a net fitness advantage over the population as a whole, despite being at a disadvantage relative to defectors within the same group. This is the only situation where Wilson's trait group formulation [3, 4] or its game theoretic counterparts [10] can explain the maintenance of cooperation [17].

3. Deterministic limit

In his seminal paper [2], Hamilton studied the asexual reproduction mode in the case there are infinitely many groups using Price's equation [25]. Here we offer an elementary derivation of a recurrence for the frequency of cooperators q_t in the offspring pool in generation t , which can be easily generalized to more complex scenarios and reproduction modes [26].

Given q_t , the proportion of groups with i cooperators among the infinitely many groups is given by the binomial distribution

$$Y_i(t) = \binom{n}{i} q_t^i (1 - q_t)^{n-i}, \quad (4)$$

and our problem is to determine how many cooperators, on average, a group with i cooperators contributes to the offspring pool. We recall that, according to Section 2, a group with i cooperators contributes m offspring to the offspring pool with m distributed by the Poisson distribution

$$p[m; \pi(i)] = \exp[-\pi(i)] \frac{[\pi(i)]^m}{m!}. \quad (5)$$

Let $Q_c(i)$ denote the probability that an offspring is a cooperator. Calculating this probability requires to be explicit about the mode of reproduction, and we will do this calculation separately for each mode in the following, but for now let us assume that $Q_c(i)$ is known. The probability that there are j cooperators among the m offspring produced by a group with i cooperators is given by the binomial distribution

$$B[m, Q_c(i)] = \binom{m}{j} [Q_c(i)]^j [1 - Q_c(i)]^{m-j}. \quad (6)$$

Therefore, a group with i cooperators contributes, on average, with

$$\sum_{m=0}^{\infty} p[m; \pi(i)] \sum_{j=0}^m B[m, Q_c(i)] j = \sum_{m=0}^{\infty} p[m; \pi(i)] m Q_c(i) = \pi(i) Q_c(i) \quad (7)$$

cooperators to the offspring pool. Note that only the mean of the distribution of the number of offspring m goes into this calculation, so any distribution with mean $\pi(i)$ gives the same result.

The proportion of cooperators in the offspring pool in generation $t + 1$ is given by the ratio between the mean number of cooperator offspring produced by the groups formed by sampling the offspring pool in generation t and the mean total number of offspring produced by these groups, i.e.

$$q_{n+1} = \frac{\sum_{i=0}^n Y_i(t) \pi(i) Q_c(i)}{\sum_{i=0}^n Y_i(t) \pi(i)}. \quad (8)$$

The sum in the denominator can be easily performed and yields $n[1 + c(r - 1)q_t]$. The last step to obtain the recurrence for q_t is to calculate $Q_c(i)$ for the different reproduction modes.

3.1. Asexual reproduction

In this case, we randomly select a parent within a group with i cooperators according to her relative fitness, and the single offspring inherits her strategy. Thus, the probability of an offspring being a cooperator is simply

$$Q_c(i) = \frac{i f_c(i)}{\pi(i)}. \quad (9)$$

Inserting this probability into Eq. (8) yields

$$q_{t+1} = q_t \frac{1 - c + r c q_t}{1 + c q_t (r - 1)}, \quad (10)$$

which is consistent with the recurrence derived using Price's equation, provided that the mean fitness in [2] is corrected to $w = 1 + q_t(K - k) = 1 + q_t c(r - 1)$. The only fixed points of recurrence (11) are $q^* = 0$ and $q^* = 1$. For $q_t \ll 1$ it is rewritten as $q_{t+1} \approx (1 - c)q_t$, which implies that $q_{t+1} < q_t$ and so $q^* = 0$

is always stable and $q^* = 1$ is always unstable. This conclusion agrees with Hamilton's that cooperation cannot progress in such a model [2].

It is instructive to briefly discuss the results for the weak altruism version of the n -prisoner's dilemma. In this case, the fitness of a cooperator is $\hat{f}_c(i) = 1 - c + irc/n$ and the fitness of a defector is $\hat{f}_d(i) = 1 + irc/n$, where i is the number of cooperators in the group of size n . The fitness of this group is still given by Eq. (3). Thus, the recurrence for the frequency of cooperators \hat{q}_t in the weak altruism scenario is

$$\hat{q}_{t+1} = \hat{q}_t \frac{1 - c + rc/n + (n-1)rc\hat{q}_t/n}{1 + c\hat{q}_t(r-1)}, \quad (11)$$

whose only fixed points are $\hat{q}^* = 0$ and $\hat{q}^* = 1$. For $\hat{q}_t \ll 1$ we have $\hat{q}_{t+1} \approx (1 - c + rc/n)\hat{q}_t$ so $\hat{q}^* = 0$ is unstable (and thus $\hat{q}^* = 1$ is stable) for $r > n$. This means that cooperators at low frequency can invade a resident population of defectors if the amplification factor r is sufficiently large.

3.2. Sexual reproduction with self-mating

In this case, we randomly select two parents according to their relative fitness, and the single offspring inherits the strategy of either parent with equal probability. Since there are no restrictions on the choice of parents, the probability that an offspring is a cooperator is

$$\begin{aligned} Q_c(i) &= \left(\frac{if_c(i)}{\pi(i)} \right)^2 + \frac{1}{2} \frac{if_c(i)}{\pi(i)} \frac{(n-i)f_d(i)}{\pi(i)} + \frac{1}{2} \frac{(n-i)f_d(i)}{\pi(i)} \frac{if_c(i)}{\pi(i)} \\ &= \frac{if_c(i)}{\pi(i)}, \end{aligned} \quad (12)$$

which is identical to the result for asexual reproduction, as expected. Thus, cooperation cannot progress also in the model where sexual reproduction with selfing is allowed.

3.3. Biparental sexual reproduction

The difference to the previous case is that the mating parents within the group of size n must be different individuals. Their single offspring inherits the strategy of either parent with equal probability. So the problem we have to solve now is to find the probability that the offspring will be a cooperator when the two parents are chosen from a group with i cooperators and $n - i$ defectors. There are three possibilities we need to consider, which are described below.

- (a) Both parents are cooperators The probability that the first selected parent is a cooperator is simply

$$\frac{if_c(i)}{if_c(i) + (n-i)f_d(i)}. \quad (13)$$

Given that the first selected parent is a cooperator, the probability that the second is also a cooperator is

$$\frac{(i-1)f_c(i)}{(i-1)f_c(i) + (n-i)f_d(i)}, \quad (14)$$

since the fitness $f_c(i)$ and $f_d(i)$ given by Eqs. (1) and (2) are computed before the selection procedure. Thus, the probability that the two selected parents are cooperators is

$$\frac{(i-1)f_c(i)}{(i-1)f_c(i) + (n-i)f_d(i)} \times \frac{if_c(i)}{if_c(i) + (n-i)f_d(i)}. \quad (15)$$

Of course, the offspring resulting from this mating is a cooperator with probability one. The probabilities of the other events are calculated using the same reasoning.

- (b) The first chosen parent is a cooperator and the second is a defector. The probability of this event is

$$\frac{(n-i)f_d(i)}{(i-1)f_c(i) + (n-i)f_d(i)} \times \frac{if_c(i)}{if_c(i) + (n-i)f_d(i)} \quad (16)$$

and the resulting offspring is a cooperator with probability 1/2.

- (c) The first chosen parent is a defector and the second is a cooperator. The probability of this event is

$$\frac{if_c(i)}{if_c(i) + (n-i-1)f_d(i)} \times \frac{(n-i)f_d(i)}{if_c(i) + (n-i)f_d(i)} \quad (17)$$

and the resulting offspring is a cooperator with probability 1/2.

Combining these probabilities, we obtain that the probability the offspring is a cooperator when two parents are chosen from a group with i cooperators and $n-i$ defectors is

$$\begin{aligned} Q_c(i) &= \frac{i(i-1)[f_c(i)]^2}{[(i-1)f_c(i) + (n-i)f_d(i)][if_c(i) + (n-i)f_d(i)]} \\ &+ \frac{1}{2} \frac{i(n-i)f_c(i)f_d(i)}{[(i-1)f_c(i) + (n-i)f_d(i)][if_c(i) + (n-i)f_d(i)]} \\ &+ \frac{1}{2} \frac{i(n-i)f_c(i)f_d(i)}{[if_c(i) + (n-i-1)f_d(i)][if_c(i) + (n-i)f_d(i)]}. \end{aligned} \quad (18)$$

Clearly, $Q_c(0) = 0$ and $Q_c(n) = 1$. Inserting $Q_c(i)$ into the recurrence (8) yields

$$q_{t+1} = F_n(q_t) \quad (19)$$

with

$$F_n(q_t) = \frac{1}{n[1 + c(r-1)q_t]} \sum_{i=0}^n \binom{n}{i} q_t^i (1-q_t)^{n-i} \pi(i) Q_c(i), \quad (20)$$

where the dependence on q_t is made explicit. Since $Q_c(0) = 0$ we have $F_n(0) = 0$ and so $q^* = 0$ is a fixed point of the recurrence (19). In addition, noting that $Q_c(n) = 1$ and $\pi(n) = n[1 + c(r - 1)]$ we obtain $F_n(1) = 1$, which implies that $q^* = 1$ is also a fixed point of the recurrence (19). Whenever these two extreme fixed points are stable, there must also be an unstable coexistence fixed point. If both extreme fixed points are unstable, then the coexistence fixed point is stable, but this situation never happens, as we will see below. We recall that the condition for the local stability of a fixed point is $|F'_n(q^*)| < 1$, where the prime indicates the derivative of $F_n(q)$, as usual [27].

Before considering the general case of group size n , it is instructive to write the recurrence (19) for $n = 2$ explicitly. We have

$$q_{t+1} = \frac{q_t}{1 + c(r-1)q_t} \left[1 + \frac{c(r-1)}{2}(1 + q_t) \right], \quad (21)$$

so that the only fixed points are $q^* = 0$ and $q^* = 1$. Noting that

$$F'_2(0) = 1 + \frac{c(r-1)}{2} > 1 \quad (22)$$

and

$$F'_2(1) = 1 - \frac{1}{2} \frac{c(r-1)}{[1 + c(r-1)]} < 1, \quad (23)$$

we conclude that the all-defectors fixed point is unstable and the all-cooperators fixed point is stable if $r > 1$. This is expected, since the advantage of the selfish strategy comes only from within-group competition, which for $n = 2$ only occurs in groups with one cooperator and one defector. However, the requirement that mating involves both individuals gives $Q_c(1) = 1/2$, so this advantage is lost for $n = 2$. Now, if we consider that a group formed by two cooperators contributes on average $2 + 2c(r - 1)$ offspring to the offspring pool, while a group formed by two defectors contributes on average only 2 offspring, the dominance of the cooperators is not surprising (see also [28]). However, for groups of size $n > 2$ the situation is not so clear-cut, as we will discuss in the following.

Let us first consider the all-defectors fixed point, $q^* = 0$, for general n . A trite calculation yields $F'_n(0) = \pi(1)Q_c(1) > 0$, which is rewritten as

$$F'_n(0) = \frac{1}{2}(1 - c) \left[1 + \frac{n - 1 + rc}{n - 1 + rc - c - rc/(n - 1)} \right], \quad (24)$$

where we have used Eqs. (3) and (18) with $i = 1$. The condition $F'_n(0) < 1$ is satisfied when

$$c > \frac{1}{2n - 3} \quad (25)$$

regardless of the value of the amplification factor r . However, if this inequality is violated, then the condition $F'_n(0) < 1$ is satisfied, provided that $r < r_s$ where

$$r_s = (n - 1) \frac{2n - 3 - c}{1 - (2n - 3)c}. \quad (26)$$

For small c , this equation reduces to $r_s = (n-1)(2n-3) + 4(n-2)(n-1)^2c$. The important result from the ‘problem of origination’ perspective [29] is that the all-defectors fixed point is unstable if inequalities (25) and $r < r_s$ are violated simultaneously. For example, for $n = 4$ the all-defectors fixed point is stable for $c > 0.2$, regardless the value of r , but for $c = 0.1$ it is unstable if $r > 29.4$. It is clear from these inequalities that increasing the group size n favors the selfish strategy.

The analysis of the all-cooperators fixed point, $q^* = 1$, is more laborious. In this case we have

$$F'_n(1) = n - \frac{c(r-1)}{1+c(r-1)} - \frac{\pi(n-1)Q_c(n-1)}{1+c(r-1)}, \quad (27)$$

which can be easily evaluated numerically using Eqs. (3) and (18) to determine the regions in the parameter space where the all-cooperators fixed point is stable, i.e., where the condition $|F'_n(1)| < 1$ is satisfied. However, in the limit $c \rightarrow 0$ we can write down an explicit expression for $F'_n(1)$, viz.,

$$F'_n(1) = 1 + c \frac{2n-3}{2(n-1)} - rc \frac{1}{2(n-1)^2} \quad (28)$$

from which it follows that the all-cooperators fixed point is stable for $r > r_a$ where $r_a = (n-1)(2n-3)$ in this limit. Note that $r_a \rightarrow r_s$ for $c \rightarrow 0$. Although for $c = 0$ the recurrence (19) reduces to $q_{t+1} = q_t$ so that the dynamics freezes at the initial conditions, a vanishingly small value of c is sufficient to lead the dynamics to $q^* = 1$ if $r > r_a$ or to $q^* = 0$ if $r < r_s$.

The results of the stability analysis are conveniently summarized in a phase diagram in the plane (c, r) as shown in Fig. 1 for $n = 4$. Only the fixed point $q^* = 1$ is stable in region C and only the fixed point $q^* = 0$ is stable in region D. In region C+D we have the bistability situation, where the outcome of the dynamics depends on the initial frequency of cooperators q_0 . In particular, the unstable coexistence fixed point $0 < q^* < 1$ determines the domains of attraction of the two stable fixed points: the dynamics leads to the all-cooperators regime when $q_0 > q^*$ and to the all-defectors regime when $q_0 < q^*$. An important feature of this phase diagram, which also holds for general group sizes $n > 2$, is that there is no stable coexistence solution. In fact, a stable coexistence fixed point would occur if $r_s < r_a$ in the limit $c \rightarrow 0$, so there would be a region where neither of the extreme fixed points is stable, but we have analytically ruled out this possibility by showing that $r_a \rightarrow r_s$ in this limit.

Another important limit where we can explicitly rewrite Eq. (27) is for $c = 1$, which corresponds to the worst-case scenario for cooperation and will be the main focus of our analysis henceforth. In this case, we find

$$F'_n(1) = n - 1 + \frac{1}{2r} \frac{n-1-r(n-2)-r^2(2n^3-10n^2+18n-11)}{n-1+r(n^2-3n+3)}, \quad (29)$$

so that the condition $|F'_n(1)| < 1$ is satisfied for $r > r_a$, where r_a is the sole positive solution of the quadratic equation

$$r_a^2 - r_a(2n^2 - 7n + 6) - (n-1) = 0. \quad (30)$$




Figure 1: Phase diagram for $n = 4$ showing the stability regions of the different fixed points. In region C, only the all-cooperators fixed point $q^* = 1$ is stable, while in region D, only the all-defectors fixed point $q^* = 0$ is stable. In region C+D both fixed points are stable. The regions C and C+D are separated by the curve $r_s = r_s(c)$ and the regions D and C+D are separated by the curve $r_a = r_a(c)$.

For large n we have $r_a \approx 2n^2$, and for the group size $n = 4$ used in drawing the phase diagram in Fig. 1 we have $r_a \approx 10.29$.

Figure 2 shows the unstable coexistence fixed point q^* for $c = 1$ and several group sizes n . In case of bistability, this fixed point determines the domains of attraction of the two stable fixed points. The coexistence and all-cooperators fixed points coincide (i.e., $q^* = 1$) at $r = r_a$, which is given for $c = 1$ by the solution of Eq. (30). We use a log-log scale in this figure to emphasize the power-law decay of q^* with increasing r . More explicitly, for $c = 1$ and large r we have an analytical expression for the unstable coexistence fixed point, viz. $q^* \approx (4n - 6)/r$, which is in perfect agreement with the numerical results shown in Fig. 2. This result is important because it shows that at maximum cost to cooperators (i.e., $c = 1$), they can take over the population even at low initial frequencies, provided that the amplification factor r is sufficiently large.

4. Simulations for finite number of groups

Here we consider the robustness of our deterministic results to demographic noise, which arises when the population is finite, i.e., when G is finite. We focus only in the case of biparental sexual reproduction, which can maintain a stable population of cooperators in the limit $G \rightarrow \infty$, as shown in the previous section.

To validate our analytical formulation, we compare in Fig. 3 the results obtained by iterating the recurrence (19) and the simulation results for $G = 10^4$ and $c = 0.05$. We choose this particular small value of the contribution to




Figure 2: Unstable coexistence fixed point q^* as a function of the amplification factor r for $c = 1$ and (from left to right) $n = 3, 4, 5, 6$. If $q_0 > q^*$ the dynamics is driven to the all-cooperators fixed point and to the all-defectors fixed point if $q_0 < q^*$. The value of r at which $q^* = 1$ signals the instability of the all-cooperators fixed point and is given by the solution of Eq. (30). The all-defectors fixed point is stable for these parameter settings.

the public goods because by varying r we can go through the three different equilibrium regimes shown in the phase diagram of Fig. 1. In addition to the good agreement between the analytical results and the individual runs of the simulations (the figure shows four runs for each parameter setting), we can see that the fluctuations due to demographic noise are higher in the bistability regime (middle panel of Fig. 3), where the dynamics take a much longer time to reach equilibrium than in the other two regimes. Moreover, the convergence to the all-defectors fixed point (left panel of Fig. 3) is typically much faster than to the all-cooperators fixed point (right panel of Fig. 3). For not too large G , in the bistability regime we can have fixation to different fixed points even starting from the same initial condition q_0 due to demographic noise. This is the problem we address next, considering a worst-case scenario for cooperation by setting $c = 1$.

To quantify the effect of demographic noise, we consider the probability of fixation of the cooperators Π , which is estimated as the fraction of 10^4 independent runs for which this strategy fixates, i.e., for which the stochastic dynamics is attracted to the all-cooperators absorbing state. In the initial population, each individual is randomly assigned to one of the strategies with equal probability, corresponding to the choice $q_0 = 0.5$ in the deterministic analysis. In the bistability regime, the relevant deterministic result is given in Fig. 2, which can be interpreted in two different ways: for fixed r there is a minimum initial frequency of cooperators $q_0 = q^*$ above which the recurrence (19) converges to the all-cooperators fixed point, and for fixed q_0 there is a minimum value of




Figure 3: Frequency of cooperators as a function of time for $r = 5$ and $q_0 = 0.5$ (left panel), $r = 15$, $q_0 = 0.5$ and $q_0 = 0.7$ (middle panel), $r = 25$ and $q_0 = 0.5$ (right panel). The jagged thin black curves are runs of the simulations for $G = 10^4$ and the smooth thick colored curves are the deterministic results. The other parameters are $n = 4$ and $c = 0.05$.

$r = r_c$ above which the recurrence (19) converges to the all-cooperators fixed point. Here we choose the later interpretation and, in particular, for $q_0 = 0.5$ and $n = 4$ we have $r_c = 20.37$. Accordingly, Fig. 4 shows the probability of fixation of the cooperators as function of the amplification factor r . As noted above, for the same parameter setting, the demographic noise can tilt the dynamics into one absorbing state or the other. The strength of this effect depends on the number of groups and on the proximity of r to r_c . In fact, the scaling assumption $\Pi \approx (r - r_c)G^{1/2}$ perfectly describes this dependence for large G , implying that the width of the transition region around r_c shrinks as $G^{-1/2}$ as G increases (see [30, 31] for other applications of curve collapsing to characterize threshold phenomena).

As already hinted at in Fig. 3, another key quantity to characterize the stochastic dynamics is the mean fixation time T_f , i.e., the mean time for the stochastic dynamics to reach the all-cooperators or the all-defectors absorbing states. Figure 5 shows that in the all-cooperators regime ($r > r_c$) the fixation time diverges with $a_g \ln G$, where the prefactor $a_g = a_g(n)$ is an increasing function of the group size n . However, in the all-defectors regime ($r < r_c$) the fixation time tends to a finite limit as G increases. Note that for finite G , the maximum of T_f does not occur at $r = r_c$, but it moves in the direction of the threshold r_c as G increases.




Figure 4: Probability of fixation Π of the cooperators as a function of the amplification factor r (left panel) for (from bottom to top at $r = 25$) $G = 400, 800, 1600, 3200$. The vertical dashed line indicates the threshold $r_c = 20.37$ beyond which the all-cooperators fixed point attracts the orbits starting at $q_0 = 0.5$ for $G \rightarrow \infty$. The right panel shows Π as a function of the scaled variable $(r - r_c)G^{1/2}$ for $G = 800, 1600, 3200$. The other parameters are $n = 4$ and $c = 1$.

Figure 6 shows the effect of the group size n on the threshold transition for finite G . Although Fig. 3 already showed that the threshold r_c increases with n , we can now see that the width of the transition region also increases with n , making the fixation of cooperators less likely for $r > r_c$ and finite G . Indeed, the right panel of Fig. 5 shows that the mean fixation time also increases with n , confirming that increasing group size makes within-group competition more favorable for defectors, reducing the chances of the cooperators to contribute to the offspring pool. Finally, we note that although in most of the results in this section we have fixed the initial frequency of cooperators at $q_0 = 0.5$, we get qualitatively similar results if we decrease q_0 , provided that we increase the range of r according to Fig. 2.

5. Discussion

There is a consensus that strong altruism, where a cooperator does not benefit from her costly altruistic behavior, cannot evolve in randomly formed, transient groups unless there is some sort of positive assortment among cooperators [2, 3, 4, 24, 32, 33] or punishment for defectors [10, 15, 16]. Note that reputation and reciprocity foster cooperation through the creation of positive assortment among players [34]. Even with these fixes, one difficulty remains: the so-called ‘origination problem’, i.e. how cooperation can take off from an initial low frequency of cooperators. However, it has been argued that this problem is an artifact of assuming, as ours and most models do, that strategies exist as




Figure 5: Mean time for fixation of cooperators or defectors T_f as a function of amplification factor r (left panel) for $n = 4$ and (from bottom to top at $r = 25$) $G = 400, 800, 1600, 3200$. The vertical dashed line indicates the threshold $r_c = 20.37$ beyond which the all-cooperators fixed point attracts the orbits starting at $q_0 = 0.5$ for $G \rightarrow \infty$. The right panel shows T_f at $r = r_c$ as function of G for (from bottom to top) $n = 3, 4, 5, 6$. The lines are the fits $T_f = b_f + a_f \ln G$ where b_f and a_f are fit parameters. The public goods contribution is $c = 1$.

discrete categories rather than as quantitative traits, as is most likely the case [29]. Alas, the quantitative strategy trait fix has only been tested in the case of weak altruism, where a cooperator also benefits from her altruistic behavior and the benefit exceeds the cost of cooperation [29]. In this case, even assuming discrete traits, there are conditions under which defectors can be invaded by rare cooperators, as we showed in Section 3.1. For the others-only variant of the n -prisoner's dilemma, and for scenarios of strong altruism in general, the problem of origination remains.

The main purpose of this paper was to highlight the importance that the mode of reproduction can have on the fate of strong altruism in populations that are frequently divided into small temporary groups, in which individuals interact locally and fitness-related processes such as mating take place. The selection dynamics resulting from the different forms of reproduction considered here were solved analytically in the limit of infinitely many groups, assuming, as usual, that individuals colonize the groups according to a binomial distribution, so that cooperators do not cluster into groups, and that local interactions follow the others-only variant of the n -prisoner's dilemma. Under biparental sexual reproduction, the results are quite striking, challenging conventional wisdom by showing that strong altruism can indeed evolve in the absence of assortment and punishment in temporary groups if the amplification factor r is sufficiently large. The basic reason is that biparental sexual reproduction may enable a resident population of cooperators to resist invasion by rare defectors, since the fitness




Figure 6: Probability of fixation of the cooperators as a function of the amplification factor r for (from left to right at $\Pi = 0.8$) $n = 3, 4, 5, 6$ (left panel). The vertical dashed lines indicate the thresholds $r_c(3) = 9.62$, $r_c(4) = 20.37$, $r_c(5) = 34.32$ and $r_c(6) = 51.83$ beyond which the all-cooperators fixed point attracts the orbits starting at $q_0 = 0.5$ for $G \rightarrow \infty$. The right panel shows the same data with the curves shifted by $r_c(n)$. The other parameters are $G = 400$ and $c = 1$.

advantage of a single defector in a group with $n - 1$ cooperators is smothered by the requirement that the recombinant offspring chromosome has a one-half chance of being a cooperator in any mating involving the defector. However, and most surprisingly, biparental sexual reproduction can also enable the invasion of a resident population of defectors by rare cooperators if the cost of cooperation c is not large, thus solving the problem of origination. These results contrast strikingly with asexual reproduction and sexual reproduction with self-mating modes, for which cooperation is always doomed and the amplification factor r can only slow the rate at which cooperators disappear from the population [2].

These findings are summarized in Fig. 1, which shows a region, labeled C, where cooperation is the only evolutionary outcome. For large group size n , this region exists for $c < 1/2n$ and $r > 2n^2$. The only modification we have made to Hamilton's original model [2] is to change asexual reproduction to biparental sexual reproduction, which is how most sexual reproduction occurs [20]. This result provides a neat solution to both the problem of origination and the conditions under which strong altruism can evolve, thus avoiding the 'inverse genetic fallacy' [35], i.e. the inappropriate attribution of mechanisms that might support cooperation (e.g., positive assortment and punishment) to the explanation of its origin. For large cooperation costs (i.e., $c \approx 1$) and large n , we find a wide region in the parameter space, labeled C+D in Fig. 1, bounded at the bottom by $r = 2n^2$, where cooperation can evolve provided that $q_0 > 4n/r$, where q_0 is the initial frequency of cooperators. These estimates for large n

are worst-case scenarios for cooperation, since increasing group size n favors defection. In addition, we have shown that the results of our analytical study for infinitely many groups are robust to the demographic noise that arises when the number of groups is finite.

How realistic are the conditions in our model for the evolution of cooperation in nature? One possible example that comes to mind is viral co-infection, which complicates the symptoms and diagnosis of disease [36], and also leads to some interesting evolutionary outcomes [37, 38, 39]. Viral co-infections can be modeled by analogy with Wilson’s trait groups [40]. In RNA viruses, recombination can occur via a copy-choice mechanism during viral replication, in which the viral RNA polymerase switches templates during strand synthesis [41]. Virus-virus interactions after co-infection can be modeled using game theory [39], as in the classic example of the RNA phage $\phi 6$, where competitive interactions were shown to follow the rules of the prisoner’s dilemma [37]. In this case, the fixation of high-multiplicity phages led to a decrease in population fitness. To escape the dilemma, it has been suggested that fitness payoffs evolve from the prisoner’s dilemma to the snowdrift game, where coexistence between cooperators and defectors is possible [42]. How general this result can be in viral co-infections is unknown, but as shown here, there may be circumstances where the payoffs are consistent with the prisoner’s dilemma and yet cooperation can still evolve.

Finally, it is an open question whether our conclusions apply to diploid sexual organisms under the more realistic assumption that altruistic and selfish strategies are quantitative traits rather than discrete behaviors. As Maynard Smith wrote [43]: “Most populations of interest have sexual diploid inheritance. In many cases this does not matter.” Apart from the obvious case where a pure phenotypic strategy is produced by a heterozygous genotype and cannot be fixed in a random mating population because of Mendelian inheritance, we believe there is no simple answer to the question of whether it does or does not matter.

Acknowledgments

JFF is partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazil), grant number 305620/2021-5. MS is funded by grant PID2021-127107NB-I00 from Ministerio de Ciencia e Innovación (Spain) and grant 2021 SGR 00526 from Generalitat de Catalunya.

Declaration of interest:

None.

References

- [1] C. Darwin, *The Descent of Man, and Selection in Relation to Sex*, John Murray, London, 1871.

- [2] W.D. Hamilton, Innate social aptitudes of man: an approach from evolutionary genetics, in: R. Fox R (ed), *ASA Studies 4: Biological Anthropology*, Malaby Press, London, 1975, pp. 133–153.
- [3] D.S. Wilson, A theory of group selection, *Proc. Nat. Acad. Sci. USA* 72 (1975) 143–146.
- [4] D.S. Wilson, Weak altruism, strong group selection, *Oikos* 59 (1990) 135–140.
- [5] J.F. Fontanari, M. Santos, The dynamics of casual groups can keep free-riders at bay, *Math. Biosc.* 372 (2024) 109188.
- [6] J.W. Pepper, Relatedness in trait group models of social evolution, *J. Theor. Biol.* 206 (2000) 355–368.
- [7] S.A. West, A.S. Griffin, A. Gardner, Social semantics: altruism, cooperation, mutualism, strong reciprocity and group selection, *J. Evol. Biol.* 20 (2007) 415–432.
- [8] S. Okasha, *Evolution and the Levels of Selection*, Oxford University Press, Oxford, 2009.
- [9] J. Fox, M. Guyer, Public Choice and cooperation in N-person Prisoner’s Dilemma, *J. Conflict Resolut.* 22 (1978) 469–481.
- [10] R. Boyd, R.J. Richerson, Punishment allows the evolution of cooperation (and anything else), in *sizable group*, *Ethol. Sociobiol.* 13 (1992))171–195.
- [11] E. Fehr, S. Gächter, Cooperation and punishment in public goods experiments, *Am. Econ. Rev.* 90 (2000) 980–994.
- [12] D.F. Zheng, H.P. Yin, C.H. Chan, P.M. Hui, Cooperative behavior in a model of evolutionary snowdrift games with N -person interactions, *Europhys. Lett.* 80 (2007) 18002.
- [13] Santos MD, Pinheiro FL, Santos FC, Pacheco JM (2012) Dynamics of N -person snowdrift games in structured populations. *J. Theor. Biol.* 315: 81–86.
- [14] M. Archetti, I. Scheuring, Review: Game theory of public goods in one-shot social dilemmas without assortment, *J. Theor. Biol.* 299 (2012) 9–20.
- [15] H. De Silva, C. Hauert, A. Traulsen, K. Sigmund, Freedom, enforcement, and the social dilemma of strong altruism, *Evol. Econ.* 20 (2010) 203–217.
- [16] K. Sigmund, H. de Silva, A. Traulsen, C. Hauert, Social learning promotes institutions for governing the commons, *Nature* 466 (2010) 861–863.
- [17] B. Charlesworth, A Note on the Evolution of Altruism in Structured Demes, *Am. Nat.* 113 (1979) 601–630.

- [18] A. Grafen, Natural selection, kin selection and group selection, in: J. R. Krebs, N. B. Davies (eds), *Behavioural Ecology: An Evolutionary Approach*, Blackwell Scientific Publications, Oxford, UK, 1984, pp. 62–84.
- [19] M. Broom, J. Rychtář, *Game-Theoretical Models in Biology*, CRC Press, Boca Raton, FL, 2013.
- [20] G.C. Williams, *Sex and Evolution*, Princeton University Press, Princeton, 1975, p. 115.
- [21] A. Robinson, Einstein said that - didn't he? *Nature* 557 (2018) 30.
- [22] A. Traulsen, J.C. Claussen, C. Hauert, Coevolutionary Dynamics: From Finite to Infinite Populations, *Phys. Rev. Lett.* 95 (2005) 238701.
- [23] J.F. Fontanari, Imitation dynamics and the replicator equation, *Europhys. Lett.* 146 (2024) 47001.
- [24] J.A. Fletcher, M. Zwick, Strong altruism can evolve in randomly formed groups. *J. Theor. Biol.* 228 (2004) 303–313.
- [25] G.R. Price, Selection and covariance, *Nature* 227 (1970) 520–521.
- [26] D. Alves, P.R.A. Campos, A.T.C. Silva, J.F. Fontanari, Group selection models in prebiotic evolution, *Phys. Rev. E* 63 (2000) 011911.
- [27] N.F. Britton, *Essential Mathematical Biology*, Springer, London, 2003.
- [28] G.C. Williams, D.C. Williams, Natural selection of individually harmful social adaptations among sibs with special reference to social insects, *Evolution* 11 (1957) 32–39.
- [29] D.S. Wilson, L. A. Dugatkin, Group selection and assortative interactions, *Am. Nat.* 149 (1997) 336–351.
- [30] K. Binder, The Monte Carlo method for the study of phase transitions: A review of some recent progress, *J. Comp. Phys.* 59 (1985) 1–55.
- [31] P.R.A. Campos, J.F. Fontanari, Finite-size scaling of the error threshold transition in finite populations, *J. Phys. A: Math. Gen.* 32 (1999) L1–L7.
- [32] L. Nunney, Group selection, altruism, and structured-deme models, *Am. Nat.* 126 (1985) 212–230.
- [33] J. Maynard Smith, The origin of altruism, *Nature* 393 (1998) 639–640.
- [34] C. Xia, J. Wang, M. Perc, Z. Wang, Reputation and reciprocity, *Phys. Life Rev.* 46 (2023) 8–45.
- [35] A.J. Field, *Altruistically Inclined?: The Behavioral Sciences, Evolutionary Theory, and the Origins of Reciprocity*, The University of Michigan Press, Ann Arbor, 2001.

- [36] Y. Du, C. Wang, Y. Zhang, Viral coinfections, *Viruses* 14 (2022) 2645.
- [37] P.E. Turner, L. Chao, Prisoner's Dilemma in an RNA virus, *Nature* 398 (1999) 441–443.
- [38] P.E. Turner, L. Chao, Escape from Prisoner's Dilemma in RNA phage $\phi 6$, *Am. Nat.* 161 (2003) 497–505.
- [39] S.F. Elena, G.P. Bernet, J.L. Carrasco, The games plant viruses play, *Curr. Opin. Virol.* 8 (2014) 62–87.
- [40] E. Szathmáry, Natural selection and dynamical coexistence of defective and complementing virus segments, *J. Theor. Biol.* 157(1992) 383–406.
- [41] T.C. Jarvis, K. Kirkegaard, The polymerase in its labyrinth: mechanisms and implications of RNA recombination, *Trends Genet.* 7 (1991) 186–191.
- [42] L. Chao, S.F. Elena, Nonlinear trade-offs allow the cooperation game to evolve from Prisoner's Dilemma to Snowdrift, *Proc. R. Soc. B* 284 (2017) 20170228.
- [43] J. Maynard Smith, *Evolution and the Theory of Games*, Cambridge University Press, Cambridge, 1982, p. 40.