An Upper Limit on the Photoproduction Cross Section of the Spin-Exotic $\pi_1(1600)$

F. Afzal⁶,⁴ C. S. Akondi⁶,¹⁰ M. Albrecht⁶,²⁷ M. Amaryan⁶,²⁴ S. Arrigo,³³ V. Arroyave,⁹ A. Asaturyan⁶,²⁷ A. Austregesilo⁰,²⁷ Z. Baldwin⁰,⁵ F. Barbosa,²⁷ J. Barlow⁰,^{10,26} E. Barriga⁰,¹⁰ R. Barsotti,¹⁵ D. Barton,²⁴ V. Baturin,²⁴ V. V. Berdnikov[®],²⁷ T. Black,²³ W. Boeglin[®],⁹ M. Boer,³¹ W. J. Briscoe[®],¹¹ T. Britton,²⁷ S. Cao,¹⁰ E. Chudakov[®],²⁷ G. Chung[®],³¹ P. L. Cole[®],¹⁷ O. Cortes,¹¹ V. Crede[®],¹⁰ M. M. Dalton[®],²⁷ D. Darulis[®],¹² A. Deur⁽⁶⁾,²⁷ S. Dobbs⁽⁶⁾,¹⁰ A. Dolgolenko⁽⁶⁾,¹⁶ M. Dugger⁽⁶⁾,¹ R. Dzhygadlo,¹³ D. Ebersole⁽⁶⁾,¹⁰ M. Edo,⁷ H. Egiyan⁽⁶⁾,²⁷ T. Erbora[®],⁹ P. Eugenio[®],¹⁰ A. Fabrizi,¹⁸ C. Fanelli[®],³³ S. Fang[®],¹⁴ J. Fitches[®],¹² A. M. Foda[®],¹³ S. Furletov[®],²⁷ L. Gan[®],²³ H. Gao,⁸ A. Gardner,¹ A. Gasparian,²² D. Glazier[®],¹² C. Gleason[®],³⁰ V. S. Goryachev[®],¹⁶ B. Grube⁰,²⁷ J. Guo⁶,⁵ L. Guo,⁹ J. Hernandez⁰,¹⁰ K. Hernandez,¹ N. D. Hoffman⁶,⁵ D. Hornidge⁰,²⁰ G. Hou,¹⁴ P. Hurck⁰,¹² A. Hurley,³³ W. Imoehl⁰,^{5,*} D. G. Ireland⁰,¹² M. M. Ito⁰,¹⁰ I. Jaegle⁰,²⁷ N. S. Jarvis⁰,⁵ T. Jeske,²⁷ M. Jing,¹⁴ R. T. Jones[©],⁷ V. Kakoyan,³⁵ G. Kalicy,⁶ V. Khachatryan,¹⁵ C. Kourkoumelis[©],² A. LaDuke,⁵ I. Larin,²⁷ D. Lawrence[©],²⁷ D. I. Lersch[©],²⁷ H. Li[©],³³ B. Liu[©],¹⁴ K. Livingston[©],¹² G. J. Lolos,²⁵ L. Lorenti,³³ V. Lyubovitskij[©],^{29,28} R. Ma,¹⁴ D. Mack,²⁷ A. Mahmood,²⁵ H. Marukyan[©],³⁵ V. Matveev[©],¹⁶ M. McCaughan⁰,²⁷ M. McCracken⁰,^{5,32} C. A. Meyer⁰,⁵ R. Miskimen⁰,¹⁸ R. E. Mitchell⁰,¹⁵ K. Mizutani⁰,²⁷ V. Neelamana⁽⁰⁾,²⁵ L. Ng⁽⁰⁾,²⁷ E. Nissen,²⁷ S. Orešić,²⁵ A. I. Ostrovidov,¹⁰ Z. Papandreou⁽⁰⁾,²⁵ C. Paudel⁽⁰⁾,⁹ R. Pedroni,²² L. Pentchev[®],²⁷ K. J. Peters,¹³ E. Prather,⁷ S. Rakshit[®],¹⁰ J. Reinhold[®],⁹ A. Remington[®],¹⁰ B. G. Ritchie[®],¹ J. Ritman[®],^{13,3} G. Rodriguez[®],¹⁰ D. Romanov[®],¹⁹ K. Saldana[®],¹⁵ C. Salgado[®],²¹ S. Schadmand[®],¹³ A. M. Schertz[®],¹⁵ K. Scheuer[®],³³ A. Schick,¹⁸ A. Schmidt[®],¹¹ R. A. Schumacher[®],⁵ J. Schwiening[®],¹³ N. Septian[®],¹⁰ P. Sharp[®],¹¹ X. Shen[®],¹⁴ M. R. Shepherd[®],¹⁵ J. Sikes,¹⁵ A. Smith[®],⁸ E. S. Smith[®],³³ D. I. Sober,⁶ A. Somov,²⁷ S. Somov,¹⁹ J. R. Stevens[®],³³ I. I. Strakovsky[®],¹¹ B. Sumner,¹ K. Suresh,³³ V. V. Tarasov[®],¹⁶ S. Taylor[®],²⁷ A. Teymurazyan,²⁵ A. Thiel[®],⁴ T. Viducic[®],²⁴ T. Whitlatch,²⁷ N. Wickramaarachchi⁰,⁶ Y. Wunderlich⁰,⁴ B. Yu⁰,⁸ J. Zarling⁰,²⁵ Z. Zhang⁰,³⁴ X. Zhou⁰,³⁴ and B. Zihlmann⁰²⁷

(The GLUEX Collaboration)

¹Polytechnic Sciences and Mathematics, School of Applied Sciences and Arts,

Arizona State University, Tempe, Arizona 85287, USA

²Department of Physics, National and Kapodistrian University of Athens, 15771 Athens, Greece

³Ruhr-Universität-Bochum, Institut für Experimentalphysik, D-44801 Bochum, Germany

⁴Helmholtz-Institut für Strahlen- und Kernphysik Universität Bonn, D-53115 Bonn, Germany

⁵Department of Physics, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

⁶Department of Physics, The Catholic University of America, Washington, D.C. 20064, USA

Department of Physics, University of Connecticut, Storrs, Connecticut 06269, USA

⁸Department of Physics, Duke University, Durham, North Carolina 27708, USA ⁹Department of Physics, Florida International University, Miami, Florida 33199, USA

¹⁰Department of Physics, Florida State University, Tallahassee, Florida 32306, USA ¹¹Department of Physics, The George Washington University, Washington, D.C. 20052, USA

¹²School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom

¹³GSI Helmholtzzentrum für Schwerionenforschung GmbH, D-64291 Darmstadt, Germany

¹⁴Institute of High Energy Physics, Beijing 100049, People's Republic of China

¹⁵Department of Physics, Indiana University, Bloomington, Indiana 47405, USA

¹⁶National Research Centre Kurchatov Institute, Moscow 123182, Russia

¹⁷Department of Physics, Lamar University, Beaumont, Texas 77710, USA

¹⁸Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003, USA

¹⁹National Research Nuclear University Moscow Engineering Physics Institute, Moscow 115409, Russia

²⁰Department of Physics, Mount Allison University, Sackville, New Brunswick E4L 1E6, Canada

²¹Department of Physics, Norfolk State University, Norfolk, Virginia 23504, USA

²²Department of Physics, North Carolina A&T State University, Greensboro, North Carolina 27411, USA ²³Department of Physics and Physical Oceanography,

University of North Carolina at Wilmington, Wilmington, North Carolina 28403, USA

²⁴Department of Physics, Old Dominion University, Norfolk, Virginia 23529, USA

²⁵Department of Physics, University of Regina, Regina, Saskatchewan S4S 0A2, Canada

²⁶Department of Mathematics, Physics, and Computer Science,

Springfield College, Springfield, Massachusetts, 01109. USA

²⁷ Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

²⁸Laboratory of Particle Physics, Tomsk Polytechnic University, 634050 Tomsk, Russia

²⁹Department of Physics, Tomsk State University, 634050 Tomsk, Russia

³⁰Department of Physics and Astronomy, Union College, Schenectady, New York 12308, USA

³¹Department of Physics, Virginia Tech, Blacksburg, VA 24061, USA

³²Department of Physics, Washington & Jefferson College, Washington, Pennsylvania 15301, USA

³³Department of Physics, William & Mary, Williamsburg, Virginia 23185, USA

Wuhan, Hubei 430072, People's Republic of China

³⁵A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute), 0036 Yerevan, Armenia

(Dated: July 4, 2024)

The spin-exotic hybrid meson $\pi_1(1600)$ is predicted to have a large decay rate to the $\omega\pi\pi$ final state. Using 76.6 pb⁻¹ of data collected with the GlueX detector, we measure the cross sections for the reactions $\gamma p \to \omega \pi^+ \pi^- p$, $\gamma p \to \omega \pi^0 \pi^0 p$, and $\gamma p \to \omega \pi^- \pi^0 \Delta^{++}$ in the range $E_{\gamma} = 8 - 10$ GeV. Using isospin conservation, we set the first upper limits on the photoproduction cross sections of the $\pi_1^0(1600)$ and $\pi_1^-(1600)$. We combine these limits with lattice calculations of decay widths and find that photoproduction of $\eta' \pi$ is the most sensitive two-body system to search for the $\pi_1(1600)$.

Quantum chromodynamics (QCD) is the fundamental theory that describes the interactions of quarks and gluons. In nature, protons and neutrons are the most common bound states of QCD, and can be modeled as color-singlet three-quark states called baryons. Even simpler hadronic states are mesons, which consist of just a quark and an anti-quark. More complicated configurations are allowed, such as hybrid mesons, in which the gluonic field contributes directly to the total angular momentum J, parity P, and charge conjugation C quantum numbers of the state. Hybrid mesons are predicted from first principles in lattice QCD (LQCD) [1, 2], as well as in phenomenological models of QCD [3, 4]. Several of the predicted hybrid mesons have spin-exotic quantum numbers, meaning their J^{PC} values are impossible for conventional mesons. LQCD predicts the lightest spinexotic nonet to have $J^{PC} = 1^{-+}$, with the isovector state, the π_1 , being the lightest [2].

Mapping out the spectrum of these hybrid states is a crucial goal of hadronic physics, since it provides important information on the role of gluons in QCD bound states. The best experimental candidate for the lightest hybrid meson is the $\pi_1(1600)$, due to the recent measurements of its decay to $\eta\pi$ and $\eta'\pi$ [5–7]. Results from LQCD, however, predict the dominant decay mode of the $\pi_1(1600)$ will be $b_1\pi$, with a branching fraction of at least 70% [8], which is supported by the fact that there have been several experimental reports of the $\pi_1(1600)$ decaying to $b_1\pi$ [9–12].

While there is considerable experimental evidence for the $\pi_1(1600)$, it has not yet been observed in the photoproduction process. The CLAS experiment has previously studied the charge-exchange reaction $\gamma p \rightarrow \pi^+\pi^+\pi^-(n)$ in the range $E_{\gamma} = 4.8 - 5.4$ GeV, and found that the dominant contributions were from the conventional $a_2^+(1320)$ and $\pi_2^+(1670)$ mesons, with no evidence for the spin-exotic $\pi_1^+(1600)$ decay to $\rho\pi$ [13]. Given the large predicted decay of the $\pi_1(1600)$ to $b_1\pi$ and the dominant decay of the b_1 to $\omega\pi$ [14, 15], in this paper we aim to study the isospin-1 contribution to $\omega\pi\pi$ photoproduction. The existing photoproduction results for $\omega\pi^+\pi^-$ suggest the isospin-0 contribution is larger than the isospin-1 contribution [16–18], which implies it will be difficult to study the isospin-1 component using only $\omega \pi^+ \pi^-$ data. However, with access to different charge combinations of the $\omega \pi \pi$ system and by exploiting isospin symmetry, it is possible to separate the isospin-0 and isospin-1 contributions. There are no known isospin-2 (i.e. flavor-exotic) states, so we assume no isospin-2 contributions to the $\omega \pi \pi$ distributions. Under this assumption, the $\omega \pi^0 \pi^0$ system is purely isospin-0 and the $\omega \pi^+ \pi^-$ system contains both isospin-0 and isospin-1. Using isospin Clebsch-Gordan coefficients, we find the isospin-0 cross section

$$\sigma(\omega\pi\pi)_{I=0} = 3\sigma(\omega\pi^0\pi^0). \tag{1}$$

The total neutral $\omega \pi \pi$ cross section can be written as $\sigma(\omega \pi \pi)_{I=0} + \sigma(\omega \pi \pi)_{I=1} = \sigma(\omega \pi^+ \pi^-) + \sigma(\omega \pi^0 \pi^0)$, so solving for $\sigma(\omega \pi \pi)_{I=1}$ gives the isospin-1 cross section

$$\sigma(\omega\pi\pi)_{I=1} = \sigma(\omega\pi^+\pi^-) - 2\sigma(\omega\pi^0\pi^0).$$
(2)

In this Letter, we measure the cross sections for

$$\gamma p \rightarrow \omega \pi^0 \pi^0 p$$
 (R1)

$$\gamma p \to \omega \pi^+ \pi^- p$$
 (R2)

$$\gamma p \rightarrow \omega \pi^- \pi^0 \Delta^{++}$$
 (R3)

where we reconstruct $\omega \to \pi^+\pi^-\pi^0$, $\pi^0 \to \gamma\gamma$ and $\Delta^{++} \rightarrow \pi^+ p$. Using Eq. 2 and R3, we isolate the cross section for photoproducing an isospin-1 $\omega \pi \pi$ system against a proton or Δ^{++} . We observe no clear structure in the mass spectrum that is consistent with a Breit-Wigner amplitude for the $\pi_1(1600)$. A comprehensive search for $\pi_1(1600) \rightarrow \omega \pi \pi$ would require a multidimensional angular analysis, which is beyond the scope of this work. However, we can use the isospin-1 $\omega \pi \pi$ spectra to set a meaningful upper limit on the $\pi_1(1600)$ photoproduction cross section, which can be used to guide our study of $\eta\pi$ and $\eta'\pi$ production. Using known resonance parameters for the $a_2(1320)$ and $\pi_1(1600)$, we determine the largest $\pi_1(1600) \rightarrow b_1\pi \rightarrow \omega\pi\pi$ contribution that is consistent with our measured $\omega \pi \pi$ data. These limits are combined with the allowed ranges of decay widths for $\pi_1(1600) \rightarrow \eta \pi$ and $\pi_1(1600) \rightarrow \eta' \pi$ from Ref. [8] to estimate the maximal $\pi_1(1600)$ contributions to the

³⁴School of Physics and Technology, Wuhan University,

photoproduced $\eta^{(\prime)}\pi$ mass spectra. In particular, we will show that the $\eta\pi$ final states can have at most a percentlevel contribution from the $\pi_1(1600)$, while our limit cannot rule out a dominant $\pi_1(1600)$ contribution to the $\eta'\pi$ systems. This leads us to the conclusion that in photoproduction, the $\eta'\pi$ final states are the most sensitive two-body systems to search for the $\pi_1(1600)$.

The GlueX experiment at Jefferson Lab has acquired a world-leading photoproduction data sample, with a goal of mapping out the hybrid meson spectrum. The linearly polarized photons are produced via the coherent bremsstrahlung process by an electron beam incident on a 50 µm thick diamond wafer, where the maximum photon polarization of about 35% occurs in the coherent peak at a photon energy of 8.8 GeV. Events with photon energies outside the coherent peak are also recorded, leading to a tagged photon energy spectrum between 7.0 GeV and 11.6 GeV. The photon beam is then incident on a liquid hydrogen target. GlueX is a nearly hermetic detector that reconstructs charged particles and photons. Particle identification is available from both dE/dx in the drift chambers and flight-time measurements in the central calorimeter and a dedicated downstream detector. Detailed information on the beamline and detector can be found elsewhere [19-26].

Events with photon beam energies between 8.0 GeV and 10.0 GeV are selected from Phase-I of the GlueX data, yielding a total luminosity of 76.6 pb^{-1} . We fully reconstruct events with the final states for reactions R1, R2, or R3. We keep events with $0.1 \, (\text{GeV}/c)^2 < -t < 0.5 \, (\text{GeV}/c)^2$, where -t is the fourmomentum transfer squared to the recoil baryon. A kinematic fit to enforce energy and momentum conservation with mass constraints on $\pi^0 \to \gamma \gamma$ is then performed and events with a resulting χ^2/ndf smaller than 5 are retained for further analysis. Beam bunches arrive everv 4 ns, so we select beam photon candidates within ± 2 ns of the beam bunch that corresponds to the reconstructed final state particles. For each event, there can be multiple beam photon candidates, as well as multiple ways to assign the final state particles. To select a unique combination for a given event, we use only the beam-photon and final-state particle combination with the smallest χ^2/ndf . For R3, we require the invariant mass $M(\pi^+ p) < 1.35 \text{ GeV}/c^2$ to select the Δ^{++} . For R2, background studies show a large contamination by $\gamma p \rightarrow b_1^- \Delta^{++}$, which we remove by requiring $M(\pi^+ p) > 1.35 \text{ GeV}/c^2$ for both π^+ combinations in the event. Background studies also show a large background for R1 from $\gamma p \to \omega \pi^0 \pi^0 \pi^0 p$, which we suppress by requiring that no additional reconstructed calorimeter showers are present in the event. After this requirement, no more than 3% of the remaining $\omega \pi^0 \pi^0$ events are due to the $\omega \pi^0 \pi^0 \pi^0$ background.

Monte Carlo (MC) samples are used to estimate the reconstruction efficiency for the $\omega \pi \pi$ events. The MC

samples are generated by modeling the decay $\pi_1(1600) \rightarrow$ $\omega\pi\pi$ as a three-body decay with a uniform distribution in phase space. We assume an exponential distribution for the four-momentum transfer squared -t, given by e^{bt} with $b = 5 \, (\text{GeV}/c)^{-2}$. The $\pi_1(1600)$ shape is modeled as a relativistic Breit-Wigner, with the mass and width taken as the pole parameters from Ref. [6], since it is the only analysis that simultaneously describes the $\eta\pi$ and $\eta' \pi$ data. This simulated data is passed through a GEANT4 simulation of the detector, and is analyzed using the same selection criteria as for the data. To determine the reconstruction efficiency, we perform a binned maximum likelihood fit on the reconstructed ω candidate masses in signal MC with a Voigtian [27], where the internal width of the ω is fixed to the value from the PDG [14]. This leaves the ω mass, the resolution, and the normalization as floating parameters in the fit to MC data. We integrate the Voigtian and divide by the number of generated events to determine our reconstruction efficiency. We then fix the mass and resolution parameters of the Voigtian to the results of the MC fits, leaving the overall normalization as the only free parameter in fits to data.

To measure the differential $\omega \pi \pi$ photoproduction cross sections $d\sigma/dm_{\omega\pi\pi}$, we divide the data into 20 equidistant bins of $\omega\pi\pi$ invariant mass $m_{\omega\pi\pi}$ from 1.20 GeV/ c^2 to 2.20 GeV/ c^2 . For each of these bins, we plot the invariant mass for each $\pi^+\pi^-\pi^0$ combination. We fit these data using a least- χ^2 fit with a Voigtian function for the ω signal plus a fourth-order polynomial for the background. These fits are then used to determine the $\omega\pi\pi$ yield $Y_{\omega\pi\pi}$ for each bin. Sample fits to the $M(\pi^+\pi^-\pi^0)$ distributions are shown in the Supplemental Materials [28].

We use the yields and reconstruction efficiencies to determine the cross sections for R1-R3 in each bin of $\omega\pi\pi$ mass. The measured cross sections are shown in Fig. 1. We expect the $a_2(1320)$ to contribute to both the $\gamma p \rightarrow \omega \pi^+ \pi^- p$ and $\gamma p \rightarrow \omega \pi^- \pi^0 \Delta^{++}$ reactions, but only the charge-exchange reaction shows a clear $a_2(1320)$ peak. This is due to the fact that the $\omega\pi^+\pi^-$ system has both isospin-1 and isospin-0 components, while the $\omega\pi^+\pi^-$ mass spectrum is dominated by isospin-0, which is consistent with the previous results from the literature [16].

As a next step, we use R1-R3 and Eq. 2 to determine the cross sections for producing an isospin-1 $\omega \pi \pi$ system recoiling against a proton or a Δ^{++} , which we show in Fig. 2. In the mass range 1.2 GeV/ $c^2 < m_{\omega\pi\pi} <$ 1.6 GeV/ c^2 , we expect contributions from non-resonant $\omega\pi\pi$, the $a_2(1320)$, the $\pi_2(1670)$, the $a_2(1700)$, and the $\pi_1(1600)$. Previous studies of $\gamma p \rightarrow \rho^0 \pi^- \Delta^{++}$ at the SLAC Hybrid facility show a large $a_2(1320)$ signal, and a substantially smaller signal near the $\pi_2(1670) \rightarrow \omega \rho$ branching fraction is just 2.7% while $\mathcal{B}(\pi_2(1670) \rightarrow$ $3\pi) = 95.8\%$ [14], implies that the $\pi_2(1670)$ should be



FIG. 1. The differential cross sections for the reactions $\gamma p \to \omega \pi^0 \pi^0 p$ (a), $\gamma p \to \omega \pi^+ \pi^- p$ (b), and $\gamma p \to \omega \pi^- \pi^0 \Delta^{++}$ (c) with 0.1 $(\text{GeV}/c)^2 < -t < 0.5 (\text{GeV}/c)^2$ as a function of $\omega \pi \pi$ invariant mass. The filled rectangles show the statistical uncertainty, and the full error bars are statistical and systematic uncertainties added in quadrature. Not included in the error bars are uncertainties due to the photon and charged-track efficiency systematic uncertainty, as well as the systematic uncertainty due to the luminosity. These uncertainties are correlated for each source across the three measurements, so they cannot be easily visualized.



FIG. 2. The isospin-1 component of the cross sections for neutral (a) and negatively charged (b) $\omega\pi\pi$ for $0.1 (\text{GeV}/c)^2 < -t < 0.5 (\text{GeV}/c)^2$. The error bars are statistical only. The cross sections are fit with an $a_2(1320)$ (green) and $\pi_1(1600)$ (magenta) Breit-Wigner shape in the range $M(\omega\pi\pi) < 1.6 \text{ GeV}/c^2$, indicated by the vertical dashed line.

a small contribution to $\omega \pi \pi$. The $a_2(1700)$ cross section has not been measured in photoproduction, but assuming that both the cross section and the branching fraction to $\omega \pi \pi$ are similar to those for the $a_2(1320)$, the $a_2(1700)$ would make up a small fraction of the events in the $\pi_1(1600)$ mass region.

We perform a binned least- χ^2 fit to these isospin-1 $\omega\pi\pi$ photoproduction cross sections with the sum of an $a_2(1320)$ and $\pi_1(1600)$ shape to determine the largest $\pi_1(1600)$ signal consistent with the data. The $a_2(1320)$ shape is a fixed-width relativistic Breit-Wigner with M = $1.3182 \text{ GeV}/c^2$ and $\Gamma = 110 \text{ MeV}$ [14], while the $\pi_1(1600)$ shape is a fixed-width relativistic Breit-Wigner using the JPAC parameters [6] from their fit to the COMPASS data [5], i.e. $M_{\pi_1(1600)} = (1564 \pm 24 \pm 86) \text{ MeV}/c^2$ and $\Gamma_{\pi_1(1600)} = (492 \pm 54 \pm 102) \text{ MeV}$. We only fit the range $m_{\omega\pi\pi} < 1.6 \text{ GeV}/c^2$ to capture the leading edge of the $\pi_1(1600)$ shape because expanding to higher masses would necessitate including a background function, as well as contributions from heavier states. The magnitude of the $a_2^0(1320)$ cross section is fixed based on the results of a partial-wave analysis to $\gamma p \to \eta \pi^0 p$ [30], while the magnitude of the $a_2^-(1320)$ cross section is a free parameter in the fit. Fixing the magnitude of the $a_2^0(1320)$ cross section is necessary because there are large fitting systematic uncertainties on the ω yields for the first three mass bins of $M(\omega \pi^0 \pi^0)$ due to a combinatorial background. For the fit of the neutral isospin-1 cross section, the only free parameter is the magnitude of the $\pi_1^0(1600)$. For the $\omega\pi^-\pi^0$ cross section, both the magnitude of $a_2^-(1320)$ and $\pi_1^-(1600)$ are allowed to float. Note that the fit model is not able to precisely match the shape of the data since no background contributions are included. We include a discussion of the systematic uncertainties in the Supplemental Materials [28].

In order to determine the upper limits on the $\pi_1(1600)$

cross sections, we need to determine how the likelihood functions vary based on the size of $\sigma(\gamma p \to \pi_1^0(1600)p)$ and $\sigma(\gamma p \to \pi_1^- \Delta^{++})$, which we denote $\sigma(\pi_1(1600))$ for simplicity. To do this, we repeat the fits to the I = 1cross sections for fixed values of $\sigma(\pi_1(1600))$. For each fit, we record the likelihood value, which gives us the likelihood as a function of $\sigma(\pi_1(1600))$. To determine the upper limit, we find the cross section corresponding to the 90th percentile in the likelihood distribution. After including the systematic uncertainties as discussed in the Supplemental Materials [28], we place 90% confidence level (C.L.) upper limits of

$$\sigma(\gamma p \to \pi_1^0(1600)p) \times \mathcal{B}(\pi_1(1600) \to b_1\pi) < 123 \text{ nb} (3)$$

and

$$\sigma(\gamma p \to \pi_1^-(1600)\Delta^{++}) \times \mathcal{B}(\pi_1(1600) \to b_1\pi) < 435 \text{ nb}$$
(4)

where we assume $\mathcal{B}(b_1 \to \omega \pi) = 100\%$. Several systematic uncertainties cancel when measuring a ratio of cross sections, including all of the luminosity systematic uncertainty and portions of the photon and tracking efficiency uncertainties. Therefore, we also report our limits normalized to the measured $a_2(1320)$ cross sections. After accounting for these reduced systematic uncertainties, we place 90% C.L. upper limits of

$$\frac{\sigma(\gamma p \to \pi_1^0(1600)p) \times \mathcal{B}(\pi_1(1600) \to b_1\pi)}{\sigma(\gamma p \to a_2^0(1320)p)} < 2.2 \quad (5)$$

and

$$\frac{\sigma(\gamma p \to \pi_1^-(1600)\Delta^{++}) \times \mathcal{B}(\pi_1(1600) \to b_1\pi)}{\sigma(\gamma p \to a_2^-(1320)\Delta^{++})} < 0.96.$$
(6)

Lattice calculations from Ref. [8] include a range of allowed values for $\mathcal{B}(\pi_1(1600) \rightarrow b_1\pi)$. We can use the lower limit on $\mathcal{B}(\pi_1(1600) \rightarrow b_1\pi)$ from their result to set an upper limit directly on the $\pi_1(1600)$ photoproduction cross section, with the caveat that it is no longer a 90% C.L. limit since the uncertainty on $\mathcal{B}(\pi_1(1600) \rightarrow b_1\pi)$ is not folded in. Using their lower bound on $\mathcal{B}(\pi_1(1600) \rightarrow b_1\pi)$ is not folded in. Using their lower bound on $\mathcal{B}(\pi_1(1600) \rightarrow b_1\pi)$, we set limits of $\sigma(\gamma p \rightarrow \pi_1^0(1600)p) < 177$ nb and $\sigma(\gamma p \rightarrow \pi_1^-(1600)\Delta^{++}) < 627$ nb. These are the first upper limits on the $\pi_1^0(1600)$ and $\pi_1^-(1600)$ photoproduction cross sections. Note that no other reaction can be used to set an upper limit on the $\pi_1(1600)$ since $b_1\pi$ is the only decay mode that has a calculated lower limit.

As mentioned above, these limits on the $\pi_1(1600)$ photoproduction cross section can be used with the calculated $\pi_1(1600)$ decay widths from LQCD [8] to determine which channels are most sensitive to the $\pi_1(1600)$ in photoproduction. To do this, we need two pieces of information: the maximum size of the $\pi_1(1600)$ signal in the channel and the total size of the photoproduction cross section for the channel. To estimate the maximum $\pi_1(1600)$ contribution in the $\eta^{(\prime)}\pi$ channels, we assume $\mathcal{B}(b_1 \to \omega \pi) = 100\%$ and multiply the measured upper limit on $\sigma(\gamma p \to \pi_1^0(1600)p) \times \mathcal{B}(\pi_1(1600) \to b_1\pi)$ by $R = \mathcal{B}(\pi_1(1600) \to \eta^{(\prime)}\pi)/\mathcal{B}(\pi_1(1600) \to b_1\pi)$. The maximum $\pi_1(1600)$ contribution comes when R is largest, so we maximize $\Gamma(\pi_1(1600) \to \eta^{(\prime)}\pi)$ and minimize $\Gamma(\pi_1(1600) \to b_1\pi)$ under the constraints that $\Gamma_{\rm tot}(\pi_1(1600)) = 492$ MeV and that the partial widths are in the ranges allowed by Ref. [8]. The corresponding values are $R = 2.8 \times 10^{-2}$ for $\eta'\pi$ and $R = 2.3 \times 10^{-3}$ for $\eta\pi$. More details on how these values were obtained are in the Supplemental Materials [28].

Next, we compare these limits to the total size of the $\eta\pi$ and $\eta'\pi$ photoproduction cross sections. We select exclusive events for the processes $\gamma p \to \eta^{(\prime)} \pi^0 p$ and $\gamma p \to$ $\eta^{(\prime)}\pi^-\Delta^{++}$ with $\Delta^{++} \to \pi^+ p$, $\eta' \to \pi^+\pi^-\eta$, and $\eta \to \gamma\gamma$. For $\gamma p \to \eta^{(\prime)} \pi^0 p$, we use the same selection criteria as $\gamma p \to \omega \pi^+ \pi^- p$, and for $\gamma p \to \eta^{(\prime)} \pi^- \Delta^{++}$ we use the same selection criteria as $\gamma p \rightarrow \omega \pi^{-} \pi^{0} \Delta^{++}$. We select the η' by requiring $|M_{\pi^+\pi^-\eta} - m_{\eta',\text{PDG}}| < 25 \text{ MeV}/c^2$, and we subtract background contributions using η' sidebands with $|M_{\pi^+\pi^-n} - m_{\eta',\text{PDG}} \pm 60 \text{ MeV}/c^2| < 25 \text{ MeV}/c^2$. In addition, we only select beam photons in the range $8.2 \text{ GeV} < E_{\gamma} < 8.8 \text{ GeV}$, which is the beam-energy range that has the maximum photon polarization. The partialwave analyses of the $\eta^{(\prime)}\pi$ systems will include the beam polarization in the fits to determine the production mechanisms involved. The cross sections in this paper do not need the polarization information, which allows us to use a wider range of energies to increase our statistics.

The reconstructed $\eta^{(\prime)}\pi^-$ invariant mass distributions are shown in Fig. 3, and the corresponding plots for $\eta^{(\prime)}\pi^0$ are in the Supplemental Materials [28]. Note that we are showing the number of reconstructed events rather than the cross section, so we convert our $a_2(1320)$ cross section and $\pi_1(1600)$ upper limit to an expected number of events using the equation $N = \sigma \mathcal{L}\epsilon \mathcal{B}$, where σ is the cross section, \mathcal{L} is the luminosity, ϵ is the reconstruction efficiency, and \mathcal{B} is the product of all required branching fractions. The figure shows the measured size of the $a_2(1320)$ signal, as well as the projected $\pi_1(1600)$ upper limit. These projections show that the $\pi_1(1600)$ will be less than 1% of the total $\eta\pi$ mass spectra, but it could be the main contribution to the $\eta'\pi$ mass spectra.

In conclusion, we have measured the cross sections for the photoproduction of three different $\omega\pi\pi$ final states off of a proton target, and used them to set an upper limit for the photoproduction cross section of the lightest hybrid meson candidate. We find that the upper limits on the $\pi_1(1600)$ cross sections are similar in size to the $a_2(1320)$ cross section. Assuming $\mathcal{B}(b_1 \to \omega\pi) = 100\%$, we place 90% C.L. upper limits of $\sigma(\gamma p \to \pi_1^0(1600)p) \times \mathcal{B}(\pi_1(1600) \to b_1\pi) < 123$ nb and $\sigma(\gamma p \to \pi_1^-(1600)\Delta^{++}) \times \mathcal{B}(\pi_1(1600) \to b_1\pi) < 435$ nb. We combine these results with lattice QCD calculations for the $\pi_1(1600)$ decay widths and find $\sigma(\gamma p \to \pi_1^0(1600)p) < 177$ nb and $\sigma(\gamma p \to \pi_1^-(1600)\Delta^{++}) <$



FIG. 3. The measured $\eta^{(\prime)}\pi^-$ invariant mass distributions (points), overlaid with the $a_2(1320)$ signal (cyan) and the $\pi_1(1600)$ upper-limit (red).

627 nb for $0.1 \,(\text{GeV}/c)^2 < -t < 0.5 \,(\text{GeV}/c)^2$. These are the first limits on $\pi_1(1600)$ photoproduction, since $\pi_1 \to b_1 \pi$ is the only decay that has a calculated lower limit on the size of its branching fraction. We also find that the best discovery potential for the $\pi_1(1600)$ in photoproduction is in the $\gamma p \to \eta' \pi^- \Delta^{++}$ reaction.

ACKNOWLEDGEMENTS

The analysis in this article was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DOE Grant No. DE-FG02-87ER40315. We would like to acknowledge the outstanding efforts of the staff of the Accelerator and the Physics Divisions at Jefferson Lab that made the experiment possible. This work was also supported in part by the U.S. Department of Energy, the U.S. National Science Foundation, NSERC Canada, the German Research Foundation, GSI Helmholtzzentrum für Schwerionenforschung GmbH, the Russian Foundation for Basic Research, the UK Science and Technology Facilities Council, the Chilean Comisión Nacional de Investigación Científica y Tecnológica, the National Natural Science Foundation of China, and the China Scholarship Council. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR23177.

 * Corresponding author: wimoehl@andrew.cmu.edu

- J. J. Dudek, R. G. Edwards, B. Joó, M. J. Peardon, D. G. Richards, and C. E. Thomas (Hadron Spectrum Collaboration), Phys. Rev. D 83, 111502(R) (2011).
- [2] J. J. Dudek, R. G. Edwards, P. Guo, and C. E. Thomas (Hadron Spectrum Collaboration), Phys. Rev. D 88, 094505 (2013).

- [3] C. A. Meyer and E. S. Swanson, Prog. Part. Nucl. Phys. 82, 21–58 (2015).
- [4] C. A. Meyer and Y. Van Haarlem, Phys. Rev. C 82, 025208 (2010).
- [5] C. Adolph *et al.* (COMPASS Collaboration) Phys. Lett. B **740**, 303–311 (2015).Erratum: Phys. Lett. B **811**, 135913 (2020).
- [6] A. Rodas *et al.* (Joint Physics Analysis Center) Phys. Rev. Lett. **122**, 042002 (2019).
- [7] B. Kopf, M. Albrecht, H. Koch, M. Küßner, J. Pychy, X. Qin and U. Wiedner, Eur. Phys. J. C 81, 12, 1056 (2021)
- [8] A. J. Woss, J. J. Dudek, R. G. Edwards, C. E. Thomas, and D. J. Wilson (Hadron Spectrum Collaboration) Phys. Rev. D 103, 054502 (2021).
- [9] D. V. Amelin *et al.* (VES Collaboration) Phys. Atom. Nucl. **62**, 445–453 (1999).
- [10] D. V. Amelin *et al.*, Phys. Atom. Nucl. **68**, 359–371 (2005).
- [11] M. Lu *et al.* (E852 Collaboration) Phys. Rev. Lett. 94, 032002 (2005).
- [12] C. A. Baker *et al.* Phys. Lett. B **563**, 140–149 (2003).
- [13] M. Nozar *et al.* (CLAS Collaboration) Phys. Rev. Lett. 102, 102002 (2009).
- [14] R. L. Workman *et al.* (Particle Data Group) Prog. Theor. Exp. Phys. **2022**, 083C01 (2022).
- [15] A. J. Woss, C. E. Thomas, J. J. Dudek, R. G. Edwards, and D. J. Wilson (Hadron Spectrum Collaboration) Phys. Rev. D 100, 054506 (2019).
- [16] M. Atkinson *et al.* (Omega Photon Collaboration) Nucl. Phys. B **229**, 269–283 (1983).
- [17] G. R. Blackett, K. Danyo, T. Handler, M. Pisharody, and G. T. Condo arXiv:hep-ex/9708032.
- [18] M. Atkinson *et al.* (Omega Photon Collaboration) Z. Phys. C **34**, 157–162 (1987).
- [19] S. Adhikari *et al.* (GlueX Collaboration) Nucl. Instrum. Meth. A 987, 164807 (2021).
- [20] F. Barbosa, C. Hutton, A. Sitnikov, A. Somov, S. Somov, and I. Tolstukhin Nucl. Instrum. Meth. A 795, 376–380 (2015).
- [21] V. V. Berdnikov, S. V. Somov, L. Pentchev, and B. Zihlmann Instrum. Exp. Tech. 58, 25–29 (2015).
- [22] M. Dugger *et al.* Nucl. Instrum. Meth. A 867, 115–127 (2017).

- [23] L. Pentchev, F. Barbosa, V. Berdnikov, D. Butler, S. Furletov, L. Robison, and B. Zihlmann Nucl. Instrum. Meth. A 845, 281–284 (2017).
- [24] T. D. Beattie *et al.* Nucl. Instrum. Meth. A **896**, 24–42 (2018).
- [25] E. Pooser *et al.* Nucl. Instrum. Meth. A **927**, 330–342 (2019).
- [26] N. S. Jarvis *et al.* Nucl. Instrum. Meth. A **962**, 163727 (2020).
- [27] C. J. Batty, S. D. Hoath, and B. L. Roberts Nucl. Instrum. Meth. 137, 179–181 (1976).
- [28] See Supplemental Material at (link to be created by Journal) for additional information about systematic uncertainties
- [29] G. T. Condo, T. Handler, W. M. Bugg, G. R. Blackett, M. Pisharody, and K. A. Danyo Phys. Rev. D 48, 3045 (1993).
- [30] M. Albrecht, "Search for Exotic Hadrons in $\eta^{(\prime)}\pi$ at GlueX" (HADRON 2023 Genova, Italy).

Supplemental Material

(Dated: July 4, 2024)

EXAMPLE FITS TO $\pi^+\pi^-\pi^0$ DISTRIBUTIONS

Figure 1 shows example fits to the ω candidate masses in data for all three reactions. The data are fit with a fourth-order polynomial to describe the background, plus a Voigtian to describe the ω signal.

UPPER LIMIT PROJECTIONS

In the main body of the paper, we present the largest possible $\pi_1(1600)$ contributions for the $\eta^{(\prime)}\pi^-$ decay mode based on our 90% C.L. upper limits. In this section, we show details on how the maximum values for $R = \mathcal{B}(\pi_1(1600) \to \eta^{(\prime)}\pi)/\mathcal{B}(\pi_1(1600) \to b_1\pi)$ were determined, as well as the corresponding projection plots for the $\eta^{(\prime)}\pi^0$ decay modes.

To estimate the maximum $\pi_1(1600)$ yield in the $\eta^{(\prime)}\pi$ decay modes, we maximize the value

$$R = \frac{\mathcal{B}(\pi_1(1600) \to \eta^{(\prime)}\pi)}{\mathcal{B}(\pi_1(1600) \to b_1\pi)}$$

$$= \frac{\Gamma(\pi_1(1600) \to \eta^{(\prime)}\pi)}{\Gamma(\pi_1(1600) \to b_1\pi)}$$
(1)

by making $\Gamma(\pi_1(1600) \rightarrow \eta^{(\prime)}\pi)$ as large as possible and $\Gamma(\pi_1(1600) \rightarrow b_1\pi)$ as small as possible. To be consistent with our $\omega\pi\pi$ analysis, we use the same value for the $\pi_1(1600)$ total width that was used to set the upper limit, which is $\Gamma_{\rm tot}(\pi_1(1600)) = 492$ MeV. This width assumption combined with the range of decay widths calculated in Ref. [8] give the values in Table I, where $\Gamma_{\rm max}(\pi_1(1600) \rightarrow \text{others})$ is the maximum partial width for all decay modes other than the $\eta^{(\prime)}\pi$ and $b_1\pi$ decay mode being considered.

We show the projected $\pi_1(1600)$ upper limits to the $\eta^{(\prime)}\pi^0$ mass distributions in Figure 2. The trends are very similar to the $\eta^{(\prime)}\pi^-$ distributions shown in the main paper: the $\pi_1(1600)$ is expected to be less than a 1% contribution to the $\eta\pi^0$ system, while it could be the main contribution to $\eta'\pi^0$.

SYSTEMATIC UNCERTAINTIES

This section details all the systematic uncertainties included in our measurement, with the results summarized in Table II.

There are several systematic effects that could potentially bias the overall normalization of our cross section estimates. Systematic uncertainties arise due to a potential mismatch of the charged-track and photon reconstruction efficiencies in data and Monte Carlo (MC). We assign conservative systematic uncertainties of 5% for the recoil proton, and 3% (5%) for each photon that enters the forward calorimeter (barrel calorimeter). For the charged pions, we determine the systematic uncertainty based on the results of a detailed study using data from the reaction $\gamma p \to \omega p$ with $\omega \to \pi^+ \pi^- \pi^0$. In this study, the ω is either reconstructed fully exclusively or with a single missing charged pion. This allows us to estimate the efficiency for reconstructing a single-charged track in data and Monte Carlo. The ratio of MC and data efficiencies are measured in bins of pion lab momentum and polar angle θ_{π} with respect to the beam axis. For the polar angle region covered by this exclusive ω production reaction ($\theta_{\pi} \leq 12^{\circ}$), we determine the weighted average bias from these efficiency ratios. For larger polar angles, we assign conservative systematic uncertainties of 3% (5%) for each charged track with $\theta_{\pi} < 20^{\circ}$ ($\theta_{\pi} > 20^{\circ}$).

We assign a conservative systematic uncertainty of 5% for the luminosity measurement, which dominantly comes from our understanding of the pair spectrometer.

To determine the systematic uncertainty from the kinematic fit, we use data on $\gamma p \to \eta' p$ with $\eta' \to \pi^+ \pi^- \eta$ and $\eta' \to \pi^0 \pi^0 \eta$ with $\eta \to \pi^+ \pi^- \pi^0$ as control samples for the $\omega \pi^+ \pi^-$ and $\omega \pi^0 \pi^0$ data, respectively. This is done because the η' provides a narrow peak with well controlled backgrounds, and has similar kinematics to the $M(5\pi)$ region that we are interested in for our upper limits. We measure the η' cross sections as a function of χ^2/ndf in the range $5 < \chi^2/\text{ndf} < 25$ and use the difference between the minimum and maximum values as an estimate for the systematic uncertainty. The systematic uncertainty determined using this method is 6.7% for $\omega \pi^+ \pi^-$, and 6.8% for $\omega \pi^0 \pi^0$. For the $\omega \pi^- \pi^0$ system, there is no such clean control sample. To determine the systematic uncertainty in this case, we directly measure $\sigma(\gamma p \to \omega \pi^- \pi^0 \Delta^{++})$ for 1.2 GeV/ $c^2 < m_{\omega\pi\pi} < 1.6 \text{ GeV}/c^2$ as a function of χ^2/ndf in the range 5 < χ^2/ndf < 20. From the difference between the minimum and maximum values, we get a systematic uncertainty of 7.5%. Note that these are conservative estimates of the systematic uncertainty since at larger χ^2/ndf , non-exclusive events can inflate the measured cross section.

For the $\omega \pi^0 \pi^0$ analysis we require that the events have no additional reconstructed calorimeter showers to suppress the background process $\gamma p \to \omega \pi^0 \pi^0 \pi^0 p$, which has an additional π^0 . To estimate the systematic uncertainty from this requirement, we measure the cross section of $\gamma p \to \eta' p$ with $\eta' \to \pi^0 \pi^0 \eta$ and $\eta \to \pi^+ \pi^- \pi^0$ before and after applying this requirement. We find that the cross section changes by 0.7%, which we assign as a systematic uncertainty.



FIG. 1. The $\pi^+\pi^-\pi^0$ invariant mass distributions in data (black points) fit with a 4th-order polynomial (dashed line) plus a Voigtian (solid line). A clear $\omega(782)$ signal is seen in all three reactions, which is accompanied by a smaller signal for the $\eta(548)$.

TABLE I. Decay widths and ratio of branching fractions of the $\pi_1(1600)$ for $\Gamma_{tot}(\pi_1(1600)) = 492$ MeV measured by JPAC [6].

Decay	$\Gamma_{\max}(\pi_1(1600) \to \eta^{(\prime)}\pi)$	$\Gamma_{\max}(\pi_1(1600) \to \text{others})$	$\Gamma_{\min}(\pi_1(1600) \to b_1\pi)$	R
$\eta\pi$	$1 { m MeV}$	$60 { m MeV}$	$431 { m MeV}$	2.3×10^{-3}
$\eta'\pi$	$12 { m MeV}$	$49 { m MeV}$	$431 { m MeV}$	2.8×10^{-2}

TABLE II. Total systematic uncertainty for each $\omega \pi \pi$ cross section. Not included in the list is the variation of the ω fit model, which varies bin-to-bin.

Source	$\sigma(\omega\pi^+\pi^-)$	$\sigma(\omega\pi^0\pi^0)$	$\sigma(\omega\pi^-\pi^0)$
Luminosity	5.0%	5.0%	5.0%
Kinematic fit	6.7%	6.8%	7.5%
Track efficiency	13.5%	9.1%	16.0%
Photon efficiency	8.1%	24.3%	16.3%
Unused shower energy	-	0.7%	-
MC substructure	2.9%	-	0.9%
MC angles	1.5%	-	0.9%
MC - t slope	2.6%	2.4%	0.3%
Proton momentum	1.3%	2.3%	2.9%
Total	18.3%	27.5%	24.8%

The nominal signal Monte Carlo simulates $\pi_1(1600) \rightarrow \omega \pi \pi$ using a uniform distribution in phase space. To determine the systematic uncertainty due to the decay model, we produce independent MC samples that include a Breit-Wigner for $b_1(1235) \rightarrow \omega \pi$ to replicate the substructure of $\pi_1^0(1600) \rightarrow b_1^{\pm}\pi^{\mp}$ and $\pi_1^-(1600) \rightarrow b_1^-\pi^0$ or $\pi_1^-(1600) \rightarrow b_1^0\pi^-$. The largest difference in the reconstruction efficiency with respect to the nominal Monte Carlo is taken as the systematic uncertainty, which gives 2.9% for $\omega \pi^+ \pi^-$ and 0.9% for $\omega \pi^- \pi^0$.

The nominal MC simulation assumes pure S-wave decays for both $\pi_1(1600) \rightarrow b_1\pi$ and $b_1 \rightarrow \omega\pi$. The $\pi_1(1600) \rightarrow b_1(1235)\pi$ decay has limited phase space, so we expect the D-wave contribution to be kinematically suppressed. To account for the possible presence of Dwaves in $b_1 \rightarrow \omega\pi$, we determine the efficiency assuming purely D-wave angular distributions for this decay. We include this efficiency by reweighting the MC data using the measured D/S amplitude ratio for the b_1 of 0.277 from the PDG [14]. This gives a systematic uncertainty of 1.5% for $\omega \pi^+ \pi^-$ and 0.9% for $\omega \pi^- \pi^0$.

The -t distributions in data and Monte Carlo may not precisely match, which could cause a mismatch of the efficiency in data and Monte Carlo. In signal Monte Carlo, all the signal reactions are simulated with a -t slope of 5 (GeV/c)⁻². As a variation, we use the efficiencycorrected -t distribution obtained from data to reweight our Monte Carlo data. This gives systematic uncertainties of 2.6% for $\omega \pi^+ \pi^-$, 2.4% for $\omega \pi^0 \pi^0$, and 0.3% for $\omega \pi^- \pi^0$.

Previous studies have shown the efficiencies for lowmomentum protons can differ in Monte Carlo and data. To study this effect, we directly measure how much the $\omega\pi\pi$ cross section for the range $1.2 \text{ GeV}/c^2 < M(\omega\pi\pi) < 1.6 \text{ GeV}/c^2$ changes when we require a minimum proton momentum that is 50 MeV/c larger than in our nominal selection. We get systematic uncertainties of 1.3%, 2.3%, and 2.9% for $\omega\pi^+\pi^-$, $\omega\pi^0\pi^0$, and $\omega\pi^-\pi^0$, respectively.

Another source of uncertainty is the fit model used to extract the ω yield. To account for this, we vary the ω signal shape from a Voigtian to shapes that are determined from Monte Carlo data and either include all possible $M(\pi^+\pi^-\pi^0)$ combinations or include only the $\pi^+\pi^-\pi^0$ combination coming from the generated ω decay. We also vary the background polynomial from order 4 to order 3 or 5. These variations are done independently, so in total there are 9 fit variations. When determining our upper limit on the $\pi_1(1600)$ cross sections, we use the fit variation that gives the largest upper limit.



FIG. 2. The measured $\eta^{(\prime)}\pi^0$ invariant mass distributions (points), overlaid with the $a_2(1320)$ signal (cyan) and the $\pi_1(1600)$ upper-limit (red).

The parameterization of the $a_2(1320)$ and $\pi_1(1600)$ are a potentially large source of uncertainty in the analysis. For the nominal shapes, we model the $a_2(1320)$ using a Breit-Wigner based on the PDG parameters [14], and the $\pi_1(1600)$ using a Breit-Wigner based on the JPAC parameters [6]. The $a_2^0(1320)$ yield is fixed based on the cross section measured in the $\eta \pi^0$ PWA and the value $\mathcal{B}(a_2(1320) \to \omega \pi \pi) = (10.6 \pm 3.2)\%$ from the PDG [14]. To account for the large uncertainty in the $a_2(1320)$ branching fraction, we also repeat the fits after varying the branching fraction up and down by its uncertainty. The fit shown in Fig. 3 of the paper uses the nominal $a_2(1320)$ branching fraction, while the $\pi_1(1600)$ upper limit is determined from a fit where the $a_2(1320)$ branching fraction is lowered by its uncertainty from the PDG, since that gives a larger upper limit. Additionally, we can vary the $\pi_1(1600)$ shape to use the PDG parameters, but since this has a smaller width, it results in a smaller upper limit.

The final source of systematic uncertainty comes from the possible interference between the decays $\pi_1^0(1600) \rightarrow$ $b_1^+\pi^-$ and $\pi_1^0(1600) \to b_1^-\pi^+$ or $\pi_1^-(1600) \to b_1^0\pi^-$ and $\pi_1^-(1600) \rightarrow b_1^- \pi^0$. To determine the size of the interference effect, we simulate these decay modes assuming totally coherent or incoherent amplitudes. Based on the isospin Clebsch-Gordan coefficients, we expect the two decay modes to destructively interfere. By determining the ratio of MC events for totally destructive interference over the incoherent case, we find that the signal could be reduced by destructive interference by up to 26% for the $\pi_1^0(1600)$ signal and 29% for the $\pi_1^-(1600)$ signal in the mass range 1.2 GeV/ $c^2 < m_{\omega\pi\pi} < 2.2$ GeV/ c^2 . The difference in the amount of interference is caused by the different amount of phase space available due to the different recoil baryons. To recover the true cross section before this $\pi_1(1600) \rightarrow b_1 \pi$ interference, we scale the likelihood function by 1.0/0.74 for the $\pi_1^0(1600)$ and by 1.0/0.71 for the $\pi_1^-(1600)$.

The total systematic uncertainties are determined by

adding the uncertainties from each source in quadrature. This gives a total systematic uncertainty of 18.3% for $\sigma(\omega\pi^+\pi^-)$, 27.5% for $\sigma(\omega\pi^0\pi^0)$, and 24.8% for $\sigma(\omega\pi^-\pi^0)$.

Three of our largest systematic uncertainties are the photon reconstruction efficiency, charged-track reconstruction efficiency, and luminosity calculation systematic uncertainties. These systematic uncertainties are correlated for each source across the three $\omega \pi \pi$ cross section measurements. To account for these large correlated uncertainties in our $\pi_1(1600)$ upper limits, we rescale our cross sections according to the size of these uncertainties. This is done both for the neutral and charged $\sigma(\omega \pi \pi)_{I=1}$. Our nominal fits shown in the paper use the cross sections before this scaling procedure, but the upper limit is determined after scaling the cross sections such that the isospin-1 cross sections give the largest $\pi_1(1600)$ upper limits. For the $\pi_1^-(1600)$, the remaining systematic uncertainties are included in the upper limit by convolving the likelihood distribution by a Gaussian whose width matches the systematic uncertainty.

To determine the neutral isospin-1 $\omega \pi \pi$ cross section, we have to take the difference of two cross sections:

$$\sigma(\omega\pi\pi)_{I=1} = \sigma(\omega\pi^+\pi^-) - 2\sigma(\omega\pi^0\pi^0)$$
(2)

The uncertainty on a difference of cross sections is given by

$$\delta_Z^2 = \left(\frac{\partial Z}{\partial A}\right)^2 \delta_A^2 + \left(\frac{\partial Z}{\partial B}\right)^2 \delta_B^2 + 2\rho_{AB} \frac{\partial Z}{\partial A} \frac{\partial Z}{\partial B} \delta_A \delta_B. \tag{3}$$

for absolute uncertainties δ , $A = \sigma(\omega \pi^+ \pi^-)$, $B = \sigma(\omega \pi^0 \pi^0)$, and $Z = A - 2B = \sigma(\omega \pi \pi)_{I=1}$. As mentioned above, the charged tracking efficiency, photon efficiency, luminosity, and kinematic fit uncertainties are correlated across the three measurements. For these systematics, we use $\rho_{AB} = 1$. For all other systematic uncertainties, we assume the sources are uncorrelated ($\rho_{AB} = 0$). For the case of correlated uncertainties, we find

$$\delta_Z = |2\delta_B - \delta_A| \tag{4}$$

while for uncorrelated uncertainties, we have

$$\delta_Z = \sqrt{\delta_A^2 + 4\delta_B^2} \tag{5}$$

Note the total uncertainty depends on the absolute size of the uncertainties, not the relative size. This means we cannot assign a global systematic uncertainty for the neutral $\sigma(\omega\pi\pi)_{I=1}$, but instead must calculate the systematic uncertainty for each mass bin. In addition, since there is no well defined global systematic uncertainty, we cannot account for the systematic uncertainties by convolving the likelihood distribution by a Gaussian in this case. Instead, we determine for each $M(\omega\pi\pi)$ bin the total systematic uncertainty to the nominal value in this bin. The effect is that the likelihood distribution is shifted instead of broadened by the systematic uncertainties.

- J. J. Dudek, R. G. Edwards, B. Joó, M. J. Peardon, D. G. Richards, and C. E. Thomas (Hadron Spectrum Collaboration), Phys. Rev. D 83, 111502(R) (2011).
- [2] J. J. Dudek, R. G. Edwards, P. Guo, and C. E. Thomas (Hadron Spectrum Collaboration), Phys. Rev. D 88, 094505 (2013).
- [3] C. A. Meyer and E. S. Swanson, Prog. Part. Nucl. Phys. 82, 21–58 (2015).
- [4] C. A. Meyer and Y. Van Haarlem, Phys. Rev. C 82, 025208 (2010).
- [5] C. Adolph *et al.* (COMPASS Collaboration) Phys. Lett. B **740**, 303–311 (2015).Erratum: Phys. Lett. B **811**, 135913 (2020).
- [6] A. Rodas et al. (Joint Physics Analysis Center) Phys. Rev. Lett. 122, 042002 (2019).
- [7] B. Kopf, M. Albrecht, H. Koch, M. Küßner, J. Pychy, X. Qin and U. Wiedner, Eur. Phys. J. C 81, 12, 1056 (2021)
- [8] A. J. Woss, J. J. Dudek, R. G. Edwards, C. E. Thomas,

and D. J. Wilson (Hadron Spectrum Collaboration) Phys. Rev. D **103**, 054502 (2021).

- [9] D. V. Amelin *et al.* (VES Collaboration) Phys. Atom. Nucl. **62**, 445–453 (1999).
- [10] D. V. Amelin *et al.*, Phys. Atom. Nucl. **68**, 359–371 (2005).
- [11] M. Lu *et al.* (E852 Collaboration) Phys. Rev. Lett. **94**, 032002 (2005).
- [12] C. A. Baker et al. Phys. Lett. B 563, 140-149 (2003).
- [13] M. Nozar *et al.* (CLAS Collaboration) Phys. Rev. Lett. 102, 102002 (2009).
- [14] R. L. Workman *et al.* (Particle Data Group) Prog. Theor. Exp. Phys. **2022**, 083C01 (2022).
- [15] A. J. Woss, C. E. Thomas, J. J. Dudek, R. G. Edwards, and D. J. Wilson (Hadron Spectrum Collaboration) Phys. Rev. D 100, 054506 (2019).
- [16] M. Atkinson *et al.* (Omega Photon Collaboration) Nucl. Phys. B **229**, 269–283 (1983).
- [17] G. R. Blackett, K. Danyo, T. Handler, M. Pisharody, and G. T. Condo arXiv:hep-ex/9708032.
- [18] M. Atkinson *et al.* (Omega Photon Collaboration) Z. Phys. C **34**, 157–162 (1987).
- [19] S. Adhikari *et al.* (GlueX Collaboration) Nucl. Instrum. Meth. A **987**, 164807 (2021).
- [20] F. Barbosa, C. Hutton, A. Sitnikov, A. Somov, S. Somov, and I. Tolstukhin Nucl. Instrum. Meth. A 795, 376–380 (2015).
- [21] V. V. Berdnikov, S. V. Somov, L. Pentchev, and B. Zihlmann Instrum. Exp. Tech. 58, 25–29 (2015).
- [22] M. Dugger *et al.* Nucl. Instrum. Meth. A 867, 115–127 (2017).
- [23] L. Pentchev, F. Barbosa, V. Berdnikov, D. Butler, S. Furletov, L. Robison, and B. Zihlmann Nucl. Instrum. Meth. A 845, 281–284 (2017).
- [24] T. D. Beattie *et al.* Nucl. Instrum. Meth. A **896**, 24–42 (2018).
- [25] E. Pooser *et al.* Nucl. Instrum. Meth. A **927**, 330–342 (2019).
- [26] N. S. Jarvis *et al.* Nucl. Instrum. Meth. A **962**, 163727 (2020).
- [27] C. J. Batty, S. D. Hoath, and B. L. Roberts Nucl. Instrum. Meth. 137, 179–181 (1976).
- [28] See Supplemental Material at (link to be created by Journal) for additional information about systematic uncertainties
- [29] G. T. Condo, T. Handler, W. M. Bugg, G. R. Blackett, M. Pisharody, and K. A. Danyo Phys. Rev. D 48, 3045 (1993).
- [30] M. Albrecht, "Search for Exotic Hadrons in $\eta^{(\prime)}\pi$ at GlueX" (HADRON 2023 Genova, Italy).