

Efficient Stochastic Template Bank using Inner Product Inequalities

KEISI KACANJA ¹, ALEXANDER H. NITZ ¹, SHICHAO WU ², MARCO CUSINATO ³, RAHUL DHURKUNDE ²,
IAN HARRY ⁴, TITO DAL CANTON ⁵ AND FRANCESCO PANNARALE ^{6,7}

¹*Department of Physics, Syracuse University, Syracuse, NY 13244, USA*

²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany and Leibniz Universität Hannover, D-30167 Hannover, Germany*

³*Departament d'Astronomia i Astrofísica, Universitat de València, Edifici d'Investigació Jeroni Munyoz, C/ Dr. Moliner, 50, E-46100 Burjassot (València), Spain*

⁴*University of Portsmouth, Institute of Cosmology and Gravitation, Portsmouth PO1 3FX, United Kingdom*

⁵*Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*

⁶*Sapienza - Università di Roma, I-00185 Roma, Italy*

⁷*INFN, Sezione di Roma, I-00185 Roma, Italy*

ABSTRACT

Gravitational wave searches are crucial for studying compact sources like neutron stars and black holes. Many sensitive modeled searches use matched filtering to compare gravitational strain data to a set of waveform models known as template banks. We introduce a new stochastic placement method for constructing template banks, offering efficiency and flexibility to handle arbitrary parameter spaces, including orbital eccentricity, tidal deformability, and other extrinsic parameters. This method can be computationally limited by the ability to compare proposal templates with the accepted templates in the bank. To alleviate this computational load, we introduce the use of inner product inequalities to reduce the number of required comparisons. We also introduce a novel application of Gaussian Kernel Density Estimation to enhance waveform coverage in sparser regions. Our approach has been employed to search for eccentric binary neutron stars, low-mass neutron stars, primordial black holes, supermassive black hole binaries. We demonstrate that our method produces self-consistent banks that recover the required minimum fraction of signals. For common parameter spaces, our method shows comparable computational performance and similar template bank sizes to geometric placement methods and stochastic methods, while easily extending to higher-dimensional problems. The time to run a search exceeds the time to generate the bank by a factor of $\mathcal{O}(10^5)$ for dedicated template banks, such as geometric, mass-only stochastic, and aligned spin cases, $\mathcal{O}(10^4)$ for eccentric and $\mathcal{O}(10^3)$ for the tidal deformable bank. With the advent of efficient template bank generation, the primary area for improvement is developing more efficient search methodologies.

1. INTRODUCTION

Gravitational waves (GWs) are distortions in spacetime. One mechanism for generating GWs is accelerating two massive compact objects such as neutron stars and black holes (Einstein 1916, 1918). When these compact objects form a binary system, their mutual gravitational attraction causes them to spiral towards each other, producing gravitational waves that travel outward at the speed of light. We observe these GWs with ground-based detectors such as the Advanced Laser

Interferometer Gravitational-wave Observatory (LIGO) and Advanced Virgo (Acernese et al. 2015; LIGO Scientific Collaboration et al. 2015). Since the commencement of the advanced detector era in 2015, Advanced LIGO and Virgo have detected $\mathcal{O}(100)$ gravitational wave sources (Abbott et al. 2019; Abbott et al. 2024, 2023; Nitz et al. 2023; Mehta et al. 2023; Olsen et al. 2022)

The initial identification of gravitational-wave sources involves conducting searches (Usman et al. 2016; Cannon et al. 2021; Aubin et al. 2021; Messick et al. 2017; Pang et al. 2020; Chu et al. 2020). Model based searches typically employ matched filtering which compares the detector data with modeled GW signals (Allen et al.

2012; Babak et al. 2013). Since the properties of a potential source are unknown, a discrete set of template waveforms, known as template bank, is used.

Template banks are carefully constructed to ensure minimal loss in the signal-to-noise ratio (SNR) (Allen et al. 2012; Sathyaprakash & Dhurandhar 1991a; Owen 1996; Owen & Sathyaprakash 1999; Balasubramanian et al. 1996; Harry et al. 2009; Brown et al. 2012; Dal Canton et al. 2014; Roy et al. 2019; Hanna et al. 2023). An ideal template bank contains at least one sufficiently similar template waveform to identify any potential signal within a designated parameter space. A valid template bank algorithm strategically fills the parameter space with templates to limit loss in SNR, and minimizes redundant templates. Additionally, it should adapt to the specific characteristics of the gravitational-wave source population studied, allowing the ability to analyze data for a diverse range of astrophysical scenarios.

Various techniques exist for constructing a template bank. The two main classes of methods are geometric (Brown et al. 2012; Babak et al. 2006; Schmidt et al. 2024), stochastic placement techniques (Harry et al. 2009; Ajith et al. 2014; Manca & Vallisneri 2010; Babak et al. 2006). There also exist hybrid methods that utilizes both approaches. (Roy et al. 2017; Roy et al. 2017). Geometric methods typically place template waveforms using a lattice technique; this generally requires a known metric, which directly quantifies the similarity of signals. Afterwards, the templates are converted into a physical parameter space. This approach guarantees that every template captures a predefined minimum percentage of the signal (Owen 1996; Prix 2007). One challenge is that the metric may only be flat in the non-physical space. This may result in over-placing templates near and outside the physical space boundaries. These templates need to be included since they may be better at recovering signals near the boundary than the templates within the physical space (Cokelaer 2007; Coogan et al. 2022; Brown et al. 2012). Consequently, since a search’s computational cost is proportional to the number of templates produced from any method, including these templates can lead to an excess in the template count, making the search more expensive to perform. This approach is not optimal for complicated boundaries, unknown metrics, and for more intricate detection studies requiring non-trivial parameter spaces.

An alternative approach is a stochastic placement technique. This method can work without an explicit metric and does not attempt to construct a lattice to place templates. Instead, templates are placed directly in the physical space and iteratively added to the bank

until the parameter space is sufficiently well covered. There are several approaches for stochastic template proposals and how templates are evaluated for inclusion in a bank. Stochastic methods typically propose templates either randomly or according to a probability distribution (Privitera et al. 2014; Indik et al. 2017). Various template acceptance strategies exist, such as rejecting templates that retrieve more than a predetermined percentage of the SNR. Alternatively, some methods gauge the density of templates; if the volume of templates surpasses a specified threshold, the template bank is deemed adequately covered (Harry et al. 2009). Unlike geometric methods, stochastic methods provide less stringent guarantees on the completeness of the template bank, sometimes resulting in banks with small holes with lower accuracy of SNR reconstruction. However, stochastic methods in general offer a higher degree of flexibility than geometric methods, and allows the ability to perform searches for many variety of circumstances such as eccentric or deformable sources.

In this paper, we present a new stochastic placement method. We show this method produces self-consistent banks that recover the SNR of any potential signal within a target search space, with losses below a chosen threshold. We show that this method has already been used to conduct various searches such as (low-mass and eccentric) neutron stars, (supermassive and primordial) black holes, and neutron-star–black-hole binaries that produce gamma-ray bursts. Finally, we show how the number of parameters in a bank scales the time to complete a search by generating different banks with varying intrinsic parameters such as spin, eccentricity, and or tidal deformability.

2. EVALUATION OF TEMPLATE BANK COVERAGE

We assess the effectiveness of a template bank by determining the fraction of SNR its best matching template can recover for a given source, known as the fitting factor (FF) (Apostolatos 1995). To quantify the similarity between template waveforms, and the ability of one GW waveform to recover the SNR of another, we define the overlap between waveforms to be

$$\mathcal{O}(h_0, h_1) = \frac{(h_0|h_1)}{\sqrt{(h_1|h_1)(h_0|h_0)}} \quad (1)$$

The noise-weighted inner product $(h_0|h_1)$ is the fraction of signal power extracted from a modeled signal h_1 with the waveform model h_0 , defined as

$$(h_0|h_1) = 4 \operatorname{Re} \int_0^\infty \frac{\tilde{h}_0(f)\tilde{h}_1^*(f)}{S_n(f)} df \quad (2)$$

Where $S_n(f)$ is the power spectral density and f is frequency. Since an overall phase and the absolute time of arrival are typically nuisance parameters in a search, we define the phase and time maximized overlap to be the match \mathcal{M} . This formalism is for a non-precessing signal. For precessing signals, refer to [Harry et al. \(2016\)](#).

$$\mathcal{M}(h_0, h_1) = \max_{\phi_c, t_c} (\mathcal{O}(h_0, h_1(\phi_c, t_c))) \quad (3)$$

The match measures how well the observed waveform correlates with the expected waveform for a given the detector sensitivity. The maximum match for a potential signal h_i with all the templates in a bank h_{tb} is the fitting factor. The same waveform approximant is used for both templates when testing the coverage of the bank.

$$FF(h_i) = \max_{h \in \{h_{tb}\}} \mathcal{M}(h_i, h) \quad (4)$$

The FF represents the fraction of the SNR that is recovered by the template bank for a given signal. For instance, consider a template bank constructed with a minimal match of 0.95. This bank is expected to recover signals with FFs of 0.95, such that up to 5% of a signal's SNR or a maximum of 14% of the total number of signals may be sacrificed. FFs less than 0.95 indicate regions within the bank where templates cannot fully recover the SNR of a reference signal, suggesting potential gaps in coverage. FFs at 0.95 and above signify that the bank is adequately populated with templates, capable of recovering at least 95% of the SNR.

3. METHODS: STOCHASTIC TEMPLATE BANK ALGORITHM

Our method uses a stochastic approach to place templates directly in the physical space and enables the generation of templates that covers a wide variety of parameter spaces includes those parameterized by mass, spin, tidal deformability, and eccentricity. If templates include detector responses, this method can also cover those extrinsic parameters. A diagram of the algorithm is illustrated in [Figure 1](#), where it is divided into two main processes: the mechanism which proposes templates (discussed in [section 3.1](#)), and the optimized procedure which determines whether a proposed template should be included into the bank (discussed in [section 3.2](#)). Overall, this method proposes templates and determines whether the inclusion of the templates sufficiently cover the parameter space to recover any potential signal and will repeat until the desired coverage is achieved.

The bank is sufficiently covered under two termination criteria: minimal match and tolerance. Toler-

ance is the fraction of accepted templates over the total amount of templates in the bank. This criterion determines how many templates must be drawn to sufficiently cover the bank. Lower tolerance values generate banks with more templates to cover the parameter space adequately; larger tolerance values do the opposite. The minimal match is the minimum percentage of SNR that at least one template in the bank can recover any fiducial signal. We use this criterion to accept or reject proposal templates. If the match between the proposal and the accepted templates is less than the minimal match, that proposed template will be added to the bank. If the template has a match greater than the minimal match condition, sufficient templates exist in the bank to recover any given signal in that region.

3.1. Template Sampling Algorithm

In this section, we detail the template sampling method, the first stage of our stochastic template bank method. We first initialize the method with a set of user-defined parameters such as properties of the source, the lower frequency cutoff, and the PSD model. The sample templates are drawn within the bounds of chirp time. Chirp time (τ_0), known as the zeroth order signal duration in time, defines the regions of template placement ([Babak et al. 2006](#)). We utilize τ_0 bins for sampling nearby templates, as significant differences in chirp time between signals would lead to higher mismatches. Moreover, stochastic methods struggle to parallelize the physical space sampling over multiple cores since splitting up the parameter space can be challenging ([Coogan et al. 2022](#)). Utilizing τ_0 boundaries facilitates faster template bank generation by parallelizing different τ_0 bins along different cores. τ_0 is defined as follows

$$\tau_0 = \frac{a_0(f_{\text{lower}})^{5/3}}{\mathcal{M}_{\text{chirp}}} \quad (5)$$

where a_0 is

$$a_0(f_{\text{lower}}) = \frac{5}{256(\pi f_{\text{lower}})^{8/3}} \quad (6)$$

and $\mathcal{M}_{\text{chirp}}$ is defined as

$$\mathcal{M}_{\text{chirp}} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} \quad (7)$$

where m_1 and m_2 is the mass of primary and secondary respectively.

To define the τ_0 boundaries, we calculate the chirp time defined above using a default lower chirp time frequency of 15Hz for the minimum and maximum mass range corresponding to the start and end τ_0 . The start τ_0 will correspond to the highest masses in the space




Figure 1. Stochastic Bank Algorithm Diagram. Points are randomly sampled from the prior distribution and within distinct bins defined by chirp time τ_0 . Waveforms are then generated with a reduced frequency model. When a template is proposed, we assess if including the proposal in the template bank will improve the coverage of the desired parameter space. This assessment is outlined in the blue box and involves comparing the proposed template with the nearest neighboring templates in chirp time. Templates are added to the bank if match between the proposed template and the accepted templates fall below the minimal match condition. If the match exceeds the minimal match condition, the bank is deemed adequately covered in that region. Match calculations are stored to streamline the process and avoid redundant computations for new proposed templates, utilizing a triangle inequality for efficient comparisons. Upon completing comparisons and reaching the minimal match threshold, tolerance checks, outlined in red, ensure the bank's coverage adequacy. If the tolerance condition is unmet, additional samples are drawn either stochastically from the prior distribution or with a Gaussian Kernel Density Estimation (KDE) method. The methods are swapped if one technique performs better than the other depicted in the beige box. Proposals are once again compared with the accepted templates. Once the tolerance condition is satisfied, accepted templates are saved, and the process iterates for subsequent bins until the entire bank is sufficiently covered. The red arrows represent the checks and repetition in method required to fulfill either tolerance or minimal match conditions.




Figure 2. The evolution of the template bank as proposed templates are added within bounded τ_0 bins. Once the first τ_0 region is sufficiently filled, the algorithm proceeds to the next region, which is overlapped with the previous bin by 50%. Points are proposed until the whole bank is sufficiently filled. The orange points correspond to the proposed points. The small colored points correspond with the accepted templates. The color bar represents the τ_0 bins explored by the method. Bank generation can be parallelized by generating it across different τ_0 bins.

and the end will correspond to the lowest mass pair. The crawl parameter determines the width of each bin. Once the τ_0 boundaries are defined, we draw the first set of points from the prior distribution of the parameters within the first τ_0 bin. The sampled templates are either accepted or rejected into the bank (elaborated in section 3.2). Sampling all the templates randomly can result in banks with a lot of holes, or lacking templates. To combat some of the gaps in the bank, we utilize a Gaussian Kernel Density Estimate (KDE). Once sufficient templates accumulate in the bank, we generate a probability density function (PDF) from the accepted proposals. From the PDF, we can identify poor regions where there are divets or tails corresponding to lack of templates and propose templates to aid in sufficiently covering the sparser regions. Another benefit for utilizing two different sampling methods is the ability to speed up proposals when one method performs worse than the other. When the fraction of accepted templates over all proposed templates, the tolerance, is worse for either sampling method, the technique will swap to drawing proposals using the alternate technique. Once the first τ_0 bin has been sufficiently filled with templates and the tolerance criteria is achieved, we begin sampling in the next τ_0 bin overlapping 50% with the previous strip. Figure 2 shows how the τ_0 strips overlap with each bin as the sampler proceeds to cover the entire parameter space.

3.2. Optimizing Template Proposal Acceptance

Once a template is proposed, we must determine if it should be added to the template bank to enhance its coverage of the target parameter space. This is done by determining the FF of our proposed template against the current set of templates in the bank to assess the similarity. Templates with FFs less than the minimal match condition are not sufficiently similar and need to be added to the bank. Templates exceeding the minimal match condition are similar and do not need to be included in the bank. To start, we generate a reduced frequency waveform model, inspired by the methodology detailed in [Ajith et al. \(2014\)](#); [Capano et al. \(2016\)](#). This model significantly reduces the computational load required for generating waveforms for the match calculations. Next, we evaluate the matches between the proposed waveform models and the closest templates in chirp time, storing these values. To rule out templates where the match is clearly above the minimal match criterion, we make use of previously stored matches. We further optimize this procedure using triangle inequality which states that the sum of any two sides of any triangle will be greater than the third side. If we compare

the distances between templates, we will automatically know that the third distance between the templates will be smaller. For example, consider the mismatch ($1 - \mathcal{M}$), which represents the distance of the templates, between waveform A and waveform B, and between waveform B and waveform C. If the sum of mismatches between A and B, B and C is smaller than the predefined 1-FF threshold, we can skip the direct calculation of the match between A and C.

$$[1-FF] \geq [1-\mathcal{M}(A, B)] + [1-\mathcal{M}(B, C)] \geq [1-\mathcal{M}(A, C)] \quad (8)$$

We iteratively compare the matches between templates until all waveforms are checked or a template exceeding the minimal match condition is found. We then return to the sampling procedure in section 3.1 to conduct tolerance checks. If these checks are unmet, we repeat sampling more potential templates until the τ_0 bin is sufficiently covered. If the checks are met, proceed to the next bin or finish executing if the final bin has been filled.

4. BANK VERIFICATION

To demonstrate that our method produces valid template banks, we choose a fiducial parameter space and use our method to generate template banks. We perform the FF calculations described in section 2 and verify the stochastic method generates self-consistent banks. We generated five separate template banks of $\mathcal{O}(10^4)$ points with fixed mass and spin range for all the banks. Masses are fixed from 2 to $10 M_{\odot}$. Spins are fixed to be between -0.2 and 0.2. We also fixed the approximant to be IMRPhenomD, the lower frequency for the waveform to be 20Hz, and the PSD to be the Advanced LIGO final design sensitivity (Buikema et al. 2020; Husa et al. 2016; Khan et al. 2016a; Husa et al. 2016; Khan et al. 2016b). To showcase the effects of tolerance choices on the bank, we generate three banks with varying tolerances, 0.05, 0.01, and 0.005, for a fixed minimal match of 0.95. We also showcase the effects of minimal match by generating the last two banks with varying minimal matches of 0.8, 0.9, and 0.95 reused from the previous bank, with a fixed tolerance of 0.01. We compare our method to another stochastic method (Ajith et al. 2014; Capano et al. 2016) by generating a bank with same parameter space, minimal match of 0.95 and a convergence value of 1000. From each of these banks, we calculated the match of the templates with random fiducial waveforms for the same approximant and parameter range defined above. We successfully reconstruct a template bank that agrees with the tolerance and minimal match conditions. Figure 3 shows




Figure 3. Cumulative distribution of fitting factors (FF) achieved by the template banks with varying minimal match (top) and varying tolerance (bottom). The minimal matches are fixed to be 0.8, 0.9 and 0.95 in the colored vertical lines and a fixed tolerance of 0.01 in the red horizontal. The tolerances for the bottom plot vary from 0.05, 0.01, and 0.005 in colored vertical dashed lines and a fixed minimal match of 0.95 in a red horizontal line. All plots achieve tolerance at the minimal match condition or before minimal match. We compare how our method performs with an existing stochastic method (Ajith et al. 2014; Capano et al. 2016) by choosing the same parameter space, minimal match of 0.95 and a convergence criteria of 1000. This bank is comparable to our method generated with a tolerance of 0.01.

the cumulative distribution of the recovered FF values for all six banks. All five banks generated with our method, are able to recover the fiducial signals before the minimal match and tolerance is met.

Table 1. The computational time to generate template banks and conducting a search for different parameter spaces. The bolded parameters corresponds to template banks generated using our method. The first three banks share the same mass-only parameter space but differ in the methods used: the geometric bank (Brown et al. 2012), an existing stochastic method (Ajith et al. 2014; Capano et al. 2016), and our method respectively. The last three rows use our method to construct template banks with spin, eccentricity and tidal deformability parameters. For searches, we assume one core processes 5000 templates in real time and estimate the time required to analyze one year of data from three detectors. The time it takes to complete a search is a factor of $\mathcal{O}(10^5)$ more than the time to complete the bank generation for the non-spinning and only spin aligned case, $\mathcal{O}(10^4)$ for the eccentric bank, and $\mathcal{O}(10^3)$ for the tidal deformable case. The non-spinning geometric technique generates banks faster than the stochastic methods for mass only banks. However, the time ratio is the largest for geometric method implicating that currently template bank generations are a negligible cost to running a search, and faster search methods should be investigated. For larger parameter spaces, the banks generated in around 30 to 60 core days. The fraction of search time over bank generation time was the lowest for these scenarios. Overall, the most time consuming process is the search. The cost of generating the banks are trivial to the cost of running the search and faster search methodologies should be developed.

Parameters	Bank Size (templates)	CPU Time To Generate Bank (core days)	CPU Time to Complete Search (core days)	Search over Bank Time Ratio	Waveform Approximant
m_1, m_2 (geometric)	5.81×10^4	0.014	1.27×10^4	9.09×10^5	Does not apply
m_1, m_2 (sbank)	5.84×10^4	0.099	1.28×10^4	1.29×10^5	TaylorF2
m_1, m_2	5.29×10^4	0.033	1.16×10^4	3.52×10^5	TaylorF2
m_1, m_2, χ_1, χ_2	4.42×10^5	0.202	9.68×10^5	4.79×10^5	TaylorF2
$m_1, m_2, \chi_1, \chi_2, e$	5.67×10^6	32.23	1.24×10^6	3.85×10^4	TaylorF2ecc
$m_1, m_2, \chi_1, \chi_2, \lambda_1, \lambda_2$	1.86×10^6	55.697	4.08×10^5	7.32×10^3	TaylorF2

5. COMPUTATIONAL SCALING

The cost of a search linearly scales with the size of the template bank. We conduct an analysis to test how the time to generate different banks compares to the time for completing the bank’s respective search. To explain the computational costs associated for searching different astrophysical populations, we conducted an analysis on six distinct banks, 4 characterized by different parameter (mass, spin, eccentricity, and deformability) generated using our method, one bank using a non-spinning geometric method (Brown et al. 2012) to compare how a geometric bank search will perform, and one mass only bank generated using an existing stochastic method (Ajith et al. 2014; Capano et al. 2016). All banks were standardized with fixed equal component masses spanning $[1, 10] M_\odot$. Additionally, we maintained a consistent minimal match threshold of 0.95 across all banks and a tolerance of 0.01 for the stochastic banks. The lower frequency for the templates was uniformly set to 20Hz, and the upper frequency was 1000Hz for each bank, and parameters such as τ_0 start, crawl, and end remained constant across the stochastic banks. For the other stochastic method, we choose the convergence criteria to be 1000 since that was the most comparable to our banks with a tolerance of 0.01 (showcased in Figure 3). For aligned spins, each of the component spins was constrained to $[-0.5, 0.5]$. For banks accounting for orbital eccentricity (e), e was set to vary from $[0,$

$0.2]$ for a reference frequency of 20 Hz, using the same spin range defined above. For the deformable bank, we set the tidal deformability λ to be $[0, 5000]$, also using the same aligned spin parameters. Except for the three banks only considering mass, we parallelized three of the banks across six distinct τ_0 bins to speed up the bank generation in wall clock time.

As we increase the number of parameters, we see the banks take a longer time to generate and the number of templates increase by roughly $\mathcal{O}(10)$ for every additional pair of parameter incorporated. Explicitly looking at the TaylorF2 waveform banks, mass only had $\mathcal{O}(10^4)$ and adding a pair of spin parameters drove the bank size up to $\mathcal{O}(10^5)$ and the tidal deformability banks was of size $\mathcal{O}(10^6)$. The eccentric bank and the tidal deformability bank took the longest to generate. The eccentric bank was faster by 23.46 core days, but had 3.04 times more templates. One reason for this difference in bank size could be due to the choice of priors. If we increase the priors to a larger range, there are more options of templates proposals and more templates are necessary to sufficiently cover the parameter space. However, the excess number of templates does not seem to limit the computational speed in the eccentric bank. One explanation for this difference can be due to the use of different waveform approximants. The eccentric waveform might have been faster in producing waveforms such that

the template comparisons are not limited by the approximant generation cost.

Given a fiducial search analyzing a year of data for a three-detector configuration, with each core capable of processing 5000 templates in real-time, we estimate and compare the time required for generating the template bank and running the search (Nitz et al. 2018). We present these results of the CPU time scaling for all banks in table 1.

We estimate the computational cost for running a non-spinning search is $\mathcal{O}(10^5)$ times greater than the bank generation costs, indicating that the speed of a template bank is negligible. We also find that the geometric bank had more templates and generated 2.3 times faster than our stochastic method. However, the search over time ratio is 2.58 times higher for geometric methods than our stochastic method depicting that faster algorithms are not beneficial towards running the search and the excess number of templates negatively effects the speed of the search. The eccentric and deformable template banks took the longest to complete at around 55 core days. In comparison to the search, the fraction of search time over bank time is $\mathcal{O}(10^4)$ and $\mathcal{O}(10^3)$ less despite having more templates. Since these banks are parallelizable, the cost of generating a bank can always be reduced in wall clock time. Overall, we find that template bank generation is already a negligible computational component of a search, even for high dimensional spaces with millions of templates. Currently, efforts to generate fast template placement algorithms are less necessary than developing more efficient search methodologies, especially for multi-parameter searches.

6. EXAMPLES

In this section, we present various applications of our stochastic method in different astrophysical searches. The stochastic method has been instrumental in constructing template banks for several types of gravitational-wave searches by utilizing the varying freedom of choice in the parameter space. Below, we discuss specific examples where this method has been successfully implemented.

6.1. Primordial Black Hole and Sub-Solar Mass Binary Neutron Star

Detecting low-mass black holes will verify the existence of Primordial Black Holes (PBHs) and provide insights on dark matter distributions (Green 2024; Framp-ton 2016; Miller et al. 2024). Due to their theorized dynamical formation, searching for these systems require additional parameters to account for the eccentric orbits of PBHs (Domènech & Sasaki 2024). Nitz & Wang

(2021a) performed a search for PBH systems using the stochastic method outlined in this paper to generate an eccentric template bank. The parameter space included eccentricity e up $[0,0.3]$ with a reference frequency of 10 Hz, primary masses ranged from $[0.1,7] M_\odot$, secondary masses ranged from $[0.1,1] M_\odot$ and used TaylorF2 (Sathyaprakash & Dhurandhar 1991b; Droz et al. 1999; Blanchet 2002; Faye et al. 2012) and TaylorF2e (Moore & Yunes 2019, 2020; Moore et al. 2018). To account for the long signal duration, our method allows the option to fix the duration of waveform models by varying the lower frequency. Nitz & Wang (2021a) used this option to set the wavelength to 512 seconds. Overall, the bank consisted of 7.8×10^6 templates, where half of the templates included orbital eccentricity.

Searching for low-mass neutron stars requires a similar template bank as the PBHs search. However, these systems requires consideration for the tidal deformability parameter instead of the eccentricity, as the eccentricity in these systems is expected to be negligible. Discovering neutron stars in the sub-solar mass range could challenge our understanding of their formation or potentially reveal a new class of stars (Doroshenko et al. 2022). Previous searches neglected the tidal deformability parameter λ , which lost up to 78.4% of the total signals (Bandopadhyay et al. 2023). This bank was constructed with tidal deformability ranging from $[0,10000]$ for both λ_1 and λ_2 to account for the loss in SNR, primary mass ranging $[0.1,2] M_\odot$, secondary mass ranging $[0.1,1] M_\odot$, both aligned spin χ_{1z} and χ_{2z} ranged from $[-0.05,0.05]$, and the approximant was chosen to be TaylorF2 (Sathyaprakash & Dhurandhar 1991b; Droz et al. 1999; Blanchet 2002; Faye et al. 2012). Similarly to the PBH search, this bank had the template waveforms fixed to 512 seconds to speed up the search. Figure 4 illustrates the varying lower frequencies of the waveforms used to maintain this duration. Overall, the bank consisted of 1.01×10^7 templates.

6.2. Eccentric NSBH and BNS Search

Measuring the orbital eccentricity of a binary provides insights into its formation history (Belczynski et al. 2001; Rodriguez et al. 2018; Romero-Shaw et al. 2019; Sedda 2020; Trani et al. 2022; Zevin et al. 2021; Dhurkunde & Nitz 2023). Searches for rare binaries with non-negligible eccentricity require additional parameters in the template bank: requiring up to 100x more templates than the typical aligned-spin, quasi-circular searches (Nitz & Wang 2021b,c; Dhurkunde & Nitz 2023). The first modeled search for eccentric spinning neutron-star binaries (BNS and NSBH) using pub-




Figure 4. Sub-solar mass neutron star search template bank with 10 million templates. Mass ranges from 0.1 to 2 for the primary mass and 0.1 to 1 for the secondary mass. The signals are fixed to 512 seconds by varying the lower frequency cutoff indicated in the color bar.

licly available data from the third observing run of the Advanced LIGO and Virgo observatories, was performed by Dhurkunde & Nitz (2023). Dhurkunde & Nitz (2023) utilized the flexible bank generation method described in this work to incorporate eccentricity in the template bank. The template bank comprised of six parameters: component masses (m_1, m_2), aligned component spins (s_{1z}, s_{2z}) and orbital eccentricity e_{20} (defined at 20 Hz), and an additional angle to account for the orientation of the elliptical orbit w.r.t an observer. Signals within the targeted search region are reliably searched using the inspiral-only TaylorF2Ecc model Moore et al. (2016), as the merger phase falls outside the sensitive range of current detectors. The template bank for the search consisted of approximately 6 million templates, and its generation took about a week to complete using 20 cores.

6.3. Binary Neutron Star Confusion Noise Cleaning

In the next decade, next-generation ground-based detectors, such as the Einstein Telescope (ET) (Hild et al. 2010; Punturo et al. 2010) and Cosmic Explorer (CE) (Reitze et al. 2019), will be available. Those detectors will not just be one order of magnitude more sensitive than the current Advanced LIGO and Virgo detectors, but also can reach 2 or 3 Hz, making the binary neutron stars’ signals last several hours or even several days. According to GWTC-3 population results (Abbott et al. 2023), Wu & Nitz (2023) simulates the mock data for those next-generation ground-based detectors and finds

that those BNS signals will form the foreground noise to reduce the detection range of detectors if not removed. This BNS confusion noise will affect Cosmic Explorer most by raising the power spectrum density around 10 Hz and introducing a noise correlation in the detector network. Wu & Nitz (2023) demonstrates a foreground cleaning method using the stochastic template bank for BNS. For bank generation, they choose $[2.4, 60] M_\odot$ as the detector-frame total mass range and $[1, 1.636]$ as the mass ratio range according to the BNS population model. With a mismatch of 0.97 using the design sensitivities of ET and CE, the foreground noise-cleaning banks contain 9.57×10^4 and 1.53×10^5 templates for CE and ET, respectively. Using these banks to detect and subtract the BNS confusion noise for each detector, they can suppress the total noise to almost the instrument noise level, allowing for near-optimal searches at the following stages. The computational cost of generating and searching with these template banks is lower compared to the full search (Lenon et al. 2021).

6.4. LISA Supermassive Black Hole Binaries Search and Inference

There are ongoing efforts to extend PyCBC to do LISA data analysis, Weaving et al. (2024) demonstrates how to use the stochastic template bank method described in this paper to find supermassive black hole binaries (SMBHBs) and use the corresponding high SNR template as the reference signal in the following heterodyning parameter estimation. Previously, people thought the template-based analysis was not viable for LISA data analysis due to the huge parameter space (Harry et al. 2008), but Weaving et al. (2024) demonstrates we can still use a sparse template bank to find all SMBHB signals in the LISA mock dataset Sangria. Different from ground-based detectors, LISA waveforms also need to take the orbital motion of spacecraft and time-delay interferometry (TDI) into account, so these make LISA waveforms much more complex. Weaving et al. (2024) uses the BBHx (Katz et al. 2020) package to generate the LISA-TDI version of the IMRPhenomD waveform and chooses an eight-dimensional parameter space (detector-frame total mass, mass ratio, two aligned spins, ecliptic latitude, ecliptic longitude, polarization angle, and inclination angle) that covers the parameters of SMBHBs in the Sangria dataset. Due to the high SNR of those SMBHB signals and almost equal sensitivities of TDI-A and TDI-E channels, they just generate the TDI-A channel’s template bank and use a mismatch threshold of 0.9 to get the SMBHB template bank consisting of around 50 templates. Finally, they

successfully detected all the SMBHB signals in the Sanguia dataset.

6.5. *Open Gravitational-Wave Catalog Search for Compact-Binary Mergers*

The fourth open gravitational-wave catalog (OGC) contains the observation of nearly 100 compact binary mergers (Nitz et al. 2023). The search covers a parameter space from neutron star binaries through heavy binary black holes (up to $\sim 1000 M_{\odot}$). The OGC search splits the analysis into four sub-regions, covering BNS, NSBH, BBH, and a focused BBH region where the largest numbers of signals are observed. The analysis based on the open source PyCBC toolkit (Usman et al. 2016; Davies et al. 2020) and makes use of the flexible template bank generation introduced in this work. This enables the use of multiple waveform approximants depending on the suitability for different parts of parameter space, e.g. the use of TaylorF2 (Sathyaprakash & Dhurandhar 1991b; Droz et al. 1999; Blanchet 2002; Faye et al. 2012) for neutron star binaries, and IMR-PhenomD (Husa et al. 2016; Khan et al. 2016b) and SEOBNRv4 for BBH signals (Taracchini et al. 2014; Bohé et al. 2016). The stochastic placement algorithm allows for the iterative generation of a larger template bank by adding to a pre-existing template bank. This allows for regions of parameter space that adapt to different requirements, e.g. a higher minimal match for the focused BBH region of 0.995 or the inclusion of tidal deformability for binary neutron star templates.

6.6. *Binaries Contain Neutron Star as Gamma-Ray Burst Progenitors*

Mergers of binary neutron star and neutron-star-black-hole systems have long been suspected to be the production sites of short duration gamma-ray bursts (Blinnikov et al. 1984; Eichler et al. 1989; Paczynski 1986, 1991; Narayan et al. 1992; Nakar 2007; Berger 2014). In the case of binary neutron stars, this hypothesis was confirmed by the simultaneous detection of GW event GW170817 and its gamma-ray burst counterpart GRB 170817A (Abbott et al. 2017). The main mechanisms invoked to launch the relativistic jet responsible for the gamma-ray emission involve neutrino pair annihilation or the presence of strong magnetic fields. Both scenarios require a remnant constituted by a central compact object accreting from a dense torus of matter surrounding it that develops immediately after the merger event. In the case of two binary neutron stars, the formation of the remnant torus is fueled by the collision of the two neutron stars, while for neutron-star-black-hole mergers the presence of the torus depends on

the parameters of the system, (see, e.g., Gonzalez et al. 2023; Kyutoku et al. 2021, for examples of numerical simulations).

When constructing template banks for searches that aim at uncovering GW signals compatible with the time and sky location of gamma-ray bursts (Harry & Fairhurst 2011; Williamson et al. 2014), the goal is to include all binary neutron star mergers and only those neutron-star-black-hole mergers that result in the formation of an accretion torus, as suggested in Pannarale & Ohme (2014). These are referred to as “EM-bright” template banks. To discriminate between neutron-star-black-hole systems that produce matter surrounding the central remnant black hole and ones that do not, the PyCBC toolkit implements a formula that predicts the remnant mass left behind in the post-merger. This formula was obtained in Foucart et al. (2018) by fitting results of numerical-relativity neutron-star-black-hole merger simulations; it returns the remnant mass M_{rem} given the following parameters of the binary system: its symmetric mass ratio, the radius of the black hole’s innermost stable circular orbit, and the neutron star compactness (see Eq. (7) in Foucart et al. 2018, for details). These banks are therefore built by prescribing priors for masses and spins of neutron stars and black holes, and applying the constraint that the remnant mass surrounding the central black hole is non-vanishing in the case of neutron-star-black-hole systems. Given a draw from the priors, there are two operations in this process that depend on the neutron star equation of state. 1) Deciding whether or not each compact object is a neutron star, so that binary neutron stars are added to the bank if necessary, and binary black holes are discarded. 2) In the case of a neutron-star-black-hole binary draw, determining the neutron star compactness corresponding to its mass in order to apply the constraint $M_{\text{rem}} > 0$. Building an EM-bright bank therefore also requires an additional input in the form of a table with mass and compactness values that represent a non-rotating¹ neutron star equilibrium configuration; this needs to be built externally by the user, see, e.g., <https://compose.obspm.fr/>. Finally, in the case of black hole spins that are not aligned to the orbital angular momentum, we replace the radius of the innermost stable circular orbit with its tilted analogue, the radius of the innermost stable spherical orbit, as detailed in Appendix A of Stone et al. (2013).

¹ Recall that the M_{rem} formula does not depend on the neutron star spin.

Two neutron-star–black-hole banks were produced: one with the EM-bright constraint and one without it. The design parameters were the same for both banks, with the exception of the remnant mass, which was fixed to be strictly greater than zero for the EM-bright bank. In this case, the 2H piecewise polytropic equation of state (Read et al. 2009) was adopted. This choice was driven by the fact that this equation of state sets a high maximum neutron star mass ($\sim 2.83 M_{\odot}$) and it favours tidal disruption because of the high neutron star compactness values it yields, compared to other equations of state. In this sense it is a conservative choice, that is, it makes the EM-bright constraint as loose as possible. The black hole mass and spin spanned from $2.83 M_{\odot}$ to $25 M_{\odot}$ and from 0.0 to 0.98, respectively. The neutron star mass ranged from $1.0 M_{\odot}$ to $2.83 M_{\odot}$, while the aligned spin ranged from 0.0 to 0.05. All priors in these intervals were taken to be uniform. Additionally, we used a minimal match of 0.97 and a lower frequency of 27.0 Hz, and we set the PSD to be the theoretical Advanced LIGO O4 sensitivity (Abbott et al. 2020). The full bank consisted of 193,235 templates, while the EM-bright bank had 143,530 templates, resulting in the exclusion of 49,705 neutron-star–black-hole templates representing the portion of parameter space where merger events do not produce a torus as a remnant.

7. CONCLUSION

We introduce an efficient and flexible method for stochastic template bank generation that produces self-consistent banks for varying tolerance and minimal match conditions. Our method has already been used for various searches for GWs from different types of compact binary sources such as eccentric BNSs and NSBHs (Dhurkunde & Nitz 2023), sub-solar mass primordial black holes (Nitz & Wang 2021a), low-mass BNSs, studies of next-generation ground-based detectors’ BNS signal detection problem (Wu & Nitz 2023; Lenon et al. 2021), LISA supermassive BBHs (Weaving et al. 2024), and the fourth open gravitational-wave catalog (Nitz et al. 2023). We find that existing methods are robust and versatile enough for various unique astrophysical scenarios.

We have demonstrated how the number of parameters in a search scales the computational cost. Compared to a non-spinning bank with only two parameters (masses only), we find an increase of up to $\mathcal{O}(10^2)$ templates as additional parameters are included in the bank. For future observatories, aligned-spin banks will get bigger by up to two orders of magnitude compared to the current banks (Dhurkunde & Nitz 2022). Fast bank generation algorithms alone will not inherently lead to

faster searches. The most pressing need for improvement is developing faster or optimizing existing search methods. Ongoing efforts to utilize various hierarchical methods could make current searches up to 20 times faster (Dhurkunde et al. 2022; Soni et al. 2022). If efforts to procure faster searches succeed, the cost of template banks will once again become considerable, and efforts to create faster methods will become critical.

In conclusion, while our method demonstrates significant improvements in template bank generation, it is important to consider the limitations and potential optimizations for utilizing our method for future optimized searches. Stochastic methods can be limited when the waveform approximant takes a long time to generate, as the time to generate a waveform linearly scales with the template bank. If the waveform generation time is comparable to the search time, geometric methods might be more appropriate. However, if a search requires a non-trivial parameter space and utilizes slow waveform approximants, optimizing stochastic bank methods becomes necessary. Our method is currently written for CPU use. This method could potentially be optimized by parallelizing the template acceptance proposals on GPUs to help speed up the generation. This optimization might help offset the additional time required for slow waveform approximants.

The bank generation code `pycbc_brute_bank`, which uses the method described in this paper, is at https://github.com/gwastro/pycbc/blob/master/bin/bank/pycbc_brute_bank.

ACKNOWLEDGMENTS

KK and AHN acknowledge support from NSF grant PHY-2309240. KK and AHN acknowledge the support from Syracuse University for providing the computational resources through the OrangeGrid High Throughput Computing (HTC) cluster. Additionally, KK expresses gratitude to Ananya Bandopadhyay and Aleya Akyüz for their valuable feedback and insightful comments, which greatly improved the quality of this paper. RD and SW acknowledges Max Planck Gesellschaft for support. MC acknowledges the support through the grant PID2021-127495NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by the European Union, and the Astrophysics and High Energy Physics programme of the Generalitat Valenciana ASFAE/2022/026 funded by MCIN and the European Union NextGenerationEU (PRTR-C17.I1). FP acknowledges support from the ICSC - Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by the European Union - NextGenerationEU and support from the Italian Min-

istry of University and Research (MUR) Progetti di ricerca di Rilevante Interesse Nazionale (PRIN) Bando 2022 - grant 20228TLHPE - CUP I53D23000630006.

REFERENCES

- Abbott, B. P., et al. 2017, *Astrophys. J. Lett.*, 848, L13, doi: [10.3847/2041-8213/aa920c](https://doi.org/10.3847/2041-8213/aa920c)
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2019, *Physical Review X*, 9, 031040, doi: [10.1103/PhysRevX.9.031040](https://doi.org/10.1103/PhysRevX.9.031040)
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2020, *Living Reviews in Relativity* 2020 23:1, 23, 1, doi: [10.1007/S41114-020-00026-9](https://doi.org/10.1007/S41114-020-00026-9)
- Abbott, R., Abbott, T. D., Acernese, F., et al. 2023, *Phys. Rev. X*, 13, 041039, doi: [10.1103/PhysRevX.13.041039](https://doi.org/10.1103/PhysRevX.13.041039)
- Abbott, R., Abbott, T. D., Acernese, F., et al. 2023, *Physical Review X*, 13, 011048, doi: [10.1103/PhysRevX.13.011048](https://doi.org/10.1103/PhysRevX.13.011048)
- Abbott, R., Abbott, T. D., Acernese, F., et al. 2024, *Phys. Rev. D*, 109, 022001, doi: [10.1103/PhysRevD.109.022001](https://doi.org/10.1103/PhysRevD.109.022001)
- Acernese, F., Agathos, M., Agatsuma, K., et al. 2015, *Classical and Quantum Gravity*, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001)
- Ajith, P., Fotopoulos, N., Privitera, S., et al. 2014, *PhRvD*, 89, 084041, doi: [10.1103/PhysRevD.89.084041](https://doi.org/10.1103/PhysRevD.89.084041)
- Allen, B., Anderson, W. G., Brady, P. R., Brown, D. A., & Creighton, J. D. E. 2012, *Phys. Rev. D*, 85, 122006, doi: [10.1103/PhysRevD.85.122006](https://doi.org/10.1103/PhysRevD.85.122006)
- Allen, B., Anderson, W. G., Brady, P. R., Brown, D. A., & Creighton, J. D. E. 2012, *PhRvD*, 85, 122006, doi: [10.1103/PhysRevD.85.122006](https://doi.org/10.1103/PhysRevD.85.122006)
- Apostolatos, T. A. 1995, *Phys. Rev. D*, 52, 605, doi: [10.1103/PhysRevD.52.605](https://doi.org/10.1103/PhysRevD.52.605)
- Aubin, F., Brighenti, F., Chierici, R., et al. 2021, *Classical and Quantum Gravity*, 38, 095004, doi: [10.1088/1361-6382/abe913](https://doi.org/10.1088/1361-6382/abe913)
- Babak, S., Balasubramanian, R., Churches, D., Cokelaer, T., & Sathyaprakash, B. S. 2006, *Classical and Quantum Gravity*, 23, 5477, doi: [10.1088/0264-9381/23/18/002](https://doi.org/10.1088/0264-9381/23/18/002)
- Babak, S., Biswas, R., Brady, P. R., et al. 2013, *Phys. Rev. D*, 87, 024033, doi: [10.1103/PhysRevD.87.024033](https://doi.org/10.1103/PhysRevD.87.024033)
- Balasubramanian, R., Sathyaprakash, B. S., & Dhurandhar, S. V. 1996, *PhRvD*, 53, 3033, doi: [10.1103/PhysRevD.53.3033](https://doi.org/10.1103/PhysRevD.53.3033)
- Bandopadhyay, A., Reed, B., Padamata, S., et al. 2023, *PhRvD*, 107, 103012, doi: [10.1103/PhysRevD.107.103012](https://doi.org/10.1103/PhysRevD.107.103012)
- Belczynski, K., Kalogera, V., & Bulik, T. 2001, *Astrophys. J.*, 572, 407, doi: [10.1086/340304](https://doi.org/10.1086/340304)
- Berger, E. 2014, *ARA&A*, 52, 43, doi: [10.1146/annurev-astro-081913-035926](https://doi.org/10.1146/annurev-astro-081913-035926)
- Blanchet, L. 2002, *Living Rev. Rel.*, 5, 3
- Blinnikov, S. I., Novikov, I. D., Perevodchikova, T. V., & Polnarev, A. G. 1984, *Soviet Astronomy Letters*, 10, 177, doi: [10.48550/arXiv.1808.05287](https://doi.org/10.48550/arXiv.1808.05287)
- Bohé, A., et al. 2016. <https://arxiv.org/abs/1611.03703>
- Brown, D. A., Harry, I., Lundgren, A., & Nitz, A. H. 2012, *PhRvD*, 86, 084017, doi: [10.1103/PhysRevD.86.084017](https://doi.org/10.1103/PhysRevD.86.084017)
- Brown, D. A., Harry, I., Lundgren, A., & Nitz, A. H. 2012, *Phys. Rev. D*, 86, 084017, doi: [10.1103/PhysRevD.86.084017](https://doi.org/10.1103/PhysRevD.86.084017)
- Buikema, A., Cahillane, C., Mansell, G. L., et al. 2020, *PhRvD*, 102, 062003, doi: [10.1103/PhysRevD.102.062003](https://doi.org/10.1103/PhysRevD.102.062003)
- Cannon, K., Caudill, S., Chan, C., et al. 2021, *SoftwareX*, 14, 100680, doi: [10.1016/j.softx.2021.100680](https://doi.org/10.1016/j.softx.2021.100680)
- Capano, C., Harry, I., Privitera, S., & Buonanno, A. 2016, *Phys. Rev. D*, 93, 124007, doi: [10.1103/PhysRevD.93.124007](https://doi.org/10.1103/PhysRevD.93.124007)
- Chu, Q., Kovalam, M., Wen, L., et al. 2020, arXiv e-prints, arXiv:2011.06787, doi: [10.48550/arXiv.2011.06787](https://doi.org/10.48550/arXiv.2011.06787)
- Cokelaer, T. 2007, *PhRvD*, 76, 102004, doi: [10.1103/PhysRevD.76.102004](https://doi.org/10.1103/PhysRevD.76.102004)
- Coogan, A., Edwards, T. D. P., Chia, H. S., et al. 2022, *PhRvD*, 106, 122001, doi: [10.1103/PhysRevD.106.122001](https://doi.org/10.1103/PhysRevD.106.122001)
- Coogan, A., Edwards, T. D. P., Chia, H. S., et al. 2022, *Phys. Rev. D*, 106, 122001, doi: [10.1103/PhysRevD.106.122001](https://doi.org/10.1103/PhysRevD.106.122001)
- Dal Canton, T., Nitz, A. H., Lundgren, A. P., et al. 2014, *PhRvD*, 90, 082004, doi: [10.1103/PhysRevD.90.082004](https://doi.org/10.1103/PhysRevD.90.082004)
- Davies, G. S., Dent, T., Tápai, M., et al. 2020, *Phys. Rev. D*, 102, 022004, doi: [10.1103/PhysRevD.102.022004](https://doi.org/10.1103/PhysRevD.102.022004)
- Dhurkunde, R., Fehrmann, H., & Nitz, A. H. 2022, *Phys. Rev. D*, 105, 103001, doi: [10.1103/PhysRevD.105.103001](https://doi.org/10.1103/PhysRevD.105.103001)
- Dhurkunde, R., & Nitz, A. H. 2022, *Phys. Rev. D*, 106, 103035, doi: [10.1103/PhysRevD.106.103035](https://doi.org/10.1103/PhysRevD.106.103035)
- . 2023. <https://arxiv.org/abs/2311.00242>
- Dhurkunde, R., & Nitz, A. H. 2023, arXiv e-prints, arXiv:2311.00242, doi: [10.48550/arXiv.2311.00242](https://doi.org/10.48550/arXiv.2311.00242)
- Domènech, G., & Sasaki, M. 2024, arXiv e-prints, arXiv:2401.07615, doi: [10.48550/arXiv.2401.07615](https://doi.org/10.48550/arXiv.2401.07615)
- Doroshenko, V., Suleimanov, V., Pühlhofer, G., & Santangelo, A. 2022, *Nature Astronomy*, 6, 1444, doi: [10.1038/s41550-022-01800-1](https://doi.org/10.1038/s41550-022-01800-1)

- Droz, S., Knapp, D. J., Poisson, E., & Owen, B. J. 1999, *PhRvD*, 59, 124016
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, *Nature*, 340, 126, doi: [10.1038/340126a0](https://doi.org/10.1038/340126a0)
- Einstein, A. 1916, *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 1, 688
- . 1918, *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 1, 154
- Faye, G., Marsat, S., Blanchet, L., & Iyer, B. R. 2012, *Class. Quant. Grav.*, 29, 175004, doi: [10.1088/0264-9381/29/17/175004](https://doi.org/10.1088/0264-9381/29/17/175004)
- Foucart, F., Hinderer, T., & Nissanke, S. 2018, *Phys. Rev. D*, 98, 081501, doi: [10.1103/PhysRevD.98.081501](https://doi.org/10.1103/PhysRevD.98.081501)
- Frampton, P. H. 2016, *Modern Physics Letters A*, 31, 1650064, doi: [10.1142/S0217732316500644](https://doi.org/10.1142/S0217732316500644)
- Gonzalez, A., et al. 2023, *Class. Quant. Grav.*, 40, 085011, doi: [10.1088/1361-6382/acc231](https://doi.org/10.1088/1361-6382/acc231)
- Green, A. M. 2024, arXiv e-prints, arXiv:2402.15211, doi: [10.48550/arXiv.2402.15211](https://doi.org/10.48550/arXiv.2402.15211)
- Hanna, C., Kennington, J., Sakon, S., et al. 2023, *Phys. Rev. D*, 108, 042003, doi: [10.1103/PhysRevD.108.042003](https://doi.org/10.1103/PhysRevD.108.042003)
- Harry, I., Bohe, A., Privitera, S., & Buonanno, A. 2016, in *APS Meeting Abstracts*, Vol. 2016, APS April Meeting Abstracts, K14.005
- Harry, I. W., Allen, B., & Sathyaprakash, B. S. 2009, *PhRvD*, 80, 104014, doi: [10.1103/PhysRevD.80.104014](https://doi.org/10.1103/PhysRevD.80.104014)
- Harry, I. W., Allen, B., & Sathyaprakash, B. S. 2009, *Phys. Rev. D*, 80, 104014, doi: [10.1103/PhysRevD.80.104014](https://doi.org/10.1103/PhysRevD.80.104014)
- Harry, I. W., & Fairhurst, S. 2011, *Phys. Rev. D*, 83, 084002, doi: [10.1103/PhysRevD.83.084002](https://doi.org/10.1103/PhysRevD.83.084002)
- Harry, I. W., Fairhurst, S., & Sathyaprakash, B. S. 2008, *Classical and Quantum Gravity*, 25, 184027, doi: [10.1088/0264-9381/25/18/184027](https://doi.org/10.1088/0264-9381/25/18/184027)
- Hild, S., Chelkowski, S., Freise, A., et al. 2010, *Classical and Quantum Gravity*, 27, 015003, doi: [10.1088/0264-9381/27/1/015003](https://doi.org/10.1088/0264-9381/27/1/015003)
- Husa, S., Khan, S., Hannam, M., et al. 2016, *PhRvD*, 93, 044006, doi: [10.1103/PhysRevD.93.044006](https://doi.org/10.1103/PhysRevD.93.044006)
- Husa, S., Khan, S., Hannam, M., et al. 2016, *Phys. Rev. D*, 93, 044006, doi: [10.1103/PhysRevD.93.044006](https://doi.org/10.1103/PhysRevD.93.044006)
- Indik, N., Haris, K., Dal Canton, T., et al. 2017, *PhRvD*, 95, 064056, doi: [10.1103/PhysRevD.95.064056](https://doi.org/10.1103/PhysRevD.95.064056)
- Katz, M. L., Marsat, S., Chua, A. J. K., Babak, S., & Larson, S. L. 2020, *PhRvD*, 102, 023033, doi: [10.1103/PhysRevD.102.023033](https://doi.org/10.1103/PhysRevD.102.023033)
- Khan, S., Husa, S., Hannam, M., et al. 2016a, *Phys. Rev. D*, 93, 044007, doi: [10.1103/PhysRevD.93.044007](https://doi.org/10.1103/PhysRevD.93.044007)
- . 2016b, *Phys. Rev. D*, 93, 044007, doi: [10.1103/PhysRevD.93.044007](https://doi.org/10.1103/PhysRevD.93.044007)
- Kyutoku, K., Shibata, M., & Taniguchi, K. 2021, *Living Rev. Rel.*, 24, 5, doi: [10.1007/s41114-021-00033-4](https://doi.org/10.1007/s41114-021-00033-4)
- Lenon, A. K., Brown, D. A., & Nitz, A. H. 2021, *PhRvD*, 104, 063011, doi: [10.1103/PhysRevD.104.063011](https://doi.org/10.1103/PhysRevD.104.063011)
- LIGO Scientific Collaboration, Aasi, J., Abbott, B. P., et al. 2015, *Classical and Quantum Gravity*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](https://doi.org/10.1088/0264-9381/32/7/074001)
- Manca, G. M., & Vallisneri, M. 2010, *PhRvD*, 81, 024004, doi: [10.1103/PhysRevD.81.024004](https://doi.org/10.1103/PhysRevD.81.024004)
- Mehta, A. K., Olsen, S., Wadekar, D., et al. 2023, <https://arxiv.org/abs/2311.06061>
- Messick, C., et al. 2017, *Phys. Rev. D*, 95, 042001, doi: [10.1103/PhysRevD.95.042001](https://doi.org/10.1103/PhysRevD.95.042001)
- Miller, A. L., Aggarwal, N., Clesse, S., et al. 2024, arXiv e-prints, arXiv:2402.19468, doi: [10.48550/arXiv.2402.19468](https://doi.org/10.48550/arXiv.2402.19468)
- Moore, B., Favata, M., Arun, K. G., & Mishra, C. K. 2016, *Phys. Rev. D*, 93, 124061, doi: [10.1103/PhysRevD.93.124061](https://doi.org/10.1103/PhysRevD.93.124061)
- Moore, B., Robson, T., Loutrel, N., & Yunes, N. 2018, *Classical and Quantum Gravity*, 35, 235006, doi: [10.1088/1361-6382/aaea00](https://doi.org/10.1088/1361-6382/aaea00)
- Moore, B., & Yunes, N. 2019, *Classical and Quantum Gravity*, 36, 185003, doi: [10.1088/1361-6382/ab3778](https://doi.org/10.1088/1361-6382/ab3778)
- . 2020, *Classical and Quantum Gravity*, 37, 225015, doi: [10.1088/1361-6382/ab7963](https://doi.org/10.1088/1361-6382/ab7963)
- Nakar, E. 2007, *Phys. Rept.*, 442, 166, doi: [10.1016/j.physrep.2007.02.005](https://doi.org/10.1016/j.physrep.2007.02.005)
- Narayan, R., Paczynski, B., & Piran, T. 1992, *ApJL*, 395, L83, doi: [10.1086/186493](https://doi.org/10.1086/186493)
- Nitz, A. H., Dal Canton, T., Davis, D., & Reyes, S. 2018, *PhRvD*, 98, 024050, doi: [10.1103/PhysRevD.98.024050](https://doi.org/10.1103/PhysRevD.98.024050)
- Nitz, A. H., Kumar, S., Wang, Y.-F., et al. 2023, *ApJ*, 946, 59, doi: [10.3847/1538-4357/aca591](https://doi.org/10.3847/1538-4357/aca591)
- Nitz, A. H., & Wang, Y.-F. 2021a, *Phys. Rev. Lett.*, 127, 151101, doi: [10.1103/PhysRevLett.127.151101](https://doi.org/10.1103/PhysRevLett.127.151101)
- . 2021b, *Phys. Rev. Lett.*, 127, 151101, doi: [10.1103/PhysRevLett.127.151101](https://doi.org/10.1103/PhysRevLett.127.151101)
- . 2021c, doi: [10.3847/1538-4357/ac01d9](https://doi.org/10.3847/1538-4357/ac01d9)
- Olsen, S., Venumadhav, T., Mushkin, J., et al. 2022, *Phys. Rev. D*, 106, 043009, doi: [10.1103/PhysRevD.106.043009](https://doi.org/10.1103/PhysRevD.106.043009)
- Owen, B. J. 1996, *Phys. Rev. D*, 53, 6749, doi: [10.1103/PhysRevD.53.6749](https://doi.org/10.1103/PhysRevD.53.6749)
- Owen, B. J., & Sathyaprakash, B. S. 1999, *Phys. Rev. D*, 60, 022002, doi: [10.1103/PhysRevD.60.022002](https://doi.org/10.1103/PhysRevD.60.022002)
- Paczynski, B. 1986, *ApJL*, 308, L43, doi: [10.1086/184740](https://doi.org/10.1086/184740)
- Paczynski, B. 1991, *Acta Astron.*, 41, 257
- Pang, P. T. H., Lo, R. K. L., Wong, I. C. F., Li, T. G. F., & Van Den Broeck, C. 2020, *PhRvD*, 101, 104055, doi: [10.1103/PhysRevD.101.104055](https://doi.org/10.1103/PhysRevD.101.104055)

- Pannarale, F., & Ohme, F. 2014, *Astrophys. J. Lett.*, 791, L7, doi: [10.1088/2041-8205/791/1/L7](https://doi.org/10.1088/2041-8205/791/1/L7)
- Privitera, S., Mohapatra, S. R. P., Ajith, P., et al. 2014, *PhRvD*, 89, 024003, doi: [10.1103/PhysRevD.89.024003](https://doi.org/10.1103/PhysRevD.89.024003)
- Prix, R. 2007, *Classical and Quantum Gravity*, 24, S481, doi: [10.1088/0264-9381/24/19/S11](https://doi.org/10.1088/0264-9381/24/19/S11)
- Punturo, M., Abernathy, M., Acernese, F., et al. 2010, *Classical and Quantum Gravity*, 27, 194002, doi: [10.1088/0264-9381/27/19/194002](https://doi.org/10.1088/0264-9381/27/19/194002)
- Read, J. S., Markakis, C., Shibata, M., et al. 2009, *Phys. Rev. D*, 79, 124033, doi: [10.1103/PhysRevD.79.124033](https://doi.org/10.1103/PhysRevD.79.124033)
- Reitze, D., Adhikari, R. X., Ballmer, S., et al. 2019, in *Bulletin of the American Astronomical Society*, Vol. 51, 35, doi: [10.48550/arXiv.1907.04833](https://doi.org/10.48550/arXiv.1907.04833)
- Rodriguez, C. L., Amaro-Seoane, P., Chatterjee, S., & Rasio, F. A. 2018, *Phys. Rev. Lett.*, 120, 151101, doi: [10.1103/PhysRevLett.120.151101](https://doi.org/10.1103/PhysRevLett.120.151101)
- Romero-Shaw, I. M., Lasky, P. D., & Thrane, E. 2019, *Mon. Not. Roy. Astron. Soc.*, 490, 5210, doi: [10.1093/mnras/stz2996](https://doi.org/10.1093/mnras/stz2996)
- Roy, S., Sengupta, A. S., & Ajith, P. 2017, arXiv e-prints, arXiv:1711.08743, doi: [10.48550/arXiv.1711.08743](https://doi.org/10.48550/arXiv.1711.08743)
- Roy, S., Sengupta, A. S., & Ajith, P. 2019, *Phys. Rev. D*, 99, 024048, doi: [10.1103/PhysRevD.99.024048](https://doi.org/10.1103/PhysRevD.99.024048)
- Roy, S., Sengupta, A. S., & Thakor, N. 2017, *Phys. Rev. D*, 95, 104045, doi: [10.1103/PhysRevD.95.104045](https://doi.org/10.1103/PhysRevD.95.104045)
- Sathyaprakash, B. S., & Dhurandhar, S. V. 1991a, *Phys. Rev. D*, 44, 3819, doi: [10.1103/PhysRevD.44.3819](https://doi.org/10.1103/PhysRevD.44.3819)
- . 1991b, *Phys. Rev. D*, 44, 3819, doi: [10.1103/PhysRevD.44.3819](https://doi.org/10.1103/PhysRevD.44.3819)
- Schmidt, S., Gadre, B., & Caudill, S. 2024, *PhRvD*, 109, 042005, doi: [10.1103/PhysRevD.109.042005](https://doi.org/10.1103/PhysRevD.109.042005)
- Sedda, M. A. 2020, *Commun. Phys.*, 3, 43, doi: [10.1038/s42005-020-0310-x](https://doi.org/10.1038/s42005-020-0310-x)
- Soni, K., Gadre, B. U., Mitra, S., & Dhurandhar, S. 2022, *Phys. Rev. D*, 105, 064005, doi: [10.1103/PhysRevD.105.064005](https://doi.org/10.1103/PhysRevD.105.064005)
- Stone, N., Loeb, A., & Berger, E. 2013, *Phys. Rev. D*, 87, 084053, doi: [10.1103/PhysRevD.87.084053](https://doi.org/10.1103/PhysRevD.87.084053)
- Taracchini, A., Buonanno, A., Pan, Y., et al. 2014, *PhRvD*, 89, 061502, doi: [10.1103/PhysRevD.89.061502](https://doi.org/10.1103/PhysRevD.89.061502)
- Trani, A. A., Rastello, S., Di Carlo, U. N., et al. 2022, *Mon. Not. Roy. Astron. Soc.*, 511, 1362, doi: [10.1093/mnras/stac122](https://doi.org/10.1093/mnras/stac122)
- Usman, S. A., Nitz, A. H., Harry, I. W., et al. 2016, *Classical and Quantum Gravity*, 33, 215004, doi: [10.1088/0264-9381/33/21/215004](https://doi.org/10.1088/0264-9381/33/21/215004)
- Weaving, C. R., Nuttall, L. K., Harry, I. W., Wu, S., & Nitz, A. 2024, *Classical and Quantum Gravity*, 41, 025006, doi: [10.1088/1361-6382/ad134d](https://doi.org/10.1088/1361-6382/ad134d)
- Williamson, A. R., Biwer, C., Fairhurst, S., et al. 2014, *Phys. Rev. D*, 90, 122004, doi: [10.1103/PhysRevD.90.122004](https://doi.org/10.1103/PhysRevD.90.122004)
- Wu, S., & Nitz, A. H. 2023, *PhRvD*, 107, 063022, doi: [10.1103/PhysRevD.107.063022](https://doi.org/10.1103/PhysRevD.107.063022)
- Zevin, M., Romero-Shaw, I. M., Kremer, K., Thrane, E., & Lasky, P. D. 2021, *Astrophys. J. Lett.*, 921, L43, doi: [10.3847/2041-8213/ac32dc](https://doi.org/10.3847/2041-8213/ac32dc)