

ROBUST Q -LEARNING FOR FINITE AMBIGUITY SETS

CÉCILE DECKER¹, JULIAN SESTER^{1,*}

July 8, 2024

¹*National University of Singapore, Department of Mathematics,
21 Lower Kent Ridge Road, 119077.*

*Corresponding author, email:jul_ses@nus.edu.sg

ABSTRACT. In this paper we propose a novel Q -learning algorithm allowing to solve distributionally robust Markov decision problems for which the ambiguity set of probability measures can be chosen arbitrarily as long as it comprises only a finite amount of measures. Therefore, our approach goes beyond the well-studied cases involving ambiguity sets of balls around some reference measure with the distance to reference measure being measured with respect to the Wasserstein distance or the Kullback–Leibler divergence. Hence, our approach allows the applicant to create ambiguity sets better tailored to her needs and to solve the associated robust Markov decision problem via a Q -learning algorithm whose convergence is guaranteed by our main result. Moreover, we showcase in several numerical experiments the tractability of our approach.

Keywords: Q -Learning, Markov Decision Processes, Model Uncertainty

1. INTRODUCTION

Markov decision problems are discrete time stochastic control problems involving an agent who can influence the environment by executing actions affecting the distribution of the realization of the subsequent state of the environment. In turn, depending on the state realization, the agent receives a feedback in form of a reward and decides on her next action. The objective of the agent is then to optimize the cumulative expected rewards obtained over an infinite time horizon by choosing the highest rewarding sequence of actions.

By construction, the described optimization problem depends crucially on the involved distributions for the state transitions. In practice, frequently in model-based approaches these distributions are estimated ([1], [32], [37]) from data or implicitly learned on-line in model-free approaches (see, e.g., [8], [39]). However chosen, the underlying model is highly prone to *model misspecification*, i.e., if the underlying distribution changes after training, the agent is trained to the wrong underlying distribution and successful execution of actions (i.e., receiving high rewards) typically will fail if the distribution change is too large. A solution to this problem is to take distributional uncertainty into account, i.e., to train the agent already by using an *ambiguity set* of probability distributions and train the agent according to the worst case measure contained in the ambiguity set, following the paradigm that during execution the distribution of the state transition might change but still is contained in a pre-defined ambiguity set. In this paper we follow this paradigm and propose a novel numerical algorithm to solve Markov decision problems while accounting for distributional uncertainty by considering arbitrary finite ambiguity sets.

1.1. Our Contribution. One frequently used numerical method to solve (non-robust) Markov decision problems that has turned out to work efficiently in practice is the so-called *Q -learning algorithm* which initially was introduced by Watkins in his PhD thesis, see [46], as well as [47] for the accompanying paper. For a general introduction to Q -learning algorithms we refer to [11] and [19], and for more example of associated applications, see among many others [3], [4], [9], [10], [16], [21], and [26].

Our contribution is to present a novel Q -learning algorithm that allows to take distributional uncertainty for the state transitions of Markov decision problems into account and which therefore

respects the so called *Knighitian uncertainty* ([20]), i.e., the uncertainty of having chosen the correct model.

We respect this model uncertainty by, instead of fixing a unique probability distribution for the state transitions, allowing the state transition between two subsequent states to be realized according to any distribution from a set of predetermined transition probabilities and then to consider the worst-case measure to compute rewards. Moreover, to account for the fact that transition probabilities may change over time, we consider a time-inhomogeneous formulation, i.e., we allow different transition probabilities in every instance of time as long as the transition probabilities are contained in the ambiguity set. To solve this *robust* formulation of a Markov decision process (as introduced in [28]) under the worst-case measure we develop a novel Q -learning algorithm.

The finite ambiguity set of transition probabilities can be designed by the applicant and its shape is, in particular, not restricted to measures that are in a certain sense close to a reference measure. This allows to consider different ambiguity sets as it were possible in the case of ambiguity sets that are defined as balls around reference measures. Put differently, using our approach, an applicant can simply include any possible scenarios in which she is interested and hence is able to control the shape and content of the ambiguity sets and hence the dynamics of the underlying Markov decision process that are deemed to be admissible. Moreover, in Section 5 we provide a proof guaranteeing convergence of our algorithm.

1.2. Related Work. While the study of distributionally robust optimization (DRO) problems and distributionally robust Markov decision problems and problems has been an active research topic already in the past decade (compare, e.g., [5], [6], [15], [23], [28], [29], [30], [34], [35], [41], [45], [48], [49], [50], and [51]), the discussion and construction of Q -learning algorithms to solve these sequential decision making problems numerically has only become an active research topic very recently.

In [6], [22], and [44] Q -learning algorithms for ambiguity sets that are constructed as balls around some reference measures with respect to the Kullback–Leibler divergence are discussed.

Moreover, in [27], the authors present a novel Q -learning algorithm for the specific case that the ambiguity set is given by a Wasserstein-ball around a reference measure.

To the best of our knowledge going beyond the aforementioned cases imposes a novelty, and has not been studied yet.

1.3. Structure. The remainder of this paper is as follows. In Section 2 we present the setting and the associated stochastic control problem. A numerical solution to this optimization problem via Q -learning is presented in Section 3 together with our main result guaranteeing convergence of the Q -learning algorithm. In Section 4 we provide numerical experiments showcasing the tractability of our Q -learning algorithm. The proofs are reported in Section 5.

2. SETTING AND SPECIFICATION OF THE PROBLEM

In this section we introduce the underlying framework which we use to establish a robust Q -learning algorithm for finite ambiguity sets.

2.1. Setting. Optimal control problems, such as Markov decision problems, are formulated using a state space comprising all possible states accessible by an underlying stochastic process. We model this state space by a d -dimensional finite Euclidean subset $\mathcal{X} \subset \mathbb{R}^d$ and eventually aim at solving a robust control problem over an infinite time horizon. Hence, we define the state space over the entire time horizon via

$$\Omega := \mathcal{X}^{\mathbb{N}} = \mathcal{X} \times \mathcal{X} \times \dots$$

equipped with the corresponding σ -algebra $\mathcal{F} := 2^{\mathcal{X}} \otimes 2^{\mathcal{X}} \otimes \dots$. Next, let $(X_t)_{t \in \mathbb{N}}$ be the state process, i.e., the stochastic process on Ω describing the states attained over time. We denote the finite set of possible actions by $A \subset \mathbb{R}^m$, where $m \in \mathbb{N}$ represents the dimension of the action space. The set of admissible policies is then given by

$$\begin{aligned} \mathcal{A} := & \{ \mathbf{a} = (a_t)_{t \in \mathbb{N}} \mid (a_t)_{t \in \mathbb{N}} : \Omega \rightarrow A; a_t \text{ is } \sigma(X_t) \text{-measurable for all } t \in \mathbb{N} \} \\ & = \{ (a_t(X_t))_{t \in \mathbb{N}} \mid a_t : \mathcal{X} \rightarrow A \text{ Borel measurable for all } t \in \mathbb{N} \}. \end{aligned}$$

In contrast to classical non-robust Markov decision problems we work under the paradigm that the precise state transition probability is not known but instead contained in ambiguity set of finitely

many transition probabilities. In the following, let $\mathcal{M}_1(\mathcal{X})$ denote the set of probability measures on $(\mathcal{X}, \mathcal{F})$, and let τ_0 denote the topology of weak convergence¹. The finite ambiguity set of probability measures in dependence of a state-action pair is then modeled by a set-valued map given by

$$(2.1) \quad \begin{aligned} \mathcal{X} \times A &\rightarrow (\mathcal{M}_1(\mathcal{X})^N, \tau_0) \\ (x, a) &\mapsto \mathcal{P}(x, a) := \left\{ \mathbb{P}^{(1)}(x, a), \dots, \mathbb{P}^{(N)}(x, a) \right\}, \quad N \in \mathbb{N}, \end{aligned}$$

where for all $k \in \{1, \dots, N\}$ and for all $(x, a) \in \mathcal{X} \times A$ we have that $\mathbb{P}^{(k)}(x, a) \in \mathcal{M}_1(\mathcal{X})$. Note that in the degenerate case where $\mathcal{P}(x, a)$ contains only a single probability distribution, we get a classical Markov decision process (i.e. a non-robust one), compare, e.g., [7].

The ambiguity set of admissible probability distributions on Ω depends on the initial state $x \in \mathcal{X}$ and the policy $\mathbf{a} \in \mathcal{A}$. Let $\delta_x \in \mathcal{M}_1(\mathcal{X})$ denote the Dirac measure at point $x \in \mathcal{X}$. Then, we define for every state-action pair $(x, \mathbf{a}) \in \mathcal{X} \times \mathcal{A}$ the underlying set of admissible probability measures of the stochastic process $(X_t)_{t \in \mathbb{N}}$ by

$$(2.2) \quad \begin{aligned} \mathfrak{P}_{x, \mathbf{a}} &:= \left\{ \delta_x \otimes \mathbb{P}_0 \otimes \mathbb{P}_1 \otimes \dots \mid \text{for all } t \in \mathbb{N} : \mathbb{P}_t : \mathcal{X} \rightarrow \mathcal{M}_1(\mathcal{X}) \text{ Borel-measurable,} \right. \\ &\quad \left. \text{and } \mathbb{P}_t \in \mathcal{P}(x_t, a_t(x_t)), \text{ for all } x_t \in \mathcal{X} \right\}, \end{aligned}$$

where the notation $\mathbb{P} = \delta_x \otimes \mathbb{P}_0 \otimes \mathbb{P}_1 \otimes \dots \in \mathfrak{P}_{x, \mathbf{a}}$ abbreviates

$$\mathbb{P}(B) = \sum_{x_0 \in \mathcal{X}} \dots \sum_{x_t \in \mathcal{X}} \dots \mathbf{1}_B((x_t)_{t \in \mathbb{N}}) \dots \mathbb{P}_{t-1}(x_{t-1}; \{x_t\}) \dots \mathbb{P}_0(x_0; \{x_1\}) \delta_x(\{x_0\}), \quad B \in \mathcal{F}.$$

The construction of ambiguity sets $\mathfrak{P}_{x, \mathbf{a}}$ provided in (2.1) and (2.2) is in the related literature often referred to as a *(s, a)-rectangular ambiguity set*, (compare [17] and [48]) and enables to obtain a dynamic programming principle of the associated Markov decision problem.

After execution of an action $a_t(X_t)$, the agent receives a feedback on the quality of the chosen action in terms of a reward $r(X_t, a_t(X_t), X_{t+1})$ where r denotes some reward function $r : \mathcal{X} \times A \times \mathcal{X} \rightarrow \mathbb{R}$.

2.2. Finite ambiguity sets and other ambiguity sets. Note that the paradigm under which we work in this paper is fundamentally different from the approaches pursued, e.g., in [6], [22], and [27] where the corresponding ambiguity sets of probability measures are defined as balls around some reference measure.

The Q-learning approaches presented in [6], [22], and [27] provide solutions to account for a potential misspecification of a reference measure, by allowing deviations from it where the size of the deviations is measured by the respective distances (Kullback–Leibler distance and Wasserstein distance). However, these approaches allow not to control of which type the distributions in the ambiguity set are but simply consider all measures that are in a certain sense close to the reference measure. If an applicant is instead interested in controlling the distributions contained in the ambiguity set, for example, by allowing only for a specific type of parametric distributions with a finite set of possible parameters or by considering an asymmetric ambiguity set, the applicant can use the Q-learning approach presented in this paper. Another relevant situation is the consideration of multiple estimated models / probability measures obtained for example via different estimation methods. To account for all of these estimations, an applicant can simply construct an ambiguity set containing all estimated probability measures.

Beyond these use cases, our approach also allows to combine different types of distributions, if desired.

The main difference therefore is that an applicant of the approach presented in this paper can control exactly what types of distributions are deemed possible whereas the approaches from [6], [22], and [27] account for a misspecification of one single reference measure.

¹For any $\mu \in \mathcal{M}_1(X)$, $(\mu_n)_{n \in \mathbb{N}} \subseteq \mathcal{M}_1(X)$ and $C_b(X, \mathbb{R})$ denoting the space of continuous and bounded functions mapping from the space X to \mathbb{R} , we have:

$$\mu_n \xrightarrow{\tau_0} \mu \Leftrightarrow \lim_{n \rightarrow \infty} \int g d\mu_n = \int g d\mu, \quad \text{for all } g \in C_b(X, \mathbb{R}).$$

2.3. Specification of the optimization problem. The *robust* optimization problem consists, for every initial state $x \in \mathcal{X}$, in maximizing the expected value of $\sum_{t=0}^{\infty} \alpha^t r(X_t, a_t(X_t), X_{t+1})$ under the worst case measure from $\mathfrak{P}_{x,\mathbf{a}}$ over all possible policies $\mathbf{a} \in \mathcal{A}$, where $\alpha \in \mathbb{R}$ is some discount factor accounting for time preferences of rewards. Therefore, the value function

$$(2.3) \quad \mathcal{X} \ni x \mapsto V(x) := \sup_{\mathbf{a} \in \mathcal{A}} \inf_{\mathbb{P} \in \mathfrak{P}_{x,\mathbf{a}}} \mathbb{E}_{\mathbb{P}} \left[\sum_{t=0}^{\infty} \alpha^t \cdot r(X_t, a_t(X_t), X_{t+1}) \right]$$

describes the expected value of $\sum_{t=0}^{\infty} \alpha^t r(X_t, a_t(X_t), X_{t+1})$ under the worst case measure from $\mathfrak{P}_{x,\mathbf{a}}$ and when executing the optimal policy $\mathbf{a} \in \mathcal{A}$ after having started in initial state $x \in \mathcal{X}$.

2.4. Dynamic Programming. We consider the one step optimization problem

$$(2.4) \quad \mathcal{T}V(x) := \max_{a \in A} \min_{\mathbb{P} \in \mathcal{P}(x,a)} \mathbb{E}_{\mathbb{P}} [r(x, a, X_1) + \alpha V(X_1)], \quad x \in \mathcal{X}$$

where $\mathcal{X} \ni x \mapsto V(x)$ is the value function defined in (2.3). Analogue to the definition of the non-robust Q -value function, see e.g. [46], we define the robust optimal Q -value function by

$$(2.5) \quad \mathcal{X} \times A \ni (x, a) \mapsto Q^*(x, a) := \inf_{\mathbb{P} \in \mathcal{P}(x,a)} \mathbb{E}_{\mathbb{P}} [r(x, a, X_1) + \alpha V(X_1)].$$

We can interpret $Q^*(x, a)$ as the *quality* of executing action a when in state x (under the worst case measure) with the best possible action leading to $V(x)$. By using the above definitions we indeed obtain the following dynamic programming equation, also famously known in the non-robust formulation as *Bellman equation*, compare e.g. [12, p.164] for a non-robust version.

Proposition 2.1. *Assume $0 < \alpha < 1$. Then, for all $x \in \mathcal{X}$ we have*

$$\max_{a \in A} Q^*(x, a) = \mathcal{T}V(x) = V(x).$$

3. ROBUST Q -LEARNING ALGORITHM

In this section we propose with Algorithm 1 a novel distributionally robust Q -learning algorithm and provide in Theorem 3.1 conditions for its convergence.

Algorithm 1 Robust Q -learning for finite ambiguity sets

Input:

State space $\mathcal{X} \subseteq \mathbb{R}^d$; Initial state $x_0 \in \mathcal{X}$; Action space $A \subseteq \mathbb{R}^m$;
 Reward function r ; Discount factor $\alpha \in (0, 1)$;
 Ambiguity set $\mathcal{X} \times A \ni (x, a) \mapsto \mathcal{P}(x, a) := \{\mathbb{P}^{(1)}(x, a), \dots, \mathbb{P}^{(N)}(x, a)\}$;
 Policy $(a_t)_t \in \mathcal{A}$; Sequence of learning rates $(\tilde{\gamma}_t)_{t \in \mathbb{N}} \subseteq [0, 1]$;

Initialize $Q_0(x, a)$ for all $(x, a) \in \mathcal{X} \times A$ to an arbitrary real value;

Set $X_0 = x_0$;

for $t = 0, 1, \dots$ **do**

Determine an index $k_t^* \in \{1, \dots, N\}$ by

$$(3.1) \quad k_t^* \in \arg \min_{k=1, \dots, N} \mathbb{E}_{\mathbb{P}^{(k)}(X_t, a_t(X_t))} \left[r(X_t, a_t(X_t), X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) \right];$$

Sample $X_{t+1} \sim \mathbb{P}^{(k_t^*)}(X_t, a_t(X_t))$;

for all $(x, a) \in \mathcal{X} \times A$ **do**

Update Q_t via

(3.2)

$$Q_{t+1}(x, a) \leftarrow \begin{cases} (1 - \tilde{\gamma}_t) \cdot Q_t(x, a) + \tilde{\gamma}_t \cdot \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) \right) & \text{if } (x, a) = (X_t, a_t(X_t)), \\ Q_t(x, a); & \text{else} \end{cases}$$

end for

end for

Output: A sequence $(Q_t(x, a))_{t \in \mathbb{N}, x \in \mathcal{X}, a \in A}$;

Given a policy $\mathbf{a} \in \mathcal{A}$ and some initial state $x_0 \in \mathcal{X}$ we define a probability measure by

$$(3.3) \quad \mathbb{P}_{x_0, \mathbf{a}} := \delta_{x_0} \otimes \mathbb{P}_0^{(k_0^*)}(\cdot, a_0(\cdot)) \otimes \mathbb{P}_1^{(k_1^*)}(\cdot, a_1(\cdot)) \otimes \cdots \in \mathfrak{P}_{x_0, \mathbf{a}},$$

where k_t^* is as defined in (3.1). Note that the probability measure defined in (3.3) is the probability measure according to which the states are sampled when applying Algorithm 1.

Our following main result now shows that the function $(Q_t)_{t \in \mathbb{N}}$ obtained as the output of Algorithm 1 converges pointwise almost surely towards the optimal robust Q -value function Q^* defined in (2.5).

Theorem 3.1. *Let $(\tilde{\gamma}_t)_{t \in \mathbb{N}} \subseteq [0, 1]$, and define for all $(x, a) \in \mathcal{X} \times A$ and for all $t \in \mathbb{N}$*

$$(3.4) \quad \gamma_t(x, a, X_t) := \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \cdot \tilde{\gamma}_t.$$

Moreover, let $0 < \alpha < 1$ and let $(x_0, \mathbf{a}) \in \mathcal{X} \times \mathcal{A}$ such that

$$(3.5) \quad \sum_{t=0}^{\infty} \gamma_t(x, a, X_t) = \infty, \quad \sum_{t=0}^{\infty} \gamma_t^2(x, a, X_t) < \infty, \quad \text{for all } (x, a) \in \mathcal{X} \times A, \quad \mathbb{P}_{x_0, \mathbf{a}} \text{ - almost surely.}$$

Then, we have for all $(x, a) \in \mathcal{X} \times A$ that

$$(3.6) \quad \lim_{t \rightarrow \infty} Q_t(x, a) = Q^*(x, a) \quad \mathbb{P}_{x_0, \mathbf{a}} \text{ - almost surely.}$$

Remark 3.2 (On the implementation).

- (a) Note that in many situations one can improve the convergence speed by replacing $\tilde{\gamma}_t$ with $\tilde{\gamma}_{\text{visits}_t(x, a)}$, where we initially set $\text{visits}_0 \equiv 0$, and then for all $(x, a) \in \mathcal{X} \times A$ and $t = 1, 2, \dots$,

$$\text{visits}_t(x, a) \leftarrow \begin{cases} \text{visits}_t(x, a) + 1 & \text{if } (x, a) = (X_t, a_t(X_t)), \\ \text{visits}_t(x, a) & \text{else,} \end{cases}$$

compare, e.g., [24]. The idea behind this modification is to account for the exploration-exploitation trade off by reducing the learning rate the more often we have visited a state and hence the more confident the algorithm is w.r.t. choosing the correct optimal action, and conversely to apply a larger learning rate if the state-action pair was visited less often until time t .

- (b) Note that applying Algorithm 1 requires to choose a pre-determined policy $\mathbf{a} \in \mathcal{A}$. Implied from usual practice in non-robust Q -learning ([13], Chapter 9), [25], or [40]) and robust Q -learning [27], a reasonable choice seems to be the ϵ -greedy policy defined by

$$\mathcal{X} \ni x \mapsto a_t(x) := \begin{cases} \operatorname{argmax}_{b \in B} Q_t(x, b) & \text{with probability } 1 - \epsilon_{\text{greedy}}, \\ a \sim \mathcal{U}(A) & \text{with probability } \epsilon_{\text{greedy}}, \end{cases}$$

for some $\epsilon_{\text{greedy}} > 0$, and where $a \sim \mathcal{U}(A)$ means that a random action a is chosen uniformly at random from the finite set A .

- (c) Due to Equation (3.1), increasing the number of probability measures N linearly increases also the computation time linearly as bloating the ambiguity set by a factor N corresponds to computing N times as many expectations of the form $\mathbb{E}_{\mathbb{P}^{(k)}(X_t, a_t(X_t))}[r(X_t, a_t(X_t), X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b)]$ in order to determine k_t^* .

4. EXAMPLES

In this section we provide two numerical examples comparing Algorithm 1 with other robust and non-robust Q -learning algorithms. To apply the numerical method from Algorithm 1, we use for all of the following examples a sequence of learning rates defined by $\tilde{\gamma}_t = \frac{1}{1+t}$ for $t \in \mathbb{N}$, a discount factor of $\alpha = 0.95$, as well as an ϵ -greedy policy with $\epsilon_{\text{greedy}} = 0.1$. Finally, we run² Algorithm 1 with 1 000 000 iterations on a processor: 13th Gen Intel(R) Core(TM) i7-13700KF 3.40 GHz. Further details of the implementation can be found under <https://github.com/CecileDecker/FiniteQLearning.git>.

²We consider the algorithm as being converged if further iterations do not change the optimal action derived from Q -learning, i.e. if $\operatorname{argmax}_{b \in A} Q_t(x, b)$ remains constant for further iterations t for all $x \in \mathcal{X}$. This was the case for all presented experiments.

4.1. **Coin Toss.** As in [28, Example 4.1] we consider an agent playing a coin toss game.

At each time step $t \in \mathbb{N}$ the agent observes the result of 10 coins where an outcome of heads corresponds to 1, and tails corresponds 0. The state X_t at time $t \in \mathbb{N}$ is then given by the sum of the 10 coins value, i.e., we have $\mathcal{X} := \{0, \dots, 10\}$.

At each time step t the agent can make a bet whether the sum of the next throw strictly exceeds the previous sum (i.e. $X_{t+1} > X_t$), or whether it is strictly smaller (i.e. $X_{t+1} < X_t$). If the agent is correct, she gets 1\$, however if the agent is wrong she has to pay 1\$. The agent also has the possibility not to play. We model this by considering the following reward function:

$$(4.1) \quad \mathcal{X} \times A \times \mathcal{X} \ni (x, a, x') \mapsto r(x, a, x') := a\mathbb{1}_{\{x < x'\}} - a\mathbb{1}_{\{x > x'\}} - |a|\mathbb{1}_{\{x = x'\}},$$

where the possible actions are given by $A := \{-1, 0, 1\}$ with $a = 1$ corresponding to betting $X_{t+1} > X_t$, $a = 0$ to not playing, and $a = -1$ to betting $X_{t+1} < X_t$. In a next step, we define two different ambiguity sets via³

$$(4.2) \quad \mathcal{P}^1(x, a) := \{\text{Bin}(10, 0.5), \text{Bin}(10, 0.6)\},$$

$$(4.3) \quad \mathcal{P}^2(x, a) := \{\text{Bin}(10, 0.5), \text{Bin}(10, 0.3)\},$$

and we aim at comparing Algorithm 1 with the Algorithm from [28] which however does not allow to build the same asymmetric sets as in (4.2) and (4.3). To build comparative ambiguity sets with Wasserstein uncertainty we define ambiguity sets such that all probability measures from \mathcal{P}^1 and \mathcal{P}^2 are contained, respectively. This leads to⁴

$$(4.4) \quad \mathcal{P}^3(x, a) := \{\mathbb{P} \in \mathcal{M}_1(\mathcal{X}) \mid W_1(\mathbb{P}, \text{Bin}(10, 0.5)) \leq 1\}$$

$$(4.5) \quad \mathcal{P}^4(x, a) := \{\mathbb{P} \in \mathcal{M}_1(\mathcal{X}) \mid W_1(\mathbb{P}, \text{Bin}(10, 0.5)) \leq 2\}$$

which are Wasserstein-balls around the *reference probability* $\text{Bin}(10, 0.5)$ with radii 1 and 2, respectively. As one can see, e.g., from [27, Equation (4.2)], \mathcal{P}^3 is the smallest Wasserstein-ball around the reference measure that contains $\text{Bin}(10, 0.6)$, whereas \mathcal{P}^4 is the smallest Wasserstein-ball containing $\text{Bin}(10, 0.3)$.

We depict the trained actions and the associated convergence time (duration of running 1 000 000 iterations of the algorithm) in Table 1.

X_t	0	1	2	3	4	5	6	7	8	9	10	Convergence time
$a_t^{\mathcal{P}^1}(X_t)$	1	1	1	1	1	0	0	-1	-1	-1	-1	19 min 54.3 sec
$a_t^{\mathcal{P}^2}(X_t)$	1	1	1	0	0	0	-1	-1	-1	-1	-1	19 min 2.5 sec
$a_t^{\mathcal{P}^3}(X_t)$	1	1	1	1	0	0	0	-1	-1	-1	-1	56 min 42.8 sec
$a_t^{\mathcal{P}^4}(X_t)$	1	1	1	0	0	0	0	0	-1	-1	-1	56 min 34.3 sec
$a_t^{\text{non-robust}}(X_t)$	1	1	1	1	1	0	-1	-1	-1	-1	-1	5.2 sec

TABLE 1. The trained actions $a_t^{\mathcal{P}^1}$, $a_t^{\mathcal{P}^2}$, $a_t^{\mathcal{P}^3}$, $a_t^{\mathcal{P}^4}$, $a_t^{\text{non-robust}}$, in dependence of the realized state X_t

We next test the profit of the actions described in Table 1 by playing 100 000 rounds of the game according to the trained policies $a_t^{\mathcal{P}^1}$, $a_t^{\mathcal{P}^2}$, $a_t^{\mathcal{P}^3}$ and $a_t^{\mathcal{P}^4}$. For simulating the 100 000 rounds we assume an underlying binomial distribution $\mathbb{P} = \text{Bin}(10, p_{\text{true}})$ with a fixed probability p_{true} for heads which we vary from 0.1 to 0.9. We depict the cumulative profits of the considered actions in Table 3.

We first observe that only in the case $p_{\text{true}} = 0.5$ the non-robust strategy is the best performing strategy. The Wasserstein-strategies are outperformed by the robust strategies taking into account either \mathcal{P}_1 or \mathcal{P}_2 in all cases for which $p_{\text{true}} \leq 0.6$, in particular for the cases $p_{\text{true}} = 0.3, 0.6$ which are by construction contained in the respective ambiguity sets.

³We denote by $\text{Bin}(n, p)$ a binomial distribution with n number of trials and p being the probability of success in a single trial.

⁴Here W_1 denotes the Wasserstein-1 distance, compare for more details, e.g., [43].

p_{true}	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
Robust, \mathcal{P}^1	-31240	-19121	-4283	15161	30484	29857	13212	-10339	-29843
Robust, \mathcal{P}^2	-24424	5231	21051	24613	22998	11715	-4984	-18896	-31185
Robust, \mathcal{P}^3	-30003	-10859	9843	21900	25395	22587	9653	-10656	-30301
Robust, \mathcal{P}^4	-24528	4263	16771	13415	9925	13366	17345	4731	-24452
Non-Robust	-31101	-18251	-521	23175	35244	23227	-1387	-18421	-31024

TABLE 2. Overall profit of the game described in Section 4.1 in dependence of different trained strategies (described in Table 1) and of the probability distribution $P = \text{Bin}(10, p_{\text{true}})$ of the simulated underlying process. We indicate with bold characters which is the best performing strategy for each choice of p_{true} .

The results imply that the choice of the *optimal* ambiguity sets depends on the scenarios that are deemed possible: If we know that⁵ $p_{\text{true}} \in \{0.5, 0.6\}$, then one should exactly consider \mathcal{P}^1 , whereas in that situation the set \mathcal{P}^3 that also covers the case $p_{\text{true}} = 0.6$ will lead to a suboptimal because too conservative strategy that also covers the case $p_{\text{true}} = 0.4$ as a possible scenario.

If an applicant however simply has a *best guess* of a reference measure and considers the cases $p_{\text{true}} = 0.4$ and $p_{\text{true}} = 0.6$ both as equally possible, then it is advisable to pursue the Wasserstein-approach outlined in [27].

4.2. Stock investing. This example is a variant of [28, Example 4.3] showcasing the use of tailored ambiguity sets which are not possible to construct and use in the same manner with Wasserstein or Entropic ambiguity sets as they are considered in [6], [22], and [44].

We consider the problem of optimally investing in the stock market, and, to this end, encode the return of a stock by 2 numeric values: either the return is positive (1) or negative (-1). The space of numerically encoded returns is therefore given by: $T := \{-1, 1\}$. At each time step t the agent can choose to buy the stock or not, or to short sell the stock. The action a_t represent this investment decision, buying the stock is encoded by 1 and not buying it by 0, whereas short-selling corresponds to -1, i.e., we have $A := \{-1, 0, 1\}$.

The agent's investment decision should depend not only on the most recent return but naturally will depend on the current *market situation*. Therefore the agent relies her investment decision on the last $h = 5$ returns. Hence, we consider the state space

$$\mathcal{X} := T^h = \{-1, 1\}^h.$$

The reward is given by the function:

$$(4.6) \quad \mathcal{X} \times A \times \mathcal{X} \ni (x, a, x') \mapsto r(x, a, x') := a \cdot x',$$

i.e., we reward a correct investment decision and penalize an incorrect investment decision.

Now, to train the agent, we construct an estimated reference measure in the following way. For the state transition, we consider the historic evolution of the (numerically encoded) returns of the underlying stock. This time series is denoted by $(R_j)_{j=1, \dots, N} \subset T^N$ for some $N \in \mathbb{N}$. We then define for some small $\gamma > 0$ a map⁶ $\mathcal{X} \times A \ni (x, a) \mapsto \tilde{\mathbb{P}}(x, a) := \sum_{i \in T} p_i(x) \cdot \delta_{\{i\}} \in \mathcal{M}_1(T)$ where for $x \in \mathcal{X}$, $i \in T$ we define:

$$(4.7) \quad p_i(x) := \frac{\tilde{p}_i(x) + \frac{\gamma}{4}}{\gamma + \sum_{j \in T} \tilde{p}_j(x)} \in [0, 1],$$

with

$$(4.8) \quad \tilde{p}_i(x) := \sum_{j=1}^{X-h+1} \mathbb{1}_{\{(\pi(x), i) = (R_j, \dots, R_{j+h-1})\}},$$

and where $\mathbb{R}^h \ni (x_1, \dots, x_h) \mapsto \pi(x_1, \dots, x_h) := (x_2, \dots, x_h) \in \mathbb{R}^{h-1}$ denotes the projection onto the last $h-1$ components. This means the data-driven construction of the measure $\tilde{\mathbb{P}}(x, a)$ relies on the relative frequency of the sequence $(\pi(x), i)$ in the time series of past realized returns $(R_j)_{j=1, \dots, N}$

⁵This would correspond to the case that the dealer in a casino does not tell whether she uses a biased coin (for which we know that $p_{\text{true}} = 0.6$ for playing the game with an agent or if she uses a fair coin.

⁶We only employ γ to avoid division by 0, hence γ can be chosen arbitrarily small.

for $i \in \{-1, 1, \}$: the more often the *market situation* $(\pi(x), i) \in \mathcal{X}$ was observed in the training data $(R_j)_{j=1, \dots, N}$ the higher the value \tilde{p}_i . Equation (4.7) is then applied to convert the frequencies \tilde{p}_i to probabilities p_i . Then, the state transition of the reference measure is partially deterministic ($h - 1$ components of subsequent states coincide) and is of the form

$$(4.9) \quad X \times A \ni (x, a) \mapsto \mathbb{P}(x, a) = \delta_{\pi(x)} \otimes \tilde{\mathbb{P}}(x, a) \in \mathcal{M}_1(\mathcal{X}).$$

To construct the probability measure from (4.9) using real data, we consider daily returns of the stock of *Alphabet Inc* (previously known as Google) in the time period from beginning of 2010 until 2024, where we depict the cumulative sum of the signs of the returns in Figure 1.




FIGURE 1. Cumulative sum of the sign of the returns and the decomposition of the data in training periods and testing periods.

We split the data into a training period ranging from January 2010 until December 2021 and two testing period starting directly thereafter until February 2023 and January 2024, respectively. We then first construct a measure \mathbb{P}_1 according to (4.9) by using all training data. To construct an ambiguity set \mathcal{P} we isolate specific periods of the training data where the market behaved very volatile (Training Period 1), bearish (Training Period 2) or bullish (Training Period 3). Including the accompanying probability measures $\mathbb{P}_2, \mathbb{P}_3, \mathbb{P}_4$ in an ambiguity set $\mathcal{P} := \{\mathbb{P}_1, \mathbb{P}_2, \mathbb{P}_3, \mathbb{P}_4\}$ then accounts for the fear of the agent to not trade profitably in these difficult market scenarios. The choice of these specific training periods is depicted in Figure 1. Using Algorithm 1, we then train different agents using the ambiguity set \mathcal{P} and using the non-robust state transitions $\mathbb{P}_i, i = 1, 2, 3, 4$, respectively. Moreover, we consider a trend following strategy (i.e., always invest according to the sign of the last return) as well as a simple buy-and-hold strategy. The results (mean average reward per trade) of the different strategies evaluated in the test period are shown in Table 3, and they show that using a robust strategy indeed enables to avoid losses in both test periods. Note moreover that in the bearish test period 1 the robust strategy in particular outperforms the non-robust agent trained on all data which is certainly due to the inclusion of \mathbb{P}_2 and \mathbb{P}_3 in the ambiguity set \mathcal{P} which are the best performing strategies in this period. Also note that in test period 2 it becomes visible that the robust approach due to its construction of a worst-case approach is better suited to avoid losses than to profit from bearish market scenarios.

5. PROOFS AND AUXILIARY RESULTS

5.1. Auxiliary Results. For any function $f : \mathcal{X} \times A \rightarrow \mathbb{R}$, we write

$$(5.1) \quad \|f\|_\infty := \sup_{x \in \mathcal{X}} \sup_{a \in A} |f(x, a)|.$$

Next, we introduce the operator \mathcal{H} operating on a function $v : \mathcal{X} \times A \rightarrow \mathbb{R}$ being defined by

$$(5.2) \quad \mathcal{X} \times A \ni (x, a) \mapsto \mathcal{H}v(x, a) := \inf_{\mathbb{P} \in \mathcal{P}(x, a)} \mathbb{E}_{\mathbb{P}} \left[r(x, a, X_1) + \alpha \max_{b \in A} v(X_1, b) \right].$$

	Test Period 1	Test Period 2
Robust (\mathcal{P})	0.0205	0.0521
Non-Robust: All Data (\mathbb{P}_1)	-0.0274	0.0729
Non-Robust: Training Period 1 (\mathbb{P}_2)	0.0377	0.0885
Non-Robust: Training Period 2 (\mathbb{P}_3)	0.0685	0.1042
Non-Robust: Training Period 3 (\mathbb{P}_4)	-0.0034	0.1458
Trend following	-0.0137	0.0781
Buy and hold	-0.0822	0.1146

TABLE 3. A comparison of the average reward per trade in the two test periods depicted in Figure 1 among different strategies. The best-performing strategy is highlighted with bold letters.

Lemma 5.1. *Let $0 < \alpha < 1$, and let the ambiguity set \mathcal{P} be defined in (2.1). Then the following fixed-point equation holds true for the optimal Q-value function defined in (2.5):*

$$\mathcal{H}Q^*(x, a) = Q^*(x, a) \text{ for all } (x, a) \in \mathcal{X} \times A.$$

Proof. This follows directly by definition of Q^* and by Proposition 2.1, compare also [27, Proof of Lemma 17]. \square

Lemma 5.2. *For any maps $q_i : \mathcal{X} \times A \rightarrow \mathbb{R}$, $i = 1, 2$, we have:*

$$\|\mathcal{H}q_1 - \mathcal{H}q_2\|_\infty \leq \alpha \|q_1 - q_2\|_\infty.$$

Proof. The proof follows is analogue to [27, Proof of Lemma 18]. \square

The following lemma is a result from stochastic approximation theory adjusted to our setting, compare [27, Lemma A.1] for the presented formulation adjusted to the underlying setting. We refer also to [36, Lemma 1], [14] [18, Theorem 1], [38, Lemma 12], and [42, Lemma 3] for further reference.

Lemma 5.3 ([36], Lemma 1). *Let $\mathbb{P}_0 \in \mathcal{M}_1(\Omega)$ be a probability measure on (Ω, \mathcal{F}) , and consider a family of stochastic processes $(\gamma_t(x, a), F_t(x, a), \Delta_t(x, a))_{t \in \mathbb{N}_0}$, $(x, a) \in \mathcal{X} \times A$, satisfying for all $t \in \mathbb{N}_0$*

$$\Delta_{t+1}(x, a) = (1 - \gamma_t(x, a)) \Delta_t(x, a) + \gamma_t(x, a) F_t(x, a) \quad \mathbb{P}_0\text{-almost surely for all } (x, a) \in \mathcal{X} \times A.$$

Let $(\mathcal{G}_t)_{t \in \mathbb{N}_0} \subseteq \mathcal{F}$ be a sequence of increasing σ -algebras such that for all $(x, a) \in \mathcal{X} \times A$ the random variables $\Delta_0(x, a)$ and $\gamma_0(x, a)$ are \mathcal{G}_0 -measurable and such that $\Delta_t(x, a)$, $\gamma_t(x, a)$, and $F_{t-1}(x, a)$ are \mathcal{G}_t -measurable for all $t \in \mathbb{N}$. Further assume that the following conditions hold.

- (i) $0 \leq \gamma_t(x, a) \leq 1$, $\sum_{t=0}^{\infty} \gamma_t(x, a) = \infty$, $\sum_{t=0}^{\infty} \gamma_t^2(x, a) < \infty$ \mathbb{P}_0 -almost surely for all $(x, a) \in \mathcal{X} \times A$, $t \in \mathbb{N}_0$.
- (ii) There exists $\delta \in (0, 1)$ such that $\|\mathbb{E}_{\mathbb{P}_0} [F_t(\cdot, \cdot) \mid \mathcal{G}_t]\|_\infty \leq \delta \|\Delta_t\|_\infty$ \mathbb{P}_0 -almost surely for all $t \in \mathbb{N}_0$.
- (iii) There exists $C > 0$ such that $\|\text{Var}_{\mathbb{P}_0} (F_t(\cdot, \cdot) \mid \mathcal{G}_t)\|_\infty \leq C(1 + \|\Delta_t\|_\infty)^2$ \mathbb{P}_0 -almost surely for all $t \in \mathbb{N}_0$.

Then, $\lim_{t \rightarrow \infty} \Delta_t(x, a) = 0$ \mathbb{P}_0 -almost surely for all $(x, a) \in \mathcal{X} \times A$.

The following result is usually referred to as Popoviciu's inequality, see [31] or [33].

Lemma 5.4. *Let Z be a random variable on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ satisfying $m \leq Z \leq M$ for some $-\infty < m \leq M < \infty$. Then, we have*

$$\text{Var}_{\mathbb{P}}(Z) \leq \frac{1}{4} (M - m)^2.$$

5.2. Proofs of the results from Section 2 and Section 3. In this section we provide the proofs of Section 2.4 and Section 3.

Proof of Proposition 2.1. Note that the equality $\max_{a \in A} Q^*(x, a) = \mathcal{T}V(x)$ directly follows by definition of the operator \mathcal{T} , and it only remains to show $\mathcal{T}V = V$. To this end, we aim at applying [28, Theorem 3.1] with $p = 0$ and verify its assumptions, i.e., we verify [28, Assumption 2.2] and [28, Assumption 2.4] by showing the following four properties.

(i) The set-valued map

$$\begin{aligned} \mathcal{X} \times A &\rightarrow (\mathcal{M}_1(\mathcal{X})^N, \tau_0) \\ (x, a) &\rightarrow \mathcal{P}(x, a) := \left\{ \mathbb{P}^{(1)}(x, a), \dots, \mathbb{P}^{(N)}(x, a) \right\} \end{aligned}$$

is nonempty, compact-valued and continuous⁷.

(ii) There exists $C_P \geq 1$ such that for all $(x, a) \in \mathcal{X} \times A, \mathbb{P} \in \mathcal{P}(x, a)$ we have

$$(5.3) \quad \int_{\mathcal{X}} (1 + \|y\|^p) \mathbb{P}(dy) \leq C_P (1 + \|x\|^p)$$

(iii) The reward function $r : \mathcal{X} \times A \times \mathcal{X} \rightarrow \mathbb{R}$ is Lipschitz-continuous in its first two components.

(iv) We have $0 < \alpha < \frac{1}{C_P}$, where C_P is defined in (ii).

To see that (i) holds, we note that \mathcal{P} is by definition non-empty and since the set $\mathcal{P}(x, a)$ is finite it is also compact for all $(x, a) \in \mathcal{X} \times A$. To show the upper hemicontinuity, we consider some $(x, a) \in \mathcal{X} \times A$ and we let $(x_n, a_n, \mathbb{P}_n)_{n \in \mathbb{N}} \subseteq \text{Gr}(\mathcal{P}) := \{(x, a, \mathbb{P}) \mid (x, a) \in \mathcal{X} \times A, \mathbb{P} \in \mathcal{P}(x, a)\}$ be a sequence such that $(x_n, a_n) \rightarrow (x, a) \in \mathcal{X} \times A$ as $n \rightarrow \infty$. Since both, \mathcal{X} and A are finite, we have $(x_n, a_n) = (x, a)$ for n large enough. Hence, $\mathbb{P}_n \in \mathcal{P}(x, a)$ for n large enough. As $\mathcal{P}(x, a)$ is finite we also have that $\mathbb{P}_n = \mathbb{P}^{(i)}(x, a)$ for infinitely many indices n for some $i \in \{1, \dots, N\}$. Hence, we can find a subsequence $(\mathbb{P}_{n_k})_{k \in \mathbb{N}}$ such that $\mathbb{P}_{n_k} \rightarrow \mathbb{P}^{(i)}(x, a) \in \mathcal{P}(x, a)$ weakly as $k \rightarrow \infty$ and the upper hemicontinuity of \mathcal{P} follows by the characterization provided in [2, Theorem 17.20].

To show the lower hemicontinuity let $(x, a) \in \mathcal{X} \times A, \mathbb{P} \in \mathcal{P}(x, a)$ and consider a sequence $(x_n, a_n)_{n \in \mathbb{N}} \subseteq \mathcal{X} \times A$ with $(x_n, a_n) \rightarrow (x, a)$ as $n \rightarrow \infty$. By definition of \mathcal{P} , there exists some $i \in \{1, \dots, N\}$ such that $\mathbb{P}^{(i)}(x, a) = \mathbb{P}$. We use this observation to define

$$\mathbb{P}_n := \mathbb{P}^{(i)}(x_n, a_n) \in \mathcal{P}(x_n, a_n), \quad n \in \mathbb{N},$$

and we obtain, since $(x_n, a_n) = (x, a)$ for n large enough that $\mathbb{P}_n \rightarrow \mathbb{P}$ weakly which shows the lower hemicontinuity of \mathcal{P} according to [2, Theorem 17.21].

To show (ii), note that $p = 0$, and define $C_P := 1$, then we have

$$\int_{\mathcal{X}} (1 + \|y\|^0) \mathbb{P}(dy) = 2 = C_P \cdot (1 + \|x\|^0)$$

for all $(x, a) \in \mathcal{X} \times A$. Next, (iii) follows since for all $x_0, x'_0, x_1 \in \mathcal{X}$ and $a, a' \in A$ we have

$$|r(x_0, a, x_1) - r(x'_0, a', x_1)| \leq \left(\max_{\substack{y_0, y'_0 \in \mathcal{X}, \\ (y_0, b) \neq (y'_0, b')}} \frac{|r(y_0, b, x_1) - r(y'_0, b', x_1)|}{\|y_0 - y'_0\| + \|b - b'\|} \right) \cdot (\|x_0 - x'_0\| + \|a - a'\|).$$

Finally, (iv) follows since $C_P = 1$ and as we have $0 < \alpha < 1$ by assumption. \square

Proof of Theorem 3.1. To establish convergence of the Q -learning algorithm from Algorithm 1, we aim at applying Lemma 5.3. To verify the assumptions of Lemma 5.3 we first note that for each $t \in \mathbb{N}$ and $(x, a) \in \mathcal{X} \times A$ the update rule from (3.2) can be written equivalently as

$$(5.4) \quad Q_{t+1}(x, a) = Q_t(x, a) + \gamma_t(x, a, X_t) \cdot \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q_t(x, a) \right);$$

where for $(x, a) \in \mathcal{X} \times A$ we recall the definition

$$(5.5) \quad \gamma_t(x, a, X_t) := \mathbf{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \cdot \tilde{\gamma}_t.$$

Next, for for each $t \in \mathbb{N}$ and $(x, a) \in \mathcal{X} \times A$ we define

$$\begin{aligned} \Delta_t(x, a) &:= Q_t(x, a) - Q^*(x, a), \\ F_t(x, a) &:= \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \right) \mathbf{1}_{\{(x, a) = (X_t, a_t(X_t))\}}. \end{aligned}$$

Moreover, we recall that as defined in Algorithm 1 we have

$$k_t^* \in \arg \min_{k=1, \dots, N} \mathbb{E}_{\mathbb{P}^{(k)}(X_t, a_t(X_t))} \left[r(X_t, a_t(X_t), X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) \right]$$

⁷Continuity of a set-valued map means the map is both upper hemicontinuous and lower hemicontinuous.

Once we have verified that the three assumptions of Lemma 5.3 are fulfilled, we obtain $\Delta_t(x, a) \rightarrow 0$ $\mathbb{P}_{x_0, \mathbf{a}}$ -almost surely as $t \rightarrow \infty$ and hence

$$Q_t(x, a) \rightarrow Q(x, a) \quad \mathbb{P}_{x_0, \mathbf{a}}\text{-almost surely as } t \rightarrow \infty$$

implying the assertion. Thus, to apply Lemma 5.3, we compute for all $(x, a) \in \mathcal{X} \times A$ that

$$\begin{aligned} \Delta_{t+1}(x, a) &= Q_t(x, a) - Q^*(x, a) + \gamma_t(x, a, X_t) \cdot \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q_t(x, a) \right) \\ &= \Delta_t(x, a) \\ &\quad + \gamma_t(x, a, X_t) \cdot \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q_t(x, a) + Q^*(x, a) - Q^*(x, a) \right) \\ &= (1 - \gamma_t(x, a, X_t)) \cdot \Delta_t(x, a) \\ &\quad + \gamma_t(x, a, X_t) \cdot \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \right) \\ &= (1 - \gamma_t(x, a, X_t)) \cdot \Delta_t(x, a) + \gamma_t(x, a, X_t) \cdot F_t(x, a), \end{aligned}$$

where we used in the last step that the indicator $\mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}}$ appears both in the definition of F_t and γ_t and that the square of the indicator function is the indicator function itself. Consider the filtration $(\mathcal{G}_t)_{t \in \mathbb{N}}$ with

$$\mathcal{G}_t := \sigma(X_1, X_2, \dots, X_t), \quad t \in \mathbb{N},$$

and $\mathcal{G}_0 := \{\emptyset, \Omega^N\}$ being the trivial sigma-algebra. Then, by definition we have for all $t \in \mathbb{N}$ and for all $(x, a) \in \mathcal{X} \times A$ that the random variables $\Delta_t(x, a)$, $\gamma_t(x, a)$, $F_{t-1}(x, a)$ are \mathcal{G}_t -measurable. Next, let $(x, a) \in \mathcal{X} \times A$, and $t \in \mathbb{N}$. Then, we have by the definition of $\mathbb{P}_{x_0, \mathbf{a}}$ in (3.3) $\mathbb{P}_{x_0, \mathbf{a}}$ -almost surely that

$$\begin{aligned} &|\mathbb{E}_{\mathbb{P}_{x_0, \mathbf{a}}}[F_t(x, a) | \mathcal{G}_t]| \\ &= \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \left| \mathbb{E}_{\mathbb{P}_{x_0, \mathbf{a}}} \left[r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \mid \mathcal{G}_t \right] \right| \\ &= \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \left| \mathbb{E}_{\mathbb{P}^{(k_t^*)}(x, a)} \left[r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \right] \right| \\ &= \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \left| \min_{i=1, \dots, N} \left(\mathbb{E}_{\mathbb{P}^{(i)}(x, a)} \left[r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) \right] \right) - Q^*(x, a) \right| \\ &= \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \left| \inf_{\mathbb{P} \in \mathcal{P}(x, a)} \mathbb{E}_{\mathbb{P}} \left[r(x, a, X_1) + \alpha \max_{b \in A} Q_t(X_1, b) \right] - Q^*(x, a) \right|. \end{aligned}$$

By using the definition of the operator \mathcal{H} from (5.2) we therefore obtain

$$|\mathbb{E}_{\mathbb{P}_{x_0, \mathbf{a}}}[F_t(x, a) | \mathcal{G}_t]| = \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} |\mathcal{H}Q_t(x, a) - Q^*(x, a)|.$$

Consequently, applying Lemma 5.1 we have

$$(5.6) \quad |\mathbb{E}_{\mathbb{P}_{x_0, \mathbf{a}}}[F_t(x, a) | \mathcal{G}_t]| = \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} |\mathcal{H}Q_t(x, a) - \mathcal{H}Q^*(x, a)|.$$

Now using (5.6) and applying Lemma 5.2 we can conclude

$$\|\mathbb{E}_{\mathbb{P}_{x_0, \mathbf{a}}}[F_t(\cdot, \cdot) | \mathcal{G}_t]\|_{\infty} \leq \|\mathcal{H}Q_t - \mathcal{H}Q^*\|_{\infty} \leq \alpha \|Q_t - Q^*\|_{\infty} = \alpha \|\Delta_t\|_{\infty}$$

showing Lemma 5.3 (ii).

Next, to show that the assumption from Lemma 5.3 (iii) is fulfilled, we observe that for all $(x, a) \in \mathcal{X} \times A$ and $t \in \mathbb{N}$ we have

$$\begin{aligned}
& \text{Var}_{\mathbb{P}_{x_0, \mathbf{a}}}(F_t(x, a) | \mathcal{G}_t) \\
&= \mathbb{1}_{\{(x, a) = (X_t, a_t(X_t))\}} \cdot \text{Var}_{\mathbb{P}^{(k_t^*)}(x, a)} \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \right) \\
(5.7) \quad &\leq \text{Var}_{\mathbb{P}^{(k_t^*)}(x, a)} \left(- \left(r(x, a, X_{t+1}) + \alpha \max_{b \in A} Q_t(X_{t+1}, b) - Q^*(x, a) \right) \right) \\
&= \text{Var}_{\mathbb{P}^{(k_t^*)}(x, a)} \left(-r(x, a, X_{t+1}) - \alpha \max_{b \in A} Q_t(X_{t+1}, b) \right) \\
&= \text{Var}_{\mathbb{P}^{(k_t^*)}(x, a)} \left(-r(x, a, X_{t+1}) - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \min_{y' \in \mathcal{X}} \max_{b' \in A} Q^*(y', b') \right).
\end{aligned}$$

We define $C_r := \max_{y_0, y_1 \in \mathcal{X}, b \in A} |r(y_0, b, y_1)| < \infty$ and compute as an upper bound for the integrand from (5.7) that

$$\begin{aligned}
& -r(x, a, X_{t+1}) - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \min_{y' \in \mathcal{X}} \max_{b' \in A} Q^*(y', b') \\
&\leq C_r - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \max_{b' \in A} Q^*(X_{t+1}, b') \\
&\leq C_r + \alpha \max_{y \in \mathcal{X}} (\max_{b' \in A} Q^*(y, b') - \max_{b \in A} Q_t(y, b)) \\
&\leq C_r + \alpha \max_{y \in \mathcal{X}} \max_{b \in A} |Q^*(y, b) - Q_t(y, b)| \\
&\leq C_r + \alpha \|\Delta_t\|_\infty =: M \in \mathbb{R}.
\end{aligned}$$

On the other hand we also may compute a lower bound as follows

$$\begin{aligned}
& -r(x, a, X_{t+1}) - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \min_{y' \in \mathcal{X}} \max_{b' \in A} Q^*(y', b') \\
&\geq -C_r - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \min_{y' \in \mathcal{X}} \max_{b' \in A} Q^*(y', b') \\
&\geq -C_r + \alpha (\min_{y' \in \mathcal{X}} \min_{b \in A} (Q^*(y', b) - Q_t(X_{t+1}, b))) \\
&\geq -C_r + \alpha (- \max_{y' \in \mathcal{X}, b \in A} |Q^*(y', b) - Q_t(X_{t+1}, b)|) \\
&\geq -C_r + \alpha \min_{y \in \mathcal{X}} (- \max_{y' \in \mathcal{X}, b \in A} |Q^*(y', b) - Q_t(y, b)|) \\
&\geq -C_r - \alpha (\max_{y', y \in \mathcal{X}, b \in A} |Q_t(y, b) - Q^*(y, b)| + |Q^*(y, b) - Q^*(y', b)|) \\
&\geq -C_r - \alpha \|\Delta_t\|_\infty - \alpha \max_{(y', y) \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| =: m \in \mathbb{R}.
\end{aligned}$$

We apply Popoviciu's inequality on variances from Lemma 5.4 with the bounds m and M computed just above, to (5.7) and obtain

$$\begin{aligned}
& \text{Var}_{\mathbb{P}_{x_0, \mathbf{a}}}(F_t(x, a) | \mathcal{G}_t) \\
& \leq \text{Var}_{\mathbb{P}^{(k_t^*)}(x, a)} \left(-r(x, a, X_{t+1}) - \alpha \max_{b \in A} Q_t(X_{t+1}, b) + \alpha \min_{y' \in \mathcal{X}} \max_{b' \in A} Q^*(y', b') \right) \\
& \leq \frac{1}{4} (M - m)^2 \\
& \leq \frac{1}{4} \left(C_r + \alpha \|\Delta_t\|_\infty + C_r + \alpha \|\Delta_t\|_\infty + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 \\
& = \frac{1}{4} \left(C_r + C_r + 2\alpha \|\Delta_t\|_\infty + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 \\
& \leq \frac{1}{2} \left(\left(2C_r + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 + 4\alpha^2 \|\Delta_t\|_\infty^2 \right) \\
& \leq \left(\left(2C_r + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 + 4\alpha^2 \|\Delta_t\|_\infty^2 \right) \\
& \leq \left(\left(2C_r + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 + 4\alpha^2 \|\Delta_t\|_\infty^2 \right) \\
& \quad + 4\alpha^2 + \left(2C_r + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 \|\Delta_t\|_\infty^2 \\
& \leq \left(4\alpha^2 + \left(2C_r + \alpha \max_{y', y \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 \right) (1 + \|\Delta_t\|_\infty^2) \\
& \leq C \cdot (1 + \|\Delta_t\|_\infty)^2,
\end{aligned}$$

with

$$(5.8) \quad C := \left(4\alpha^2 + \left(2C_r + \alpha \max_{(y', y) \in \mathcal{X}, b \in A} |Q^*(y, b) - Q^*(y', b)| \right)^2 \right).$$

This shows that also Lemma 5.3 (iii) is fulfilled, and hence, with an application of Lemma 5.3 we obtain $Q_t(x, a) \rightarrow Q(x, a)$ $\mathbb{P}_{x_0, \mathbf{a}}$ -almost surely as $t \rightarrow \infty$ for all $(x, a) \in \mathcal{X} \times A$. \square

REFERENCES

- [1] Victor Aguirregabiria and Pedro Mira. Swapping the nested fixed point algorithm: A class of estimators for discrete Markov decision models. *Econometrica*, 70(4):1519–1543, 2002.
- [2] Charalambos D. Aliprantis and Kim C. Border. *Infinite dimensional analysis*. Springer, Berlin, third edition, 2006. A hitchhiker's guide.
- [3] Andrea Angiuli, Nils Detering, Jean-Pierre Fouque, and Jimin Lin. Reinforcement learning algorithm for mixed mean field control games. *arXiv preprint arXiv:2205.02330*, 2022.
- [4] Andrea Angiuli, Jean-Pierre Fouque, and Mathieu Lauriere. Reinforcement learning for mean field games, with applications to economics. *arXiv preprint arXiv:2106.13755*, 2021.
- [5] Nicole Bäuerle and Alexander Glauner. Distributionally robust Markov decision processes and their connection to risk measures. *Mathematics of Operations Research*, 2021.
- [6] Nicole Bäuerle and Alexander Glauner. Q-learning for distributionally robust Markov decision processes. In *Modern Trends in Controlled Stochastic Processes*, pages 108–128. Springer, 2021.
- [7] Nicole Bäuerle and Ulrich Rieder. *Markov decision processes with applications to finance*. Springer Science & Business Media, 2011.
- [8] Sinan Çalıřır and Meltem Kurt Pehlivanođlu. Model-free reinforcement learning algorithms: A survey. In *2019 27th signal processing and communications applications conference (SIU)*, pages 1–4. IEEE, 2019.
- [9] Jay Cao, Jacky Chen, John Hull, and Zissis Poulos. Deep hedging of derivatives using reinforcement learning. *The Journal of Financial Data Science*, 3(1):10–27, 2021.
- [10] Arthur Charpentier, Romuald Elie, and Carl Remlinger. Reinforcement learning in economics and finance. *Computational Economics*, pages 1–38, 2021.
- [11] Jesse Clifton and Eric Laber. Q-learning: Theory and applications. *Annual Review of Statistics and Its Application*, 7:279–301, 2020.
- [12] Avinash K Dixit. *Optimization in economic theory*. Oxford University Press, USA, 1990.
- [13] Matthew F Dixon, Igor Halperin, and Paul Bilokon. *Machine learning in Finance*, volume 1170. Springer, 2020.
- [14] Aryeh Dvoretzky. *On stochastic approximation*. University of California Press, 1956.

- [15] Laurent El Ghaoui and Arnab Nilim. Robust solutions to Markov decision problems with uncertain transition matrices. *Operations Research*, 53(5):780–798, 2005.
- [16] Jian Huang, Junyi Chai, and Stella Cho. Deep learning in finance and banking: A literature review and classification. *Frontiers of Business Research in China*, 14(1):1–24, 2020.
- [17] Garud N Iyengar. Robust dynamic programming. *Mathematics of Operations Research*, 30(2):257–280, 2005.
- [18] Tommi Jaakkola, Michael I Jordan, and Satinder P Singh. On the convergence of stochastic iterative dynamic programming algorithms. *Neural computation*, 6(6):1185–1201, 1994.
- [19] Beakcheol Jang, Myeonghwi Kim, Gaspard Harerimana, and Jong Wook Kim. Q-learning algorithms: A comprehensive classification and applications. *IEEE access*, 7:133653–133667, 2019.
- [20] Frank Hyneman Knight. *Risk, uncertainty and profit*, volume 31. Houghton Mifflin, 1921.
- [21] Petter N Kolm and Gordon Ritter. Modern perspectives on reinforcement learning in finance. *Modern Perspectives on Reinforcement Learning in Finance (September 6, 2019)*. *The Journal of Machine Learning in Finance*, 1(1), 2020.
- [22] Zijian Liu, Qinxun Bai, Jose Blanchet, Perry Dong, Wei Xu, Zhengqing Zhou, and Zhengyuan Zhou. Distributionally robust q -learning. In *International Conference on Machine Learning*, pages 13623–13643. PMLR, 2022.
- [23] Shie Mannor, Ofir Mebel, and Huan Xu. Robust MDPs with k -rectangular uncertainty. *Mathematics of Operations Research*, 41(4):1484–1509, 2016.
- [24] Jarryd Martin, Suraj Narayanan Sasikumar, Tom Everitt, and Marcus Hutter. Count-based exploration in feature space for reinforcement learning. *arXiv preprint arXiv:1706.08090*, 2017.
- [25] Volodymyr Mnih, Koray Kavukcuoglu, David Silver, Andrei A Rusu, Joel Veness, Marc G Bellemare, Alex Graves, Martin Riedmiller, Andreas K Fidjeland, Georg Ostrovski, et al. Human-level control through deep reinforcement learning. *nature*, 518(7540):529–533, 2015.
- [26] Mohammad Bagher Naghibi-Sistani, MR Akbarzadeh-Tootoonchi, MH Javidi-Dashte Bayaz, and Habib Rajabi-Mashhadi. Application of Q-learning with temperature variation for bidding strategies in market based power systems. *Energy Conversion and Management*, 47(11-12):1529–1538, 2006.
- [27] Ariel Neufeld and Julian Sester. Robust q -learning algorithm for Markov decision processes under Wasserstein uncertainty. *Automatica*, 2024+.
- [28] Ariel Neufeld, Julian Sester, and Mario Šikić. Markov decision processes under model uncertainty. *Mathematical Finance*, 33(3):618–665, 2023.
- [29] Brian Ning, Franco Ho Ting Lin, and Sebastian Jaimungal. Double deep Q-learning for optimal execution. *Applied Mathematical Finance*, 28(4):361–380, 2021.
- [30] Kishan Panaganti and Dileep Kalathil. Sample complexity of robust reinforcement learning with a generative model. In *International Conference on Artificial Intelligence and Statistics*, pages 9582–9602. PMLR, 2022.
- [31] Tiberiu Popoviciu. Sur les équations algébriques ayant toutes leurs racines réelles. *Mathematica*, 9(129-145):20, 1935.
- [32] John Rust. Structural estimation of Markov decision processes. *Handbook of econometrics*, 4:3081–3143, 1994.
- [33] Rajesh Sharma, Madhu Gupta, and Girish Kapoor. Some better bounds on the variance with applications. *Journal of Mathematical Inequalities*, 4(3):355–363, 2010.
- [34] Nian Si, Fan Zhang, Zhengyuan Zhou, and Jose Blanchet. Distributionally robust batch contextual bandits. *arXiv preprint arXiv:2006.05630*, 2020.
- [35] Nian Si, Fan Zhang, Zhengyuan Zhou, and Jose Blanchet. Distributionally robust policy evaluation and learning in offline contextual bandits. In *International Conference on Machine Learning*, pages 8884–8894. PMLR, 2020.
- [36] Satinder Singh, Tommi Jaakkola, Michael L Littman, and Csaba Szepesvári. Convergence results for single-step on-policy reinforcement-learning algorithms. *Machine learning*, 38(3):287–308, 2000.
- [37] Sorawoot Srisuma and Oliver Linton. Semiparametric estimation of Markov decision processes with continuous state space. *Journal of Econometrics*, 166(2):320–341, 2012.
- [38] Csaba Szepesvári and Michael L Littman. Generalized Markov decision processes: Dynamic-programming and reinforcement-learning algorithms. In *Proceedings of International Conference of Machine Learning*, volume 96, 1996.
- [39] Jean Tarbouriech, Shubhanshu Shekhar, Matteo Pirodda, Mohammad Ghavamzadeh, and Alessandro Lazaric. Active model estimation in Markov decision processes. In *Conference on Uncertainty in Artificial Intelligence*, pages 1019–1028. PMLR, 2020.
- [40] Michel Tokic and Günther Palm. Value-difference based exploration: adaptive control between epsilon-greedy and softmax. In *Annual conference on artificial intelligence*, pages 335–346. Springer, 2011.
- [41] Kerem Uğurlu. Robust optimal control using conditional risk mappings in infinite horizon. *Journal of Computational and Applied Mathematics*, 344:275–287, 2018.
- [42] Hado van Hasselt and Marco A Wiering. Convergence of model-based temporal difference learning for control. In *2007 IEEE International Symposium on Approximate Dynamic Programming and Reinforcement Learning*, pages 60–67. IEEE, 2007.
- [43] Cédric Villani. *Optimal transport: old and new*, volume 338. Springer, 2008.
- [44] Shengbo Wang, Nian Si, Jose Blanchet, and Zhengyuan Zhou. A finite sample complexity bound for distributionally robust q -learning. In *International Conference on Artificial Intelligence and Statistics*, pages 3370–3398. PMLR, 2023.
- [45] Yue Wang and Shaofeng Zou. Policy gradient method for robust reinforcement learning. *arXiv preprint arXiv:2205.07344*, 2022.

- [46] Christopher JCH Watkins. Learning from delayed rewards. *Ph. D. thesis, King's College, University of Cambridge*, 1989.
- [47] Christopher JCH Watkins and Peter Dayan. Q-learning. *Machine learning*, 8(3-4):279–292, 1992.
- [48] Wolfram Wiesemann, Daniel Kuhn, and Berç Rustem. Robust Markov decision processes. *Mathematics of Operations Research*, 38(1):153–183, 2013.
- [49] Huan Xu and Shie Mannor. Distributionally robust Markov decision processes. *Mathematics of Operations Research*, 37(2):288–300, 2012.
- [50] Wenhao Yang, Liangyu Zhang, and Zhihua Zhang. Toward theoretical understandings of robust Markov decision processes: Sample complexity and asymptotics. *The Annals of Statistics*, 50(6):3223–3248, 2022.
- [51] Zhengqing Zhou, Zhengyuan Zhou, Qinxun Bai, Linhai Qiu, Jose Blanchet, and Peter Glynn. Finite-sample regret bound for distributionally robust offline tabular reinforcement learning. In *International Conference on Artificial Intelligence and Statistics*, pages 3331–3339. PMLR, 2021.