

# Deriving Improved Plasma Fluid Equations from Collisional Kinetic Theory

Y. S. Dimant\*

*Boston University, Center for Space Physics, Boston, MA, United States*

Correspondence\*:  
Corresponding Author  
dimant@bu.edu

## ABSTRACT

Developing a quantitative understanding of wave plasma processes in the lower ionosphere requires a reasonably accurate theoretical description of the underlying physical processes. For such highly collisional plasma environment as the E-region ionosphere, kinetic theory represents the most accurate theoretical description of wave processes. For the analytical treatment, however, the collisional kinetic theory is extremely complicated and succeeds only in a limited number of physical problems. To date, most research applied oversimplified fluid models that lack a number of critical kinetic aspects, so that the coefficients in the corresponding fluid equations are often accurate only to an order of magnitude. This paper presents the derivation for the highly collisional, partially magnetized case relevant to E-region conditions. It provides a more accurate reduction of the ion and, especially, electron kinetic equations to the corresponding 5-moment fluid equations by using a new set of analytic approximations. This derivation results in more accurate fluid-model set of equations appropriate for most E-region problems. The results of this paper could be used for a routine practical analysis when working with actual data. The improved equations can also serve as a basis for more accurate plasma fluid computer simulations.

**Keywords:** E-region ionosphere, magnetized plasma, plasma-neutral collisions, kinetic theory, fluid equations, 5-moment description

## 1 INTRODUCTION

At altitudes of the equatorial and high-latitude E-region ionospheres, the ionosphere is highly collisional in such a way that ions are almost demagnetized by their frequent collisions with the surrounding neutral molecules while electrons still remain strongly magnetized. Strong DC electric fields perpendicular to the geomagnetic field cause their electrojets and give rise to plasma instabilities whose nonlinear development produces plasma density irregularities observed by radars and rockets.

Developing a quantitative understanding of wave plasma processes in the lower ionosphere requires an accurate theoretical description of the underlying physical processes. For such dissipative environment, collisional plasma kinetic theory represents the most accurate theoretical description of wave processes. Particle-in-cell (PIC) simulations present the most advanced way to apply the kinetic approach, but such massive computer simulations (Oppenheim and Dimant, 2004; Oppenheim et al., 2008; Oppenheim and Dimant, 2013; Oppenheim et al., 2020) are usually quite costly. In many cases,

simple estimates and parameter dependencies provided by an analytic approach will suffice. For the analytical treatment, however, the collisional kinetic theory is extremely complicated and succeeds only in a limited number of physical problems. To date, most research applied an oversimplified fluid model that lacks many critical kinetic aspects. These models mostly apply to weakly collisional conditions. The coefficients in the simple fluid equations are often accurate only to an order of magnitude because they were not obtained using the full kinetic theory. This paper presents the derivation of improved fluid equations for the highly collisional, partially magnetized case relevant to E-region conditions, starting from a more consistent kinetic approach. It provides more accurate values for the fluid-model coefficients.

There are different approaches to analytical description of low-frequency plasma processes in the E-region ionosphere, including both the kinetic theory and fluid models. Traditionally, the kinetic theory of the FB instability applied an oversimplified BGK collision operator (Bhatnagar et al., 1954). This operator does not follow from an accurate Boltzmann collision operator, but represents an artificial construct. It simplifies dramatically the analytical treatment, satisfying the particle number conservation, as well as the momentum and energy balances (albeit under certain conditions, see below). This simplified approach is reasonably applicable to the description of the heavy ions, but it is totally unacceptable to the description of the light electrons (Dimant and Sudan, 1995a).

More accurate approaches to the kinetic description of electrons under conditions of the E-region wave processes, such as the Farley-Buneman (FB) instability, have been developed by a few research groups. Stubbe (1990) modified the BGK terms to allow for the different rates of the electron energy and momentum losses. This simple modification, however, does not follow from the Boltzmann operator and cannot be trusted. Later, two independent research groups developed more sophisticated and accurate approaches. Kissack and collaborators (Kissack et al., 1995, 1997, 2008a,b) applied Grad's method (Grad, 1949; Rodbard et al., 1995), while Dimant and Sudan (1995a) used an expansion in Legendre polynomials with respect to the angles in the velocity space (Gurevich, 1978; Allis, 1982). The latter kinetic approach has allowed the authors to predict a new electron thermal-driven instability in the lower E/upper D regions (Dimant and Sudan, 1995b,c), which has been later explained in terms of a much simpler fluid model (Dimant and Sudan, 1997). This effect has been verified by others (Robinson, 1998; St. -Maurice and Kissack, 2000). Later, a similar thermal-instability process has been also suggested for ions (Kagan and Kelley, 2000; Dimant and Oppenheim, 2004; Dimant et al., 2023).

This paper presents a consistent reduction of the ion and electron kinetic equations to the 5-moment fluid equations by using a new set of analytic approximations. This derivation results in a more accurate fluid model appropriate for most E-region plasma problems.

The paper is organized as follows. Section 2 introduces the collisional kinetic equation and reviews the generic procedure for obtaining the moment equations. The collisional parts there are not specified and remain in the general integral form. Section 3 describes ion momentum equation obtained using the BGK collision model. The most important part is section 4. It derives low-frequency electron-fluid equations using a kinetic theory based on the efficient isotropization of the electron distribution function in the velocity space (Gurevich, 1978; Dimant and Sudan, 1995a). This requires a more detailed and sophisticated treatment. Section 4.1 derives the moment equations where the heat conductivity and frictional heating are given in terms of a still unspecified small directional part of the velocity distribution function. To illustrate major ideas of closing the derivation, section 4.2 describes the simplest case of the constant (i.e., velocity-independent) kinetic collision parameters. Section 4.3 presents the general results obtained in detail in the Appendix. Compared to the simplest electron-fluid equations from section 4.2, the general momentum and thermal-balance equations include a larger number of coefficients, as well

as additional heat-conductivity terms. By their look, the latter may appear collisionless, but they have arisen exclusively due to the velocity dependence of the kinetic electron-neutral collision frequency. All these results could be used for a routine practical analysis when working with actual data. The improved equations can serve as a basis for more accurate plasma fluid computer simulations.

## 2 GENERAL KINETIC TREATMENT

This section discusses a general approach to deriving the fluid model from the kinetic theory for any plasma particles. To avoid confusion, throughout this paper we will use the following nomenclature. We denote various kinds of particles (charged or neutral) by Latin subscripts:  $p$ ,  $q$ , etc., that stand for electrons ( $e$ ), ions of various kinds ( $i$ ) and neutrals ( $n$ ), while denoting vector components by Greek subscripts:  $\alpha$ ,  $\beta$ , etc.

Non-relativistic kinetics of charged particles of the kind  $p$  with the velocity  $\vec{v}_p$  at a given location  $\vec{r}$  and time  $t$  is described by the Boltzmann kinetic equation,

$$\partial_t f_p + \nabla \cdot (\vec{v}_p f_p) + \partial_{\vec{v}_p} \cdot \left[ \frac{q_p}{m_p} \left( \vec{E} + \vec{v}_p \times \vec{B} \right) f_p \right] = \left( \frac{df_p}{dt} \right)_{\text{col}}, \quad (1)$$

where  $f_p(\vec{v}_p, \vec{r}, t)$  is the single-particle velocity distribution function. The left-hand side (LHS) of Eq. (1) describes the collisionless (Vlasov) dynamics of the  $p$ -species charged particles in smoothed over many particles electric ( $\vec{E}$ ) and magnetic ( $\vec{B}$ ) fields (for simplicity, we ignore here a gravity force);  $q_p$  and  $m_p$  are the  $p$ -particle charge and mass, respectively. The LHS of Eq. (1) is intentionally written in a conservative (divergence) form which is more convenient for deriving the moment equations.

The right-hand side (RHS) of Eq. (1), term  $(df_p/dt)_{\text{col}} = \sum_q S_{pq}$ , is the collisional operator describing binary collisions of the  $p$ -particles with all available kinds of charged and neutral particles denoted by  $q$  (including the  $p$ -particles themselves). In the general case, the partial components  $S_{pq}$  represent integral operators that involve products of  $f_p(\vec{v}_p, \vec{r}, t)$  by  $f_q(\vec{v}_q, \vec{r}, t)$ . The partial operator  $S_{qq}$  is quadratically nonlinear, while  $S_{pq}$  with  $p \neq q$  are linear with respect to  $f_p$ .

The binary collisions can be either elastic or inelastic. Elastic collisions conserve the total kinetic energy, momentum and angular momentum of the colliding pair. The corresponding partial collisional operator,  $S_{pq}$ , can be described by the well-known Boltzmann collision integral (Shkarofsky et al., 1966; Gurevich, 1978; Lifshitz and Pitaevskii, 1981; Schunk and Nagy, 2009; Khazanov, 2011). During an inelastic collision of a charged particle with a neutral particle, a fraction of the total kinetic energy goes to excitation (de-excitation) of the neutral particle (or ion) or to release of electrons via ionization. Inelastic processes in the lower ionosphere often involve molecular dissociation, recombination with ions and electron attachment, accompanied by photon radiation or absorption. The complete kinetic description of all these processes is complicated. In many cases, however, inelastic collisions are close to elastic and one can continue using Boltzmann's integral with minor modifications (Gurevich, 1978; Shkarofsky et al., 1966). Kinetic Eq. (1) with Boltzmann's collision integral *per se* represents a significant simplification over the full multi-particle kinetics, but it still remains quite difficult for a mathematical treatment and requires further simplifications.

Being interested in fluid-model equations that follow from kinetic Eq. (1), right below we review the conventional approach to deriving equations for the lowest-order moments of the distribution function. The material of this section will serve as a guide for more specific derivations of the following sections.

The three lowest-order velocity moments include the  $p$ -species particle density,

$$n_p(\vec{r}, t) \equiv \int f_p d^3 v_p, \tag{2}$$

mean fluid velocity,

$$\vec{V}_p(\vec{r}, t) \equiv \langle \vec{v}_p \rangle = \frac{1}{n_p} \int \vec{v}_p f_p d^3 v_p, \tag{3}$$

and effective temperature,

$$T_p(\vec{r}, t) = \frac{m_p}{3} \left\langle \left( \vec{v}_p - \vec{V}_p \right)^2 \right\rangle = \frac{m_p}{3n_p} \int \left( \vec{v}_p - \vec{V}_p \right)^2 f_p d^3 v_p. \tag{4}$$

The derivations below will also involve other velocity-averaged quantities defined by

$$\langle \dots \rangle \equiv \frac{1}{n_p} \int (\dots) f_p d^3 v_p. \tag{5}$$

Integrations in Eqs. (2)–(5) are performed over the entire 3-D velocity space.

First, we consider the particle-number balance. Integrating Eq. (1) over the particle velocities with  $f_p \rightarrow 0$  as  $v_p \equiv |\vec{v}_p| \rightarrow \infty$ , we easily obtain the continuity equation for the  $p$ -particle fluid,

$$\partial_t n_p + \nabla \cdot \left( n_p \vec{V}_p \right) = \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p. \tag{6}$$

The RHS of Eq. (6) includes various particle sources and losses, like ionization, recombination and electron attachment. The collisions between the charged particles of the same species usually conserve the average particle number and hence do not contribute to the RHS of Eq. (6).

Second, we obtain the momentum-balance equation that involves the mean fluid drift velocity,  $\vec{V}_p$ . Integrating Eq. (1) with the weighting function  $m_p \vec{v}_p$ , for a given vector-component  $\alpha$  of the momentum density, we obtain

$$\begin{aligned} & m_p \partial_t (n_p V_{p\alpha}) + \sum_{\beta=1}^3 \partial_{x_\beta} \mathbf{P}_{p\alpha\beta} + m_p \sum_{\beta=1}^3 \partial_{x_\beta} (n_p V_{p\alpha} V_{p\beta}) \\ & = q_p \left[ E_\alpha + \frac{1}{c} \left( \vec{V}_p \times \vec{B} \right)_\alpha \right] n_p + m_p \int v_{p\alpha} \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p, \end{aligned} \tag{7}$$

where  $\mathbf{P}_p$  is the total pressure tensor with vector components defined as

$$\mathbf{P}_{p\alpha\beta} \equiv m_p \int (v_{p\alpha} - V_{p\alpha}) (v_{p\beta} - V_{p\beta}) f_p d^3 v_p. \tag{8}$$

It combines the isotropic pressure,  $P_p \delta_{mn}$  ( $\delta_{mn} = 1$  if  $m = n$  and  $\delta_{mn} = 0$  otherwise),  $P_p = n_p T_p$ , with the viscosity tensor,  $\Pi_{p\alpha\beta} \equiv \mathbf{P}_{p\alpha\beta} - P_p \delta_{\alpha\beta}$ . Equation (7) includes momentum changes due to various average forces and those caused by particle density variations. To exclude the latter and separate the net effect of the total force, we multiply Eq. (6) by  $m_p \vec{V}_p$  and subtract the resultant equation from Eq. (7).

This yields the conventional momentum-balance equation,

$$m_p n_p \frac{D_p \vec{V}_p}{Dt} = q_p n_p \left[ \vec{E} + \frac{1}{c} (\vec{V}_p \times \vec{B}) \right] - \nabla \cdot \mathbf{P}_p + \vec{R}_p, \quad (9)$$

where  $D_p/Dt \equiv \partial_t + \vec{V}_p \cdot \nabla$  is the convective (also called substantial or material) derivative for the average  $p$ -particle flow and

$$\vec{R}_p \equiv m_p \int (\vec{v}_p - \vec{V}_p) \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p. \quad (10)$$

Here and below, the “dot”-products of a vector,  $\vec{a}$ , with a two-component tensor,  $\mathbf{A}$ , depending on the multiplier order, denote vectors with the components  $(\vec{a} \cdot \mathbf{A})_\alpha \equiv \sum_{\beta=1}^3 a_\beta \mathbf{A}_{\beta\alpha}$  or  $(\mathbf{A} \cdot \vec{a})_\alpha \equiv \sum_{\beta=1}^3 \mathbf{A}_{\alpha\beta} a_\beta$ . The tensor divergence,  $\nabla \cdot \mathbf{P}_p = \nabla P_p + \nabla \cdot \mathbf{\Pi}_p$ , represents a vector which uses the obvious symmetry  $\mathbf{P}_{p\alpha\beta} = \mathbf{P}_{p\beta\alpha}$  following from Eq. (8). The RHS of Eq. (9) includes all smooth forces acting on the average particle flow of the charged particles, such as the total Lorentz force, pressure gradient, and total friction,  $\vec{R}_p$ . The latter is associated with collisions of the given  $p$ -particles with all other charged or neutral particles. It includes no momentum exchange between the same-species particles because their mutual collisions automatically conserve the total momentum,  $\int \vec{v}_p S_{pp} d^3 v_p = 0$ .

Third, to obtain the total energy-balance equation, we integrate Eq. (1) with the weighting function  $m_p v_p^2/2$  and obtain

$$\partial_t \mathcal{E}_p + \nabla \cdot \int \frac{m_p v_p^2}{2} \vec{v}_p f_p d^3 v_p = \vec{j}_p \cdot \vec{E} + \frac{m_p}{2} \int v_p^2 \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p, \quad (11)$$

where  $\mathcal{E}_p$  is the  $p$ -species average kinetic-energy density and  $\vec{j}_p$  is their electric-current density,

$$\mathcal{E}_p \equiv \int \frac{m_p v_p^2}{2} f_p d^3 v_p, \quad \vec{j}_p \equiv q_p n_p \vec{V}_p. \quad (12)$$

Before proceeding, we separate from the kinetic particle velocity  $\vec{v}_p$  the mean drift velocity  $\vec{V}_p$ , so that (11) becomes

$$\begin{aligned} & \partial_t \left[ n_p \left( \frac{m_p V_p^2}{2} + \frac{3T_p}{2} \right) \right] \\ & + \nabla \cdot \left[ n_p \left( \frac{m_p V_p^2}{2} + \frac{5T_p}{2} \right) \vec{V}_p + \mathbf{\Pi}_p \cdot \vec{V}_p + \frac{n_p m_p}{2} \left\langle (\vec{v}_p - \vec{V}_p)^3 \right\rangle \right] \\ & = \vec{j}_p \cdot \vec{E} + \vec{V}_p \cdot \vec{R}_p + \frac{m_p}{2} \int (\vec{v}_p - \vec{V}_p)^2 \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p \\ & + \frac{m_p V_p^2}{2} \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p. \end{aligned} \quad (13)$$

where  $(\vec{v}_p - \vec{V}_p)^3 = |\vec{v}_p - \vec{V}_p|^2 (\vec{v}_p - \vec{V}_p)$ . Equation (13) describes dynamic variations of the total energy density. It includes a part associated with the average fluid motion,  $n_p m_p V_p^2/2$ , and the internal

thermal energy,  $n_p T_p$ . To extract the equation exclusively for the particle temperature,  $T_p$ , we multiply Eq. (6) by  $(m_p V_p^2/2 + 3T_p/2)$ , take the scalar product of Eq. (9) with  $\vec{V}_p$  and subtract the resultant two equations from Eq. (13). This yields

$$\begin{aligned} & \frac{3n_p}{2} \frac{D_p T_p}{Dt} + n_p T_p \nabla \cdot \vec{V}_p + \mathbf{\Pi}_p \cdot \nabla \cdot \vec{V}_p + \nabla \cdot \left[ \frac{n_p m_p}{2} \left\langle (\vec{v}_p - \vec{V}_p)^3 \right\rangle \right] \\ & = \frac{m_p}{2} \int (\vec{v}_p - \vec{V}_p)^2 \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p + \left( \frac{m_p V_p^2}{2} - \frac{3T_p}{2} \right) \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p, \end{aligned} \tag{14}$$

where  $\mathbf{\Pi}_p \cdot \nabla \cdot \vec{V}_p \equiv \sum_{\alpha, \beta=1}^3 \mathbf{\Pi}_{p\alpha\beta} \nabla_\alpha \vec{V}_{p\beta}$ . Note that after this step the electric field has been eliminated from the energy balance equation. This is a crucial step in deriving the proper form of the frictional heating, as described below.

Typically, equations like Eq. (14) represent the final form of the thermal-balance equation. These equations are most convenient for calculations. In order to clarify the physical meaning of some terms, however, it is helpful to recast Eq. (14) in a slightly different form. Rewriting the continuity Eq. (6) as

$$\frac{D_p n_p}{Dt} + n_p \nabla \cdot \vec{V}_p = \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p,$$

we recast the two first terms in the LHS of Eq. (14) as

$$\begin{aligned} \frac{3n_p}{2} \frac{D_p T_p}{Dt} + n_p T_p \nabla \cdot \vec{V}_p &= \frac{3n_p}{2} \frac{D_p T_p}{Dt} - T_p \frac{D_p n_p}{Dt} + T_p \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p \\ &= n_p T_p \frac{D_p s_p}{Dt} + T_p \int \left( \frac{df_p}{dt} \right)_{\text{col}} d^3 v_p, \end{aligned} \tag{15}$$

where  $s_p \equiv \ln(T_p^{3/2}/n_p) = \ln(P_p^{3/2}/n_p^{5/2})$  represents the specific entropy of the  $p$ -species fluid (Braginskii, 1965) (for a single-atomic gas, the adiabatic coefficient  $\gamma = 5/3$ ). This recast allows interpreting  $n_p T_p \nabla \cdot \vec{V}_p$  as the adiabatic heating (cooling) term. The two remaining terms in the LHS of Eq. (14) describe the work performed by viscous forces and the fluid heat conductance. All these processes are collisionless.

All collisional processes in the thermal balance Eq. (14) are described by its RHS. After rearranging the last term in Eq. (15) to the RHS of Eq. (14), the last term there becomes  $(m_p V_p^2/2 - 5T_p/2) \int (df_p/dt)_{\text{col}} d^3 v_p$ . All integral terms involving  $(df_p/dt)_{\text{col}}$  describe the frictional heating and thermal inflows (outflows) associated with possible emergence (disappearance) of  $p$ -particles as a result of ionization, recombination, etc. For the general form of  $(df_p/dt)_{\text{col}}$ , calculating the frictional heating is not an easy task. Below we employ two different kinds of further approximation: one is more appropriate for heavy single-charged ions (Section 3), while the other is suitable for light electrons (Section 4).

Before proceeding further, we emphasize that in general no truncated chain of moment equations is closed because starting from the momentum equation every further moment equation involves higher-order moments. To allow the moment equation chain to be rigorously truncated, the most appropriate is the near-equilibrium case when the particle distribution function, along with its small perturbations, remains reasonably close to Maxwellian (Dimant and Sudan, 1995a; Kissack et al., 1995). This case

allows describing the particle kinetics using a restricted number of spatially and temporarily varying parameters, such as the particle density, temperature, and average drift velocity (5-moment equations). In real situations, however, this is not always the case, that is why inconsistencies in the fluid description often happen (e.g., García-Colín et al., 2004; Velasco et al., 2002). Higher-order sets of equations allow more serious deviations from Maxwellian, but still with a restricted number of additional fluid parameters. Fluid models that include restricted numbers of equations using approximate closures, such as the 5 or 13-moment models (Schunk and Nagy, 2009), can be successfully employed in situations when there are no sharp gradients, extreme fields, abundant superthermal particles, or extremely large temperature differences between different species of the colliding particles. These conditions are usually met at the equatorial electrojet. If they cannot be met, then the adequate description of plasma dynamics may require a direct solution of the corresponding collisional kinetic equation.

### 3 BGK COLLISION KINETICS AND THE FLUID MODEL FOR IONS

In this section, we consider the ion kinetics in low-ionized, highly collisional plasma. Bearing in mind mainly the lower-E/upper-D regions of the ionosphere or similar media, we can neglect coulomb collisions between the charged particles compared to their much more frequent collisions with the neutrals. For the ion-neutral collision integral, we can use the simple Bhatnagar-Gross-Krook (BGK) model (Bhatnagar et al., 1954). Disregarding ionization-recombination processes, we can write the simplest BGK collision operator in the neutral frame of reference as

$$\left(\frac{df_i}{dt}\right)_{\text{col}}^{\text{BGK}} = \nu_i (f_{\text{eff}} - f_i), \quad (16)$$

where  $f_{\text{eff}}$  is a fictitious Maxwellian function, normalized to the locally varying ion density,  $n_i(\vec{r}, t)$ , with the constant neutral temperature  $T_n$ :

$$f_{\text{eff}}(\vec{v}_i, \vec{r}, t) \equiv n_i(\vec{r}, t) \left(\frac{m_i}{2\pi T_n}\right)^{3/2} \exp\left(-\frac{m_i v_i^2}{2T_n}\right). \quad (17)$$

For the BGK model, it is essential that the ion-neutral collision frequency,  $\nu_i$ , is assumed constant. The model collision term in the form of Eq. (16) conserves the local number of particles. Applied to both ions and neutrals, the BGK model also conserves the total momentum of the two colliding particles.

The BGK model does not follow from Boltzmann's collision integral under any rigorous approximations, but it is a reasonable fit for single-charged ions that collide, predominantly elastically, with the surrounding neutrals of the same (or close) mass. Recent 2-D hybrid computer simulations of the Farley-Buneman instability that used for ions this kinetic equation (Kovalev et al., 2008) have demonstrated a good agreement with similar results of the more accurate fully-kinetic PIC or hybrid simulations (Janhunen, 1995; Oppenheim et al., 2008, 1996, 1995; Oppenheim and Dimant, 2004). There are two major reasons why this oversimplified model works well for the predominantly ion-neutral collisions typical for the lower ionosphere. First, within a 1000 K temperature range, the ion-neutral collision frequency is nearly velocity-independent (Maxwell molecule collisions) (Schunk and Nagy, 2009). Second, collisions of ions with neutral particles of the same or close mass have roughly equal rates of the average momentum and energy transfer, described by the single parameter  $\nu_i$ . Both these factors distinguish dramatically the ion-neutral collisions from the electron-neutral ones, as we discuss in the following section.

For the distribution function of single-charged positive ions,  $f_i(\vec{v}_i, \vec{r}, t)$ , the BGK kinetic equation in the conservative (divergence) form is given by

$$\partial_t f_i + \nabla \cdot (\vec{v}_i f_i) + \partial_{\vec{v}_i} \cdot \left[ \left( \frac{e\vec{E}}{m_i} + \Omega_i \vec{v}_i \times \hat{b} \right) f_i \right] = \nu_i (f_{\text{eff}} - f_i). \tag{18}$$

In this section, we derive the 5-moment ion fluid-model equations for  $n_i = \int f_i d^3 v_i$ ,  $\vec{V}_i = \langle \vec{v}_i \rangle = \int \vec{v}_i f_i d^3 v_i$ , and  $T_i = m_i \langle \delta \vec{v}_i \rangle^2 / 3 = (m_i / 3 n_i) \int \delta v_i^2 f_i d^3 v_i$ , where  $\delta \vec{v}_i \equiv \vec{v}_i - \vec{V}_i$ . We will largely follow the major steps of the previous section, but will do this specifically for the ions in the BGK approximation. This will help us to better understand the physical meaning of the algebraic manipulations and intermediate results.

Integrating Eq. (18) over velocities, we obtain the ion continuity equation,

$$\partial_t n_i + \nabla \cdot (n_i \vec{V}_i) = \frac{D_i n_i}{Dt} + n_i \nabla \cdot \vec{V}_i = 0. \tag{19}$$

Further, integrating Eq. (18) over ion velocities with the weighting function  $\vec{v}_i$ , we obtain the total momentum balance equation,

$$\partial_t (n_i \vec{V}_i) + \partial_{x_m} \int v_{i_k} v_{i_m} f d^3 v_i - n_i \left( \frac{e\vec{E}}{m_i} + \Omega_i \vec{V}_i \times \hat{b} \right) = -\nu_i n_i \vec{V}_i. \tag{20}$$

Using Eq. (19) and recalling the definition of the pressure in Eq. (8), we obtain the final momentum equation,

$$m_i n_i \frac{D_i \vec{V}_i}{Dt} = m_i n_i \left( \frac{e\vec{E}}{m_i} + \Omega_i \vec{V}_i \times \hat{b} \right) - \nabla \cdot \mathbf{P}_i - \nu_i n_i m_i \vec{V}_i. \tag{21}$$

Integrating Eq. (18) with the weighting function  $m_i v_i^2 / 2$ , we obtain

$$\begin{aligned} \partial_t \left( \frac{m_i n_i V_i^2}{2} + \frac{3 n_i T_i}{2} \right) + \nabla \cdot \int \vec{v}_i f_i \frac{m_i v_i^2}{2} d^3 v_i \\ = n_i e \vec{E} \cdot \vec{V}_i + \nu_i \left[ \frac{3}{2} n_i (T_n - T_i) - \frac{m_i n_i V_i^2}{2} \right]. \end{aligned} \tag{22}$$

Using the easily derivable relation

$$\int \vec{v}_i f_i \frac{m_i v_i^2}{2} d^3 v_i = \vec{V}_i \left( \frac{m_i n_i V_i^2}{2} + \frac{5 n_i T_i}{2} \right) + \vec{q}_i,$$

where the vector components of  $\vec{q}_i$  are given by

$$q_{i\alpha} \equiv m_i \int \delta v_{i\alpha} \frac{\delta v_i^2}{2} f_i d^3 v_i + \sum_{\beta=1}^3 V_{i\beta} \mathbf{\Pi}_{i\alpha\beta}, \tag{23}$$



we obtain

$$\begin{aligned} & \frac{D_i}{Dt} \left( \frac{m_i n_i V_i^2}{2} + \frac{3n_i T_i}{2} \right) \\ & + \left( \frac{m_i n_i V_i^2}{2} + \frac{3n_i T_i}{2} \right) \nabla \cdot \vec{V}_i + \nabla \cdot (n_i T_i \vec{V}_i + \vec{q}_i) \\ & = n_i e \vec{E} \cdot \vec{V}_i + \nu_i \left[ \frac{3}{2} n_i (T_n - T_i) - \frac{m_i n_i V_i^2}{2} \right]. \end{aligned} \tag{24}$$

Taking a scalar product of Eq. (21) with  $\vec{V}_i$ , we obtain

$$n_i \frac{D_i}{Dt} \left( \frac{m_i V_i^2}{2} \right) = n_i e \vec{E} \cdot \vec{V}_i - \vec{V}_i \cdot \nabla \cdot \mathbf{P}_i - \nu_i m_i n_i V_i^2. \tag{25}$$

Using Eq. (19) and  $\mathbf{P}_i = n_i T_i \mathbf{I} + \mathbf{\Pi}_i$ , we rewrite Eq. (25) as

$$\begin{aligned} & \frac{D_i}{Dt} \left( \frac{n_i m_i V_i^2}{2} \right) \\ & = n_i e \vec{E} \cdot \vec{V}_i - \vec{V}_i \cdot \nabla (n_i T_i) - \vec{V}_{i\alpha} \nabla_\beta \mathbf{\Pi}_{i\alpha\beta} - \nu_i m_i n_i V_i^2 - \frac{n_i m_i V_i^2}{2} \nabla \cdot \vec{V}_i. \end{aligned} \tag{26}$$

Rewriting Eq. (19) as

$$T_i \frac{D_i}{Dt} \left( \frac{3n_i}{2} \right) + \frac{3n_i T_i}{2} \nabla \cdot \vec{V}_i = 0 \tag{27}$$

we subtract both Eqs. (26) and (27) from Eq. (24) and obtain

$$\begin{aligned} & \frac{3n_i}{2} \frac{D_i T_i}{Dt} + n_i T_i \cdot \nabla \vec{V}_i + \nabla_\alpha q_\alpha - V_{i\beta} \nabla_\alpha \mathbf{\Pi}_{i\alpha\beta} \\ & = \frac{\nu_i m_i n_i V_i^2}{2} + \frac{3}{2} \nu_i n_i (T_n - T_i). \end{aligned}$$

Using

$$\nabla_\alpha q_{i\alpha} - V_{i\beta} \nabla_\alpha \mathbf{\Pi}_{i\alpha\beta} = m_i \nabla_\alpha \int \delta v_{i\alpha} \frac{\delta v_i^2}{2} f_i d^3 v_i + \mathbf{\Pi}_{i\alpha\beta} \nabla_\alpha V_{i\beta}$$

and  $n_i \nabla \cdot \vec{V}_i = -D_i n_i / dt$ , we obtain the final ion balance equation

$$\begin{aligned} & \frac{3n_i}{2} \frac{D_i T_i}{Dt} - T_i \frac{D_i n_i}{Dt} + \nabla \cdot \int m_i \delta \vec{v}_i \frac{\delta v_i^2}{2} f_i d^3 v_i + \mathbf{\Pi}_{i\alpha\beta} \nabla_\alpha V_{i\beta} \\ & = \frac{\nu_i m_i n_i V_i^2}{2} + \frac{3}{2} \nu_i n_i (T_n - T_i). \end{aligned} \tag{28}$$

The two last terms in the LHS of the ion thermal balance Eq. (28) describe the thermal conduction. The thermal flux given by the integral term and  $\mathbf{\Pi}_{i\alpha\beta}$  should be determined from higher-order moment equations. Alternatively, Eq. (28) could be closed using proper approximations like Maxwellian  $f_i$ .

The first term in the RHS of Eq. (28) describes the total ion frictional heating. This term has originated from a subtraction of the frictional energy loss described by

$$\nu_i \left( \int \frac{m_i v^2}{2} f_i d^3 v_i - \frac{3n_i T_i}{2} \right) = \frac{\nu_i m_i n_i V_i^2}{2},$$

from the total work performed during the collisional braking of the ion fluid as a whole,  $\nu_i m_i n_i V_i^2$ , given by the last term in Eq. (25). This means that when the ion fluid is frictionally braked only the half of the released energy heats the ions, while the other half goes directly to the colliding neutrals. Incidentally, this is precisely the total amount of frictional heating for ions and neutrals of the equal mass,  $m_n = m_i$ , following from a more detailed kinetic theory (Schunk and Nagy, 2009; Dimant and Oppenheim, 2004),

$$\frac{\nu_i m_i m_n n_i V_i^2}{m_i + m_n} = \frac{\nu_i m_i n_i V_i^2}{2}.$$

Thus, the BGK model of ion-neutral collisions describes correctly the ion frictional heating if the colliding partners have the same or close masses. This means that one can successfully employ for ions moment equations (27)–(28) under moderate conditions when the ion distribution function is reasonably close to Maxwellian. Such conditions mostly apply to the equatorial E region, rather than to the high-latitude ionosphere.

## 4 COLLISIONAL KINETICS AND THE FLUID MODEL FOR ELECTRONS

This section is the central piece of this paper. It derives the electron fluid equations from an approximate, but rigorous, kinetic theory based on characteristics of the actual physical conditions and wave processes in the E-region ionosphere. For electrons, the oversimplified BGK collision model (employed above for ions) can apply only to plasma processes whose characteristic wave frequencies exceed substantially the electron collision frequencies. However, for low-frequency processes in the highly collisional E/D-region ionosphere, where the opposite condition usually holds (see Dimant and Sudan, 1995a, and references therein), the electron BGK collision model is totally unsuitable. The main reason is that the rate of electron-neutral collisional exchange of momentum,  $\nu_{en}$ , is a few orders of magnitude larger than the corresponding rate of the energy exchange,  $\delta_{en} \nu_{en}$  (Gurevich, 1978). This means that during collisions with heavy neutrals the light electrons scatter over angles in the velocity space much more frequently than they change their kinetic energy. In low-frequency processes of the lower ionosphere, this leads to an efficient isotropization of the electron distribution function. The BGK model, however, completely ignores this feature. In addition, the BGK model does not cover the clearly pronounced velocity dependence of the kinetic electron-neutral collision frequency  $\nu_{en}(v_e)$  (Gurevich, 1978; Schunk and Nagy, 2009). This velocity dependence plays an important role for some E-region instabilities, (see, e.g., Dimant and Sudan, 1997, and references therein), and it modifies the instability and wave characteristics.

### 4.1 General kinetic approach and momentum equations

In a weakly ionized plasma of the lower ionosphere, collisions of an electron with other charged particles, including other electrons,  $\nu_{ee}, \nu_{ei}$ , are usually negligible compared to electron neutral collisions,  $\nu_e \approx \nu_{en}$ . At altitudes above 75 km, strongly magnetized electrons, involved in low-frequency processes with  $\omega \ll \nu_e \ll \Omega_e$ , have an almost isotropic velocity distribution whose speed dependence can deviate

significantly from Maxwellian. For such processes, an adequate kinetic description is by expanding the velocity distribution function  $f_e(\vec{r}, t, \vec{v}_e)$  in Legendre polynomials with respect to angles in the velocity space (Shkarofsky et al., 1966; Gurevich, 1978; Khazanov, 2011). To the first-order accuracy with respect to a small anisotropy of  $f_e(\vec{r}, t, \vec{v}_e)$ , one can represent the total electron distribution function as a combination of the major isotropic part,  $F_0(\vec{r}, t, v_e)$ , where  $v_e \equiv |\vec{v}_e|$ , and a relatively small directional part determined by a single vector-function  $\vec{f}_1(\vec{r}, t, v_e)$  (Gurevich, 1978; Dimant and Sudan, 1995a),

$$f_e(\vec{r}, t, \vec{v}_e) \approx F_0(\vec{r}, t, v_e) + \frac{\vec{f}_1(\vec{r}, t, v_e) \cdot \vec{v}_e}{v_e} = F_0 + |\vec{f}_1| \cos \theta, \quad (29)$$

where  $\theta$  is the angle between  $\vec{f}_1$  and  $\vec{v}_e$ . Here we assume that  $|\vec{f}_1| \ll F_0$ , along with the similar inequalities for the speed derivatives (see below). The major isotropic part,  $F_0$ , determines scalar velocity-averaged characteristics of the electron fluid, such as the electron density and temperature, while the small directional part,  $\vec{f}_1$ , determines vector characteristics, such as the average drift velocity and various fluxes. In this approximation, any higher-order anisotropies are neglected. For electrons in the highly collisional E-region ionosphere, the higher-order anisotropies usually play no role (see below).

The assumption of  $|\vec{f}_1| \ll F_0$  is well justified for electrons within the kinetic energy range  $\mathcal{E}_e < 2$  eV ( $v_e < 1000$  km/s). This range usually includes both the thermal bulk of electrons ( $\mathcal{E}_e \lesssim 0.03$  eV for the cold E-region ionosphere) and a significant fraction of superthermal electrons. In this energy rate, the ratio of the mean, mostly inelastic, collisional energy loss to that of the predominantly elastic momentum loss,  $\delta_{en}(v_e)\nu_{en}(v_e)/\nu_{en}(v_e) = \delta_{en}(v_e)$ , is usually quite small:  $\delta_{en}(v_e) \sim (2-4) \times 10^{-3}$  (Gurevich, 1978) (although it is two orders of magnitude larger than the corresponding purely elastic rate,  $\delta_{en}^{elas} \approx 2m_e/m_n$ ). The ratio of  $|\vec{f}_1|$  to  $F_0$  is typically  $\sim \sqrt{\delta_{en}}$ , so that the directional part of the electron distribution function in Eq. (29) turns out to be automatically small compared to the major isotropic part,  $|\vec{f}_1| \ll F_0$ . However, this brings up the following question. If there were an imposed DC electric field,  $\vec{E} \perp \vec{B}$ , so strong that the corresponding  $\vec{E} \times \vec{B}$ -drift velocity,  $\vec{V}_{dr} = \vec{E} \times \vec{B}/B^2$ , would be comparable to the mean electron thermal speed,  $v_{Th}^e = (T_e/m_e)^{1/2}$ , then the condition of  $|\vec{f}_1| \ll F_0$  would become invalid. As a matter of fact, however, such strong field would heat electrons so much that the heated thermal velocity  $v_{Th}^e$  would automatically exceed  $\vec{V}_{dr}$ . If the new electron temperature is  $\lesssim 23,000$  K (corresponding to 2 eV) then the approximation (29) still holds. This is a significant difference of electrons from heavy ions with  $\delta_{in} \simeq 1$ .

The kinetic equation with the general electron-neutral collision operator,  $(df_e/dt)_{coll}$ , leads to the two coupled equations for  $F_0(\vec{r}, t, v_e)$  and  $\vec{f}_1(\vec{r}, t, v_e)$  (Gurevich, 1978; Dimant and Sudan, 1995a):

$$\partial_t F_0 + \frac{v_e}{3} \nabla \cdot \vec{f}_1 - \frac{e}{3m_e v_e^2} \frac{\partial}{\partial v_e} \left( v_e^2 \vec{E} \cdot \vec{f}_1 \right) = S_0, \quad (30a)$$

$$\partial_t \vec{f}_1 - \Omega_e \hat{b} \times \vec{f}_1 + v_e \nabla F_0 - \frac{e \vec{E}}{m_e} \frac{\partial F_0}{\partial v_e} = \vec{S}_1, \quad (30b)$$

where

$$S_0 \equiv \frac{1}{2} \int_{-1}^1 \left( \frac{df_e}{dt} \right)_{col} d(\cos \vartheta), \quad \vec{S}_1 \equiv \frac{3}{2} \int_{-1}^1 \left( \frac{df_e}{dt} \right)_{col} \frac{\vec{f}_1}{|\vec{f}_1|} (\cos \vartheta) d(\cos \vartheta) \quad (31)$$

(note that the expressions for  $S_{0,1}$  in Dimant and Sudan (1995a) missed the correct normalization factors). Bearing in mind moderately fast wave processes,  $\tau_{rec}^{-1} \ll \omega \ll \nu_e$ , where  $\tau_{rec}$  is an effective recombination lifetime at a given altitude, we will ignore ionization-recombination processes, as we did above for

the ions. The kinetic description of electrons based on Eq. (30) differs dramatically from any kinetic description based on the BGK collision model.

Fluid equations based on Eq. (30), usually implying a nearly-Maxwellian velocity distribution, have been successfully explored by a number of researchers (see, e.g., Gurevich, 1978; Dimant and Sudan, 1995a, and references therein). However, the form of major fluid equations presented in Gurevich (1978), Chapter 5, does not clearly show the basic structure of generic Eqs. (9) and (14) or similar ion Eqs. (21) and (28). By this, we mean that Gurevich’s equations show neither explicit adiabatic heating and cooling, nor frictional heating  $\propto V_{e,i}^2$ . Adiabatic terms proportional to  $(\gamma_{e,i} - 1)$  in Gurevich (1978) Eqs. (5.3) and (5.4) and the corresponding terms in the following equations appear to have been introduced “by hand” and are actually extraneous. One can verify that these adiabatic terms have already been implicitly distributed among other terms of the temperature balance equations within the corresponding fluxes given by Gurevich (1978) equations (5.8)–(5.11), so that they are accounted in Eqs. (5.3)–(5.4) twice.

The explicit adiabatic terms show up naturally in the kinetic approaches based on small perturbations of the distribution function shifted by the average particle drift velocity. These approaches clearly differ from that based on perturbations of the non-shifted velocity distribution, as in Eq. (29) resulting in Eq. (30). For relatively small drift velocities, however, the two different approaches should yield the same results. Below we demonstrate that the kinetic approach based on Eqs. (29)–(30) does reproduce in a rigorous and natural way the electron fluid equations with the correct adiabatic heating and cooling, frictional heating, etc. We will also calculate kinetic corrections associated with the general velocity dependence of the electron-neutral collision frequency and non-Maxwellian velocity distribution. The Appendix contains details of these calculations.

In accord with the low-frequency condition of  $\omega \ll \nu_e \ll \Omega_e$ , we neglect in Eq. (30b) the electron inertia term  $\partial_t \vec{f}_1$  and use a standard approximation  $\vec{S}_1 \approx -\nu_e(v_e) \vec{f}_1$  (Gurevich, 1978; Dimant and Sudan, 1995a). This allows us to close this set of equations in a simple way. As a result, we obtain

$$-\frac{e\vec{E}}{m_e} \frac{\partial F_0}{\partial v_e} - \Omega_e \hat{b} \times \vec{f}_1 + v_e \nabla F_0 = -\nu_e(v_e) \vec{f}_1. \tag{32}$$

Resolving this vector equation with respect to  $\vec{f}_1$ , we obtain

$$\vec{f}_1(v_e) = -\mathbf{N}(v_e) \cdot \vec{K} F_0, \tag{33}$$

where the kinetic electron mobility tensor  $\mathbf{N}(v)$  and the differential vector operator  $\vec{K}$  are given by

$$\mathbf{N}(v_e) \approx \begin{bmatrix} \frac{\nu_e(v_e)}{\Omega_e^2} & \frac{1}{\Omega_e} & 0 \\ -\frac{1}{\Omega_e} & \frac{\nu_e(v_e)}{\Omega_e^2} & 0 \\ 0 & 0 & \frac{1}{\nu_e(v_e)} \end{bmatrix}, \tag{34}$$

$$\vec{K} \equiv v_e \nabla - \frac{e\vec{E}}{m_e} \frac{\partial}{\partial v_e}. \tag{35}$$

Here and elsewhere, we neglect second-order small terms  $\sim \nu_e^2$  compared to  $\Omega_e^2$  and represent all tensors in the matrix form for the Cartesian system  $\hat{x}, \hat{y}, \hat{z}$  with the  $\hat{z}$ -axis along  $\vec{B}$ . We can write Eqs. (33)–(34)

explicitly in terms of the parallel ( $\parallel$ ) and perpendicular ( $\perp$ ) to  $\vec{B}$  components as

$$f_{1\parallel} = -\frac{1}{\nu_e(v_e)} K_{\parallel} F_0, \quad \vec{f}_{1\perp} = -\left( \frac{\nu_e(v) \vec{K}_{\perp}}{\Omega_e^2} + \frac{\hat{b}}{\Omega_e} \times \vec{K}_{\perp} \right) F_0, \quad (36)$$

where  $\hat{b} = \hat{z}$  is the unit vector along  $\vec{B}$ . The spatial derivatives in Eq. (33) or (36) express the drift-diffusion approximation in the collisional kinetic theory, while the velocity derivatives describe electron energy variations caused by the electric field  $\vec{E}$ .

Furthermore, expressing the RHS of Eq. (30a) as (Gurevich, 1978)

$$S_0 = \frac{1}{2v_e^2} \frac{\partial}{\partial v_e} \left[ v_e^2 \delta_{en} \nu_e \left( v_e F_0 + \frac{T_n}{m_e} \frac{\partial F_0}{\partial v_e} \right) \right], \quad (37)$$

we obtain (Dimant and Sudan, 1995a)

$$\partial_t F_0 + \frac{1}{3v_e^2} \vec{K} \cdot (v_e^2 \vec{f}_1) = \frac{1}{2v_e^2} \frac{\partial}{\partial v_e} \left[ v_e^2 \delta_{en} \nu_e \left( v_e F_0 + \frac{T_n}{m_e} \frac{\partial F_0}{\partial v_e} \right) \right]. \quad (38)$$

Expressing here  $\vec{f}_1$  in terms of  $F_0$  via Eq. (33) or (36), we obtain a closed kinetic equation for the major isotropic distribution function,  $F_0(\vec{r}, t, v)$ . Its solution, with the use of (33) or (36), provides both parts of the distribution function, so that its scalar and vector moments can be calculated by a straightforward speed integration. According to Eq. (29), the lowest-order moments of the electron distribution function, such as the electron density, mean drift velocity, and temperature, are given by

$$n_e \equiv 4\pi \int_0^{\infty} F_0 v_e^2 dv_e, \quad \vec{V}_e \equiv \frac{4\pi}{3n_e} \int_0^{\infty} \vec{f}_1 v_e^3 dv_e, \quad T_e \equiv \frac{4\pi m_e}{3n_e} \int_0^{\infty} F_0 v_e^4 dv_e. \quad (39)$$

A direct solution of the kinetic Eq. (38) would be the most accurate and general way to describing the electron behavior (Dimant and Sudan, 1995a). However, the goal of this paper is to obtain a set of the lowest-order fluid equations in order to properly describe E-region plasma processes, even if this set of equations is not fully closed due to possible deviations of the electron velocity distribution function from Maxwellian.

As above, we start from the particle conservation. Using the definitions of Eq. (39) and integrating Eq. (38) over  $v_e$  with the weighting function  $4\pi v_e^2$ , we obtain the standard electron continuity equation,

$$\partial_t n_e + \nabla \cdot (n_e \vec{V}_e) = 0. \quad (40)$$

Next, we integrate Eq. (32) with the weighting function  $4\pi v_e^3 / (3n_e)$ . Applying the integration by parts, we obtain an equation

$$\frac{e\vec{E}}{m_e} - \Omega_e \hat{b} \times \vec{V}_e + \frac{\nabla(n_e T_e)}{m_e n_e} + \frac{4\pi}{3n_e} \int_0^{\infty} \vec{f}_1 \nu_e v_e^3 dv_e = 0, \quad (41)$$

that describes the momentum balance of the inertialess electron fluid. Equation (41) includes the Lorentz force, pressure gradient, and collisional friction. As we show in the Appendix, in the general case of a

velocity-dependent collision frequency,  $\nu_e(v)$ , the last term in the LHS of Eq. (41), in addition to the collisional friction, may also include an anisotropic addition to the total pressure gradient.

Taking a scalar product of Eq. (41) with  $m_e n_e \vec{V}_e$ , we obtain the expression

$$\vec{V}_e \cdot \left[ n_e e \vec{E} + \nabla (n_e T_e) \right] + \frac{4\pi m_e}{3} \vec{V}_e \cdot \int_0^\infty \vec{f}_1 \nu_e v_e^3 dv_e = 0 \quad (42)$$

analogous to Eq. (25) for ions. This expression represents the total work done by the electric field and other forces on the average electron flow. We will employ this expression right below.

Now we derive an equation describing the total energy balance. Integrating Eq. (38) with the weighting function  $2\pi m_e v_e^4$ , we obtain

$$\begin{aligned} \partial_t \left( \frac{3n_e T_e}{2} \right) + \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e + n_e e \vec{E} \cdot \vec{V}_e \\ = -2\pi m_e \int_0^\infty \left( v_e F_0 + \frac{T_n}{m_e} \frac{dF_0}{dv_e} \right) \delta_{en} \nu_e v_e^3 dv_e. \end{aligned} \quad (43)$$

Using Eq. (42), we eliminate from Eq. (43) the work done by the electric field on the average flow,  $n_e e \vec{E} \cdot \vec{V}_e$ , and obtain

$$\begin{aligned} \partial_t \left( \frac{3n_e T_e}{2} \right) - \vec{V}_e \cdot \nabla (n_e T_e) + \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e \\ = \frac{4\pi m_e}{3} \vec{V}_e \cdot \int_0^\infty \vec{f}_1 \nu_e v_e^3 dv_e - 2\pi m_e \int_0^\infty \left( v_e F_0 + \frac{T_n}{m_e} \frac{dF_0}{dv_e} \right) \delta_{en} \nu_e v_e^3 dv_e. \end{aligned} \quad (44)$$

Here we have rearranged the terms between the two sides of the equation in such a way that all terms proportional to the collision frequency remain in the RHS, while all other terms are put in the LHS. After so doing, it may be tempting to interpret the first term in the RHS of Eq. (44) as the electron frictional heating. In the general case of velocity-dependent  $\nu_e(v)$ , however, this interpretation would not be perfectly accurate, as we show in Appendix and section 4.3 below.

Equation (44) is not yet the final form of the thermal-balance equation. It needs to be further transformed into a form similar to Eq. (14) or (28). In Appendix, we develop this recast for the general case of velocity-dependent  $\nu_e(v)$ . However, right below we proceed with the simplest model of constant  $\nu_e$  and  $\delta_{en}$ . This model is inaccurate for electron-neutral collisions of the lower ionosphere (Gurevich, 1978; Schunk and Nagy, 2009), but it will allow us to clarify basic ideas of closing Eq. (44).

## 4.2 Constant collisional parameters

For constant  $\nu_e$  and  $\delta_{en}$ , using the definitions of Eq. (39) and integrating the last term of Eq. (44) by parts, we obtain

$$\begin{aligned} \partial_t \left( \frac{3n_e T_e}{2} \right) - \vec{V}_e \cdot \nabla (n_e T_e) + \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e \\ = m_e \nu_e n_e V_e^2 + \frac{3}{2} \delta_{en} \nu_e n_e (T_n - T_e), \end{aligned} \quad (45)$$

Using Eq. (33), we rewrite the third term in the LHS as

$$\frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e = -\frac{5}{2m_e} \nabla \cdot \mathbf{N} \cdot \left[ \nabla (\lambda n_e T_e^2) + n_e T_e e \vec{E} \right]. \tag{46}$$

Here the double-dot product involving a tensor means  $\nabla \cdot \mathbf{N} \cdot \nabla \dots = \sum_{\alpha, \beta=1}^3 \partial_{x_\alpha} (\mathbf{N}_{\alpha\beta} \partial_{x_\beta} \dots)$  (and similar for  $\nabla \cdot \mathbf{N} \cdot \vec{E}$ ) and we have also introduced a dimensionless parameter of order unity,  $\lambda$ ,

$$\lambda \equiv \frac{4\pi m_e^2}{15 n_e T_e^2} \int_0^\infty v_e^6 F_0 dv_e = \frac{m_e \int_0^\infty v_e^6 F_0 dv_e}{5 T_e \int_0^\infty v_e^4 F_0 dv_e} = \frac{3 \left( \int_0^\infty F_0 v_e^2 dv_e \right) \int_0^\infty F_0 v_e^6 dv_e}{5 \left( \int_0^\infty F_0 v_e^4 dv_e \right)^2}. \tag{47}$$

Note that for the Maxwellian isotropic part of the electron distribution function,

$$F_0 = n_e \left( \frac{m_e}{2\pi T_e} \right)^{3/2} \exp \left( -\frac{m_e v_e^2}{2T_e} \right), \tag{48}$$

we have  $\lambda = 1$ .

Using Eq. (39) and (35), we obtain

$$\vec{V}_e = -\frac{4\pi}{3n_e} \mathbf{N} \cdot \int_0^\infty v_e^3 \vec{K} F_0 dv_e = -\mathbf{N} \cdot \left[ \frac{e \vec{E}}{m_e} + \frac{\nabla (n_e T_e)}{m_e n_e} \right]. \tag{49}$$

Multiplying Eq. (49) by  $m_e n_e T_e$ , we can rewrite it as

$$-\mathbf{N} \cdot (n_e T_e e \vec{E}) = m_e n_e T_e \vec{V}_e + \mathbf{N} \cdot T_e \nabla (n_e T_e).$$

This relation allows us to eliminate the electric field from Eq. (46), so that the latter becomes

$$\begin{aligned} & \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e \\ &= \frac{5}{2m_e} \nabla \cdot \left\{ \mathbf{N} \cdot \left[ (1 - \lambda) T_e^2 \nabla n_e - (2\lambda - 1) n_e T_e \nabla T_e \right] + m_e n_e T_e \vec{V}_e \right\}. \end{aligned} \tag{50}$$

Using Eqs. (40) and (50), after a simple algebra,

$$\begin{aligned} & \partial_t \left( \frac{3n_e T_e}{2} \right) - \vec{V}_e \cdot \nabla (n_e T_e) + \frac{5}{2} \nabla \cdot (n_e T_e \vec{V}_e) \\ &= \partial_t \left( \frac{3n_e T_e}{2} \right) - \vec{V}_e \cdot \nabla (n_e T_e) + \frac{5}{2} \nabla \cdot (n_e T_e \vec{V}_e) \\ & \quad - \frac{5T_e}{2} \left[ \partial_t n_e + \nabla \cdot (n_e \vec{V}_e) \right] = \frac{3n_e}{2} \frac{D_e T_e}{Dt} - T_e \frac{D_e n_e}{Dt}, \end{aligned} \tag{51}$$

we obtain the sought-for temperature-balance equation in a more standard form,

$$\frac{3n_e}{2} \frac{D_e T_e}{Dt} - T_e \frac{D_e n_e}{Dt} - \nabla \cdot \vec{q}_e = m_e \nu_e n_e V_e^2 + \frac{3}{2} \delta_{en} \nu_e n_e (T_n - T_e). \tag{52}$$

Here the electron thermal flux density,  $\vec{q}_e$ , is given by

$$\vec{q}_e = \frac{5T_e}{2m_e} \mathbf{N} \cdot [(2\lambda - 1) n_e \nabla T_e + (\lambda - 1) T_e \nabla n_e] = \vec{q}_{e\parallel} + \vec{q}_{eP} + \vec{q}_{eH}, \quad (53)$$

where its explicit parallel, Pedersen, and Hall components are given by

$$\begin{aligned} \vec{q}_{e\parallel} &= \frac{5T_e [(2\lambda - 1) n_e \nabla_{\parallel} T_e + (\lambda - 1) T_e \nabla_{\parallel} n_e]}{2m_e \nu_e}, \\ \vec{q}_{eP} &= \frac{5T_e \nu_e [(2\lambda - 1) n_e \nabla_{\perp} T_e + (\lambda - 1) T_e \nabla_{\perp} n_e]}{2m_e \Omega_e^2}, \\ \vec{q}_{eH} &= \frac{\Omega_e}{\nu_e} (\hat{b} \times \vec{q}_{eP}) = \frac{5T_e \hat{b} \times [(2\lambda - 1) n_e \nabla_{\perp} T_e + (\lambda - 1) T_e \nabla_{\perp} n_e]}{2m_e \Omega_e}. \end{aligned} \quad (54)$$

The two first terms in the LHS of Eq. (52), as well as the similar ones in Eq. (14) or (28), describe adiabatic heating or cooling of the electron fluid, while  $\nabla \cdot \vec{q}_e$  describes the heat conductivity. Note that the Hall component of  $\vec{q}_e$  can contribute into electron heat conductance only as a quadratically-nonlinear effect and, since  $\nabla \cdot \vec{q}_{eH} \propto (\nabla_{\perp} n_e \times \nabla_{\perp} T_e)$  only if the gradients of  $n_e$  and  $T_e$  are not parallel.

As mentioned above, for Maxwellian  $F_0(v_e)$  we have  $\lambda = 1$ , so that the term in  $\vec{q}_e$  proportional to  $\nabla n_e$  disappears. This fact can be understood as follows. If the major part of the distribution function remains Maxwellian then it is determined only by two space-dependent parameters: the density,  $n_e$ , and the temperature,  $T_e$ . If there is a density gradient but no temperature gradient, then electrons of all energies will diffuse from denser regions to less dense ones with no redistribution of the temperature and hence with no heat conductivity.

If the electron velocity distribution deviates from Maxwellian (this happens, e.g., when a low-ionized plasma heated by strong electric fields is embedded in an abundant cold neutral atmosphere with a significantly different temperature (Milikh and Dimant, 2003)) then the situation is more complicated. The effective electron temperature  $T_e$ , which is proportional to the mean electron chaotic energy, can be uniformly distributed, but the details of the electron energy distribution may differ significantly in different regions of space. The energy transport is stronger for electrons with higher energies than it is for lower-energy electrons. Hence, if there are spatial gradients of high-energy distribution tails then more energetic particles provide stronger energy redistribution. This may make, e.g., some less dense regions to be on average more energetic than the denser regions, even if they had initially equal effective temperatures. Moreover, it is even possible to imagine a situation when electron heat is transferred from cooler regions to hotter ones, leading to a further electron temperature elevation in the latter. This counter-intuitive, but theoretically possible, effect should not surprise because a strongly non-Maxwellian, i.e., strongly non-equilibrium plasma cannot be adequately described by the conventional equilibrium thermodynamics.

### 4.3 Velocity-dependent parameters

In the actual lower ionosphere, the electron-neutral kinetic collision frequency,  $\nu_e$ , and the energy loss fraction,  $\delta_{en}$ , have clearly pronounced velocity dependencies (Gurevich, 1978; Schunk and Nagy, 2009). This does not allow  $\nu_e(v_e)$  and  $\delta_{en}(v_e)$  to be factored out from the integrals in Eqs. (41)–(44), making the derivation of the general momentum and temperature-balance equations more complicated than that described in section 4.2. Such derivation is developed in detail in the Appendix, while here we only present the results.



In the general case of velocity-dependent  $\nu_e$  and  $\delta_{en}$ , electron continuity Eq. (40) stays the same. The other two moment equations have the same basic structure as (49) and (52), but they contain additional terms and include a large number of dimensionless factors of order unity listed in Eqs. (63)–(68) below.

The general inertialess expression for the average electron drift velocity  $\vec{V}_e$  is given by

$$\vec{V}_e = -\frac{1}{m_e} \mathbf{M} \cdot \left[ e\vec{E} + \frac{\nabla_{\perp}(n_e T_e)}{n_e} + \frac{\beta_{\parallel}}{\alpha_{\parallel}} \frac{\nabla_{\parallel}(n_e T_e)}{n_e} \right], \tag{55}$$

where

$$\mathbf{M} \equiv \frac{4\pi}{3n_e} \int_0^{\infty} \frac{d(v_e^3 \mathbf{N}(v_e))}{dv_e} F_0(v_e) dv_e = \begin{bmatrix} \frac{\alpha_P \langle \nu_e \rangle_e}{\Omega_e^2} & \frac{1}{\Omega_e} & 0 \\ -\frac{1}{\Omega_e} & \frac{\alpha_P \langle \nu_e \rangle_e}{\Omega_e^2} & 0 \\ 0 & 0 & \alpha_{\parallel} \left\langle \frac{1}{\nu_e} \right\rangle_e \end{bmatrix}, \tag{56}$$

an the tensor  $\mathbf{N}$  is given by Eq. (34) and  $\langle \dots \rangle_e$  denotes the velocity average over the major (isotropic) part of the electron distribution function,

$$\langle \dots \rangle_e = \frac{4\pi \int_0^{\infty} (\dots) F_0(v_e) v_e^2 dv_e}{n_e} = \frac{\int_0^{\infty} (\dots) F_0(v_e) v_e^2 dv_e}{\int_0^{\infty} F_0(v_e) v_e^2 dv_e}. \tag{57}$$

The electron current density is given by  $\vec{j}_e = en_e \vec{V}_e$ , so that the electric conductivity tensor is given by  $\sigma_e = (n_e e^2 / m_e) \mathbf{M}$ . The corresponding diagonal terms represent the Pedersen ( $\propto \alpha_P$ ) and parallel ( $\propto \alpha_{\parallel}$ ), while the antisymmetric off-diagonal terms ( $\propto 1/\Omega_e$ ) represent the Hall conductivity.

The general thermal-balance equation is given by

$$\begin{aligned} & \frac{3n_e}{2} \frac{D_e T_e}{Dt} - T_e \frac{D_e n_e}{Dt} \\ & + \frac{5}{2} \left( \frac{\rho_{\parallel} - \beta_{\parallel}}{\alpha_{\parallel}} \right) n_e T_e \nabla_{\parallel} \cdot \vec{V}_{e\parallel} + \left( \frac{5\rho_{\parallel} - 3\alpha_{\parallel} - 2\beta_{\parallel}}{2\alpha_{\parallel}} \right) \vec{V}_{e\parallel} \cdot \nabla_{\parallel} (n_e T_e) - \nabla \cdot \vec{q}_e \\ & = \alpha_P \langle \nu_e \rangle_e m_e n_e \left( V_{e\perp}^2 + \frac{V_{\parallel}^2}{\alpha_{\parallel} \xi} \right) - 2\pi m_e \int_0^{\infty} v_e^3 \delta_{en} \nu_e \left( v_e F_0 + \frac{T_n}{m_e} \frac{dF_0}{dv_e} \right) dv_e, \end{aligned} \tag{58}$$

where

$$\vec{q}_e \equiv \vec{q}_{eP} + \vec{q}_{eH} + \vec{q}_{e\parallel} = \mathbf{X} \cdot \frac{\nabla T_e}{T_e} + (\mathbf{X} - \mathbf{\Lambda}) \cdot \frac{\nabla n_e}{n_e}, \tag{59}$$

is the thermal-flux density with

$$\mathbf{X} = \frac{5n_e T_e}{2m_e} \begin{bmatrix} \frac{\chi_P \langle \nu_e \rangle}{\Omega_e^2} & \frac{\chi_H}{\Omega_e} & 0 \\ -\frac{\chi_H}{\Omega_e} & \frac{\chi_P \langle \nu_e \rangle}{\Omega_e^2} & 0 \\ 0 & 0 & \frac{\chi_{\parallel}}{\langle \nu_e \rangle} \end{bmatrix}, \tag{60a}$$

$$\mathbf{\Lambda} = \frac{5n_e T_e}{2m_e} \begin{bmatrix} \frac{\mu_P \langle \nu_e \rangle}{\Omega_e^2} & \frac{\lambda}{\Omega_e} & 0 \\ -\frac{\lambda}{\Omega_e} & \frac{\mu_P \langle \nu_e \rangle}{\Omega_e^2} & 0 \\ 0 & 0 & \frac{\mu_{\parallel}}{\langle \nu_e \rangle} \end{bmatrix}. \tag{60b}$$

The explicit Pedersen, Hall, and parallel components of  $\vec{q}_e$  are given by

$$\begin{aligned} \vec{q}_{eP} &\equiv \frac{5T_e \langle \nu_e \rangle [\chi_P n_e \nabla_{\perp} T_e + (\chi_P - \mu_P) T_e \nabla_{\perp} n_e]}{m_e \Omega_e^2}, \\ \vec{q}_{eH} &\equiv \frac{5T_e \hat{b} \times [\chi_H n_e \nabla_{\perp} T_e + (\chi_H - \lambda) T_e \nabla_{\perp} n_e]}{m_e \Omega_e}, \\ \vec{q}_{e\parallel} &\equiv \frac{5T_e [\chi_{\parallel} n_e \nabla_{\parallel} T_e + (\chi_{\parallel} - \mu_{\parallel}) T_e \nabla_{\parallel} n_e]}{m_e \langle \nu_e \rangle_e}, \end{aligned} \tag{61}$$

$$\chi_P \equiv 2\mu_P + \alpha_P - \beta_P - \rho_P, \quad \chi_H \equiv 2\lambda - 1, \quad \chi_{\parallel} \equiv 2\mu_{\parallel} - \beta_{\parallel}, \tag{62}$$

In addition to  $\lambda$  defined by (47), Eqs. (55)–(62) include

$$\alpha_P \equiv \frac{\int_0^{\infty} \frac{d(v^3 \nu_e(v_e))}{dv_e} F_0 dv_e}{3 \int_0^{\infty} \nu_e(v_e) F_0 v_e^2 dv_e}, \quad \alpha_{\parallel} \equiv \frac{\int_0^{\infty} \frac{d(v_e^3 / \nu_e(v_e))}{dv} F_0 dv_e}{3 \int_0^{\infty} \frac{1}{\nu_e(v_e)} F_0 v^2 dv_e}, \tag{63}$$

$$\rho_{\parallel} \equiv \frac{\left( \int_0^{\infty} \frac{d}{dv_e} \left( \frac{v_e^5}{\nu_e(v_e)} \right) F_0 dv_e \right) \left( \int_0^{\infty} v_e^2 F_0 dv_e \right)}{5 \left( \int_0^{\infty} v_e^4 F_0 dv_e \right) \left( \int_0^{\infty} \frac{v_e^2}{\nu_e(v_e)} F_0 dv_e \right)}, \tag{64a}$$

$$\rho_P \equiv \frac{\left( \int_0^{\infty} \frac{d}{dv_e} (\nu_e(v_e) v_e^5) F_0 dv_e \right) \left( \int_0^{\infty} v_e^2 F_0 dv_e \right)}{5 \left( \int_0^{\infty} v_e^4 F_0 dv_e \right) \left( \int_0^{\infty} \nu_e(v_e) v_e^2 F_0 dv_e \right)}, \tag{64b}$$

$$\beta_P \equiv \frac{\left( \int_0^{\infty} \nu_e(v_e) F_0 v_e^4 dv_e \right) \int_0^{\infty} F_0 v_e^2 dv_e}{\left( \int_0^{\infty} \nu_e(v_e) F_0 v_e^2 dv_e \right) \int_0^{\infty} F_0 v_e^4 dv_e}, \tag{65a}$$

$$\beta_{\parallel} \equiv \frac{\left( \int_0^{\infty} \frac{F_0 v_e^4}{\nu_e(v_e)} dv_e \right) \int_0^{\infty} F_0 v_e^2 dv_e}{\left( \int_0^{\infty} \frac{F_0 v_e^2}{\nu_e(v_e)} dv_e \right) \int_0^{\infty} F_0 v_e^4 dv_e}, \tag{65b}$$

$$\delta_P \equiv \frac{\left( \int_0^{\infty} \nu_e^2 F_0 v_e^4 dv_e \right) \left( \int_0^{\infty} F_0 v_e^2 dv_e \right)^2}{\left( \int_0^{\infty} \nu_e F_0 v_e^2 dv_e \right)^2 \left( \int_0^{\infty} F_0 v_e^4 dv_e \right)}, \tag{66}$$

$$\mu_P \equiv \frac{3 \left( \int_0^\infty \nu_e(v_e) F_0 v_e^6 dv_e \right) \left( \int_0^\infty F_0 v_e^2 dv_e \right)^2}{5 \left( \int_0^\infty \nu_e(v_e) F_0 v_e^2 dv_e \right) \left( \int_0^\infty F_0 v_e^4 dv_e \right)^2}, \tag{67a}$$

$$\mu_{\parallel} \equiv \frac{3 \left( \int_0^\infty \frac{F_0 v_e^6}{\nu_e(v_e)} dv_e \right) \left( \int_0^\infty F_0 v_e^2 dv_e \right)^2}{5 \left( \int_0^\infty \frac{F_0 v_e^2}{\nu_e(v_e)} dv_e \right) \left( \int_0^\infty F_0 v_e^4 dv_e \right)^2}, \tag{67b}$$

$$\xi \equiv \left\langle \frac{1}{\nu_e} \right\rangle_e \langle \nu_e \rangle_e = \frac{\left( \int_0^\infty \nu_e(v_e) F_0(v_e) v_e^2 dv_e \right) \left( \int_0^\infty \nu_e^{-1}(v_e) F_0(v_e) v_e^2 dv_e \right)}{\left( \int_0^\infty F_0(v_e) v_e^2 dv_e \right)^2}. \tag{68}$$


For constant  $\nu_e$  and arbitrary  $F_0(v)$ , we have  $\alpha_{P,\parallel} = \rho_{P,\parallel} = \beta_{P,\parallel} = \delta_P = \xi = 1$ ,  $\mu_{P,\parallel} = \lambda$ ,  $\chi_{P,H,\parallel} = 2\lambda - 1$  and  $\mathbf{M} = \mathbf{N}$ , so that Eqs. (58)–(61) reduce to Eqs. (52)–(54). If, additionally,  $F_0(v)$  is Maxwellian then we have even simpler parameters:  $\mu_{P,\parallel} = \chi_{P,H,\parallel} = \lambda = 1$ .

In a broad range of electron energies,  $\mathcal{E}_e \lesssim 0.3$  eV, the velocity dependence of  $\nu_e$  in the lower ionosphere can be approximated by a simple power-law dependence,  $\nu_e \propto v_e^{2\alpha}$ , with  $\alpha \approx 5/6$  [(Gurevich, 1978), Sect. 2.3.1, Fig. 7] or, practically to the same accuracy, with  $\alpha = 1$  (Dimant and Sudan, 1995a). For the general power-law dependent  $\nu_e \propto v_e^{2\alpha}$  with  $\alpha$  in the range between 0 and 1 and Maxwellian  $F_0(v_e)$ , Eqs. (63)–(68) simplify dramatically,

$$\begin{aligned} \alpha_P = \beta_P &= 1 + \frac{2\alpha}{3}, & \alpha_{\parallel} = \beta_{\parallel} &= 1 - \frac{2\alpha}{3}, \\ \rho_{\parallel} = \mu_{\parallel} &= \frac{(3 - 2\alpha)(5 - 2\alpha)}{15}, & \rho_P = \mu_P &= \frac{(3 + 2\alpha)(5 + 2\alpha)}{15}, \\ \delta_P &= \frac{\sqrt{\pi}\Gamma(5/2 + 2\alpha)}{3\Gamma^2(3/2 + \alpha)}, & \xi &= \frac{1 - 4\alpha^2}{\sin \frac{\pi(2\alpha+1)}{2}} = \frac{4\alpha^2 - 1}{\sin \frac{\pi(2\alpha-1)}{2}}. \end{aligned} \tag{69}$$

The case of  $\alpha = 1/2$  corresponds to hard-sphere collisions. In this case, the indeterminate expression for  $\xi$  yields  $4/\pi \approx 1.273$ . For  $\alpha = 5/6$  (Gurevich, 1978), we have  $\alpha_P = \beta_P \approx 1.556$ ,  $\alpha_{\parallel} = \beta_{\parallel} \approx 0.444$ ,  $\rho_{\parallel} = \mu_{\parallel} \approx 0.296$ ,  $\rho_P = \mu_P \approx 2.074$ ,  $\delta_P \approx 3.095$ , and  $\xi \approx 2.053$ . For  $\alpha = 1$  (Dimant and Sudan, 1995a), all these factors deviate from unity even further, e.g.,  $\alpha_{\parallel} = \beta_{\parallel} \approx 0.333$ ,  $\delta_P \approx 3.889$ , and  $\xi = 3$ . Thus the quantitative effect of the velocity dependence of  $\nu_e(v)$  is significant and should not be ignored. Figure 1 shows the coefficients given by Eq. (69) for general values of the power-law exponent  $\alpha$  within the physically realistic range of  $0 \leq \alpha \leq 1$ .

Comparison of general energy-balance Eq. (58) with Eq. (52) shows that the velocity dependence of the collision parameters results not only in more complicated heat conductivity, frictional heating, and cooling, but in additional terms associated with the plasma motion and gradients in the parallel to  $\vec{B}$  direction. It is important that these seemingly collisionless terms originate fully from electron-neutral collisions due to the velocity distribution of  $\nu_e(v)$ . Similar effects in the Hall and Pedersen directions are inconsequential and are neglected here. One should not, however, neglect Hall and Pedersen components in the heat conductivity because the plasma temperature and density gradients in those directions can be much sharper than those in the parallel direction.



**Figure 1.** Fluid-model coefficients for the power-law dependent  $e$ - $n$  collision frequency,  $\nu_{en}(v_e) \propto v_e^{2\alpha}$ .

Now we discuss the last (cooling) term in the RHS of Eq. (58). For general velocity-dependent  $\delta_{en}(v)$ , but a Maxwellian distribution function  $F_0 \propto \exp[-m_e v^2 / (2T_e)]$ , using (39), we reduce this term to

$$-2\pi m_e \int_0^\infty v_e^3 \delta_{en} \nu_e \left( v_e F_0 + \frac{T_n}{m_e} \frac{dF_0}{dv_e} \right) dv_e = \frac{3}{2} \frac{\langle \delta_{en} v_e^2 \nu_e \rangle_e n_e}{\langle v_e^2 \rangle_e} (T_n - T_e). \tag{70}$$

For general  $F_0$ , but constant  $\delta_{en}$ , we can rewrite the cooling term in Eq. (58) as

$$-2\pi m_e \int_0^\infty v_e^3 \delta_{en} \nu_e \left( v_e F_0 + \frac{T_n}{m_e} \frac{dF_0}{dv_e} \right) dv_e = \frac{3}{2} \delta_{en} \langle \nu_e \rangle_e n_e (\alpha_P T_n - \beta_P T_e), \tag{71}$$

where we integrated by parts and used Eqs. (63) and (65a). Equation (71) shows that for general non-Maxwellian  $F_0$  the cooling term is not necessarily proportional to the temperature difference  $(T_n - T_e)$ . However, for the power-law dependent  $\nu_e \propto v_e^{2\alpha}$  and Maxwellian  $F_0$ , according to Eq. (69), we have  $\beta_P = \alpha_P = 1 + 2\alpha/3$ . In this case, the structure of the cooling term proportional to  $\alpha_P \langle \nu_e \rangle_e n_e$  matches that of the frictional heating term for a purely perpendicular field,  $\alpha_P \langle \nu_e \rangle_e m_e n_e V_{e\perp}^2$ , as seen from the first term in the RHS of Eq. (58).

## 5 DISCUSSION

When applying a fluid model, either for analytic calculations or simulations, it is important to have the corresponding equations with accurate parameters applicable to the relevant physical conditions. These equations and parameters are usually derived from the kinetic theory, so that their accuracy is determined by the accuracy of the underlying kinetic approach.

Based on two different approximate kinetic approaches, this paper derives the fluid-model equations that describe low-frequency plasma processes in the highly dissipative E-region ionosphere. The treatment is restricted to collisions of the plasma particles, ions or electrons, with the neutral molecules only; no Coulomb collisions are considered. The neglect of Coulomb collisions at the E-region ionosphere is usually well justified, although sometimes electron-electron collisions may play a role, resulting in a more efficient maxwellization of the electron distribution function (Dimant and Sudan, 1995a). Such maxwellization makes the fluid model (as opposed to the pure kinetic theory) even more applicable. For the plasma particle collisions with neutrals (elastic or inelastic), we assume the known cross-sections as functions of the colliding particle velocities. These velocity dependencies of the collisional cross-sections can be taken from the literature (e.g., for the dominant electron-nitrogen collisions, see Itikawa, 2006; Song et al., 2023, and references therein). Assuming these known cross-sections, we can always calculate the velocity dependencies of the kinetic collision frequencies,  $\nu_p(\vec{v}_p)$  ( $p = i, e, n$ ) that are used in this paper.

The resultant fluid-model parameters are expressed in general integral forms through the known velocity dependencies of  $\nu_p$ . For the most important plasma processes, such as the small- to medium-scale cross-field plasma instabilities (the thermal Farley-Buneman and gradient drift instabilities), closed 5-moment multi-fluid models are usually sufficient for the accurate fluid description. Given the plasma species  $p$ , the 5-moment set of the unknowns includes the particle density ( $n_p$ ), temperature ( $T_p$ ), and the three components of the mean drift velocity,  $\vec{V}_p$ .

For the ionospheric ions ( $p = i$ ), we have employed the well-known (and fairly simple) BGK collisional model. For the heavy ions, the applicability of the BGK collision operator can be justified by the fact that within the thermal bulk and around, the kinetic ion-neutral collision frequency  $\nu_{in}$  is approximately constant, i.e., velocity-independent; this approximation corresponds to so-called Maxwell molecule collisions (Schunk and Nagy, 2009). Additionally, in the E-region the ion masses ionosphere are fairly close to the neutral-molecule masses,  $m_i \simeq m_n$ . As we demonstrate in section 3, in the case of  $m_i = m_n$  the oversimplified BGK model results even in a quantitatively accurate frictional heating and cooling terms, see the RHS of equation (28). We should bear in mind, however, that under sufficiently strong electric field,  $E \gtrsim m_i \nu_i V_{Ti} / e$ , the ion distribution function can be significantly distorted with an appreciable deviation from Maxwellian. Although the major ion-fluid terms in these cases remain valid, the entire 5-moment model cannot necessarily be easily closed, and hence its validity may be questionable. The factor of strong electric field is usually of importance for the high-latitude E region under conditions of severe magnetospheric perturbations (geomagnetic storms or substorms), while at the equatorial E region the electric fields are typically much weaker, so that the closed 5-moment ion fluid model is usually more applicable.

The central part of this paper is the derivation of the 5-moment fluid equations for electrons ( $p = e$ ). For the light electrons, unlike the ions, the simple BGK model cannot serve even as a crude approximation. As we explained in section 4, the reasons for the total BGK-model inapplicability are the two major facts: (1) the mean rate of the collisional loss of the electron energy is much less than the corresponding loss of the electron momentum, so that the electron behavior cannot be described by a single collisional parameter; (2) the kinetic collisional frequency has a pronounced electron velocity dependence,  $\nu_e(v_e)$ . The first fact leads to a strong isotropization of the electron velocity distribution in the velocity space, while the speed dependence of the electron velocity distribution is effectively decoupled from its angular dependence. The second fact leads to noticeable modifications of the electron fluid coefficients and even to occurrence of additional thermal-diffusion terms. As a result, in the general case the fluid-model

coefficients acquire additional dimensionless multipliers whose values are determined by some integral relations over the entire electron distribution function, see Eqs. (59)–(68). For the Maxwellian function, and especially for the power-law dependencies of the  $\nu_e$ -speed dependence,  $\nu_e(v_e) \propto v_e^{2\alpha}$ , these general integral relations reduce to simple algebraic ones, see Eq. (69) and Figure 1. From that figure we see that some numerical multipliers can deviate from the unity quite significantly, although they still remain in the same order of magnitude. A better quantitative knowledge of these fluid coefficients is important for accurate calculations and predictions of physical characteristics of various plasma wave and other processes. As our future knowledge of the speed dependence of the kinetic collision rates becomes more precise, using the more general integral relationships obtained here one can improved the values of the corresponding fluid-model coefficients.

The kinetic approach for the electron description employed in this paper is based on the expansion of the electron velocity distribution in Legendre polynomials in the velocity space and keeping the two first terms of such expansion, see Eq.(29): the dominant isotropic part,  $F_0(\vec{r}, t, v_e)$ , and a small directional part,  $|f_1(\vec{r}, t, v_e)| \cos \theta$ , where  $\theta$  is the angle between  $\vec{f}_1$ . This approach is analogous to that employed by Gurevich (1978), see also Dimant and Sudan (1995a), although, as we explained in section 4.1, Gurevich's fluid equations for electrons, Gurevich (1978, Chapter 5), derived through this kinetic approach, differ from ours. Gurevich's equations are written in a form that does not include explicit adiabatic and frictional heating terms. Also, as we mentioned above, the “adiabatic” terms proportional to  $(\gamma_{e,i} - 1)$  in Gurevich's Eqs. (5.3) and (5.4) are extraneous, while their correct equivalent has already been implicitly distributed within the other terms in Gurevich's Eqs. (5.8)–(5.11) (and hence included twice).

An alternative kinetic approach to build an electron fluid description is based on Grad's method (Kissack et al., 1995, 1997, 2008a,b). The latter assumes a finite number of parameters that characterize the velocity distribution, implying that the electron velocity distribution is reasonably close to Maxwellian. Our approach is much more general in terms of the  $v_e$ -dependence, but it restricts the angular distribution of the electron distribution function to the simplest linear deviation from the isotropy (see above). This approximation allows calculating vector fluxes like  $n_e \vec{V}_e$  or energy fluxes (see below), but higher-order tensor characteristics like an anisotropic pressure, etc., may require an accuracy beyond its field of applicability. Note, however, that high-order tensor characteristics for electrons are not expected to be significant due to the relatively high rate of electron distribution function isotropization associated with a small value of  $\delta_{en} \sim (2-4) \times 10^{-3}$  within the low-energy electron energy range,  $\mathcal{E}_e \equiv m_e v_e^2 / 2 < 2$  eV (Gurevich, 1978). Note also that under physical conditions when the two methods are applicable both techniques actually provide reasonable close quantitative results. At the same time, our kinetic approach provides much simpler, and hence much more practical, algebraic expressions applicable to various small- and medium-scale E-region processes.

## 6 CONCLUSION

Based on relevant physical conditions, we have derived improved fluid equations for the E-region ionosphere. In this derivation, for the E-region ions and electrons we have used two different approximate kinetic approaches.

For the ions, we have employed the simple BGK collision operator (section 3). This resulted in full 5-moment set of the continuity, momentum, and energy balance equations, see Eqs. (19), (21), and (28). Though these equations look conventional, our derivation has demonstrated that for the E-region ions

with almost equal masses of the ions and neutrals, the BGK collision operator leads to the quantitatively accurate frictional and cooling rates.

The central part of this paper is the derivation of the electron fluid equations. For the electrons, the BGK collisional operator is inapplicable and we have employed the kinetic approach based on the expansion of the electron distribution function,  $f_e(\vec{v}_e)$ , in Legendre polynomials over the angles in the velocity space. Due to physical conditions resulting in efficient isotropization of  $f_e(\vec{v}_e)$  allowed us to restrict the treatment to the two highest terms of the Legendre expansion: the dominant isotropic part,  $F_0(v_e)$  and a small directional part  $\vec{f}_1 \cdot \vec{v}_e/v_e$ . The former is responsible for calculating the scalar fluid quantities, such as the electron density and temperature, while the directional part allows one to calculate the electron flux (or electric current) density. An important factor in our derivations is the fact that the electron-neutral collisional frequencies are strongly velocity-dependent. Assuming these velocity dependencies to be known, we have derived the full set of the 5-moment equations: the continuity equation, the momentum equation, and the thermal balance equation. Since the E-region electrons in all relevant low-frequency processes are essentially inertialess, the momentum equation reduces to an explicit expression for the electron mean drift velocity given by Eq. (55). The most non-trivial result is the thermal balance equation given in the general case by Eq. (58), where the parameters are given by Eqs. (47), (63)–(62). For the Maxwellian distribution function and the power-law speed dependence of the electron-neutral collision frequency,  $\nu_e(v_e) \propto v_e^{2\alpha}$ , the integral relationships for the fluid-model parameters reduce to simple algebraic expressions given by Eq. (69), see also Fig. 1.

Comparison of general energy-balance Eq. (58) with the corresponding equation for the velocity-independent electron collision frequency, see Eq. (52), shows that the velocity dependence of the collision parameters results in a more complicated heat conductivity, frictional heating, and cooling, as well as in additional terms associated with the plasma motion and gradients in the parallel to  $\vec{B}$  direction. These terms look collisionless, but they originate exclusively from the velocity distribution of  $\nu_e(v)$ . Similar effects in the Hall and Pedersen directions are inconsequential and have been neglected. However, one should not neglect the Hall and Pedersen components in the heat conductivity because the plasma temperature and density gradients in those directions can be much sharper than those in the parallel direction.

The results of this paper could be used for a routine practical analysis when working with actual data. The improved equations can also serve as a basis for more accurate plasma fluid computer simulations. In the general case, the applicability of the closed 5-moment equations is restricted by reasonably moderate conditions of the equatorial E region. For the high-latitude E-region ionosphere, the accurate description may require using a fully kinetic treatment.

## ACKNOWLEDGMENT

The author is deeply grateful to Prof. Oppenheim for his long-term support, extensive help, and multiple discussions. This work was funded by NSF Grants 1755350, 1007789, 0442075 and NASA Grants 80NSSC21K1322, 80NSSC19K0080.

## APPENDIX

**MOMENT EQUATIONS FOR VELOCITY-DEPENDENT  $\nu_e(v_e)$**

In this Appendix, we derive Eqs. (55) and (58). The key step of this derivation is expressing the electric field,  $\vec{E}$ , in terms of the electron average drift velocity,  $\vec{V}_e$ , and the  $n_e, T_e$ -gradients.

For velocity-dependent  $\nu_e(v)$ , which cannot be factored out of the corresponding integrals, we use the following approach. Using Eq. (33) and integrating by parts, we obtain from Eq. (39)

$$\vec{V}_e = -\frac{4\pi}{3n_e} \int_0^\infty v^4 \mathbf{N} \cdot \nabla F_0 dv - \mathbf{M} \cdot \frac{e\vec{E}}{m_e}, \tag{A.1}$$

where  $\mathbf{M}$  is given by Eq. (56) with  $\alpha_{P,\parallel}$  defined by Eq. (63). By inverting the matrix  $\mathbf{M}$ , we obtain

$$\frac{e\vec{E}}{m_e} = -\mathbf{M}^{-1} \cdot \left( \vec{V}_e + \frac{4\pi}{3n_e} \int_0^\infty v^4 \mathbf{N}(v) \cdot \nabla F_0 dv \right). \tag{A.2}$$

Using Eq. (33), we obtain

$$\vec{f}_1(u) = -\mathbf{N}(u) \cdot \left[ u \nabla F_0 + \mathbf{M}^{-1} \cdot \left( \vec{V}_e + \frac{4\pi}{3n_e} \int_0^\infty \mathbf{N}(v) \cdot \nabla F_0(v) v^4 dv \right) \frac{\partial F_0}{\partial u} \right]. \tag{A.3}$$

To complete the derivation of the momentum and temperature-balance equations using Eqs. (41) and (44), we need to calculate there the two integrals involving  $\vec{f}_1$ . Both integrals have the same structure described by  $\int_0^\infty \vec{f}_1 g(v) v^3 dv$ , where the function  $g(v)$  is either  $\nu_e(v)$  or  $v^2$ . Using integration by parts, changing the variables, and re-ordering the integration wherever needed, we obtain

$$\begin{aligned} \int_0^\infty \vec{f}_1 g(v_e) v_e^3 dv_e &= \int_0^\infty \frac{d \left( g(u) u^3 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \cdot \vec{V}_e \right)}{du} F_0(u) du \\ &+ \int_0^\infty \left[ \frac{4\pi}{3n_e} \int_0^\infty \frac{d \left( g(u) u^3 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \right)}{du} F_0(u) du - g(v) \mathbf{I} \right] \\ &\cdot \mathbf{N}(v) \cdot \nabla F_0(v_e) v_e^4 dv_e, \end{aligned}$$

where  $\mathbf{I}$  is the unit tensor:  $\mathbf{I}_{\alpha\beta} = 1$  if  $\alpha = \beta$  and  $\mathbf{I}_{\alpha\beta} = 0$  otherwise. As a result, for the integrals in Eqs. (41) and (44) we obtain

$$\begin{aligned} &\frac{4\pi m_e}{3} \int_0^\infty \nu_e \vec{f}_1 v_e^3 dv_e \\ &= \frac{4\pi m_e}{3} \int_0^\infty \frac{d \left( \nu_e(u) u^3 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \cdot \vec{V}_e \right)}{du} F_0(u) du \\ &+ \frac{4\pi m_e}{3} \int_0^\infty \left[ \frac{4\pi}{3n_e} \int_0^\infty \frac{d \left( \nu_e(u) u^3 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \right)}{du} F_0(u) du - \nu_e(v) \mathbf{I} \right] \\ &\cdot \mathbf{N}(v) \cdot \nabla F_0(v_e) v_e^4 dv_e, \end{aligned} \tag{A.4}$$



and

$$\begin{aligned} \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e &= \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \frac{d(u^5 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \cdot \vec{V}_e)}{du} F_0(u) du \\ &+ \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \left[ \frac{4\pi}{3n_e} \int_0^\infty \frac{d(u^5 \mathbf{N}(u) \cdot \mathbf{M}^{-1})}{du} F_0(u) du - v^2 \mathbf{I} \right] \\ &\cdot \mathbf{N}(v) \cdot \nabla F_0(v_e) v_e^4 dv_e. \end{aligned} \tag{A.5}$$

First, we calculate the integrals in the RHS of Eq. (A.4). After a simple algebra, we obtain

$$\begin{aligned} \frac{4\pi}{3n_e} \int_0^\infty \frac{d(\nu_e(u) u^3 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \cdot \vec{V}_e)}{du} F_0(u) du \\ \approx \alpha_P \langle \nu_e \rangle \vec{V}_{e\perp} + \frac{(\alpha_P^2 - \mu) \langle \nu_e \rangle^2}{\Omega_e} (\hat{b} \times \vec{V}_{e\perp}) + \frac{\langle \nu_e \rangle \vec{V}_{\parallel}}{\alpha_{\parallel} \xi}, \end{aligned} \tag{A.6}$$

where

$$\mu \equiv \frac{\left( \int_0^\infty \frac{d(\nu_e^2(v_e) v_e^3)}{dv_e} F_0 dv_e \right) \int_0^\infty F_0 v_e^2 dv_e}{3 \left( \int_0^\infty \nu_e v_e^2 F_0 dv_e \right)^2}, \tag{A.7}$$

and similar for the second term in the RHS of (A.5),

$$\begin{aligned} \frac{4\pi}{3n_e} \int_0^\infty \nu_e \vec{f}_1 v_e^3 dv_e &\approx \alpha_P \langle \nu_e \rangle \vec{V}_{e\perp} + \frac{(\alpha_P^2 - \mu) \langle \nu_e \rangle^2}{\Omega_e} (\hat{b} \times \vec{V}_{e\perp}) + \frac{\langle \nu_e \rangle \vec{V}_{\parallel}}{\alpha_{\parallel} \xi} \\ &- \frac{[\alpha_P (\alpha_P - \beta_P) - \mu + \delta_P] \langle \nu_e \rangle^2 \nabla_{\perp} (n_e T_e)}{\Omega_e^2 m_e n_e} \\ &+ \frac{(\alpha_P - \beta_P) \langle \nu_e \rangle \hat{b} \times \nabla_{\perp} (n_e T_e)}{\Omega_e m_e n_e} + \left( \frac{\beta_{\parallel}}{\alpha_{\parallel}} - 1 \right) \frac{\nabla_{\parallel} (n_e T_e)}{m_e n_e}. \end{aligned} \tag{A.8}$$

Here and below, all parameters have been defined by Eqs. (63)–(68). Substituting Eq. (A.8) to Eq. (41) and neglecting the second-order small terms  $\sim \nu_e^2/\Omega_e^2$  wherever applicable, we obtain

$$\frac{e\vec{E}}{m_e} - \Omega_e \hat{b} \times \vec{V}_{e\perp} + \langle \nu_e \rangle_e \left( \alpha_P \vec{V}_{e\perp} + \frac{\vec{V}_{e\parallel}}{\alpha_{\parallel} \xi} \right) + \frac{\nabla_{\perp} (n_e T_e)}{m_e n_e} + \frac{\beta_{\parallel}}{\alpha_{\parallel}} \frac{\nabla_{\parallel} (n_e T_e)}{m_e n_e} = 0. \tag{A.9}$$

Expressing the combination of the second and third terms in the LHS of Eq. (A.9) as a matrix-by-vector product,

$$-\Omega_e \hat{b} \times \vec{V}_{e\perp} + \langle \nu_e \rangle_e \left( \alpha_P \vec{V}_{e\perp} + \frac{\vec{V}_{e\parallel}}{\alpha_{\parallel} \xi} \right) = \begin{bmatrix} \alpha_P \langle \nu_e \rangle_e & \Omega_e & 0 \\ -\Omega_e & \alpha_P \langle \nu_e \rangle_e & 0 \\ 0 & 0 & \frac{\langle \nu_e \rangle_e}{\alpha_{\parallel} \xi} \end{bmatrix} \vec{V}_e,$$

and resolving the resultant matrix equation for  $\vec{V}_e$  to the first-order accuracy with respect to  $\nu_e/\Omega_e$ ,

$$\begin{bmatrix} \alpha_P \langle \nu_e \rangle_e & \Omega_e & 0 \\ -\Omega_e & \alpha_P \langle \nu_e \rangle_e & 0 \\ 0 & 0 & \frac{\langle \nu_e \rangle_e}{\alpha_{\parallel} \xi} \end{bmatrix}^{-1} \approx \begin{bmatrix} \frac{\alpha_P \langle \nu_e \rangle_e}{\Omega_e^2} & -\frac{1}{\Omega_e} & 0 \\ \frac{1}{\Omega_e} & \frac{\alpha_P \langle \nu_e \rangle_e}{\Omega_e^2} & 0 \\ 0 & 0 & \frac{\alpha_{\parallel} \xi}{\langle \nu_e \rangle_e} \end{bmatrix} = \mathbf{M},$$

we arrive at the explicit expression for the electron fluid drift velocity  $\vec{V}_e$  given by Eq. (55).

Further, we calculate the integrals in Eq. (A.5). To the first-order accuracy with respect to  $\nu_e/\Omega_e$ , we obtain

$$\begin{aligned} & \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \frac{d}{dv} \left( v_e^5 \mathbf{N} \cdot \mathbf{M}^{-1} \cdot \vec{V}_e \right) F_0 dv_e \\ & \approx \frac{5}{2} \left\{ \nabla_{\perp} \cdot \left( \vec{V}_{e\perp} n_e T_e \right) + \nabla_{\parallel} \left( \frac{\rho_{\parallel} V_{e\parallel} n_e T_e}{\alpha_{\parallel}} \right) \right. \\ & \left. + \frac{\langle \nu_e \rangle_e (\rho_P - \alpha_P)}{\Omega_e} \nabla_{\perp} \cdot \left[ \hat{b} \times \left( \vec{V}_{e\perp} n_e T_e \right) \right] \right\}. \end{aligned} \tag{A.10}$$

and

$$\begin{aligned} & \frac{4\pi}{3n_e} \int_0^\infty \frac{d \left( u^5 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \right)}{du} F_0(u) du \\ & \approx \frac{5T_e}{m_e} \begin{bmatrix} 1 & \frac{(\alpha_P - \rho_P) \langle \nu_e \rangle_e}{\Omega_e} & 0 \\ \frac{(\rho_P - \alpha_P) \langle \nu_e \rangle_e}{\Omega_e} & 1 & 0 \\ 0 & 0 & \frac{\rho_{\parallel}}{\alpha_{\parallel} \langle 1/\nu_e \rangle_e} \end{bmatrix}. \end{aligned} \tag{A.11}$$

where the dimensionless parameters of order unity  $\rho_{\parallel, P}$  are defined by Eq. (64). Using Eqs. (A.10) and (A.11) and proceeding with calculations similar to those above, we obtain

$$\begin{aligned} & \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \left[ \frac{4\pi}{3n_e} \int_0^\infty \frac{d \left( u^5 \mathbf{N}(u) \cdot \mathbf{M}^{-1} \right)}{du} F_0(u) du - v^2 \mathbf{I} \right] \\ & \cdot \mathbf{N}(v) \cdot \nabla F_0(v_e) v_e^4 dv_e \\ & = \nabla_{\perp} \cdot \frac{5 \langle \nu_e \rangle_e \left[ (\beta_P + \rho_P - \alpha_P) T_e \nabla_{\perp} (n_e T_e) - \mu_P \nabla_{\perp} (n_e T_e^2) \right]}{m_e \Omega_e^2} \\ & + \nabla_{\perp} \cdot \frac{5 \hat{b} \times \left[ T_e \nabla_{\perp} (n_e T_e) - \lambda \nabla_{\perp} (n_e T_e^2) \right]}{m_e \Omega_e} \\ & + \nabla_{\parallel} \cdot \frac{5 \left[ \beta_{\parallel} T_e \nabla_{\perp} (n_e T_e) - \mu_{\parallel} \nabla_{\perp} (n_e T_e^2) \right]}{m_e \langle \nu_e \rangle_e} \end{aligned} \tag{A.12}$$

Separating the temperature and density gradients, after introducing linear combinations of the numerical coefficients  $\chi_{P,H,\parallel}$ , as defined in (62), we obtain from (A.10) and (A.12)

$$\begin{aligned} \frac{2\pi m_e}{3} \nabla \cdot \int_0^\infty \vec{f}_1 v_e^5 dv_e &= \frac{5}{2} \left[ \nabla_\perp \cdot \left( n_e T_e \vec{V}_{e\perp} \right) + \frac{\rho_\parallel}{\alpha_\parallel} \nabla_\parallel \left( n_e T_e V_{e\parallel} \right) \right. \\ &\left. + \langle \nu_e \rangle_e (\alpha_P - \rho_P) \frac{\nabla_\perp \cdot \left( \hat{b} \times n_e T_e \vec{V}_{e\perp} \right)}{\Omega_e} \right] - \nabla \cdot \vec{q}_e, \end{aligned} \quad (\text{A.13})$$

where parameters  $\beta_{P,\parallel}$ ,  $\mu_{P,\parallel}$  and tensors  $\mathbf{X}$ ,  $\mathbf{\Lambda}$  are defined by (65), (67), and (60). The last term in (A.13) can be written explicitly as

$$\vec{q}_e \equiv \vec{q}_{eP} + \vec{q}_{eH} + \vec{q}_{e\parallel} = \mathbf{X} \cdot \frac{\nabla T_e}{T_e} + (\mathbf{X} - \mathbf{\Lambda}) \cdot \frac{\nabla n_e}{n_e} \quad (\text{A.14})$$

where

$$\begin{aligned} \vec{q}_{eP} &\equiv \frac{5T_e \langle \nu_e \rangle [\chi_{P\parallel} n_e \nabla_\perp T_e + (\chi_P - \mu_P) T_e \nabla_\perp n_e]}{m_e \Omega_e^2} \\ \vec{q}_{eH} &\equiv \frac{5T_e \hat{b} \times [\chi_{H\parallel} n_e \nabla_\perp T_e + (\chi_H - \lambda) T_e \nabla_\perp n_e]}{m_e \Omega_e} \\ \vec{q}_{e\parallel} &\equiv \frac{5T_e [\chi_{\parallel\parallel} n_e \nabla_\parallel T_e + (\chi_\parallel - \mu_\parallel) T_e \nabla_\parallel n_e]}{m_e \langle \nu_e \rangle}. \end{aligned} \quad (\text{A.15})$$

From Eqs. (44), (51), (A.8), (A.13), neglecting terms that are first- and second-order small with respect to  $\nu_e/\Omega_e$ , we obtain thermal-balance Eq. (58).

## REFERENCES

- Allis, W. P. (1982). Semidivergence of the Legendre expansion of the Boltzmann equation. *Phys. Rev. A* 26, 1704–1712. doi:10.1103/PhysRevA.26.1704
- Bhatnagar, P. L., Gross, E. P., and Krook, M. (1954). A Model for Collision Processes in Gases. I. Small Amplitude Processes in Charged and Neutral One-Component Systems. *Phys. Rev.* 94, 511–525. doi:10.1103/PhysRev.94.511
- Braginskii, S. I. (1965). Transport Processes in a Plasma. *Reviews of Plasma Physics* 1, 205
- Dimant, Y. S. and Oppenheim, M. M. (2004). Ion thermal effects on E-region instabilities: linear theory. *J. Atmos. Solar-Terr. Phys.* 66, 1639–1654. doi:10.1016/j.jastp.2004.07.006
- Dimant, Y. S., Oppenheim, M. M., Evans, S., and Martinez-Sykora, J. (2023). Unified fluid theory of the collisional thermal Farley-Buneman instability including magnetized multi-species ions. *Phys. Plasmas* 30, 102101. doi:10.1063/5.0155500
- Dimant, Y. S. and Sudan, R. N. (1995a). Kinetic theory of low-frequency cross-field instability in a weakly ionized plasma. I. *Phys. Plasmas* 2, 1157–1168. doi:10.1063/1.871394
- Dimant, Y. S. and Sudan, R. N. (1995b). Kinetic theory of low-frequency cross-field instability in a weakly ionized plasma. II. *Phys. Plasmas* 2, 1169–1181. doi:10.1063/1.871395
- Dimant, Y. S. and Sudan, R. N. (1995c). Kinetic theory of the Farley-Buneman instability in the E region of the ionosphere. *J. Geophys. Res.* 100, 14605–14624. doi:10.1029/95JA00794

- Dimant, Y. S. and Sudan, R. N. (1997). Physical nature of a new cross-field current-driven instability in the lower ionosphere. *J. Geophys. Res.* 102, 2551–2564. doi:10.1029/96JA03274
- García-Colín, L. S., Velasco, R. M., and Uribe, F. J. (2004). Inconsistency in the Moment's method for solving the Boltzmann equation. *Journal of Non Equilibrium Thermodynamics* 29, 257–277. doi:10.1515/JNETDY.2004.054
- Grad, H. (1949). On the kinetic theory of rarefied gases. *Communications on Pure and Applied Mathematics* 2, 331–407. doi:https://doi.org/10.1002/cpa.3160020403
- Gurevich, A. V. (1978). *Nonlinear phenomena in the ionosphere*, vol. 10 (New York: Springer Verlag). doi:10.1007/978-3-642-87649-3
- Itikawa, Y. (2006). Cross Sections for Electron Collisions with Nitrogen Molecules. *Journal of Physical and Chemical Reference Data* 35, 31–53. doi:10.1063/1.1937426
- Janhunen, P. (1995). On recent developments in E-region irregularity simulations and a summary of related theory. *Ann. Geophys.* 13, 791–806. doi:10.1007/s00585-995-0791-7
- Kagan, L. M. and Kelley, M. C. (2000). A thermal mechanism for generation of small-scale irregularities in the ionospheric E region. *J. Geophys. Res.* 105, 5291–5302. doi:10.1029/1999JA900415
- Khazanov, G. V. (2011). *Kinetic Theory of the Inner Magnetospheric Plasma* (Springer, New York)
- Kissack, R. S., Kagan, L. M., and St.-Maurice, J. P. (2008a). Thermal effects on Farley-Buneman waves at nonzero aspect and flow angles. I. Dispersion relation. *Phys. Plasmas* 15, 022901. doi:10.1063/1.2834275
- Kissack, R. S., Kagan, L. M., and St.-Maurice, J. P. (2008b). Thermal effects on Farley-Buneman waves at nonzero aspect and flow angles. II. Behavior near threshold. *Phys. Plasmas* 15, 022902. doi:10.1063/1.2834276
- Kissack, R. S., St.-Maurice, J. P., and Moorcroft, D. R. (1997). The effect of electron-neutral energy exchange on the fluid Farley-Buneman instability threshold. *J. Geophys. Res.* 102, 24091–24116. doi:10.1029/97JA02036
- Kissack, R. S., St.-Maurice, J. P., and Moorcroft, D. R. (1995). Electron thermal effects on the Farley-Buneman fluid dispersion relation. *Phys. Plasmas* 2, 1032–1055. doi:10.1063/1.871383
- Kovalev, D. V., Smirnov, A. P., and Dimant, Y. S. (2008). Modeling of the Farley-Buneman instability in the E-region ionosphere: a new hybrid approach. *Ann. Geophys.* 26, 2853–2870. doi:10.5194/angeo-26-2853-2008
- Lifshitz, E. M. and Pitaevskii, L. P. (1981). *Physical kinetics* (Elsevier)
- Milikh, G. M. and Dimant, Y. S. (2003). Model of anomalous electron heating in the E region: 2. Detailed numerical modeling. *J. Geophys. Res.* 108, 1351. doi:10.1029/2002JA009527
- Oppenheim, M., Dimant, Y., Longley, W., and Fletcher, A. C. (2020). Newly Discovered Source of Turbulence and Heating in the Solar Chromosphere. *Astrophys. J.* 891, L9. doi:10.3847/2041-8213/ab75bc
- Oppenheim, M., Otani, N., and Ronchi, C. (1995). Hybrid simulations of the saturated Farley-Buneman instability in the ionosphere. *Geophys. Res. Lett.* 22, 353–356. doi:10.1029/94GL03277
- Oppenheim, M., Otani, N., and Ronchi, C. (1996). Saturation of the Farley-Buneman instability via nonlinear electron ExB drifts. *J. Geophys. Res.* 101, 17273–17286. doi:10.1029/96JA01403
- Oppenheim, M. M., Dimant, Y., and Dyrud, L. P. (2008). Large-scale simulations of 2-D fully kinetic Farley-Buneman turbulence. *Ann. Geophys.* 26, 543–553. doi:10.5194/angeo-26-543-2008
- Oppenheim, M. M. and Dimant, Y. S. (2004). Ion thermal effects on E-region instabilities: 2D kinetic simulations. *J. Atmos. Solar-Terr. Phys.* 66, 1655–1668. doi:10.1016/j.jastp.2004.07.007

- Oppenheim, M. M. and Dimant, Y. S. (2013). Kinetic simulations of 3-D Farley-Buneman turbulence and anomalous electron heating. *J. Geophys. Res.* 118, 1306–1318. doi:10.1002/jgra.50196
- Robinson, T. R. (1998). The effects of small scale field aligned irregularities on E-region conductivities: implications for electron thermal processes. *Advances in Space Research* 22, 1357–1360. doi:10.1016/S0273-1177(98)80034-3
- Rodbard, M. G., Bezerra, J., A. G., and Kremer, G. M. (1995). A combined Chapman-Enskog and Grad method. II. Ionized gases. *Phys. Plasmas* 2, 642–648. doi:10.1063/1.871416
- Schunk, R. and Nagy, A. (2009). *Ionospheres: Physics, Plasma Physics, and Chemistry* (Cambridge: University Press). doi:10.1017/CBO9780511635342
- Shkarofsky, J. P., Johnston, T. W., and Bachynski, M. P. (1966). *The Particle Kinetics of Plasmas* (Reading: Addison-Wesley)
- Song, M.-Y., Cho, H., Karwasz, G. P., Kokoouline, V., and Tennyson, J. (2023). Cross Sections for Electron Collisions with N<sub>2</sub>, N<sub>2</sub><sup>\*</sup>, and N<sub>2</sub><sup>+</sup>. *Journal of Physical and Chemical Reference Data* 52, 023104. doi:10.1063/5.0150618
- St. -Maurice, J. P. and Kissack, R. S. (2000). The role played by thermal feedback in heated Farley-Buneman waves at high latitudes. *Ann. Geophys.* 18, 532–546. doi:10.1007/s00585-000-0532-x
- Stubbe, P. (1990). The concept of a kinetic transport theory. *Phys. Fluids B* 2, 22–33. doi:10.1063/1.859535
- Velasco, R. M., Uribe, F. J., and García-Colín, L. S. (2002). Inconsistencies in moment methods. *Phys. Rev. E* 66, 032103. doi:10.1103/PhysRevE.66.032103