Star Formation in Extreme Environments: A 200 pc High Velocity Gas Stream in the Galactic Centre

V. S. Veena^{1,2}, W.-J. Kim², Álvaro Sánchez-Monge^{3,4}, P. Schilke², K. M. Menten¹, G. A. Fuller^{2,5}, M. C. Sormani⁶, F. Wyrowski¹, W. E. Banda-Barragán^{7,8}, D. Riquelme^{9,10}, P. Tarrío^{11,12}, and P. de Vicente¹²

- ¹ Max-Planck-Institut für Radioastronomie, auf dem Hügel 69, 53121 Bonn, Germany e-mail: vadamattom@mpifr-bonn.mpg.de
- ² I. Physikalisches Institut, Universität zu Köln, Zülpicher Str. 77, 50937 Köln, Germany
- ³ Institut de Ciències de l'Espai (ICE, CSIC), Carrer de Can Magrans s/n, E-08193, Bellaterra, Barcelona, Spain
- ⁴ Institut d'Estudis Espacials de Catalunya (IEEC), Barcelona, Spain
- Jodrell Bank Centre for Astrophysics, Department of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, UK
- ⁶ Università dell'Insubria, via Valleggio 11, 22100 Como, Italy
- ⁷ Escuela de Ciencias Físicas y Nanotecnología, Universidad Yachay Tech, Hacienda San José S/N, 100119 Urcuquí, Ecuador
- 8 Hamburger Sternwarte, University of Hamburg, Gojenbergsweg 112, 21029 Hamburg, Germany
- ⁹ Departamento de Astronomía, Universidad de La Serena, Av. Cisternas 1200, La Serena, Chile
- ¹⁰ Instituto Multidisciplinario de Investigación y Postgrado, Universidad de La Serena, Raúl Bitrán 1305, La Serena, Chile
- ¹¹ Centro de Desarrollos Tecnológicos, Observatorio de Yebes (IGN), 19141 Yebes, Guadalajara, Spain
- Observatorio Astronómico Nacional (OAN-IGN), 28014, Madrid, Spain

Received; accepted

ABSTRACT

Context. The expanding molecular ring (EMR) manifests itself as a parallelogram in the position-velocity diagram of spectral line emission from the Central Molecular Zone (CMZ) surrounding the Galacic centre (GC). It is a high velocity ($|V_{LSR}| > 100 \text{ km s}^{-1}$) extended molecular gas structure. The formation of the EMR is believed to be associated with the bar driven inflow onto the nuclear region of the Galaxy. The physical and chemical properties, as well as the evolution of the EMR and its connection to other GC clouds and the CMZ as a whole, are not yet fully comprehended.

Aims. Using multiwavelength data, we investigate the gas kinematics, star formation activity, and the presence of shocked gas in a 200 pc long high velocity gas stream ($V_{LSR} \sim +150 \text{ km s}^{-1}$) with a double helix morphology named the *helix stream*, that is located 15–55 pc above the CMZ ($l \sim 0^{\circ} - 1.5^{\circ}$; $b \sim 0.05^{\circ} - 0.4^{\circ}$) and is kinematically associated with the EMR/parallelogram.

Methods. To study the kinematics of the helix stream, we used 13 CO (J=2-1) data from the SEDIGISM survey and 12 CO (J=1-0) archival data from the Nobeyama telescope. Additional multiwavelength archival data from infrared to radio wavelengths are used to investigate the star formation activity. We carried out molecular line observations using the IRAM 30m, Yebes 40m, and APEX 12m telescopes. The detection of four rotational transitions of the SiO molecule (J=1-0, 2-1, 5-4, 7-6) indicate the presence of shocks. We derived the SiO column densities and abundances in different regions of the helix stream using the rotational diagram method. We also performed non-local thermodynamic equilibrium (non-LTE) modelling of the SiO emission to analyse the excitation conditions of the shocked gas.

Results. The presence of clumps with submillimeter continuum emission from dust and a candidate H π region signify the ongoing star formation activity within the helix stream. The cloud is massive $(2.5 \times 10^6 \, \mathrm{M_\odot})$ and highly turbulent $(\Delta V_{\mathrm{mean}} = 18 \, \mathrm{km \, s^{-1}})$. We find evidence of cloud-cloud collisions towards the eastern edge $(l \sim 1.3^\circ)$, suggesting a dynamic interaction with the CMZ. An expanding shell is detected within the cloud with radius of 6.7 pc and an expansion velocity of 35 km s⁻¹. The shell might be powered by several supernovae or a single hypernova. The relative abundance of SiO within the helix stream with respect to H₂ implies extensive shock processes occurring on large scales $(X(\mathrm{SiO}) \sim 10^{-9})$. The helical or cork-screw velocity structure observed within the individual strands of the helix stream indicates twisting and turning motions occurring within the cloud.

Conclusions. We propose that the helix stream is the continuation of the near side bar lane, that is overshooting after "brushing" the CMZ and interacting with it at the location of the G1.3 cloud. This interpretation finds support both from numerical simulations and prior observational studies of the CMZ. Our findings carry profound implications for understanding star formation in extreme conditions and they elucidate the intricate properties of gas and dust associated with nuclear inflows in barred spiral galaxies.

Key words. Galaxy: center / Galaxy: evolution / ISM: molecules / ISM: kinematics and dynamics / Radio lines: ISM

1. Introduction

The central part of the Milky Way, the Galactic centre region (GC), located at a distance of 8.2 kpc (Reid et al. 2019; GRAVITY Collaboration et al. 2019), provides an exceptional astrophysical setting to explore the characteristics of radiation

and matter within galactic nuclei. It contains the compact nonthermal radio source Sgr A* that is coincident with the Galactic super-massive black hole (Young et al. 2023). The GC hosts a variety of phenomena such as star formation, large-scale gas flows, magnetic loops, and galactic winds (Kakiuchi et al. 2018; Ponti et al. 2021; Veena et al. 2023; Yusef-Zadeh et al. 2024). The in-




Fig. 1. (Top) 13 CO (2–1) integrated intensity map in the velocity range -200 to +200 km s $^{-1}$. The box in red indicates the area containing the high velocity stream (right panel) and × marks the positions of Sgr A*. (Bottom) 13 CO (2–1) integrated intensity map of the high velocity stream in the velocity range +100 to +200 km s $^{-1}$. The five positions used for SiO observations are marked as × and are labelled.

ner 1 kpc of the GC harbours a large molecular complex that extends ~400 pc and is known as the central molecular zone (CMZ, Morris & Serabyn 1996). The molecular gas in the CMZ exhibits complex structures and kinematics as well as peculiar velocity features. This region contains the largest concentration of high-density molecular gas in the Galaxy (Ferrière et al. 2007). The morphology, kinematics, and physical properties and the chemistry of the molecular gas in the CMZ have been investigated in great detail (e.g., Bally et al. 1987; Morris & Serabyn 1996; Jones et al. 2012, 2013; Ginsburg et al. 2016; Le Petit et al. 2016; Battersby et al. 2020; Henshaw et al. 2023).

The mass accretion onto the Galactic nucleus is primarily governed by inflows from kiloparsec scales (e.g., Englmaier & Gerhard 1999). In barred galaxies, the central bar generates a non-axisymmetric gravitational field inducing non-circular gas motions, with gravitational torques crucial for driving mass inflows (Sorensen et al. 1976; Binney et al. 1991; Hatchfield et al. 2021). "Dust lanes" are structures associated with galactic bars and can be observed in external galaxies seen face-on (Sandage 1961; Dalcanton et al. 2004; Stuber et al. 2023). Theoretical models, combined with observational evidence, indicate that these dust lanes serve as pathways through which gas streams onto the CMZ from distances of ~3 kpc (Sormani & Barnes 2019, and references therein). Identifying these structures, including accretion sites within the CMZ, is challenging due to the solar system's location within a Galactic spiral arm. Clas-

sic characterisation methods involve position-velocity (PV) diagrams (e.g., Marshall et al. 2008).

There exists an extended high velocity gas structure in the GC ($|l| \le \pm 2.0^{\circ}$), known as the expanding molecular ring (EMR), that is vertically extended to $\sim b \pm 0.6^{\circ}$ (Scoville 1972; Oort 1977; Bally et al. 1987; Oka et al. 1998). The EMR is recognized in PV diagrams as a collection of high-velocity arcs resembling a parallelogram at $|V_{LSR}| > 100 \text{ km s}^{-1}$ (Sofue 2017) and was initially thought to be a radially expanding ring of molecular gas. However, the modern and more widely accepted interpretation is that the EMR has its origins in the non-circular motions driven by the Galactic bar that cause infalling gas from dust lanes to overshoot the CMZ (see Henshaw et al. 2023, and references therein). Although the EMR and the CMZ appear to be located close to each other in the GC, the study of the star formation activity and the physical and chemical properties of the EMR have been limited by the fact that many observational studies of the GC region do not extend enough in latitude or do not have the high sensitivity needed to detect low density emission from the EMR, as its molecular lines have a brightness that is an order of magnitude lower than lines in the CMZ.

In this work, we investigate the kinematics and star formation activity of the helix stream (Section 3.1) that is spatially and kinematically associated with the EMR. For our analysis, we use multiwavelength data ranging from infrared to radio wavelengths. The organization of the paper is as follows: The details of data are given in Section 2. Section 3 describes the results of




Fig. 2. ¹³CO (2–1) intensity map in the velocity range +130 to +190 km s⁻¹showing the double helix morphology of the helix stream.

our multiwavelength analysis, whereas in Section 4 we discuss our findings. In Section 5 we present our conclusions.

2. Data

2.1. SEDIGISM survey

The SEDIGISM (Structure, Excitation, and Dynamics of the Inner Galactic Interstellar Medium) project is a spectral line survey of the Milky Way in the J = 2-1 rotational transition of the ¹³CO and C¹⁸O isotopologues of the carbon monoxide molecule (CO), which have rest frequencies of 220.398684 and 219.560354 GHz, respectively (Schuller et al. 2017). It was carried out with the Atacama Pathfinder Experiment (APEX) 12-metre submillimetre telescope (Güsten et al. 2006) and covers 78 deg² in longitude, l, of the inner region of the Galaxy ($-60^{\circ} < l < +18^{\circ}$ with a longitude, b, coverage of $|b| < 0.5^{\circ}$). The survey has a larger latitude coverage in the GC region ($|b| < 1.0^{\circ}$). The effective spectral and angular resolutions are of 0.25 km s⁻¹ and 28", respectively. The typical root mean square (rms) noise of the survey is 0.8 K in main beam brightness temperature, $T_{\rm MB}$. In order to study the ¹³CO emission towards the GC region, we focused on data within the longitude range $l: 0^{\circ} - 1.5^{\circ}$, $|b| < 1.0^{\circ}$. We smoothed the original data cube to a velocity resolution of 1 km s⁻¹ for our analysis. The resultant rms noise is 0.4 K.

2.2. SiO observations with IRAM 30 m telescope

In order to investigate the presence of shocks within the cloud, we use the SiO J=2-1 line with a rest frequency of 86.84699 GHz. Observations in the 3 mm band were carried out using the Institut de Radioastronomie Millimétrique (IRAM) 30-metre telescope during the period 2021 September 09-11 (Project ID: 057-21) with the Eight MIxer Receiver (EMIR; Carter et al. 2012). The standard position switching mode observations towards 5 positions along the cloud (see Fig. 1(Bottom) and Table 1) were covered the frequency range 85.7–109.2 GHz. The half power beam width (HPBW) of the IRAM telescope is 29" at this 87 GHz. We used the FTS200 backend with a spectral resolution of 200 kHz (equivalent to 0.6 km s⁻¹at 3 mm) and, as for the Yebes and Yebes data (see below), the data reduction was performed using the Continuum and Line Analysis Single-dish Software (CLASS) which is part of the GILDAS software (Pety 2005). In order to convert antenna temperatures (T_A^*) to main beam temperatures $T_{\rm MB}$, we used the relation $T_{\text{MB}} = F_{\text{eff}}/B_{\text{eff}} \times T_{\text{A}}^*$, where $F_{\text{eff}} = 0.95$ is the forward efficiency and $B_{\text{eff}} = 0.81$ is the main beam efficiency.

2.3. Yebes 40 m SiO observations

In order to probe the J=1-0 transition of SiO (ν_0 =43.42385 GHz), observations in 7 mm were carried out using the Yebes 40 m telescope on September 27, 2021 (Project ID: 21A017) with the HEMT (High Electron Mobility Transistors) receiver. It allows broad-band observations with an instantaneous bandwidth of 18 GHz in two linear polarisations. The backends used are Fast Fourier Spectrometers (FFTs) with 2.5 GHz bandwidths and a spectral resolution of 38 kHz (Tercero et al. 2021). The HPBW of the Yebes telescope is 40.6" at this frequency. The position switching mode observations towards 5 positions (Fig. 1(Bottom) and Table 1) covered the frequency range 31.5-50 GHz. The spectra were measured in units of antenna temperature and in order to convert this to $T_{\rm MB}$, we used a beam efficiency factor of 0.55.

2.4. APEX SiO observations

To probe the J=5-4 ($v_0=217.10492$ GHz) and J=7-6(ν_0 =303.9268 GHz) transitions of SiO, we used the APEX telescope (Project IDs: M9518C_109, M9518A_109). The position switching observations were carried out towards 5 positions (Fig. 1(Bottom) and Table 1) using the nFLASH230 receiver and the 7 pixel Large APEX sub-Millimetre Array (LASMA). The 2 sideband nFLASH230 receiver is connected to two FFT backends, providing a simultaneous coverage of two sidebands with a bandwidth of 7.9 GHz each. The separation between the centre of the two sidebands is 16 GHz. LASMA pixels consists of sideband separating mixers (2SB) which provides an IF bandwidth of 4-8 GHz for each of the two sidebands. The HPBW are 28.5'' and 20'' for the SiO (5–4) and (7–6) lines, respectively. In order to convert antenna temperatures to T_{MB} , we used beam efficiency factors of 0.83 and 0.74 for the nFLASH230 and LASMA receivers.

3. Results

3.1. A high velocity molecular cloud in the GC with double helix morphology

Fig. 1(Top) shows the ^{13}CO (2–1) integrated intensity map of the GC region in the velocity range from -200 to $+200~\text{km}\,\text{s}^{-1}$. The image shows emission that extends predominantly along, mostly slightly below, the Galactic plane, which represents the CMZ. On the positive latitude end of the distribution, diffuse emission is observed that arches toward higher latitudes, up to $\sim 0.4^{\circ}$. Fig. 1(Bottom) presents the zoomed-in view of the above




Fig. 3. 12 CO(1–0) integrated intensity map (Oka et al. 1998) of the helix stream (shown in Fig. 1(Right) and Fig. 2) in the velocity range $100-200 \text{ km s}^{-1}$. The white straight line AB used to construct the PV diagram is marked. The five positions used for SiO observations are marked as × and are labelled.

Table 1. List of 5 positions used for spectral line observations with IRAM, APEX and Yebes telescopes

Position	RA(J2000) (hms)	Dec(J2000) (° ' ")	l (Deg)	b (Deg)
P1	17:47:14.23	-27:39:23.22	1.278	0.361
P2	17:46:46.15	-27:50:54.03	1.061	0.349
P3	17:45:32.49	-28:20:59.21	0.492	0.320
P4	17:45:11.09	-28:38:30.31	0.202	0.235
P5	17:45:19.04	-28:46:22.86	0.105	0.142

mentioned region, integrated within the smaller velocity range +100 to $+200~km~s^{-1}$. Bright, elongated emission extending over $\sim 1.4^\circ$ (in longitude) is observed. The width of the emission feature is $\sim 0.15^\circ$ (in latitude) and the corresponding aspect ratio is 9.3, implying that the structure is filament-like in morphology. It has a tilt with an inclination angle $\sim 8.5^\circ$, with respect to the Galactic plane. The CO cloud exhibits multiple clumpy substructures. The high velocity of the cloud (>100 km s $^{-1}$) cannot be explained with a simple Galactic rotation model (Sofue 2017). Its unique location in the GC and large velocities suggest its association with the EMR. Assuming that the cloud is at the distance of the GC (D=8.2±1 kpc), we estimate its extent to be 200 pc \times 21.5 pc. The cloud is located 15 - 55 pc above the Galactic mid-plane.

To investigate further the intricate substructure of the cloud, we created nine intensity maps integrated within 10 km s⁻¹ velocity intervals, which are presented in Fig. A.1. Several clumpy features are observed. The lowest velocity features correspond to the emission from the edges of the filament ($100-130 \text{ km s}^{-1}$). The emission from the central region is evident in higher velocity images $(+130 \text{ to } +190 \text{ km s}^{-1})$. In order to analyse the morphology of the emission from the central region, which can be visualised better with a lower dynamic range image, we generated an additional integrated intensity map of emission in the high velocity range (i. e., $+130 \text{ to } +190 \text{ km s}^{-1}$). The resultant image is presented in Fig. 2. The emission has a remarkable "double helix" morphology containing two helically wound strands. The origin of this double helix morphology is explored in Section 4.3. The cloud will be addressed as the *helix stream* in the following sections.

3.2. Kinematics of the helix stream

3.2.1. Position-velocity diagram

In order to study the velocity structure and large-scale kinematics of the helix, we used ¹²CO maps from the GC CO survey

by Oka et al. (1998), who mapped the GC region in the ¹²CO (J=1-0) transition using the 45 m telescope at Nobeyama Radio Observatory. The spectral resolution and grid spacing are 0.65 km s⁻¹ and 34", respectively. The advantage of using ¹²CO (1–0) emission (critical density $\approx 2 \times 10^3 \text{ cm}^{-3}$) to study the large-scale cloud kinematics is that we are sensitive to regions of low density (and low column density) and thus can also probe the velocity structure of the diffuse envelope surrounding the helix stream. However, it is important to note that in denser (and higher column density) regions, opacity effects may occur, potentially affecting the velocity measurements. Fig. 3 shows the integrated intensity map of the ¹²CO emission integrated over the same velocity range as in Fig. 1(Right). The overall morphology and spatial extent of the ¹²CO(1–0) emission distribution is similar to that of the ${}^{13}CO(2-1)$ emission except for the fact that the ¹²CO emission appears more extended and clumpier compared to the sharp emission features observed in the latter, only tracing the denser regions. To explore the velocity distribution of the region, we construct a PV diagram along the major axis of the cloud (see Fig. 3).

From the PV diagram (Fig. 4(Left)), we identify a wide variety of emission features with velocities ranging from ~0 to +200 km s⁻¹. The emission corresponding to the helix stream is seen as an elongated feature in the velocity range ~ 100 – 200 km s⁻¹. Towards the eastern edge of the cloud ($l \sim 1.3^{\circ}$), several clumpy features are observed in the velocity range from ~ 70 to $100 \, \mathrm{km \, s^{-1}}$. These are connected to the high velocity emission with bridge-like features. Such bridge-like features have been found to exist in several Galactic molecular clouds and are believed to be signatures of cloud-cloud collisions (e.g., Haworth et al. 2015; Sormani et al. 2019). The bridge feature corresponds to the region in the 2D intensity map (Fig .1(Right)) where the filament slightly arches downwards towards the CMZ. As the 100 km s⁻¹ clumps are further connected to lower velocity features consistent with velocities of CMZ clouds (0-100 km s⁻¹), this indicate a possible interaction between the CMZ and the helix stream. Another prominent feature in the PV diagram is a ring-like structure extending $\sim 0.26^{\circ}$ (36 pc), centred at $l \sim 0.42^{\circ}$ (marked as a dashed ellipse in Fig. 4(Left)). Such ring or arc-like features are associated with expanding shells of gas within the cloud (Arce et al. 2011) and are the signatures of possible stellar feedback. We explore the origin of the expanding shell in Section 4.1. Apart from velocity sub-structures such as the bridge, the ring and the 100 km s⁻¹clump, a large scale monotonous velocity variation of $\sim 100 \, \mathrm{km \, s^{-1}}$ along the filament is seen, which corresponds to a gradient of $0.5 \text{ km s}^{-1} \text{ pc}^{-1}$.

3.2.2. Spectral decomposition using GAUSSPY+

To analyse the velocity distribution of the dense gas in detail, we employed pixel-by-pixel spectral decomposition of the ¹³CO (2–1) data using the GAUSSPY+ package (Lindner et al. 2015; Riener et al. 2019). We chose data of the ¹³CO (2–1) line instead of ¹²CO(1–0) as ¹³CO is less abundant than ¹²CO by factors of ~20–30 in the GC, and its lines are less affected by optical depth effects. GAUSSPY+ is based on GaussPy, a python based Autonomous Gaussian Decomposition algorithm, which is used to analyse complex spectra arising in the interstellar medium (ISM) by decomposing them into multiple Gaussian components. It involves a technique called derivative spectroscopy, and automatically determines the initial guesses for the Gaussian components for individual spectra. In the later stages of the fitting routine, GAUSSPY+ also carries out an automated spatial refitting based




Fig. 4. (Left) ¹²CO(1–0) PV diagram of the helix stream along the cut AB indicated in Fig. 3 overlaid with the position-velocity loci of Hi-GAL prestellar (blue dots) and protostellar (red triangles) clumps discussed in Section 3.4. (Right) The ¹³CO (2–1) PPV plot of the helix stream generated from spectral decomposition using GAUSSPY+. Each velocity component is colour coded according to its velocity dispersion. Ellipse marks the shell feature discussed in Section 4.1.

on neighbouring fit solutions. We adopted the default parameters for the spectral decomposition provided by Riener et al. (2019) except for the smoothing parameters $\alpha 1$ and $\alpha 2$ for which we generated a training set consisting of 5000 random spectra from the data cube. Using the training set, we estimated $\alpha 1$ and $\alpha 2$ to be 1.23 and 4.55, respectively. The spectra towards the bridge region were fitted with up to 8 components. However, on average, three components were fitted to a typical spectrum.

In Fig. 4(Right), all the identified components above a signal-to noise ratio of 5 and velocities exceeding 70 km s⁻¹ are presented as a position-position-velocity (PPV) plot. The data are colour coded according to the velocity dispersion of the corresponding component. The overall morphology of the features observed in the PPV diagram is consistent with that of the ¹²CO PV diagram, though the extended diffuse emission is not as evident as in the ¹²CO PV diagram owing to the fact that the ¹³CO emission traces higher column density gas compared to the former. The bridge feature seen in the ¹²CO PV diagram is also observed in the PPV plot. The ring-like feature identified in the PV diagram is clearly evident in the ¹³CO PPV plot as a spherical shell with a central cavity.

Fig. 5 presents a histogram of velocity dispersions (based on Full Width at Half Maximum (FWHM) of the $^{13}\mathrm{CO}$ emission line) from the spectral decomposition using GAUSSPY+. The velocity dispersion is found to range from 1.0 to 25 km s $^{-1}$ with a mean of 7.8 km s $^{-1}(\sigma_{\mathrm{median}}=7.2~\mathrm{km\,s^{-1}})$. As seen in the plot, the histogram of the velocity dispersion follows a lognormal distribution. The observed velocity dispersions in the helix stream are consistent with velocity dispersion measurements of clouds in the CMZ region ($\sigma_{\mathrm{median}}=9.8~\mathrm{km\,s^{-1}}$; Henshaw et al. 2016). Highest velocity dispersion (i.e., $\sigma>15~\mathrm{km\,s^{-1}}$) is observed mainly towards the 100 km s $^{-1}$ eastern clump, the connecting bridge, and the shell-feature at $l\sim0.42^\circ$. As the observed line width could have contributions from thermal as well as non-thermal motions, we proceed to estimate the magnitudes of each component using the following expressions (e.g., Myers 1983)

$$\sigma_{\rm T} = \sqrt{\frac{k_{\rm B} T_{\rm kin}}{\mu m_{\rm H}}} \tag{1}$$

$$\sigma_{\rm NT} = \sqrt{(\sigma)^2 - (\sigma_{\rm T})^2},\tag{2}$$

where $\sigma_{\rm T}$ and σ correspond to the thermal and the observed dispersion, $T_{\rm kin}$ is the kinetic temperature, μ is the molecular weight of $^{13}{\rm CO}$ that is 29, $k_{\rm B}$ is the Boltzmann constant and $m_{\rm H}$, the mass of the hydrogen atom. Following the approach of Henshaw et al. (2016), we assume a fiducial temperature range of 60–100 K typical for the CMZ. This corresponds to thermal velocity dispersion ranging between 0.1–0.2 km s⁻¹ and is thus mostly neglible. The obtained non-thermal $\sigma_{\rm NT}$ dispersion can be used to estimate the Mach number using the expression

$$\mathcal{M}_{3D} \approx \sqrt{3} \frac{\sigma_{\rm NT}}{c_{\rm s}}$$
 (3)

where $c_s = \sqrt{k_B T_{kin}/\mu_p m_H}$ is the isothermal sound speed and μ_n =2.33 is the mean weight for molecular hydrogen gas. For the same temperature range of 60–100 K, c_s is found to range from $0.5-0.6 \,\mathrm{km}\,\mathrm{s}^{-1}$. For the mean velocity dispersion of 7.8 km s⁻¹, the Mach number \mathcal{M}_{3D} ranges between 23–27 at a scale of 1 pc (equivalent to the angular resolution of the ¹³CO data). This suggests that on parsec-scales, the gas motions within the helix stream are on the average, highly supersonic. However, there is a caveat that the obtained Mach number could be an upper limit, in case of ordered gas flows and substructures present at sub-parsec scales that are unresolved. Towards the circular shell and the eastern bridge seen in the PV and PPV plots, the Mach numbers exceed 50. This could be explained as increased turbulence as a result of expanding gas motions within the cavity and/or cloud-cloud collisions along the bridge. In all cases, our estimates of \mathcal{M}_{3D} suggest that the gas harbours strong turbulence.

3.2.3. Size-velocity dispersion relation

Spectral line observations of molecular clouds reveal a power-law correlation between the velocity dispersion and the projected size, with a power-law index ranging between 0.2 to 0.6 (e.g., Larson 1981; Solomon et al. 1987; Caselli & Myers 1995). This is known as the Larson's scaling relationship. In this section, we examine the velocity dispersion-size relationship of the helix stream derived from ¹³CO (2–1) data. For this, we first identify the relevant structures within the cloud using the dendrogrambased structure identification python package ASTRODENDRO (Rosolowsky et al. 2008). The dendrogram is a powerful technique to analyse hierarchical structures within a molecular cloud.




Fig. 5. Histogram of velocity dispersion of spectra decomposed using GAUSSPY+. Blue line corresponds to the lognormal fit to the histogram.

In order to compute the dendrogram, three parameters are required: (i) min_value, the minimum intensity threshold taken as $3\sigma_{\rm rms}$ where $\sigma_{\rm rms}$ is the rms noise, (ii) min_delta that is the minimum height of the identified structure, taken to be $1.5\sigma_{\rm rms}$, and (iii) min npix, the minimum number of pixels within a structure, considered to be 6, equivalent to two beams in area. The structures are categorised into trunks (largest continuous structures), branches (intermediate structures), or leaves (smallest substructures). For each structure, we compute the size R, extracted from the geometric mean of the semi-major and semi-minor axis (radius parameter in ASTRODENDRO) i.e., $R = \eta \sqrt{\sigma_{maj}\sigma_{min}}$ where η , the factor that relates the radius of a spherical cloud to its one-dimensional rms size, is assumed to be 1.91 (e.g., Solomon et al. 1987; Mazumdar et al. 2021). The velocity dispersion σ corresponds to the v_{rms} parameter, that is the intensity-weighted second moment of velocity. Using ASTRODENDRO, we identify 635 distinct structures within the cloud.


The σ -R plot of the helix stream is presented in Fig. 6(Top). The largest structures identified within the cloud have R \sim 10–20 pc and the corresponding velocity dispersions are in the range of 9–12 km s $^{-1}$, whereas a systematic decrease is observed in the velocity dispersions of smaller structures. For comparison, we have also plotted data corresponding to other Galactic clouds (Caselli & Myers 1995; Oka et al. 2001b; Heyer et al. 2009). From the plot, it appears that σ is correlated with R. To quantify this, we fitted the data with a power-law of the form

$$\left(\frac{\sigma}{\mathrm{km}\,\mathrm{s}^{-1}}\right) = \alpha \left(\frac{\mathrm{R}}{\mathrm{pc}}\right)^{\beta}$$

From the χ^2 fit, we estimated the power-law index β and coefficient α as 0.7 ± 0.01 and 1.90 ± 0.05 , respectively. The obtained index is steeper than the classical power-law index of 0.5 (Larson 1981) and is comparable to that of CMZ clouds (~ 0.66 , Shetty et al. 2012; Kauffmann et al. 2017).

3.2.4. Cloud stability and dynamics

In order to investigate whether the helix stream is stable against gravitational collapse, we examined the modified Larson's relationship, known as the Heyer relationship (Heyer et al. 2009). The surface densities (Σ) of the dendrogram structures are calculated and plotted against $\sigma/R^{0.5}$ in the bottom panel of Fig. 6. We find a systematic variation of $\sigma/R^{0.5}$ with Σ . The structures within the helix stream deviate from the loci of gravitationally






Fig. 6. (Top) Comparison of size versus velocity dispersion in the helix stream (red dots) with values of Galactic centre clouds from Oka et al. (2001b, green ×), the Orion A and B giant molecular clouds (magenta stars; Caselli & Myers 1995) and other Galactic giant molecular clouds (blue crosses; Heyer et al. 2009). The dotted line corresponds to the best-fit result from Solomon et al. (1987), $\sigma = 0.7R^{0.5}$. The dashed line corresponds to the best-fit result for the helix stream $\sigma = 2.97 R^{0.70}$. (Bottom) Heyer's relation (surface density, Σ , versus $\sigma/R^{0.5}$) for different Galactic clouds. Dots correspond to the structures within the helix stream, squares correspond to structures within the helix stream where follow-up molecular observations are carried out, crosses represent Galactic giant molecular clouds (Heyer et al. 2009), and triangles represent high-density cores in the GC 'dust-ridge' clouds (Barnes et al. 2019). Black curves indicate solutions of pressure-bounded virial equilibrium and dotted lines correspond to α_{vir} =1.0, 3.0, and 10.0 where there is negligible external pressure.

bound clouds where the virial parameter, $\alpha_{\rm vir}$, equals 1. The stability of a cloud can be defined based on the virial critical parameter ($\alpha_{\rm crit}$). Unbound, sub-critical clouds are characterised by $\alpha_{\rm vir} > \alpha_{\rm crit}$, whereas gravitationally bound, supercritical clouds are characterised by $\alpha_{\rm vir} < \alpha_{\rm crit}$. According to Kauffmann et al. (2013), for non-magnetised clouds, $\alpha_{\rm crit} \gtrsim 2$, whereas for strongly magnetised clouds, $\alpha_{\rm crit} \ll 2$. Heyer et al. (2009) estimated the mean $\alpha_{\rm vir}$ as 1.9, whereas for the helix stream, it ranges between 2.7 and 36.5, with a mean of 9.5, implying that a majority of the structures within the helix stream are highly subcritical and unbound. Field et al. (2011) suggested that the external pressure ($P_{\rm e}$) introduces an additional confining force that




Fig. 7. ¹³CO integrated intensity map overlaid with Hi-GAL prestellar (green asterisks) and protostellar (magenta asterisks) clumps. Red triangle corresponds to the location of the radio source GPSR5 0.431+0.262. Blue circle roughly corresponds to the location of ring-like feature observed in the PV diagram. The five positions used for SiO observations are marked as ×.

Table 2. Properties of Herschel Hi-GAL compact sources identified within the helix stream

Source ID	GLON (Deg)	GLAT (Deg)	Diameter (pc)	Distance ^a (kpc)	Velocity (km s ⁻¹)	$ ext{Mass}^b \ (10^2 ext{M}_\odot)$	$T_{ ext{dust}}^{b}$ (K)	$L_{\rm Bol}$ $(10^3 {\rm L}_{\odot})$	Σ (g cm ⁻²)	Class ^c
HIGALBM0.1020+0.1502	0.1020	0.1502	1.9	8.4	93.7	55.5	16.4	14.2	0.39	2
HIGALBM0.1017+0.1750	0.1017	0.1749	1.1	8.4	113.8	50.5	14.9	4.4	1.19	1
HIGALBM0.1129+0.1423	0.1129	0.1423	1.3	8.4	88.9	14.3	15.9	2.3	0.22	1
HIGALBM0.3521+0.2479	0.3521	0.2479	1.0	8.4	112.4	9.3	15.5	2.5	0.27	2
HIGALBM0.3736+0.2481	0.3736	0.2481	1.0	8.4	110.4	43.2	12.0	1.1	1.13	1
HIGALBM0.3944+0.2428	0.3944	0.2428	1.1	8.4	105.9	14.3	15.0	2.0	0.34	2
HIGALBM0.4322+0.2101	0.4322	0.2100	0.9	8.4	114.1	1.5	28.2	7.3	0.05	1
HIGALBM0.4677+0.3197	0.4677	0.3197	0.8	8.4	154.2	4.1	16.7	0.9	0.16	2
HIGALBM0.5138+0.3377	0.5138	0.3377	1.4	8.3	162.9	9.4	14.8	1.4	0.12	2
HIGALBM0.6988+0.3696	0.6988	0.3696	1.4	8.4	153.8	2.2	17.9	0.9	0.03	2
HIGALBM0.9817+0.3846	0.9817	0.3846	1.2	8.4	162.1	17.1	13.6	1.0	0.31	1
HIGALBM1.0573+0.3538	1.0573	0.3538	0.7	8.4	171.6	55.7	10.4	0.5	3.02	1
HIGALBM1.0620+0.3648	1.0620	0.3648	0.4	8.4	166.8	2.5	18.8	1.0	0.48	2
HIGALBM1.2317+0.3524	1.2317	0.3524	0.9	8.4	174.1	12.2	14.8	1.5	0.38	2
HIGALBM1.2876+0.2928	1.2876	0.2928	1.5	8.6	110.6	33.3	13.0	1.8	0.41	2
HIGALBM1.2882+0.2743	1.2882	0.2743	1.0	8.6	103.1	51.4	10.8	0.7	1.47	1
HIGALBM1.3052+0.2733	1.3052	0.2733	0.7	8.6	116.6	5.7	14.9	0.7	0.31	2
HIGALBM1.3390+0.2649	1.3390	0.2649	1.2	8.6	108.8	21.9	13.0	1.9	0.43	2
HIGALBM1.3435+0.2596	1.3435	0.2596	1.2	8.6	102.7	137.4	10.3	1.3	2.64	1
			-							

Notes. (a) Mège et al. (2021), (b) Obtained by fitting the SEDs in the spectral range $160 \, \mu\text{m} \le \lambda \le 500 \, \mu\text{m}$ with a modified blackbody function, (c) 1: prestellar, 2: protostellar

is not accounted under the simple virial equilibrium assumption, which only considers gravitational and kinetic energies. Based on their analysis, the data from Heyer et al. (2009) that appears to systematically deviate from the simple virial equilibrium can be in pressure-bounded virial equilibrium (PVE), if the external force $P_{\rm e}/k_{\rm B}$ is in the range 10^4 – 10^6 K cm⁻³. The solutions to the PVE, as proposed by Field et al. (2011), can be written as


$$\frac{\sigma^2}{R} = \frac{1}{3} (\pi \Gamma G \Sigma + \frac{4P_e}{\Sigma}) \tag{4}$$

where Γ is related to the density structure of the cloud. Here, for simplicity, we assume $\Gamma=0.73$, corresponding to a centrally concentrated density structure (e.g., Field et al. 2011; Walker et al. 2018). The solutions to the PVE appear to be V-shaped in the Σ versus $\sigma/R^{0.5}$ plot, observed in the bottom panel of Fig. 6.

Assuming that the helix stream is in PVE, the external pressure has to be of the order of $P_{\rm e}/k_{\rm B}\sim 10^5-10^7~{\rm K~cm^{-3}}$. In Section 3.2.2, we find that there is highly supersonic turbulence within the helix stream. This turbulent pressure could be responsible for the PVE within the helix stream.

3.3. Column density and mass

To estimate the mass and column densities of the cloud, we followed the X-factor method, to convert the integrated intensities of ^{13}CO (2–1) emission to H₂ column density. For this, we used the mean conversion factor $X_{^{13}\text{CO}(2-1)}=1.08\times10^{21}$ (K km s $^{-1}$) estimated by Schuller et al. (2017). The integrated intensity map in the velocity range +100 to +200 km s $^{-1}$ is converted to a column density map. The peak column density is estimated to be 8.7×10^{22} cm $^{-2}$, whereas the average column density along the






Fig. 8. (Top) Mass-radius relation of 19 Hi-GAL clumps. The shaded area represents range of masses consistent with low mass star formation, satisfying the criteria M≤870 M_{\odot} (r/pc)^{1.33} (Kauffmann & Pillai 2010). The dotted lines indicate surface density thresholds of 1 and 0.05 g cm⁻², respectively. (Bottom) Bolometric luminosity-mass relation of 19 Hi-GAL clumps. Dotted lines represent (L/M)=0.1, 1.0, and 10.0 L_{\odot}/M_{\odot} , respectively.

filament is $2.0\times10^{22}~\text{cm}^{-2}$. The total column density $N(H_2)$ can be used to estimate the mass of the helix stream using the equation

$$M = N(H_2) \mu_g m_H A \tag{5}$$

where M is the total mass of the cloud, μ_g is the mean weight of molecular gas taken as 2.86 assuming that the gas is 70% molecular hydrogen by mass (Ward-Thompson et al. 2010), m_H is the mass of the hydrogen atom, and A is the area of the cloud calculated by adding the area of individual pixels within the 5σ contour of ¹³CO emission. The total mass of the helix stream is thus estimated to be 2.5×10^6 M $_{\odot}$. We would like to caution that the estimated mass could be affected by optical depth effects towards the peak positions of the high density clumps as well as a lower X₁₃CO(2-1) factor in the GC region (e.g., Dahmen et al. 1998; Gramze et al. 2023). We also estimated column densities and mass of the cloud using the H₂ column density map produced by the point process mapping (PPMAP) method (Marsh et al. 2017). The PPMAP uses Hi-GAL continuum data in the wavelength range 70-500 μ m and it takes complete account of the point spread functions (PSFs) of the telescope used, resulting in a spatial resolution of 12". Dust temperatures and column densities are determined by fitting a spectral energy distribution (SED) assuming an opacity of 0.1 cm²g⁻¹ at 300 μ m, a power-law index β of 2, and a dust to gas ratio of 100. The PPMAP derived dust temperature and column density maps of the helix stream region are presented in Fig. A.2. The peak column density is 1.2×10^{23} cm⁻² and the mean column density is 2.3×10^{22} cm⁻², similar to the column density estimates based on ^{13}CO emission. We estimate the total mass of the cloud from the column density using equation 5. Using the PPMAP method we find the mass of the cloud to be 4.2×10^6 M $_{\odot}$. This estimate is an upper limit as the source is in the GC region and there could be contributions from dust associated with the ISM along the line of sight. In addition, there could be uncertainties due to the assumed dust to gas ratio. The agreement between the mass estimates from X-factor and PPMAP methods, each falling in the order of 10^6 M $_{\odot}$, gives confidence in our results, despite methodological variations and potential uncertainties.

3.4. Dust clumps associated with the helix stream

In order to study the properties of the cold dust clumps associated with the cloud, we use the Herschel Hi-GAL compact source catalogue (Elia et al. 2017, 2021) which is a bandmerged catalogue of cold dust clumps identified in the Herschel wavelength bands (70–500 μ m). The catalogue comprises sources that are detected in at least three consecutive Herschel bands. The distances to the clumps are estimated based on methods described in Mège et al. (2021) and physical properties are computed using a single temperature graybody fit to the SEDs. Clumps are classified into three different categories: starless clumps, prestellar clumps, and protostellar clumps. Starless clumps are objects that are gravitationally unbound. Prestellar clumps are gravitationally bound clumps that are not associated with 70 μ m sources. Protostellar clumps are those clumps with star formation activity, i.e., indicated by the presence of 70 μ m sources.

From the Hi-GAL catalogue, we found 19 clumps within the 5σ contour of the ¹³CO intensity map. We selected only clumps whose velocities range between 70 and 200 km s⁻¹, corresponding to velocities of observed kinematic features in the PV diagram. The distribution of Hi-GAL clumps along the helix stream is shown in Fig. 7 and properties of individual clumps are listed in Table 2. Of 19 clumps, 8 are in the prestellar phase and 11 are in the protostellar phase. Masses of clumps range between 1.5 and 137.4×10^2 M_{\odot} and luminosities range between 0.7 and 14.2×10^3 M_{\odot}. The radius-mass relation of these sources is presented in Fig. 8(Top). Of 19 clumps, 17 satisfy the criteria for massive star formation defined by Kauffmann & Pillai (2010), i.e., $M > 870 M_{\odot} (r/pc)^{1.33}$. The surface densities of 5 prestellar clumps exceed the surface density threshold of 1 g cm⁻² for massive star formation, defined by Krumholz & McKee (2008). In Fig. 8 (Bottom), we present the mass-luminosity relationship of the Hi-GAL clumps, which is considered to be a good diagnostic of a clump's evolutionary (Molinari et al. 2008). The bolometric luminosity to mass ((L/M)) ratio of all, except for one clump, lies in the range $0.1-10.0 L_{\odot}/M_{\odot}$. Of these, 8 clumps have $(L/M) < 1.0 L_{\odot}/M_{\odot}$. Overall, the low (L/M) ratio of the Hi-GAL clumps suggests that they are in relatively early evolutionary phases (Giannetti et al. 2017) in which the gas is still accumulated and compressed ($(L/M) \lesssim 2.0 L_{\odot}/M_{\odot}$) or the young stellar objects gaining mass $(2.0 L_{\odot}/M_{\odot} \lesssim (L/M) \lesssim 40.0 L_{\odot}/M_{\odot})$. The distribution of 19 Hi-GAL clumps in the position-velocity space, as depicted in Fig. 4(Left), reveals a notable concentration towards the bridge and shell features in the PV diagram. This spatial clustering suggests a scenario of triggered star formation, wherein the interaction of molecular gas within these




Fig. 9. SiO (1–0), (2–1), (5–4) and (7–6) and H¹³CO⁺(2–1) spectra toward positions P1, P2, P3, P4 and P5. Gaussian fits to the SiO and H¹³CO⁺(2–1) spectra are shown in red and blue, respectively.

regions leads to enhanced densities and the initiation of gravitational collapse. Such triggering could be driven by external factors such as shock fronts, turbulence, or feedback effects.

3.5. Massive star formation within the expanding CO shell

In Section 3.2, we identified ring/shell-like structures in the PV and PPV plots that are associated with a shell-like feature in the integrated intensity map. Here we explore the origin of the CO shell and its implications on the star formation activity in the helix stream. The approximate centre of the cavity in the intensity plot is at $(l,b)=(0.42^{\circ},0.26^{\circ})$ with an angular diameter of $\sim 9.0'$, which translates to a diameter of 22 pc (see Fig. 7). As mentioned in Section 3.2.1, an expanding shell of gas would create a ring-like feature in the PV and PPV plots (Arce et al. 2011). The velocity of the circular feature in the PV diagram ranges between $100-170~{\rm km\,s^{-1}}$, which translates to an expansion velocity of $\sim 35~{\rm km\,s^{-1}}$.

As mentioned in Section 3.4, the distribution of Hi-GAL sources reveals a clustering of sources in the region. Of these, two are protostellar clumps. In addition, we also searched for thermal radio sources within the expanding shell. We find a source GPSR5 0.431+0.262 from the VLA survey for compact radio sources near the GC by Lazio & Cordes (2008), marked in Fig. 7. The object is detected at 1 GHz and 5 GHz bands (angular diameter of 1.9"). The radio spectral index measurements by Lazio & Cordes (2008) show that it is thermal in nature with a spectral index $\alpha = 0.2$. The angular size comparable to the image resolution of the data $(2.4'' \times 1.3'')$ suggests that the source remains unresolved. At the GC distance, this gives an upper limit diameter of 0.1 pc. Assuming the object to be an H II region, we estimate the Lyman continuum photon rate of the H II region under the assumptions of optically thin emission and negligible absorption by the dust (Mezger & Henderson 1967; Schmiedeke et al. 2016)

$$\left(\frac{N_{Lyc}}{s^{-1}}\right) = 4.771 \times 10^{42} \left(\frac{S_{\nu}}{Jy}\right) \left(\frac{T_e}{K}\right)^{-0.45} \left(\frac{\nu}{GHz}\right)^{0.1} \left(\frac{d}{pc}\right)^2$$
 (6)

Here S_{ν} is the flux density at frequency ν taken as 26.7 mJy at 5 GHz (Becker et al. 1994), T_{e} is the electron temperature, and d is

the distance to the source that is 8.2 kpc. For estimating the electron temperature in the region, we use the electron temperature gradient curve across the Galactic disk (Quireza et al. 2006) and consider an approximate value of 7000 K at the GC. The resultant Lyman continuum photon rate is found to be $1.87 \times 10^{47} \, \text{s}^{-1}$. Considering that the H II region is excited by a single zero age main sequence (ZAMS) star, we find the spectral type (Panagia 1973) of the source to be B0–B0.5.

The source is also detected in the infrared bands (see Fig. A.3) as a compact source. However, no nebulous infrared emission is detected within the interior of the shell (see Fig. A.3). HII regions have characteristic infrared colors, that are useful for identifying these objects (e.g. Wood & Churchwell 1989; Hughes & MacLeod 1989; Anderson et al. 2012). According to Makai et al. (2017), the infrared colors $\log_{10}(F_{24 \mu m}/F_{12 \mu m}) \ge 0$ and $\log_{10}(F_{70 \mu m}/F_{12 \mu m}) \ge 1.2$, and $\log_{10}(F_{24 \mu m}/F_{12 \mu m}) \ge 0$ and $\log_{10}(F_{160 \mu m}/F_{70 \mu m}) \le 0.67$ reliably identify H II regions, independent of their size. To determine the infrared flux densities of the source, we use the MIPSGAL (Gutermuth & Heyer 2015) and the Hi-GAL (Molinari et al. 2016) point source catalogues. For our source, we find $\log_{10}(F_{24 \mu m}/F_{12 \mu m}) = 0.9$, $\log_{10}(F_{70 \mu m}/F_{12 \mu m}) = 1.2$, and $\log_{160}(F_{70 \mu m}/F_{12 \mu m}) = 0.4$, consistent with the HII region scenario. The identification of a potential H_{II} region and the presence of clustered Hi-GAL clumps suggest that the shell's origin is likely due to feedback effects resulting from ongoing star formation activity. We will explore this scenario in detail in Section 4.1.

3.6. SiO emission: evidence of large scale shocks

To investigate the presence of shocks, in particular in view of the high Mach numbers discussed above, we carried out SiO observations toward 5 positions along the helix stream. SiO is a salient tracer of shocks as it is produced through the sputtering of Si-bearing material in grains (Schilke et al. 1997). The target positions are marked in Fig. 7. Four rotational transitions of SiO are observed and the details of these transitions are presented in Table 3. The resultant SiO spectra are presented in Fig. 9. SiO emission is observed toward all five positions and all positions except P2 have two velocity components. Line parameters were extracted by fitting Gaussian components to the




Fig. 10. Rotation diagrams for J = 1-0, 2-1, 5-4 and 7-6 transitions toward positions P1, P2, P3, P4 and P5.

individual spectra and the results are tabulated in Table. A.1. A systematic velocity gradient is observed as we move from P1 to P5. For the J = (2 - 1) transition, the line widths are in the range $10.3-30.9 \text{ km s}^{-1}$ with a mean value of 18.6 km s^{-1} . Observed line widths are generally comparable to high velocity components observed in other Galactic clouds ($\Delta V \ge 14 \text{ km s}^{-1}$; Csengeri et al. 2016). To compare the properties of shocked gas emission with those of general dense gas, we examined H¹³CO⁺ emission towards the same five positions (Fig. 9). The H¹³CO⁺ molecule serves as an excellent tracer for dense gas, given its critical density of $\sim 10^5$ cm⁻³ and the optically thin nature of its emission. The (2-1) transition of $H^{13}CO^{+}$ has a rest frequency of 86.75429 GHz, which is covered within the same spectral setup used for the SiO (2-1) observations. Line parameters were extracted by fitting Gaussian components to the individual spectra and the results are tabulated in Table. A.2. Across all positions except P3, SiO line widths are generally broader than H¹³CO⁺ line widths. Specifically, the mean SiO (2-1) line width across all positions is 18.6 km s^{-1} , while the mean $H^{13}CO^{+}$ (1-0) line width is 13.1 km s⁻¹. The broader line width of SiO could be driven by moderate to high-velocity shocks and/or outflows. The exception at position P3 may reflect local variations in shock activity or gas dynamics.

3.6.1. SiO column density and abundance

In order to estimate the SiO column density and abundance in the helix stream, we follow the simplistic assumption that the emission is optically thin and assume local thermodynamic equilibrium (LTE) conditions. To compensate for difference in beam sizes and different beam dilution effects, we applied a correction factor $\theta_{1-0}^2/\theta_{line}^2$ to (2–1), (5–4) and (7–6) intensities, where θ_{1-0} is the beam FWHM of (1–0) transition and θ_{line} is the FWHM corresponding to the transition in consideration. Under these assumptions, the total SiO column density N(SiO) and the measured integrated line intensity *I* in K km s⁻¹ are related as (e.g., Armijos-Abendaño et al. 2015)

Table 3. Details of observed SiO transitions

Transition	Frequency (GHz)	$E_{ m up} \ (m K)$	$ Log_{10}(A_{ij}) \\ (s^{-1}) $	$S \mu^2 $ (D ²)	rms (mK)
1–0	43.4238	2.08	-5.51584	9.63	35
2-1	86.8469	6.25	-4.53354	19.26	20
5–4	217.1049	31.26	-3.28429	48.14	12
7–6	303.9268	58.35	-2.83461	67.38	4

$$\frac{N_{\rm u}}{g_{\rm u}} = \frac{N({\rm SiO})}{Q_{\rm rot}} e^{-E_{\rm u}/kT_{\rm rot}} = \frac{1.67 \times 10^{14} I}{\nu \mu^2 S}$$
 (7)

where $N_{\rm u}$ and $g_{\rm u}$ are the molecular column density and degeneracy of the upper energy level, $Q_{\rm rot}$ is the SiO rotational partition function, $E_{\rm u}/k$ is the energy of the upper energy level (in K), $T_{\rm rot}$ is the rotation temperature in K, ν is the transition frequency in GHz, μ is the dipole moment in Debye and S is the line strength. We can construct a rotation diagram (or "Boltzmann plot" Goldsmith & Langer 1999) by taking the natural logarithm of equation (7),

$$\ln \frac{1.67 \times 10^{14} I}{\nu \mu^2 S} = \ln \frac{N(\text{SiO})}{Q_{rot}} - \frac{E_u}{k T_{rot}}$$
(8)

The SiO rotation diagrams for all five positions are presented in Fig. 10. For positions exhibiting multiple velocity components, we summed all components to estimate rotation temperatures and column densities. We find rotation temperatures in the range 8.5–9.6 K and beam averaged column densities in the range $1.1 - 3.4 \times 10^{13}$ cm⁻² with a mean of 2.1×10^{13} cm⁻². The column densities are listed in Table 4. We would like to caution that the temperature and column density estimates based on rotation diagram method could suffer from optical depth and





Fig. 11. SiO column density (N(SiO)) as a function of molecular hydrogen column density (N(H_2)). Dotted lines correspond to SiO abundances (X(SiO)) with respect to N(H_2). Red triangles correspond to helix stream, blue dots represent data from Csengeri et al. (2016), magenta crosses from Zhu et al. (2020) and green squares correspond to data from Kim et al. (2023).

beam dilution effects. It is worth noting that temperatures derived from rotation diagrams (8–10 K) are significantly lower than the typical kinetic temperatures expected in the CMZ (25–200 K; Krieger et al. 2017). This discrepancy is likely due to SiO lines becoming increasingly sub-thermal with increasing J (and therefore increasing critical density), which is consistent with the presence of shocks.

In order to estimate the abundance of SiO, X(SiO), relative to molecular hydrogen $(N(SiO)/N(H_2))$, we used the H₂ column density map produced by the PPMAP method. The column densities and abundances are listed in Table 4. The SiO abundances range from 4.8×10^{-10} to 1.5×10^{-9} with a mean of 7.7×10^{-10} . Interestingly, we see an enhanced abundance towards P1 (1.5×10^{-9}) that is almost three times that of the mean abundance toward other positions (5.9×10^{-10}) . The integrated intensity ratios of SiO with respect to H¹³CO⁺ also shows a similar trend where the ratio towards P1 is thrice that of mean ratio towards other positions (see Table 4). P1 corresponds to the region where signatures of cloud-cloud collision are observed in the PV diagram and the PPV plot. Hence, the enhanced SiO abundance in this region could indicate SiO enrichment from cloudcloud collision. In Fig. 11, we compare the SiO and H₂ column densities of the helix stream with SiO data from other Galactic clouds. For this, we used the SiO column densities from Csengeri et al. (2016), Zhu et al. (2020) and Kim et al. (2023). We use the corresponding molecular hydrogen column density data from the ATLASGAL catalogue (Urquhart et al. 2018, 2022). The sample consists of objects in different evolutionary stages such as infrared dark clouds, massive starless and star forming clumps, high mass protostellar objects, and ultracompact H II regions. From the figure, it is evident that the mean SiO abundance in the helix stream be higher than that of spiral arm clouds (1.1×10^{-10}) . However, the SiO abundance of the helix stream is similar to that of other GC clouds ($\sim 10^{-9}$, e.g., Martín-Pintado et al. 1997; Armijos-Abendaño et al. 2020).






Fig. 12. RADEX predictions for the SiO (5-4)/(2-1) integrated intensity ratio plotted as a function of the thermal pressure $(n(H_2)T)$ for two different line widths $13~{\rm km\,s^{-1}(top)}$ and $30~{\rm km\,s^{-1}(bottom)}$. The calculations were carried out for SiO column density of $10^{13}~{\rm cm^{-2}}$ and kinetic temperatures of 25, 50, 100 and 250 K. The observed range of line ratios for the narrow components (represented by blue dashed lines) and broad components (represented by red dashed lines) are also plotted.

3.6.2. Excitation conditions using radiative transfer modelling

Given the limitations of the rotation diagram method in estimating the physical conditions of the helix stream, particularly due to the sub-thermal excitation of SiO lines and the presence of shocks, we employ the non-LTE modelling for a more reliable analysis. For this, we use the non-LTE radiative transfer code RADEX (van der Tak et al. 2007) that utilises the large velocity gradient (LVG) approximation. To determine SiO column densities, we compared the intensity ratios of the SiO (2–1) and (5–4) lines, taking advantage of their comparable spatial resolutions (~ 29"). In Fig. 12, we present RADEX non-LTE predictions illustrating the SiO (5-4)/(2-1) intensity ratio as a function of thermal pressure $(n(H_2)T)$ across a range of kinetic temperatures. We kept the SiO column density fixed at 10^{13} cm⁻². Two separate plots are provided: one for $\Delta V = 13$ km/s, representing observed narrow components, and another for $\Delta V = 30 \text{ km/s}$ to account for broader components. From the plots, we find that the (5-4)/(2-1)line ratio as a function of thermal pressure remains nearly constant despite changes in kinetic temperature. The thermal pressures lie in the range $1.8\text{--}7.3 \times 10^6$ K cm⁻³. This pressure estimate from the RADEX analysis is consistent with the external pressure required for pressure-bounded equilibrium in the helix stream, evident from Fig. 6(Bottom), and as discussed in Section 3.2.4. The relatively constant behaviour of the (5--4)/(2--1) line ratio with respect to thermal pressure suggests a decoupling of this ratio from changes in kinetic temperature which is indicative of non-LTE conditions in the helix stream. If we assume kinetic temperature of 50 K, typical for the CMZ, we get corresponding H_2 volume densities in the range $3.6\text{--}14\times10^4$ cm⁻³.

Table 4. SiO column densities and relative abundance

	N(SiO) cm ⁻²	N(H ₂) cm ⁻²	X(SiO)	$\frac{I_{SiO\ (2-1)}}{I_{H^{13}CO^{+}(2-1)}}$
P1	3.4×10^{13}	2.2×10^{22}	1.5×10^{-9}	12.4
P2	1.3×10^{13}	2.7×10^{22}	4.8×10^{-10}	4.6
P3	1.1×10^{13}	2.1×10^{22}	5.2×10^{-10}	3.5
P4	1.7×10^{13}	2.3×10^{22}	7.4×10^{-10}	4.2
P5	3.2×10^{13}	5.1×10^{22}	6.3×10^{-10}	3.8

4. Discussion

4.1. Origin of the expanding CO shell

Circular or shell-like cavities extending to parsec scales, termed as bubbles, are commonly observed within Galactic high mass star forming regions (Weaver et al. 1977) as well as in external galaxies (Barnes et al. 2023; Watkins et al. 2023). These bubbles are believed to be three-dimensional structures formed by radiation, spherical outflows or winds emanating from high mass stars (Churchwell et al. 2006). Such bubbles are often traced by infrared dust emission, atomic (HI) emission or molecular line emission (e.g., Cappa et al. 2003; Churchwell et al. 2007; Beaumont & Williams 2010; Barnes et al. 2023). The velocity data and the morphological distribution of the molecular gas provide insights into the geometry of the shell and the thickness of the parent cloud. Shells can be classified into momentum driven or pressure driven based on the mechanism responsible for the expansion. Momentum driven shells are formed by stellar winds with sufficient strength where expansion is triggered by the transfer of wind mechanical energy into the ambient material (Castor et al. 1975). In regions where weak or negligible stellar winds exist, the energy generated within an H II region could lead to a pressure difference between the ionised and the ambient neutral gas. This triggers a pressure-driven expansion of the H_{II} region. As a consequence, a shock front forms at the boundary between the ionised gas and the ambient neutral gas and the shell traces the boundary of the shock (Garay & Lizano 1999). The expanding shell feature within the helix stream appears as a complete ring in the PV and PPV plots revealing blueshifted and red-shifted emission corresponding to front and back hemispheres of the shell. This implies that the centre of the shell coincides with the centre of the cloud. The azimuthally averaged radial profile of the shell is presented in Fig. 13. We estimated the radius and the thickness of the shell by fitting a Gaussian to the radial profile (e.g., Arce et al. 2011). The peak of the Gaussian corresponds to the radius of the shell whereas the FWHM corresponds to the thickness of the shell. We find the radius of




Fig. 13. Azimuthally averaged radial profile of the CO shell. Dotted line corresponds to the Gaussian fit to the intensity profile.

the shell to be 2.8', equivalent to 6.7 pc, and the shell thickness to be 3.8', that is 8.9 pc. As there is no diffuse, large scale ionised or warm dust emission present within the shell (see Fig. A.3), we assume the source to be momentum driven, powered by stellar winds. Following the approach of Arce et al. (2011), we test this hypothesis by estimating the wind mass loss rate (\hat{m}_w) required to generate the observed shell using the expression

$$\dot{m}_w = \frac{P_{\text{shell}}}{v_w \, \tau_w} \tag{9}$$

where P_{shell} is the total shell momentum, v_w is the wind velocity and τ_w is the wind timescale. The momentum of the shell is given by $P_{\text{shell}} = M_{\text{shell}} V_{\text{exp}}$ where M_{shell} is the mass of the shell and $V_{\rm exp}$ is the expansion velocity. Using the column density map based on PPMAP method, we find the mass of the shell as 1.4×10^5 M_{\odot}. Using the shell expansion velocity of 35 km s⁻¹ from the PV diagram (Fig. 4), we estimate P_{shell} as $4.9 \times 10^6 \text{ M}_{\odot} \text{ km s}^{-1}$. Considering the radio source GPSR5 0.431+0.262 to be the source powering the stellar wind, we assume $v_w = 2000 \text{ km s}^{-1}$ for a typical O or B-type star (Prinja & Massa 1998; Chen et al. 2013). Assuming a wind timescale of 1 Myr, we find the mass loss rate as $2.5 \times 10^{-3} \text{ M}_{\odot} \text{yr}^{-1}$, notably exceeding the expected rate for O–B stars $(10^{-7} - 10^{-6} \text{ M}_{\odot} \text{yr}^{-1};$ Chen et al. 2013) by a factor of $\sim 10^3 - 10^4$. This discrepancy strongly indicates that the observed shell formation is instigated by multiple sources or a highly energetic event. The momentum injected to the ISM by a single supernova explosion is $\sim 10^5 \,\mathrm{M_\odot} \,\mathrm{km} \,\mathrm{s}^{-1}$ (Koo et al. 2020). This suggests that the shell might have been formed by multiple supernovae or a single hypernova, which releases ~ 10 times more kinetic energy than a supernova (Iwamoto et al. 1998). This possibility aligns with similar observations in circumnuclear regions, such as the CO 0.13-0.13 cloud in the GC (Oka et al. 2001a) and the centre of the NGC 253 galaxy (Sakamoto et al. 2006), where hypernova events have been proposed as potential candidates for driving molecular bubble/cavity expansions. We also detect seven X-ray sources from the Chandra catalogue (Muno et al. 2006) within the shell. These sources could potentially be cataclysmic variables, accreting neutron stars or black holes, young isolated pulsars, or Wolf-Rayet/O stars in colliding wind binaries. However, the specific identification of these sources and their association with the observed shell lie beyond the scope of the present study.




Fig. 14. Schematic view of the inner few kiloparsecs of the GC based on Fig. 3 of Henshaw et al. (2023), showing near and far bar lanes, the CMZ, the G1.3 cloud complex, and the Sgr E complex. The relative position of the helix stream is also shown in the figure. Dashed blue and pink lines correspond to overshot gas from the near and far bar lanes, respectively. Dashed arrow indicate the direction of the observer's line of sight.

4.2. Origin of SiO emission

SiO emission could trace high velocity shocks ($v_{\text{shock}} \sim 20$ – 50 km s⁻¹) from protostellar outflows, medium to high velocity irradiated shocks ($v_{\text{shock}} \sim 10 - 20 \text{ km s}^{-1} \text{to } v_{\text{shock}} \sim 50 \text{ km s}^{-1}$) in hot cores, photon dominated regions and supernova remnants, and very low velocity shocks ($v_{\text{shock}} < 10 \text{ km s}^{-1}$) in star forming regions and infrared dark clouds (Hatchell et al. 2001; Schilke et al. 2001; Gusdorf et al. 2008; Nguyen-Lu'o'ng et al. 2013; Csengeri et al. 2016; Cosentino et al. 2022; Kim et al. 2023). Previous studies identified large scale SiO emission in the GC clouds associated with the SgrA region and the CMZ (e.g., Martín-Pintado et al. 1997; Riquelme et al. 2010; Jones et al. 2012). Typical SiO abundances are found to be $X_{SiO} \sim 10^{-9}$. The enhanced SiO emission observed in the SgrA region exhibits a correlation with the 6.4 keV Fe line, as reported by Amo-Baladrón et al. (2009). This correlation is attributed to two potential mechanisms: fluorescence in an X-ray reflection nebula or the impact of low-energy cosmic rays followed by electronic relaxation. SiO mapping studies toward the SgrB2 star forming region by Armijos-Abendaño et al. (2020) reveal shocked gas with a turbulent substructure, consistent with a large-scale cloudcloud collision. The average SiO abundance is $\sim 10^{-9}$ and this along with observed SiO (2-1) line intensities and gas temperature of ~ 30 K agree with models of grain sputtering by C-type shocks.

The consistent detection of SiO emission towards all five positions within the helix stream suggests the existence of a pervasive, large-scale shock distributed along the 200 pc structure. Among the many studies focused on SiO in the GC and the CMZ region, the helix stream holds a unique distinction. Its location ~ 50 pc above the Galactic plane and the CMZ and its high LSR velocities sets it apart as an intriguing and singular feature within the GC. The width of the SiO line is related to the shock velocity and depending on the projection effects and turbulence, the shock velocity could be higher or lower than the line width (Louvet et al. 2016). Considering the SiO (2–1) line, we observe two components at all positions except P2. Specifically, positions

P1, P4, and P5 display components with FWHM>20 km s⁻¹. At P4 and P5, the FWHMs of the broader components range between 31–34 km s⁻¹, three times larger than that of the narrowest component. Narrow components (<20 km s⁻¹) identified toward all five positions have integrated intensities in the range 1.1–3.7 K km s⁻¹ and is consistent with chemical models of low velocity shocks with $n_{\rm H}=10^4~{\rm cm}^{-3}~(\nu_{\rm shock}<20~{\rm km\,s}^{-1};$ Louvet et al. 2016). Presence of broad components toward P1, P4 and P5 indicate the presence of moderate velocity shocks. This variation in shock velocities suggests diverse local conditions within the helix stream, indicating a complex environment with different shock strengths and possibly varied physical properties across different regions of the cloud.

The observed signatures of cloud-cloud collision towards the eastern edge of the cloud, coupled with indications of high supersonic turbulence, offer compelling evidence suggesting the formation of SiO associated with these events. These collisions are known to trigger shocks and compressive processes within the interstellar medium, creating conditions favourable to the formation of SiO. The presence of enhanced SiO abundance observed specifically at position P1 further supports the scenario of cloud-cloud collisions. This enhancement is consistent with the notion that the collision zone may have triggered sputtering of grains, elevating SiO levels. Regarding the broader components observed at positions P4 and P5, multiple factors could contribute to their origin. One plausible explanation could be the presence of heightened turbulence within these regions. Increased turbulence levels, possibly induced by dynamic processes such as collisions or shocks, could result in broader line widths. Alternatively, the broad components at P4 and P5 might also arise from localised star formation activities such as outflows, jets, and winds from young stellar objects as we observe multiple Hi-GAL clumps near P5. Follow up high resolution mapping studies, combined with simulations are crucial in unravelling the intricate interplay between cloud collisions, turbulence, outflows or jets of embedded sources in shaping observed SiO properties within the cloud.

4.3. Formation of the helix stream

In barred spiral galaxies, the central bar significantly shapes the dynamics of the gas reservoir within the galactic nucleus, crucially influencing their formation and evolution by funnelling matter from the spiral arms into the nuclear region (e.g., Combes & Gerin 1985; Hummel et al. 1990; Sellwood & Wilkinson 1993; Oh et al. 2012). Numerical simulations show that the gas that flows in and around the bar depends on periodic orbits and there exists two main classes of orbits: x_1 and x_2 orbits (Contopoulos & Grosbol 1989; Binney et al. 1991; Athanassoula 1992). The x_1 orbits are elongated, with their major axis aligned along the bar direction whereas the x_2 orbits are innermost orbits, relatively less elongated and aligned perpendicular to the bar. The gas in outer regions follows the x_1 orbits, whereas the gas in the CMZ follows the x_2 orbits. The gas plunges from x_1 to x_2 orbits along the bar lanes. According to Henshaw et al. (2023), the high velocity emission from the EMR/parallelogram originates from dense shocked overshooting gas that has been falling along the dust lane and is in the process of transitioning from x_1 to x_2 orbits. The shocked streams of gas also known as dust lanes are often observed in barred galaxies (Stuber et al. 2023) and fuel the star formation, accretion onto the supermassive black hole, and large-scale outflows (Sormani & Barnes 2019). The timeaveraged mass inflow rate along the dust lanes is $\sim 2.7 \text{ M}_{\odot} \text{yr}^{-1}$ out of which, up to 30% accretes onto the CMZ, while the rest




Fig. 15. Integrated intensity map of the ¹³CO emission in the velocity range 100, 200 km s⁻¹(same as Fig. 1(Right)) overlaid with the ¹³CO integrated intensity contours in the velocity range 150, 200 km s⁻¹, highlighting the helix morphology. Dashed green and cyan curves corresponds to two strands of the helix, strand 1 and strand 2.

overshoots and accretes later (Sormani & Barnes 2019; Hatchfield et al. 2021).

In the GC, there exists a population of compact clouds known as extended velocity features (EVFs) that are characterised by extreme velocity dispersions ($\Delta V > 100 \text{ km s}^{-1}$; Sormani et al. 2019). According to Sormani et al. (2019) and Liszt (2006, 2008), these are either (a) material which originates from collisions between the material in the dust lane and the material that has overshot from dust lanes on the opposite side (e.g., Gramze et al. 2023), or (b) material that originates from the collision between the dust lane and the CMZ. Sormani et al. (2019) identified an EVF at $l=1.3^{\circ}$, which is proposed to be of the second category, i.e., collision between the dust lane and the CMZ.

The G1.3 cloud ($l \sim 1.28^{\circ}$, $b \sim 0.07^{\circ}$) is located at the edge of the CMZ and has high SiO abundance (2.6×10^{-9}) ; Riquelme et al. 2018). Busch et al. (2022) investigated the morphology, physical properties as well as the chemical composition of the G1.3 cloud. They found signatures of cloud-cloud collisions, consistent with gas accretion from the near-side dust lane onto the CMZ region. Two prominent velocity components (~100, 180 km s⁻¹) are observed, connected by an emission bridge. The eastern edge of the helix stream is at $l = 1.3^{\circ}$, where we also observe signatures of cloud-cloud collisions in the PV and the PPV plots. It is located 40 pc above the G1.3 cloud. The similarity in velocity components (\sim 127, 177 km s⁻¹) and estimated SiO abundance towards position P1 at the eastern edge (1.9×10^{-9}) suggests that the helix stream is physically connected to the G1.3 cloud. The helix stream likely originates from turbulent shocked gas resulting from the interaction between the near dust lane and the CMZ, overshot to higher latitudes (see Fig. 14), while the other part collided with the CMZ, forming the G1.3 cloud. The helix stream extends beyond the Galactic plane, with the eastern edge indicating the bar lane brushing against the CMZ, forming a velocity bridge, and the rest of the stream continuing along the bar lane trajectory. On the opposite side, the Sgr E region is believed to be at the intersection of the far dust lane and the CMZ. Wallace et al. (2022) identified CO filaments at much smaller scales (\sim 2 pc) in the Sgr E region with aspect ratio \sim 5 : 1, and explained them as gas being stretched as it is rapidly accreted by the gravitational field of the Galactic bar while falling toward the CMZ.

The origin of the double helix morphology of the cloud is unclear. In order to examine the velocity structure of the helix stream in detail, we generated additional PV diagrams along two strands of the helix as depicted in Fig. 15. The resultant PV diagrams along strand 1 and strand 2 are illustrated in Fig. 16. Across both diagrams, we identify two cloud components. The primary component, indicative of the velocity of the helix stream, exhibits velocities exceeding 150 km s⁻¹, while the

secondary component, manifesting as clumpy features, aligns with the bridge feature (Fig. 4), associated with the cloud-cloud collision with the CMZ, with velocities lower than 140 km s⁻¹. The PV diagrams unveil a distinct helical or cork-screw morphology at scales of 6 pc and 14 pc for strand 1 and strand 2, respectively, implying twisting and turning motions within the two strands of the helix stream. Similar twisting motions have been observed in few Galactic filamentary clouds (e.g., González Lobos & Stutz 2019; Álvarez-Gutiérrez et al. 2021). Beyond these smaller scale helical features, the PV diagram towards strand 1 also exhibits a large-scale wave-like morphology with a characteristic wavelength of approximately ~86 pc, signifying the complex kinematics of the helix stream.

In addition to the helix stream, few other clouds within the GC region exhibit a helix or double-helix morphology. The double helix nebula (DHN) is a 25 pc infrared nebula that is ~100 pc above the GC and is interpreted as a torsional Alfvén wave propagating vertically away from the Galactic disc, possibly driven by the rotation of the magnetised circumnuclear disc (Morris et al. 2006). The GC molecular tornado (GCT) is a helical-spur of molecular gas observed in CO at $l = 1.2^{\circ}$, $V_{LSR} = 70 \text{ km s}^{-1}$, extending 170 pc vertically in the plane and is believed to be formed by magnetic squeezing mechanism where a vertical magnetic tube/flux squeezed by the molecular gas in Galactic rotation (Sofue 2007). The pigtail cloud is another helical CO cloud extending ~ 30 pc vertically at $l = -0.7^{\circ}$, $V_{LSR} \sim -70$ to -30 km s⁻¹ and is proposed to be associated with a twisted and coiled magnetic tube resulting from interaction of clouds in the x_1 and x_2 orbits (Matsumura et al. 2012). Unlike these clouds, the helix stream is aligned nearly parallel to the Galactic plane. Interestingly, the GCT cloud is spatially located close to the eastern edge of the helix stream. From Fig. 4(Left), we observe bridging emission features from 70 km s⁻¹, the velocity of GCT to $180\ km\ s^{-1}$, the velocity of the helix stream. Its spatial proximity to the GCT, coupled with bridging emission features observed between their velocities, strongly suggests a physical connection between the two. This connection implies a significant linkage between the helix stream and the magnetically-induced GCT, potentially indicating a common mechanism shaping both clouds. A comparison of the ¹³CO emission from the helix stream with the dust polarisation measured with the Planck 353 GHz data (Planck Collaboration et al. 2020) is presented in Fig. A.4. The observed dust polarisation, which is directly linked to the magnetic field of the cloud, reveals that the polarisation vectors are parallel to the high-intensity ridges within the helix stream. Consequently, the orientation of the magnetic field (rotated 90° with respect to the dust polarisation angle) is perpendicular to the high-intensity emission. However, the observed dust polarisation could potentially be contaminated by line-of-sight dust emission. Therefore, detailed follow-up polarisation studies are necessary to accurately disentangle the true magnetic field structure and its relation to the observed morphology of the helix stream.

An alternate proposition for the origin of the double helix morphology is that the two strands of the double helix represent two interacting high velocity gas streams that are gravitationally wound to each other, while moving along their respective trajectories and revolving around each other. Under this assumption, we can roughly estimate the mass of the streams by applying the Kepler's third law,

$$\frac{P^2}{a^3} = \frac{4\pi^2}{G(M_1 + M_2)}\tag{10}$$




Fig. 16. (Left) PV diagram along the strand 1 of the helix stream shown as green curve in Fig. 15. (Right) PV diagram along the strand 2 of the helix stream shown as cyan curve in Fig. 15.

where P is the orbital period, a is the separation between the two streams of gas and M_1 and M_2 correspond to the masses of the individual streams and G is the gravitational constant. We calculate P by dividing the 2D wavelength of the helix (\sim 30 pc) with the velocity of the streams. The line of sight velocity of the streams is $\sim\!\!170~km\,s^{-1} and$ assuming that the total velocity vectors point in the same direction as the bar major axis (~30° with respect to the line of sight), we find the stream velocity as 80 km s⁻¹. Using these, we estimate P to be 0.36 Myr. Assuming $M_1=M_2$, and a=4.5 pc (half of the separation between the two strand of the helix), we find the total mass of the system as 5.9×10^6 M_{\odot}. We find that the calculated mass using Kepler's third law is comparable to the observed mass estimate for the helix stream ($\sim 2-5 \times 10^6 \ M_{\odot}$). Large-scale oscillatory gas flows with wavelengths ranging from 0.3-400 pc are observed toward molecular clouds in Galactic and extragalactic environments. Henshaw et al. (2020) proposed that they are likely to be formed via gravitational instabilities. Alves et al. (2020) identified a coherent 2.7 kpc sinusoidal wavy chain of gas clouds in the Solar neighbourhood, known as the Radcliffe Wave. Followup studies by Konietzka et al. (2024) show that the Radcliffe Wave is a coherent oscillating structure, with phase velocities ranging from 5–40 km s⁻¹. Gravitational perturbations and feedback mechanisms have been suggested as the possible origins of the Radcliffe wave. Considering that the helix stream could also have its morphology and velocity structure shaped by gravitational instabilities, it is possible that the strands of the double helix can be approximated as two superposed Radcliffe Wavelike structures.

The intricate morphology and kinematics of the helix stream, along with its association with other helical clouds and the application of Kepler's third law, provide compelling evidence for a multifaceted origin involving gravitational interactions, magnetic fields, and possibly other physical processes. Our study revealing active massive star formation within this cloud for the first time not only challenges existing paradigms but also emphasise the significance of clouds with high line of sight velocities in the evolution of the Galaxy. Furthermore, the helix stream stands as an invaluable local template, offering insights into understanding sub-parsec scale mass inflow dynamics within the nuclear regions of barred galaxies.

5. Conclusion

We carried out a comprehensive multiwavelength study of the high velocity helix stream in the GC region, that is associated with the EMR/parallelogram. Using CO data, we performed a kinematic analysis of the cloud, followed by an investigation of the star formation activity using data from infrared to radio wavelengths. We explored the presence of shocked gas in the region using SiO molecular line data. The outcomes of this study can be summarised as follows:

- − The helix stream is a highly elongated (aspect ratio: 9.3) 200 pc high velocity cloud (100–200 km s⁻¹) positioned at a vertical distance of ~15–55 pc above the CMZ. The mass of the cloud is estimated to be $2.5 \times 10^6 \, \mathrm{M}_{\odot}$.
- The ¹²CO PV and ¹³CO PPV plots of the cloud display a bridging feature towards the eastern edge connecting the helix stream to several clumpy features in the velocity range 70–100 km s⁻¹, a signature of cloud-cloud collision.
- The σ R plot of the helix stream is fitted with a power-law index of 0.7, that is steeper than the classical power-law index of 0.5. The obtained index is similar to that of other GC clouds. We find the gas motion within the cloud to be supersonic ($\mathcal{M}_{3D} > 20$).
- The PV and PPV plots also reveal a ring-like feature at $l \sim 0.42^\circ$, associated with an expanding shell of molecular gas. We find the expansion velocity to be 35 km s⁻¹. The shell radius and thickness are 6.7 pc and 8.9 pc, respectively. The large momentum of the expanding shell suggests its formation associated with multiple supernovae or a single hypernova.
- A thermal radio source is identified towards the centre of the expanding CO shell. The infrared colours of the radio source is consistent with that of an H π region, revealing the massive star formation activity within the shell. The ZAMS spectral type of the ionising source is estimated to be B0–B0.5.

- We find 19 Hi-GAL dust clumps within the cloud indicating the ongoing star formation activity within the cloud. There are 8 prestellar clumps and 11 protostellar clumps. Of these clumps, 17 satisfy criteria for massive star formation.
- Detection of SiO emission within the helix stream indicate the presence of a large-scale pervasive shock in the region. Mean SiO abundance is $\sim 10^{-9}$. An enhancement in SiO abundance is observed towards the eastern edge, suggesting additional SiO production from a cloud-cloud collision.
- The PV diagrams along the individual strands of the helix stream reveal twisting and turning motions within the cloud at scales of 6–14 pc. In addition, the PV diagrams also reveal larger scale (~86 pc) wavy patterns.
- The helix stream is likely formed by the dust lane-CMZ interaction and represents the turbulent shocked gas that is overshot to high latitudes after "brushing" the CMZ at the location of G1.3 cloud. This interpretation finds support in the observed kinematic signatures, as well as large Mach numbers and enhanced SiO abundance within the cloud. The distinctive double helix structure of the cloud may arise from the combined influences of turbulence, gravity, and magnetic squeezing.

Acknowledgements. We thank the referee for the valuable comments and suggestions that improved the quality of the paper. We thank the staff of APEX 12m telescope, IRAM 30m telescope and Yebes 40m telescope for their excellent support. This publication is based on data acquired with the Atacama Pathfinder Experiment (APEX) under programmes 092.F-9315, 193.C-0584, 109.A-9518, and 109.C-9518. APEX is a collaboration among the Max-Planck-Institut fur Radioastronomie, the European Southern Observatory, and the Onsala Space Observatory. The processed data products are available from the SEDIGISM survey database located at https://sedigism.mpifr-bonn.mpg.de/index.html, which was constructed by James Urquhart and hosted by the Max Planck Institute for Radio Astronomy. This work is based on observations carried out with the Yebes 40 m telescope (21A017). The 40 m radio telescope at Yebes Observatory is operated by the Spanish Geographic Institute (IGN; Ministerio de Transportes y Movilidad Sostenible). This work is based on observations carried out under project number 057-21 with the IRAM 30m telescope. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). This work is based in part on observations made with the Spitzer Space Telescope, which was operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This publication also made use of data products from Herschel. Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. This work has made use of 353 GHz dust polarisation data, based on observations obtained with Planck (http://www.esa.int/Planck), an ESA science mission with instruments and contributions directly funded by ESA Member States, NASA, and Canada. MCS acknowledges financial support from the European Research Council under the ERC Starting Grant "GalFlow" (grant 101116226). D. Riquelme acknowledges the financial support of DIDULS/ULS, through the project PAAI 2023.

References

Álvarez-Gutiérrez, R. H., Stutz, A. M., Law, C. Y., et al. 2021, ApJ, 908, 86 Alves, J., Zucker, C., Goodman, A. A., et al. 2020, Nature, 578, 237 Amo-Baladrón, M. A., Martín-Pintado, J., Morris, M. R., Muno, M. P., &

Rodríguez-Fernández, N. J. 2009, ApJ, 694, 943

Anderson, L. D., Zavagno, A., Barlow, M. J., García-Lario, P., & Noriega-Crespo, A. 2012, A&A, 537, A1

Arce, H. G., Borkin, M. A., Goodman, A. A., Pineda, J. E., & Beaumont, C. N. 2011, ApJ, 742, 105

Armijos-Abendaño, J., Banda-Barragán, W. E., Martín-Pintado, J., et al. 2020, MNRAS, 499, 4918

Armijos-Abendaño, J., Martín-Pintado, J., Requena-Torres, M. A., Martín, S., & Rodríguez-Franco, A. 2015, MNRAS, 446, 3842

Athanassoula, E. 1992, MNRAS, 259, 345

```
Bally, J., Stark, A. A., Wilson, R. W., & Henkel, C. 1987, ApJS, 65, 13
Barnes, A. T., Longmore, S. N., Avison, A., et al. 2019, MNRAS, 486, 283
Barnes, A. T., Watkins, E. J., Meidt, S. E., et al. 2023, ApJ, 944, L22
Battersby, C., Keto, E., Walker, D., et al. 2020, ApJS, 249, 35
Beaumont, C. N. & Williams, J. P. 2010, ApJ, 709, 791
Becker, R. H., White, R. L., Helfand, D. J., & Zoonematkermani, S. 1994, ApJS,
   91, 347
```

Binney, J., Gerhard, O. E., Stark, A. A., Bally, J., & Uchida, K. I. 1991, MNRAS, 252, 210

Busch, L. A., Riquelme, D., Güsten, R., et al. 2022, A&A, 668, A183 Cappa, C. E., Arnal, E. M., Cichowolski, S., Goss, W. M., & Pineault, S. 2003, in A Massive Star Odyssey: From Main Sequence to Supernova, ed. K. van der Hucht, A. Herrero, & C. Esteban, Vol. 212, 596

Carter, M., Lazareff, B., Maier, D., et al. 2012, A&A, 538, A89

Caselli, P. & Myers, P. C. 1995, ApJ, 446, 665 Castor, J., McCray, R., & Weaver, R. 1975, ApJ, 200, L107 Chen, Y., Zhou, P., & Chu, Y.-H. 2013, ApJ, 769, L16

Churchwell, E., Povich, M. S., Allen, D., et al. 2006, ApJ, 649, 759 Churchwell, E., Watson, D. F., Povich, M. S., et al. 2007, ApJ, 670, 428

Combes, F. & Gerin, M. 1985, A&A, 150, 327

Contopoulos, G. & Grosbol, P. 1989, A&A Rev., 1, 261

Cosentino, G., Jiménez-Serra, I., Tan, J. C., et al. 2022, MNRAS, 511, 953 Csengeri, T., Leurini, S., Wyrowski, F., et al. 2016, A&A, 586, A149

Dahmen, G., Huttemeister, S., Wilson, T. L., & Mauersberger, R. 1998, A&A, 331, 959

Dalcanton, J. J., Yoachim, P., & Bernstein, R. A. 2004, ApJ, 608, 189 Elia, D., Merello, M., Molinari, S., et al. 2021, MNRAS, 504, 2742

Elia, D., Molinari, S., Schisano, E., et al. 2017, MNRAS, 471, 100 Englmaier, P. & Gerhard, O. 1999, MNRAS, 304, 512

Ferrière, K., Gillard, W., & Jean, P. 2007, A&A, 467, 611

Field, G. B., Blackman, E. G., & Keto, E. R. 2011, MNRAS, 416, 710

Garay, G. & Lizano, S. 1999, PASP, 111, 1049

Giannetti, A., Leurini, S., Wyrowski, F., et al. 2017, A&A, 603, A33

Ginsburg, A., Henkel, C., Ao, Y., et al. 2016, A&A, 586, A50

Goldsmith, P. F. & Langer, W. D. 1999, ApJ, 517, 209

González Lobos, V. & Stutz, A. M. 2019, MNRAS, 489, 4771 Gramze, S. R., Ginsburg, A., Meier, D. S., et al. 2023, ApJ, 959, 93

GRAVITY Collaboration, Abuter, R., Amorim, A., et al. 2019, A&A, 625, L10 Gusdorf, A., Pineau Des Forêts, G., Cabrit, S., & Flower, D. R. 2008, A&A, 490,

Güsten, R., Nyman, L. Å., Schilke, P., et al. 2006, A&A, 454, L13

Gutermuth, R. A. & Heyer, M. 2015, AJ, 149, 64 Hatchell, J., Fuller, G. A., & Millar, T. J. 2001, A&A, 372, 281

Hatchfield, H. P., Sormani, M. C., Tress, R. G., et al. 2021, ApJ, 922, 79

Haworth, T. J., Shima, K., Tasker, E. J., et al. 2015, MNRAS, 454, 1634 Henshaw, J. D., Barnes, A. T., Battersby, C., et al. 2023, in Astronomical Society of the Pacific Conference Series, Vol. 534, Protostars and Planets VII, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura, 83

Henshaw, J. D., Kruijssen, J. M. D., Longmore, S. N., et al. 2020, Nature Astronomy, 4, 1064

Henshaw, J. D., Longmore, S. N., Kruijssen, J. M. D., et al. 2016, MNRAS, 457,

Heyer, M., Krawczyk, C., Duval, J., & Jackson, J. M. 2009, ApJ, 699, 1092 Hughes, V. A. & MacLeod, G. C. 1989, AJ, 97, 786

Hummel, E., van der Hulst, J. M., Kennicutt, R. C., & Keel, W. C. 1990, A&A, 236, 333

Iwamoto, K., Mazzali, P. A., Nomoto, K., et al. 1998, Nature, 395, 672 Jones, P. A., Burton, M. G., Cunningham, M. R., et al. 2012, MNRAS, 419, 2961 Jones, P. A., Burton, M. G., Cunningham, M. R., Tothill, N. F. H., & Walsh, A. J. 2013, MNRAS, 433, 221

Kakiuchi, K., Suzuki, T. K., Fukui, Y., et al. 2018, MNRAS, 476, 5629

Kauffmann, J. & Pillai, T. 2010, ApJ, 723, L7

Kauffmann, J., Pillai, T., & Goldsmith, P. F. 2013, ApJ, 779, 185 Kauffmann, J., Pillai, T., Zhang, Q., et al. 2017, A&A, 603, A89

Kim, W. J., Urquhart, J. S., Veena, V. S., et al. 2023, A&A, 679, A123

Konietzka, R., Goodman, A. A., Zucker, C., et al. 2024, Nature, 628, 62 Koo, B.-C., Kim, C.-G., Park, S., & Ostriker, E. C. 2020, ApJ, 905, 35

Krieger, N., Ott, J., Beuther, H., et al. 2017, ApJ, 850, 77

Krumholz, M. R. & McKee, C. F. 2008, Nature, 451, 1082

Larson, R. B. 1981, MNRAS, 194, 809

Lazio, T. J. W. & Cordes, J. M. 2008, ApJS, 174, 481 Le Petit, F., Ruaud, M., Bron, E., et al. 2016, A&A, 585, A105

Lindner, R. R., Vera-Ciro, C., Murray, C. E., et al. 2015, AJ, 149, 138

Liszt, H. S. 2006, A&A, 447, 533

Liszt, H. S. 2008, A&A, 486, 467

Louvet, F., Motte, F., Gusdorf, A., et al. 2016, A&A, 595, A122

Makai, Z., Anderson, L. D., Mascoop, J. L., & Johnstone, B. 2017, ApJ, 846, 64 Marsh, K. A., Whitworth, A. P., Lomax, O., et al. 2017, MNRAS, 471, 2730

Marshall, D. J., Fux, R., Robin, A. C., & Reylé, C. 2008, A&A, 477, L21 Martín-Pintado, J., de Vicente, P., Fuente, A., & Planesas, P. 1997, ApJ, 482, L45

Article number, page 16 of 20

```
Matsumura, S., Oka, T., Tanaka, K., et al. 2012, ApJ, 756, 87
```

Mazumdar, P., Wyrowski, F., Urquhart, J. S., et al. 2021, A&A, 656, A101

Mège, P., Russeil, D., Zavagno, A., et al. 2021, A&A, 646, A74

Mezger, P. G. & Henderson, A. P. 1967, ApJ, 147, 471

Molinari, S., Pezzuto, S., Cesaroni, R., et al. 2008, A&A, 481, 345

Molinari, S., Schisano, E., Elia, D., et al. 2016, A&A, 591, A149

Morris, M. & Serabyn, E. 1996, ARA&A, 34, 645

Morris, M., Uchida, K., & Do, T. 2006, Nature, 440, 308

Muno, M. P., Bauer, F. E., Bandyopadhyay, R. M., & Wang, Q. D. 2006, ApJS, 165, 173

Myers, P. C. 1983, ApJ, 270, 105

Nguyen-Lu'o'ng, Q., Motte, F., Carlhoff, P., et al. 2013, ApJ, 775, 88

Oh, S., Oh, K., & Yi, S. K. 2012, ApJS, 198, 4

Oka, T., Hasegawa, T., Sato, F., Tsuboi, M., & Miyazaki, A. 1998, ApJS, 118, 455

Oka, T., Hasegawa, T., Sato, F., Tsuboi, M., & Miyazaki, A. 2001a, PASJ, 53, 779

Oka, T., Hasegawa, T., Sato, F., et al. 2001b, ApJ, 562, 348

Oort, J. H. 1977, ARA&A, 15, 295

Panagia, N. 1973, AJ, 78, 929

Pety, J. 2005, in SF2A-2005: Semaine de l'Astrophysique Française, ed. F. Casoli, T. Contini, J. M. Hameury, & L. Pagani, 721

Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, A&A, 641, A12

Ponti, G., Morris, M. R., Churazov, E., Heywood, I., & Fender, R. P. 2021, A&A, 646, A66

Prinja, R. K. & Massa, D. L. 1998, in Astronomical Society of the Pacific Conference Series, Vol. 131, Properties of Hot Luminous Stars, ed. I. Howarth, 218

Quireza, C., Rood, R. T., Bania, T. M., Balser, D. S., & Maciel, W. J. 2006, ApJ, 653, 1226

Reid, M. J., Menten, K. M., Brunthaler, A., et al. 2019, ApJ, 885, 131

Riener, M., Kainulainen, J., Henshaw, J. D., et al. 2019, A&A, 628, A78

Riquelme, D., Amo-Baladrón, M. A., Martín-Pintado, J., et al. 2018, A&A, 613, A42

Riquelme, D., Bronfman, L., Mauersberger, R., May, J., & Wilson, T. L. 2010, A&A, 523, A45

Rosolowsky, E. W., Pineda, J. E., Kauffmann, J., & Goodman, A. A. 2008, ApJ, 679, 1338

Sakamoto, K., Ho, P. T. P., Iono, D., et al. 2006, ApJ, 636, 685

Sandage, A. 1961, The Hubble Atlas of Galaxies

Schilke, P., Pineau des Forêts, G., Walmsley, C. M., & Martín-Pintado, J. 2001, A&A, 372, 291

Schilke, P., Walmsley, C. M., Pineau des Forets, G., & Flower, D. R. 1997, A&A, 321, 293

Schmiedeke, A., Schilke, P., Möller, T., et al. 2016, A&A, 588, A143

Schuller, F., Csengeri, T., Urquhart, J. S., et al. 2017, A&A, 601, A124

Scoville, N. Z. 1972, ApJ, 175, L127

Sellwood, J. A. & Wilkinson, A. 1993, Reports on Progress in Physics, 56, 173
Shetty, R., Beaumont, C. N., Burton, M. G., Kelly, B. C., & Klessen, R. S. 2012, MNRAS, 425, 720

Sofue, Y. 2007, PASJ, 59, 189

Sofue, Y. 2017, MNRAS, 470, 1982

Solomon, P. M., Rivolo, A. R., Barrett, J., & Yahil, A. 1987, ApJ, 319, 730

Sorensen, S. A., Matsuda, T., & Fujimoto, M. 1976, Ap&SS, 43, 491

Sormani, M. C. & Barnes, A. T. 2019, MNRAS, 484, 1213

Sormani, M. C., Treß, R. G., Glover, S. C. O., et al. 2019, MNRAS, 488, 4663

Stuber, S. K., Schinnerer, E., Williams, T. G., et al. 2023, A&A, 676, A113

Tercero, F., López-Pérez, J. A., Gallego, J. D., et al. 2021, A&A, 645, A37

Urquhart, J. S., König, C., Giannetti, A., et al. 2018, MNRAS, 473, 1059

Urquhart, J. S., Wells, M. R. A., Pillai, T., et al. 2022, MNRAS, 510, 3389

van der Tak, F. F. S., Black, J. H., Schöier, F. L., Jansen, D. J., & van Dishoeck, E. F. 2007, A&A, 468, 627

Veena, V. S., Riquelme, D., Kim, W. J., et al. 2023, A&A, 674, L15

Walker, D. L., Longmore, S. N., Zhang, Q., et al. 2018, MNRAS, 474, 2373

Wallace, J., Battersby, C., Mills, E. A. C., et al. 2022, ApJ, 939, 58

Ward-Thompson, D., Kirk, J. M., André, P., et al. 2010, A&A, 518, L92 Watkins, E. J., Kreckel, K., Groves, B., et al. 2023, A&A, 676, A67

Weaver, R., McCray, R., Castor, J., Shapiro, P., & Moore, R. 1977, ApJ, 218,

Wood, D. O. S. & Churchwell, E. 1989, ApJ, 340, 265

Young, A., Gillessen, S., de Zeeuw, T., et al. 2023, A&A, 670, A36

Yusef-Zadeh, F., Zhao, J.-H., Arendt, R., et al. 2024, MNRAS, 530, 235

Zhu, F.-Y., Wang, J.-Z., Liu, T., et al. 2020, MNRAS, 499, 6018

Appendix A: Additional figures and tables




Fig. A.1. SEDIGISM 13 CO intensity maps in 10 km s $^{-1}$ velocity intervals from +110 to +200 km/s. The velocity labels in individual panels correspond to central velocities of each interval.




Fig. A.2. (Left) Column density and (Right) dust temperature maps using PPMAP method overplotted with 5σ contours of ^{13}CO integrated emission.




Fig. A.3. Mid-infrared 8 μ m (Left) and 24 μ m (Right) maps of the expanding CO shell overlaid with 13 CO emission contours. Contour levels are 18 and 23 Kkm/s. The thermal radio source GPSR5 0.431+0.262 is marked as a circle.

Table A.1. SiO line parameters

		SiO (1–0)			SiO (2-1)			SiO (5-4)			SiO (7–6)	
	$\int_{C} T_{MB} dv$ (K km s ⁻¹)	V_{LSR} $(km s^{-1})$	ΔV (km s ⁻¹)	$\int T_{MB} dv$ (K km s ⁻¹)	$\frac{V_{LSR}}{(km \ s^{-1})}$	ΔV (km s ⁻¹)	$\int T_{MB} dv$ (K km s ⁻¹)	V_{LSR} (km s^{-1})	ΔV (km s ⁻¹)	$\int T_{MB} dv$ (K km s ⁻¹)	V_{LSR} $(km s^{-1})$	ΔV (km s ⁻¹)
P1	9.6 ± 0.3 5.8 ± 0.2	126.4 ± 0.3 177.3 ± 0.3	23.9 ± 0.8 16.1 ± 0.7	5.5 ± 0.1 3.7 ± 0.1	$126.5 \pm 0.2 \\ 177.0 \pm 0.2$	23.6 ± 0.5 18.3 ± 0.5	0.5 ± 0.06 0.4 ± 0.05	127.8 ± 1.4 176.8 ± 1.1	22.2 ± 2.6 16.1 ± 2.4	0.14 ± 0.01 0.01 ± 0.02	129.8 ± 0.8 181.3 ± 0.8	14.3 ± 1.7 12.9 ± 2.1
P2	P2 6.6 ± 0.2	168.5 ± 0.2	168.5 ± 0.2 15.4 ± 0.6 3.1 ± 0.1	3.1 ± 0.1	168.4 ± 0.2	16.5 ± 0.5	0.3 ± 0.03	168.3 ± 0.7	11.1 ± 2.0	168.4 ± 0.2 16.5 ± 0.5 0.3 ± 0.03 168.3 ± 0.7 11.1 ± 2.0 0.07 ± 0.01 167.4 ± 1.1	167.4 ± 1.1	9.9 ± 2.3
P3	4.8 ± 0.4	P3 4.8 ± 0.4 156.2 ± 1.6 37.6 ± 3.9	37.6 ± 3.9	1.8 ± 0.1 1.1 ± 0.1	149.2 ± 0.3 163.3 ± 0.7	10.3 ± 0.8 11.8 ± 1.4	0.2 ± 0.02	0.2 ± 0.02 150.2 ± 0.5	6.2 ± 1.4	0.04 ± 0.01	0.04 ± 0.01 146.2 ± 1.3 11.0 ± 3.3	11.0 ± 3.3
P4	4.8 ± 0.5 1.8 ± 0.5	133.7 ± 0.9 163.9 ± 2.9	19.9 ± 2.6 21.9 ± 7.1	1.5 ± 0.2 2.8 ± 0.3	$132.1 \pm 0.3 \\ 145.7 \pm 1.4$	10.5 ± 1.0 30.9 ± 2.0	0.2 ± 0.02 0.3 ± 0.1	131.3 ± 0.9 140.2 ± 3.3	7.1 ± 2.1 15.8 ± 4.9	0.06 ± 0.02	138.0 ± 1.9 14.9 ± 6.2	14.9 ± 6.2
P5	5.2 ± 0.7 11.1 ± 0.9	94.9 ± 0.3 114.4 ± 1.2	12.9 ± 0.9 31.5 ± 2.3	3.1 ± 0.4 8.5 ± 0.5	95.2 ± 0.3 116.0 ± 1.0	11.5 ± 0.8 34.2 ± 2.3	0.3 ± 0.04 1.6 ± 0.08	96.1 ± 0.4 112.8 ± 0.8	6.8 ± 0.9 31.8 ± 1.7	0.03 ± 0.01 0.22 ± 0.01	98.8 ± 0.5 107.5 ± 1.9	3.9 ± 1.0 28.5 ± 1.9

Table A.2. Parameters of H¹³CO⁺(2–1) lines

	$\int T_{MB} dv$	V_{LSR}	ΔV
	$(K \text{ km s}^{-1})$	$(km s^{-1})$	$(km s^{-1})$
D.1	0.0 0.1	1010 10	21 = 22
P1	0.8 ± 0.1	121.8 ± 1.3	21.7 ± 3.3
	0.07 ± 0.06	138.9 ± 2.3	5.8 ± 3.5
	0.2 ± 0.05	179.6 ± 1.1	7.9 ± 2.8
P2	0.7 ± 0.1	167.6 ± 0.7	12.8 ± 1.7
Da	06.01	1507 . 00	156.04
P3	0.6 ± 0.1	150.7 ± 0.9	15.6 ± 2.4
	0.2 ± 0.1	169.7 ± 1.5	9.9 ± 3.2
P4	0.6 ± 0.1	131.1 ± 4.1	13.4 + 4.1
P4	0.0 = 0.1		10
	0.5 ± 0.1	149.4 ± 4.1	16.1 ± 4.1
P5	2.0 ± 0.4	95.4 ± 1.3	13.8 ± 2.4
	1.1 ± 0.4	112.5 ± 2.4	13.9 ± 5.4




Fig. A.4. ¹³CO integrated intensity map of the helix stream overlaid with Planck 353 GHz polarisation vectors (white). Note that the orientation of the magnetic field is perpendicular to the polarisation vectors.