

A SET OF q -ORTHOGONAL FUNCTIONS

DANDAN CHEN AND SIYU YIN

ABSTRACT. In 1981 Gasper considered q -analogous functions of $C_k^{(\alpha,\beta)}(e^{i\theta})$ which contain the q -orthogonality relation for the continuous q -ultraspherical polynomials. We resolve the remaining question posed by Gasper regarding the q -orthogonality relation. Furthermore, we provide an evaluation of a new q -beta integral with seven parameters.

1. INTRODUCTION

In his work on the Dirichlet problem for the Heisenberg group, Greiner [8] showed that each L_α -spherical harmonic is a unique linear combination of functions of the form

$$e^{in\psi} \sin^{|n|/2} \theta H_k^{(\alpha,n)}(e^{i\theta})$$

where $k \in \mathbb{N}$ and $n \in \mathbb{Z}$. The function $H_k^{(\alpha,n)}(e^{i\theta})t^k$ is defined by the generating function

$$(1 - te^{-i\theta})^{(n-|n|+\alpha-1)/2} (1 - te^{i\theta})^{-(n+|n|+\alpha+1)/2} = \sum_{k=0}^{\infty} H_k^{(\alpha,n)}(e^{i\theta})t^k.$$

Greiner [8] raised the question of whether the functions $H_k^{(\alpha,n)}(e^{i\theta})t^k$ are orthogonal or bi-orthogonal with respect to some complex valued weight functions. Unfortunately, the theory of polynomials orthogonal on the unit circle cannot be applied to $C_k^{(\alpha,\beta)}(e^{i\theta})$ since these functions are not polynomials in powers of $z = e^{i\theta}$, and the weight function $\omega^{(\alpha,\beta)}(\theta)$ is not real valued when $\alpha \neq \beta$.

Gasper [6] illustrated that the more general class of functions $C_k^{(\alpha,\beta)}(e^{i\theta})$ indeed satisfies the orthogonality relation. He also considered q -(basic) analogs of $C_k^{(\alpha,\beta)}(e^{i\theta})$ which contain the continuous q -ultraspherical polynomials. Gasper provided conditions under which these functions were orthogonal and explored more general functions $C_k^{(\alpha,\beta,\gamma)}(e^{i\theta}; q)$ and $\omega^{(\alpha,\beta,\gamma)}(\theta; q)$, defined as follows:

$$\sum_{k=0}^{\infty} C_k^{(\alpha,\beta,\gamma)}(e^{i\theta}; q)t^k = \frac{(\alpha e^{-i\theta}t; q)_\infty (\beta e^{i\theta}t; q)_\infty}{(e^{-i\theta}t; q)_\infty (\gamma e^{i\theta}t; q)_\infty},$$

2010 *Mathematics Subject Classification.* 05A30, 33D05, 33D15, 33D45, 42C05.

Key words and phrases. q -beta integral, q -orthogonal functions, q -ultraspherical polynomials, Askey-Wilson polynomials.

$$\omega^{(\alpha, \beta, \gamma)}(\cos \theta; q) = \frac{(e^{-2i\theta}; q)_\infty (\gamma e^{2i\theta}; q)_\infty}{(\alpha e^{-2i\theta}; q)_\infty (\beta e^{2i\theta}; q)_\infty}.$$

Here and later, we adopt the standard q -series notation:

$$(a; q)_0 := 1, \quad (a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty := \prod_{k=0}^{\infty} (1 - aq^k)$$

for $|q| < 1$ and $n \in \mathbb{N}$. However, Gasper was unable to derive a similar orthogonal equation as in equation (3.6) from [6] without imposing additional restrictions on the parameters.

The continuous q -ultraspherical polynomials, also known as the Rogers polynomials or the Rogers-Askey-Ismail polynomials, constitute a family of orthogonal polynomials introduced by L. J. Rogers in 1895 [13]. Rogers utilized these polynomials to establish the Rogers-Ramanujan identities and described some essential properties of these polynomials, but he did not prove their orthogonality. The orthogonality of these polynomials was first proved by R. Askey and M. E. H. Ismail in 1980 [2]. More Recently, the first author and Liu [5] provided a new proof of q -orthogonality relation for the continuous q -ultraspherical polynomials and evaluated a new q -beta integral with five parameters.

Motivated by the first author and Liu's method in [5], we aim to address the remaining question posed by Gasper [6] by considering the more general situation of q -orthogonal functions: $C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)$ and $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q)$, which are defined as

$$(1.1) \quad \sum_{n=0}^{\infty} \Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q) \frac{t^n}{(q; q)_n} = \frac{(\alpha xt, \beta yt; q)_\infty}{(\gamma xt, \delta yt; q)_\infty},$$

and

$$(1.2) \quad \sum_{n=0}^{\infty} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) t^n = \frac{(\alpha t e^{i\theta}, \beta t e^{-i\theta}; q)_\infty}{(\gamma t e^{i\theta}, \delta t e^{-i\theta}; q)_\infty},$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{C}$.

Theorem 1.1. *For $\max\{|q|, |\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|\} < 1$, we have*

$$(1.3) \quad \int_0^{2\pi} C_m^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(\alpha, \beta, \gamma, \delta)}(\cos \theta|q) d\theta \\ = \frac{2\pi \left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q\right)_\infty}{\left(q, \frac{\alpha\beta}{\gamma\delta}; q\right)_\infty} \left(\frac{1}{1 - \frac{\alpha q^n}{\gamma}} + \frac{1}{1 - \frac{\beta q^n}{\delta}} \right) \frac{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_n (\gamma\delta)^n}{(q; q)_n} \delta_{m,n},$$

where

$$\omega^{(\alpha, \beta, \gamma, \delta)}(\cos \theta|q) = \frac{\left(\frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q\right)_\infty}{\left(\frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q\right)_\infty}$$

and $\delta_{m,n} = 1$ for $m = n$; $\delta_{m,n} = 0$ for else.

Letting $\gamma = \delta = 1$ in Theorem 1.1, we find that it simplifies directly to the result presented in [5, Theorem 1.3]. Based on Theorem 1.1, we derive the following new q -beta integral formula with seven parameters $\alpha, \beta, \gamma, \delta, s, t$ and q . More formulas about q -beta integral can be seen in [1].

Theorem 1.2. *For $\max\{|q|, |\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|, |\gamma s|, |\gamma t|, |\delta s|, |\delta t|\} < 1$, we have*

$$\begin{aligned} & \int_0^{2\pi} \frac{(\alpha te^{i\theta}, \beta te^{-i\theta}, \alpha se^{i\theta}, \beta se^{-i\theta}, \frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_\infty}{(\gamma te^{i\theta}, \delta te^{-i\theta}, \gamma se^{i\theta}, \delta se^{-i\theta}, \frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q)_\infty} d\theta \\ &= \frac{2\pi (\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q)_\infty}{(q, \frac{\alpha\beta}{\gamma\delta}; q)_\infty} \sum_{n=0}^{\infty} \left(\frac{1}{1 - \frac{\alpha}{\gamma} q^n} + \frac{1}{1 - \frac{\beta}{\delta} q^n} \right) \frac{(\frac{\alpha\beta}{\gamma\delta}; q)_n}{(q; q)_n} (\gamma\delta st)^n. \end{aligned}$$

Furthermore, taking $\alpha = a\gamma$ and $\beta = a\delta$ in Theorem 1.1, we obtain the following q -integral formula by using the orthogonality relation for the continuous q -ultraspherical polynomials and Rogers ${}_6\phi_5$ summation formula.

Theorem 1.3. *For $\max\{|q|, |a|, |b|\} < 1$, we have*

$$\begin{aligned} & \int_0^\pi C_m^{(b\gamma, b\delta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta \\ &= \begin{cases} \frac{(\gamma\delta)^n (1-aq^n) (\frac{b}{a}; q)_{\frac{m-n}{2}} (b; q)_{\frac{m+n}{2}} (a\gamma\delta)^{\frac{m-n}{2}}}{(1-a) h_n(a|q) (q; q)_{\frac{m-n}{2}} (aq; q)_{\frac{m+n}{2}}}, & \text{if } m \equiv n \pmod{2}, \\ 0, & \text{if } m \not\equiv n \pmod{2}, \end{cases} \end{aligned}$$

where

$$h_n(a|q) = \frac{(q, a^2; q)_\infty (q; q)_n (1 - aq^n)}{2\pi (a, aq; q)_\infty (a^2; q)_n (1 - a)}.$$

The paper is organized as follows. In Section 2, we first collect some useful identities on basic hypergeometric series and the asymptotic properties of the q -functions $C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)$. Then in Section 3 we demonstrate the proofs of Theorems 1.1–1.3 by using the orthogonality relation for the continuous q -ultraspherical polynomials and Rogers ${}_6\phi_5$ summation formula, and we also obtain the recurrence relation between $C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)$ and $C_m^{(b\gamma, b\delta, \gamma, \delta)}(e^{i\theta}; q)$ without the details.

2. PRELIMINARIES

In this section, we first collect some useful identities on basic hypergeometric series. The basic hypergeometric series or the q -hypergeometric series ${}_r+1\phi_r$ are defined by [7]

$${}_r+1\phi_r \left[\begin{matrix} a_1, a_2, \dots, a_{r+1} \\ b_1, b_2, \dots, b_r \end{matrix}; q, z \right] = \sum_{k=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_k z^k}{(q, b_1, \dots, b_r; q)_k}.$$

We will also need the ${}_{r+1}W_r(a_1; a_2, \dots, a_{r-1}; q, z)$ very-well-poised series [7]:

$${}_{r+1}W_r(a_1; a_2, \dots, a_{r-1}; q, z) = {}_{r+1}\phi_r \left[\begin{matrix} a_1, q\sqrt{a_1}, -q\sqrt{a_1}, a_2, \dots, a_{r-1} \\ \sqrt{a_1}, -\sqrt{a_1}, qa_1/a_2, \dots, qa_1/a_{r-1} \end{matrix}; q, z \right].$$

The q -binomial theorem [7, p. 92] is defined as

$$(2.1) \quad \frac{(az; q)_\infty}{(z; q)_\infty} = \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} z^n, \quad |z| < 1.$$

The q -binomial coefficient is denoted by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}.$$

The continuous q -ultraspherical polynomials are defined as [2]:

$$(2.2) \quad \frac{(\beta r e^{i\theta}, \beta r e^{-i\theta}; q)_\infty}{(r e^{i\theta}, r e^{-i\theta}; q)_\infty} = \sum_{n=0}^{\infty} C_n(\cos \theta; \beta | q) r^n.$$

By (2.2), we can easily find that $C_0(\cos \theta; \beta | q) = 1$, $C_1(\cos \theta; \beta | q) = 2 \cos \theta (1 - \beta) / (1 - q)$ and the recurrence relation

$$(1 - q^{n+1}) C_{n+1}(\cos \theta; \beta | q) = 2 \cos \theta (1 - \beta q^n) C_n(\cos \theta; \beta | q) - (1 - \beta^2 q^{n-1}) C_{n-1}(\cos \theta; \beta | q).$$

From (2.2) and the q -binomial theorem (2.1), it follows that

$$C_n(\cos \theta; \beta | q) = \sum_{k=0}^n \frac{(\beta; q)_k (\beta; q)_{n-k}}{(q; q)_k (q; q)_{n-k}} e^{i(n-2k)\theta} = \sum_{k=0}^n \frac{(\beta; q)_k (\beta; q)_{n-k}}{(q; q)_k (q; q)_{n-k}} \cos(n-2k)\theta.$$

F. Jackson defined the q -integral of the function $f(x)$ from a to b by [7, p. 23] [10]

$$(2.3) \quad \int_a^b f(x) d_q x = (1-q)b \sum_{n=0}^{\infty} q^n f(bq^n) - (1-q)a \sum_{n=0}^{\infty} q^n f(aq^n).$$

The generating function of $\Phi_n^{(\alpha, \beta)}$ [12, Proposition 3.2] is that

$$(2.4) \quad \sum_{n=0}^{\infty} \Phi_n^{(\alpha, \beta)}(x, y | q) \frac{t^n}{(q; q)_n} = \frac{(\alpha x t, \beta y t; q)_\infty}{(x t, y t; q)_\infty}.$$

Proposition 2.1. *For $\max\{|\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|, |q|\} < 1$, the q -function $C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)$ has some asymptotic properties:*

- (1) $|C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)| \leq C_n^{(\alpha, \beta, \gamma, \delta)}(1; q)$,
- (2) $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}, e^{-i\theta} | q) = (q; q)_n C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)$,
- (3) $\limsup_{n \rightarrow \infty} C_n^{(\alpha, \beta, \gamma, \delta)}(1; q)^{1/n} = \max\{|\gamma|, |\delta|\}$.

Proof. (1) We observe that $(q; q)_n > 0$ when $|q| < 1$, where $n \in \mathbb{N}$ and $|e^{i(n-2k)\theta}| = 1$. Given that $|\frac{\alpha}{\gamma}| < 1$ and $|\frac{\beta}{\delta}| < 1$, we have $|C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)| \leq C_n^{(\alpha, \beta, \gamma, \delta)}(1; q)$.

(2) By comparing the coefficients of the powers of t in equation (1.1) and equation (1.2), we complete the proof of (2).

(3) From (1.2), we get

$$\frac{(\alpha t, \beta t; q)_\infty}{(\gamma t, \delta t; q)_\infty} = \sum_{n=0}^{\infty} C_n^{(\alpha, \beta, \gamma, \delta)}(1; q) t^n.$$

According to the definition of the radius of convergence for power series centered at the origin, we determine that the radius of convergence R of the above series is given by $R = \min\{\frac{1}{|\gamma|}, \frac{1}{|\delta|}\}$, thus we finish the proof of (3). \square

Proposition 2.2. *Let $\max\{|q|, |\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|\} < 1$. Then for any nonnegative integer k , and any t such that $|t| < \min\{\frac{1}{|\gamma|}, \frac{1}{|\delta|}\}$, the series*

$$\sum_{n=0}^{\infty} C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(q; q)_{n+k}}{(\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}} t^n$$

converges uniformly and absolutely on $[0, 2\pi]$.

Proof. Given the condition $\max\{|q|, |\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|\} < 1$, we apply the triangular inequality to establish the following chain of inequalities:

$$\left| \frac{(q; q)_{n+k}}{(\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}} \right| \leq \frac{(-|q|; |q|)_\infty}{(|\frac{\alpha\beta}{\gamma\delta}|; |q|)_{n+k}} \leq \frac{(-|q|; |q|)_\infty}{(|\frac{\alpha\beta}{\gamma\delta}|; |q|)_\infty}.$$

Utilizing Proposition 2.1, we derive the inequality

$$\begin{aligned} & \left| \sum_{n=0}^{\infty} C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(q; q)_{n+k}}{(\alpha\beta; q)_{n+k}} t^n \right| \\ & \leq \frac{(-|q|; |q|)_\infty}{(|\frac{\alpha\beta}{\gamma\delta}|; |q|)_\infty} \sum_{n=0}^{\infty} C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(1; q) C_n^{(\alpha, \beta, \gamma, \delta)}(1; q) |t|^n. \end{aligned}$$

By applying the root test, we determine that the right-hand side of the inequality converges when $|t| < \min\{\frac{1}{|\gamma|}, \frac{1}{|\delta|}\}$. Thus, the series in Proposition 2.2 converges uniformly and absolutely on the interval $[0, 2\pi]$. \square

Proposition 2.3. [12, Proposition 5.1] *If there are no zero factors in the denominator of the integral and then we have*

$$\Phi_n^{(a,b)}(x, y|q) = \frac{(ab; q)_n (a, b, by/x, ax/y; q)_\infty}{(1-q)y(q, ab, x/y, qy/x; q)_\infty} \int_x^y \frac{(qz/x, qz/y; q)_\infty z^n}{(bz/x, az/y; q)_\infty} d_q z.$$

Comparing (1.1) with (2.4), we obtain that

$$(2.5) \quad \Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q) = \Phi_n^{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}\right)}(\gamma x, \delta y|q).$$

Utilizing the relationship between $\Phi_n^{(\alpha, \beta)}(x, y|q)$ and $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q)$ presented in (2.5), we obtain the following q -integral representation for $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q)$.

Proposition 2.4. *If there are no zero factors in the denominator of the integral, then we have*

$$(2.6) \quad \Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q) = \frac{(\frac{\alpha\beta}{\gamma\delta}; q)_n (\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\beta y}{\gamma x}, \frac{\alpha x}{\delta y}; q)_\infty}{(1-q)\delta y (q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma x}{\delta y}, \frac{q\delta y}{\gamma x}; q)_\infty} \int_{\gamma x}^{\delta y} \frac{(\frac{qz}{\gamma x}, \frac{qz}{\delta y}; q)_\infty z^n}{(\frac{\beta z}{\gamma\delta x}, \frac{\alpha z}{\gamma\delta y}; q)_\infty} d_q z.$$

3. PROOFS OF THEOREM 1.1 – THEOREM 1.3

Firstly, we shall give the proof of Theorem 1.1 by q -beta integral and the relation between $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(x, y|q)$ and $C_k^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)$.

Proof of Theorem 1.1. Assuming $k \in \mathbb{N}$, we begin by dividing both sides of (2.6) by $(\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}$ and replacing n by $n+k$, we have

$$\frac{\Phi_{n+k}^{(\alpha, \beta, \gamma, \delta)}(x, y|q)}{(\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}} = \frac{(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\beta y}{\gamma x}, \frac{\alpha x}{\delta y}; q)_\infty}{(1-q)\delta y (q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma x}{\delta y}, \frac{q\delta y}{\gamma x}; q)_\infty} \int_{\gamma x}^{\delta y} \frac{(\frac{qz}{\gamma x}, \frac{qz}{\delta y}; q)_\infty z^{n+k}}{(\frac{\beta z}{\gamma\delta x}, \frac{\alpha z}{\gamma\delta y}; q)_\infty} d_q z.$$

Next, we multiply both sides by $\Phi_n^{(a, b, c, d)}(u, v|q)t^n/(q; q)_n$ and sum the resulting equation over n from 0 to ∞ , leading to the following result:

$$\begin{aligned} & \sum_{n=0}^{\infty} \Phi_{n+k}^{(\alpha, \beta, \gamma, \delta)}(x, y|q) \Phi_n^{(a, b, c, d)}(u, v|q) \frac{t^n}{(q; q)_n (\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}} \\ &= \sum_{n=0}^{\infty} \Phi_n^{(a, b, c, d)}(u, v|q) \frac{t^n}{(q; q)_n} \frac{(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\beta y}{\gamma x}, \frac{\alpha x}{\delta y}; q)_\infty}{(1-q)\delta y (q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma x}{\delta y}, \frac{q\delta y}{\gamma x}; q)_\infty} \int_{\gamma x}^{\delta y} \frac{(\frac{qz}{\gamma x}, \frac{qz}{\delta y}; q)_\infty z^{n+k}}{(\frac{\beta z}{\gamma\delta x}, \frac{\alpha z}{\gamma\delta y}; q)_\infty} d_q z \\ &= \frac{(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\beta y}{\gamma x}, \frac{\alpha x}{\delta y}; q)_\infty}{(1-q)\delta y (q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma x}{\delta y}, \frac{q\delta y}{\gamma x}; q)_\infty} \int_{\gamma x}^{\delta y} \frac{(\frac{qz}{\gamma x}, \frac{qz}{\delta y}; q)_\infty (autz, bvtz; q)_\infty z^k}{(\frac{\beta z}{\gamma\delta x}, \frac{\alpha z}{\gamma\delta y}; q)_\infty (cutz, dtz; q)_\infty} d_q z. \end{aligned}$$

Setting $x = u = e^{i\theta}$, $y = v = e^{-i\theta}$, $a = \alpha$, $b = \beta$, $c = \gamma$, $d = \delta$ in the aforementioned equation, we obtain that, for $\max\{|q|, |\frac{\alpha}{\gamma}|, |\frac{\beta}{\delta}|\} < 1$ and $|t| < \min\{\frac{1}{|\gamma|}, \frac{1}{|\delta|}\}$,

$$\begin{aligned} & \frac{(q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_\infty}{(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q)_\infty} \sum_{n=0}^{\infty} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{-i\theta}; q) \frac{(q; q)_{n+k} t^n}{(\frac{\alpha\beta}{\gamma\delta}; q)_{n+k}} \\ &= \frac{1 - \frac{\delta}{\gamma} e^{-2i\theta}}{(1-q)\delta e^{-i\theta}} \int_{\gamma e^{i\theta}}^{\delta e^{-i\theta}} \frac{(\frac{qz}{\gamma} e^{-i\theta}, \frac{qz}{\delta} e^{i\theta}, \alpha t z e^{i\theta}, \beta t z e^{-i\theta}; q)_\infty z^k}{(\frac{\beta z}{\gamma\delta} e^{-i\theta}, \frac{\alpha z}{\gamma\delta} e^{i\theta}, \gamma t z e^{i\theta}, \delta t z e^{-i\theta}; q)_\infty} d_q z \end{aligned}$$

$$\begin{aligned}
&= \left(1 - \frac{\delta}{\gamma} e^{-2i\theta}\right) \delta^k e^{-ik\theta} \sum_{n=0}^{\infty} \frac{(q^{n+1}, \frac{\delta}{\gamma} e^{-2i\theta} q^{n+1}, \alpha \delta t q^n, \beta \delta t e^{-2i\theta} q^n; q)_{\infty}}{\left(\frac{\beta}{\gamma} e^{-2i\theta} q^n, \frac{\alpha}{\gamma} q^n, \gamma \delta t q^n, \delta^2 t e^{-2i\theta} q^n; q\right)_{\infty}} q^{n(k+1)} \\
&\quad + \left(1 - \frac{\gamma}{\delta} e^{2i\theta}\right) \gamma^k e^{ik\theta} \sum_{n=0}^{\infty} \frac{(q^{n+1}, \frac{\gamma}{\delta} e^{2i\theta} q^{n+1}, \beta \gamma t q^n, \alpha \gamma t e^{2i\theta} q^n; q)_{\infty}}{\left(\frac{\alpha}{\delta} e^{2i\theta} q^n, \frac{\beta}{\delta} q^n, \gamma \delta t q^n, \gamma^2 t e^{2i\theta} q^n; q\right)_{\infty}} q^{n(k+1)}.
\end{aligned}$$

Let us consider the right-hand side of the above equation. We define that

(3.1)

$$\left(1 - \frac{\delta}{\gamma} e^{-2i\theta}\right) \delta^k e^{-ik\theta} \sum_{n=0}^{\infty} \frac{(q^{n+1}, \frac{\delta}{\gamma} e^{-2i\theta} q^{n+1}, \alpha \delta t q^n, \beta \delta t e^{-2i\theta} q^n; q)_{\infty}}{\left(\frac{\beta}{\gamma} e^{-2i\theta} q^n, \frac{\alpha}{\gamma} q^n, \gamma \delta t q^n, \delta^2 t e^{-2i\theta} q^n; q\right)_{\infty}} q^{n(k+1)} := \sum_{n=k}^{\infty} \lambda_n e^{-in\theta},$$

(3.2)

$$\left(1 - \frac{\gamma}{\delta} e^{2i\theta}\right) \gamma^k e^{ik\theta} \sum_{n=0}^{\infty} \frac{(q^{n+1}, \frac{\gamma}{\delta} e^{2i\theta} q^{n+1}, \beta \gamma t q^n, \alpha \gamma t e^{2i\theta} q^n; q)_{\infty}}{\left(\frac{\alpha}{\delta} e^{2i\theta} q^n, \frac{\beta}{\delta} q^n, \gamma \delta t q^n, \gamma^2 t e^{2i\theta} q^n; q\right)_{\infty}} q^{n(k+1)} := \sum_{n=k}^{\infty} \mu_n e^{in\theta},$$

where $\{\lambda_n\}_{n=k}^{\infty}$ and $\{\mu_n\}_{n=k}^{\infty}$ are independent of θ . Then we arrive at

$$\begin{aligned}
&\frac{(q, \frac{\alpha\beta}{\gamma\delta}, \frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_{\infty}}{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}, \frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q\right)_{\infty}} \sum_{n=0}^{\infty} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(q; q)_{n+k} t^n}{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_{n+k}} \\
&= \sum_{n=k}^{\infty} \lambda_n e^{-in\theta} + \sum_{n=k}^{\infty} \mu_n e^{in\theta}.
\end{aligned}$$

By Proposition 2.2, we know that the above series on the left-hand side converges uniformly and absolutely. From the well-known fact in complex function theory, $\int_0^{2\pi} e^{in\theta} d\theta = 2\pi \delta_{n,0}$, we have, for $k > 0$,

$$(3.3) \quad \sum_{n=0}^{\infty} \left\{ \int_0^{2\pi} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(\frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_{\infty}}{\left(\frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q\right)_{\infty}} d\theta \right\} \frac{(q; q)_{n+k} t^n}{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_{n+k}} = 0.$$

Comparing the coefficients of powers of t on the both sides of (3.3), we obtain the orthogonality that, for $k > 0$,

$$(3.4) \quad \int_0^{2\pi} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_{n+k}^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(\frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_{\infty}}{\left(\frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q\right)_{\infty}} d\theta = 0.$$

On the other hand, for $k = 0$,

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left\{ \int_0^{2\pi} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{(\frac{\gamma}{\delta} e^{2i\theta}, \frac{\delta}{\gamma} e^{-2i\theta}; q)_{\infty}}{\left(\frac{\alpha}{\delta} e^{2i\theta}, \frac{\beta}{\gamma} e^{-2i\theta}; q\right)_{\infty}} d\theta \right\} \frac{(q; q)_n t^n}{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_n} \\
&= 2\pi \frac{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q\right)_{\infty}}{\left(q, \frac{\alpha\beta}{\gamma\delta}; q\right)_{\infty}} (\lambda_0 + \mu_0).
\end{aligned}$$

Let us begin by considering λ_0 and μ_0 . Inspecting the series in (3.1) and (3.2), we obtain the constant term of the Fourier expansion of this series :

$$\lambda_0 = \sum_{n=0}^{\infty} \frac{(q^{n+1}, \alpha\delta tq^n; q)_{\infty}}{\left(\frac{\alpha}{\gamma}q^n, \gamma\delta tq^n; q\right)_{\infty}} q^n,$$

$$\mu_0 = \sum_{n=0}^{\infty} \frac{(q^{n+1}, \beta\gamma tq^n; q)_{\infty}}{\left(\frac{\beta}{\delta}q^n, \gamma\delta tq^n; q\right)_{\infty}} q^n.$$

Applying the q -binomial theorem (2.1) to the right-hand side of the above equation, we simplify λ_0 and μ_0 to:

$$\lambda_0 = \sum_{n=0}^{\infty} \frac{(\gamma\delta t)^n}{1 - \frac{\alpha}{\gamma}q^n} \quad \text{and} \quad \mu_0 = \sum_{n=0}^{\infty} \frac{(\gamma\delta t)^n}{1 - \frac{\beta}{\delta}q^n}.$$

Thus we have

$$\sum_{n=0}^{\infty} \left\{ \int_0^{2\pi} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q)^2 \frac{\left(\frac{\gamma}{\delta}e^{2i\theta}, \frac{\delta}{\gamma}e^{-2i\theta}; q\right)_{\infty}}{\left(\frac{\alpha}{\delta}e^{2i\theta}, \frac{\beta}{\gamma}e^{-2i\theta}; q\right)_{\infty}} d\theta \right\} \frac{(q; q)_n t^n}{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_n}$$

$$= 2\pi \frac{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q\right)_{\infty}}{\left(q, \frac{\alpha\beta}{\gamma\delta}; q\right)_{\infty}} \sum_{n=0}^{\infty} \left(\frac{1}{1 - \frac{\alpha}{\gamma}q^n} + \frac{1}{1 - \frac{\beta}{\delta}q^n} \right) (\gamma\delta t)^n.$$

Equating the coefficients of powers of t on both sides of the above equation, we deduce that

$$\int_0^{2\pi} \left\{ C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \right\}^2 \frac{\left(\frac{\gamma}{\delta}e^{2i\theta}, \frac{\delta}{\gamma}e^{-2i\theta}; q\right)_{\infty}}{\left(\frac{\alpha}{\delta}e^{2i\theta}, \frac{\beta}{\gamma}e^{-2i\theta}; q\right)_{\infty}} d\theta$$

$$= 2\pi \frac{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q\right)_{\infty} \left(\frac{\alpha\beta}{\gamma\delta}; q\right)_n}{\left(q, \frac{\alpha\beta}{\gamma\delta}; q\right)_{\infty} (q; q)_n} \left(\frac{1}{1 - \frac{\alpha}{\gamma}q^n} + \frac{1}{1 - \frac{\beta}{\delta}q^n} \right) (\gamma\delta)^2.$$

Combining the above equation with (3.4), we complete the proof of Theorem 1.1. \square

Next, we demonstrate the proof of theorem 1.2 by using Theorem 1.1.

Proof of Theorem 1.2. Multiplying both sides of (1.3) by $s^m t^n$, we have

$$\int_0^{2\pi} C_m^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) \frac{\left(\frac{\gamma}{\delta}e^{2i\theta}, \frac{\delta}{\gamma}e^{-2i\theta}; q\right)_{\infty}}{\left(\frac{\alpha}{\delta}e^{2i\theta}, \frac{\beta}{\gamma}e^{-2i\theta}; q\right)_{\infty}} s^m t^n d\theta$$

$$= 2\pi \frac{\left(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q\right)_{\infty}}{\left(q, \frac{\alpha\beta}{\gamma\delta}; q\right)_{\infty}} \left(\frac{1}{1 - \frac{\alpha}{\gamma}q^n} + \frac{1}{1 - \frac{\beta}{\delta}q^n} \right) \frac{\left(\frac{\alpha\beta}{\gamma\delta}; q\right)_n (\gamma\delta)^n}{(q; q)_n} s^m t^n \delta_{m,n}.$$

Summing the above equation over m and n from 0 to ∞ respectively, we deduce that

$$\int_0^{2\pi} \left(\sum_{m=0}^{\infty} C_m^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) s^m \right) \left(\sum_{n=0}^{\infty} C_n^{(\alpha, \beta, \gamma, \delta)}(e^{i\theta}; q) t^n \right) \frac{\left(\frac{\gamma}{\delta}e^{2i\theta}, \frac{\delta}{\gamma}e^{-2i\theta}; q\right)_{\infty}}{\left(\frac{\alpha}{\delta}e^{2i\theta}, \frac{\beta}{\gamma}e^{-2i\theta}; q\right)_{\infty}} d\theta$$

$$= 2\pi \frac{(\frac{\alpha}{\gamma}, \frac{\beta}{\delta}; q)_{\infty}}{(q, \frac{\alpha\beta}{\gamma\delta}; q)_{\infty}} \sum_{n=0}^{\infty} \left(\frac{1}{1 - \frac{\alpha}{\gamma}q^n} + \frac{1}{1 - \frac{\beta}{\delta}q^n} \right) \frac{(\frac{\alpha\beta}{\gamma\delta}; q)_n}{(q; q)_n} (\gamma\delta st)^n.$$

Applying (1.2) to the left-hand side of the above equation, we complete the proof of Theorem 1.2. \square

Now, we shall provide the proof of Theorem 1.3 concerning the orthogonal relation of the q -functions $C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)$.

Proof of Theorem 1.3. Let $\alpha = a\gamma$ and $\beta = a\delta$ in (1.2). Then, the generating function of $C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)$ is given by

$$(3.5) \quad \sum_{n=0}^{\infty} C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) t^n = \frac{(a\gamma t e^{i\theta}, a\delta t e^{-i\theta}; q)_{\infty}}{(\gamma t e^{i\theta}, \delta t e^{-i\theta}; q)_{\infty}}.$$

Furthermore, the orthogonal relation holds for $\max\{|q|, |a|\} < 1$ as follows:

$$\begin{aligned} & \int_0^{2\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta \\ &= \frac{4\pi (a, a; q)_{\infty}}{(q, a^2; q)_{\infty}} \frac{1}{1 - aq^n} \frac{(a^2; q)_n (\gamma\delta)^n}{(q; q)_n} \delta_{m,n} \end{aligned}$$

by letting $\alpha = a\gamma$ and $\beta = a\delta$ in Theorem 1.1. After some simplify calculations, the left-hand side of the above formula becomes

$$\begin{aligned} & \int_0^{2\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta \\ &= 2 \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta. \end{aligned}$$

Recall [7, p. 186], for $|q| < 1, |\beta| < 1$,

$$\int_0^{\pi} C_m(\cos \theta; \beta | q) C_n(\cos \theta; \beta | q) W_{\beta}(\cos \theta | q) d\theta = \frac{\delta_{m,n}}{h_n(\beta | q)},$$

where

$$h_n(\beta | q) = \frac{(q, \beta^2; q)_{\infty} (q; q)_n (1 - \beta q^n)}{2\pi (\beta, \beta q; q)_{\infty} (\beta^2; q)_n (1 - \beta)}.$$

More details can be found in [3, 4, 9, 11]. Using the definition of $h_n(\beta | q)$ above, we can get the orthogonal relation, for $\max\{|q|, |a|\} < 1$,

$$(3.6) \quad \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta = \frac{(\gamma\delta)^m \delta_{m,n}}{h_n(a | q)},$$

where

$$h_n(a | q) = \frac{(q, a^2; q)_{\infty} (q; q)_n (1 - aq^n)}{2\pi (a, aq; q)_{\infty} (a^2; q)_n (1 - a)}.$$

By (3.5), we have the following summation of $C_n^{(\alpha, \alpha, \gamma, \gamma)}(e^{i\theta}; q)(1 - aq^n)t^n$ over n from 0 to ∞ :

$$(3.7) \quad \sum_{n=0}^{\infty} C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)(1 - aq^n)t^n = \frac{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}; q)_{\infty}}{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_{\infty}}(1 - a)(1 - a\gamma\delta t^2).$$

Combining (3.6) with (3.7), we obtain that

$$\begin{aligned} & \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)\omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta|q) \frac{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}; q)_{\infty}}{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_{\infty}}(1 - a\gamma\delta t^2)d\theta \\ &= \frac{2\pi(\gamma\delta)^m(a, aq; q)_{\infty}(a^2; q)_m t^m}{(q, a^2; q)_{\infty}(q; q)_m}. \end{aligned}$$

Then let $t \rightarrow tq^k$ in the equation above, we have

$$(3.8) \quad \begin{aligned} & \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)\omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta|q) \frac{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}; q)_{\infty}}{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_{\infty}} \frac{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_k}{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}; q)_k} (1 - a\gamma\delta t^2 q^{2k})d\theta \\ &= \frac{2\pi(\gamma\delta)^m(a, aq; q)_{\infty}(a^2; q)_m (tq^k)^m}{(q, a^2; q)_{\infty}(q; q)_m}. \end{aligned}$$

Multiplying both sides of (3.8) by

$$\frac{(a\gamma\delta t^2, \frac{aq}{b}; q)_k}{(1 - a\gamma\delta t^2)(q, b\gamma\delta t^2; q)_k} b^k$$

and then summing the resulting equation over k from 0 to ∞ , we get that

$$\begin{aligned} & \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)\omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta|q) {}_6W_5(a\gamma\delta t^2; \gamma te^{i\theta}, \delta te^{-i\theta}, \frac{aq}{b}; q, b) \frac{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}; q)_{\infty}}{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_{\infty}} d\theta \\ &= \frac{2\pi(\gamma\delta)^m(a, aq; q)_{\infty}(a^2; q)_m t^m}{(q, a^2; q)_{\infty}(q; q)_m(1 - a\gamma\delta t^2)} \sum_{k=0}^{\infty} \frac{(a\gamma\delta t^2, \frac{aq}{b}; q)_k}{(q, b\gamma\delta t^2; q)_k} (bq^m)^k. \end{aligned}$$

From the Rogers ${}_6\phi_5$ summation formula [7, Eq. (2.7.1)], for $|aq/bcd| < 1$,

$$(3.9) \quad {}_6W_5(a; b, c, d; q, \frac{aq}{bcd}) = \frac{(aq, aq/bc, aq/bd; q)_{\infty}}{(aq/b, aq/c, aq/d, aq/bcd; q)_{\infty}},$$

we arrive at

$${}_6W_5(a\gamma\delta t^2; \gamma te^{i\theta}, \delta te^{-i\theta}, \frac{aq}{b}; q, b) = \frac{(a\gamma\delta t^2 q, aq, b\gamma te^{i\theta}, b\delta te^{-i\theta}; q)_{\infty}}{(a\gamma tqe^{i\theta}, a\delta tqe^{-i\theta}, b\gamma\delta t^2, b; q)_{\infty}}.$$

Then we can obtain that

$$(3.10) \quad \int_0^{\pi} C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)\omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta|q) \frac{(b\gamma te^{i\theta}, b\delta te^{-i\theta}; q)_{\infty}}{(\gamma te^{i\theta}, \delta te^{-i\theta}; q)_{\infty}} d\theta$$

$$= \frac{2\pi(\gamma\delta)^m t^m (a, b; q)_\infty (a^2; q)_m}{(q, a^2; q)_\infty (q; q)_m} \sum_{k=0}^{\infty} \frac{(\frac{aq}{b}; q)_k (b\gamma\delta t^2 q^k; q)_\infty}{(q; q)_k (a\gamma\delta t^2 q^k; q)_\infty} (bq^m)^k.$$

Considering the summation on the right-hand side of the formula above, by q -binomial theorem (2.1), we have

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(\frac{aq}{b}; q)_k (b\gamma\delta t^2 q^k; q)_\infty}{(q; q)_k (a\gamma\delta t^2 q^k; q)_\infty} (bq^m)^k &= \sum_{k=0}^{\infty} \frac{(\frac{aq}{b}; q)_k}{(q; q)_k} (bq^m)^k \sum_{s=0}^{\infty} \frac{(\frac{b}{a}; q)_s}{(q; q)_s} (a\gamma\delta t^2 q^k)^s \\ &= \sum_{s=0}^{\infty} \frac{(\frac{b}{a}; q)_s}{(q; q)_s} (a\gamma\delta t^2)^s \sum_{k=0}^{\infty} \frac{(\frac{aq}{b}; q)_k}{(q; q)_k} (bq^{m+s})^k \\ &= \sum_{s=0}^{\infty} \frac{(\frac{b}{a}; q)_s}{(q; q)_s} (a\gamma\delta t^2)^s \frac{(aq^{m+s+1}; q)_\infty}{(bq^{m+s}; q)_\infty} \\ &= \frac{(a; q)_\infty}{(b; q)_\infty} \sum_{s=0}^{\infty} \frac{(\frac{b}{a}; q)_s (b; q)_{m+s}}{(q; q)_s (a; q)_{m+s+1}} (a\gamma\delta t^2)^s. \end{aligned}$$

Hence, we deduce that

$$\begin{aligned} &\int_0^\pi C_m^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \frac{(b\gamma t e^{i\theta}, b\delta t e^{-i\theta}; q)_\infty}{(\gamma t e^{i\theta}, \delta t e^{-i\theta}; q)_\infty} \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta \\ &= \frac{2\pi(a, a; q)_\infty (a^2; q)_m (\gamma\delta)^m t^m}{(q, a^2; q)_\infty (q; q)_m} \sum_{n=0}^{\infty} \frac{(\frac{b}{a}; q)_n (b; q)_{m+n}}{(q; q)_n (a; q)_{m+n+1}} (a\gamma\delta t^2)^n. \end{aligned}$$

With the help of (3.5), it follows that

$$\begin{aligned} &\sum_{m=0}^{\infty} \left\{ \int_0^\pi C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) C_m^{(b\gamma, b\delta, \gamma, \delta)}(e^{i\theta}; q) \omega^{(a\gamma, a\delta, \gamma, \delta)}(\cos \theta | q) d\theta \right\} t^m \\ &= \frac{2\pi(a, a; q)_\infty (a^2; q)_n (\gamma\delta)^n}{(q, a^2; q)_\infty (q; q)_n} \sum_{j=0}^{\infty} \frac{(\frac{b}{a}; q)_j (b; q)_{n+j}}{(q; q)_j (a; q)_{n+j+1}} (a\gamma\delta)^j t^{n+2j}. \end{aligned}$$

Comparing the coefficients of powers of t in the above equation, we complete the proof of Theorem 1.3. \square

Besides, we also obtain the recurrence relationship between $C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q)$ and $C_m^{(b\gamma, b\delta, \gamma, \delta)}(e^{i\theta}; q)$.

Proposition 3.1. *For $\max\{|a|, |b|\} < 1$, we have*

(3.11)

$$C_m^{(b\gamma, b\delta, \gamma, \delta)}(e^{i\theta}; q) = \sum_{\substack{n=0 \\ n \equiv m \pmod{2}}}^m (1 - aq^n) C_n^{(a\gamma, a\delta, \gamma, \delta)}(e^{i\theta}; q) \frac{(\frac{b}{a}; q)_{\frac{m-n}{2}} (b; q)_{\frac{m+n}{2}}}{(q; q)_{\frac{m-n}{2}} (a; q)_{\frac{m+n}{2}+1}} (a\gamma\delta)^{\frac{m-n}{2}}.$$

Acknowledgements. The first author was supported in part by the National Natural Science Foundation of China (Grant No. 12201387).

REFERENCES

- [1] W.A. Al-Salam and A. Verma, Some remarks on q -beta integral. Proc. Amer. Math. Soc. 85, 360–362 (1982).
- [2] R. Askey and M. Ismail, The Rogers q -ultraspherical polynomials, Approximation Theory. III (E. W. Cheney, ed.), Academic Press, New York, 1980, pp. 175–182.
- [3] R. Askey and M. Ismail, A generalization of the ultraspherical polynomials, Studies in Pure Mathematics (P. Erdős, ed.), Birkhäuser, Basel, 1983, pp. 55–678.
- [4] R. Askey and J. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, Mem. Amer. Math. Soc. 54 (1985), no. 319, iv+55
- [5] D. Chen and Z.-G. Liu, New proof of some q -orthogonal functions, preprint.
- [6] G. Gasper, *Orthogonality of certain functions with respect to complex valued weights*. Canadian J. Math. **33** (1981), no.5, 1261–1270. MR0638380
- [7] G. Gasper and M. Rahman, Basic Hypergeometric Series, Cambridge Univ. Press, Cambridge, MA, 2004.
- [8] P.C. Greiner, Spherical harmonics on the Heisenberg group, Canadian Math. Bull. 23 (1980), 383-396.
- [9] Mourad E. H. Ismail, Classical and quantum orthogonal polynomials in one variable, with two chapters by Walter Van Assche, with a foreword by Richard A. Askey, reprint of the 2005 original, Encyclopedia of Mathematics and its Applications, vol. 98, Cambridge University Press, Cambridge, 2009.
- [10] F. Jackson, On q -definite integrals, Q. J. Pure Appl. Math. 50 (1910) 101-112.
- [11] R. Koekoek, P.A. Lesky and R.F. Swarttouw, Hypergeometric orthogonal polynomials and their q -analogues, with a foreword by Tom H. Koornwinder, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2010.
- [12] Z.-G. Liu, On a system of q -partial differential equations with applications to q -series. Analytic number theory, modular forms and q -hypergeometric series, 445–461, Springer Proc. Math. Stat., 221, Springer, Cham, 2017.
- [13] L.J. Rogers, Third memoir on the expansion of certain infinite products, Proc. London Math. Soc. 26 (1895) 15–32.

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, PEOPLE'S REPUBLIC OF CHINA

NEWTOUCH CENTER FOR MATHEMATICS OF SHANGHAI UNIVERSITY, SHANGHAI, PEOPLE'S REPUBLIC OF CHINA

Email address: mathcdd@shu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, PEOPLE'S REPUBLIC OF CHINA

Email address: siyuyin0113@126.com