

A GLUING CONSTRUCTION OF D_k ALF GRAVITATIONAL INSTANTONS AND EXISTENCE OF NON-HOLOMORPHIC MINIMAL SPHERES

XUWEN ZHU

ABSTRACT. This note extends the construction of D_k ALF gravitational instantons in Singer–Schroers [SS21] to a new case where the nonlinear superposition is given by the D_1 Atiyah–Hitchin metric and $k - 1$ copies of A_0 Taub-NUT metrics. We then give a general class of ALF spaces such that each of them contains a non-holomorphic minimal sphere. Together with Foscolo’s construction [Fos19] this gives a large class of $K3$ surfaces containing non-holomorphic minimal spheres.

1. INTRODUCTION

A gravitational instanton is a complete noncompact 4-dimensional hyperKähler manifold with L^2 curvature decay at infinity. There has been a long history in the study of gravitational instantons and a lot of recent developments have been made. Gravitational instantons arise as building blocks for degeneration of $K3$ surfaces [GW00, Fos19, HSVZ22, CVZ20], which makes them powerful tools in probing $K3$ surfaces near degeneration limits.

The gravitational instantons have been classified into six types, ALE / F / G / H / G^* / H^* , based on the end structure at infinity [Min10, CC21, SZ22]. In this paper we will discuss the ALF gravitational instantons, which is modeled at infinity by quotients of circle bundles over \mathbb{R}^3 and has cubic volume growth. There are two families of ALF spaces depending on the monodromy at infinity being cyclic or dihedral, A_k and D_k , labelled by non-negative integers k . In this paper we will discuss the construction of D_k type ALF spaces via a nonlinear superposition of D_1 and $k - 1$ copies of A_0 ALF spaces, and using this we will show the existence of non-holomorphic minimal spheres in these spaces as well as in $K3$ surfaces.

This gluing construction extends the work by Schroers and Singer [SS21], who obtained D_k ALF gravitational instantons by gluing the moduli space \mathcal{M}_2^0 of centered $SU(2)$ monopoles (which is the only D_0 ALF space) with k copies of Taub-NUT space (each of which is an A_0 ALF

space). As commented in their paper, this construction can be extended to other cases. In this paper we work on the case where the center piece is replaced by the Atiyah–Hitchin metric AH [AH88], which is defined to be the double cover of \mathcal{M}_2^0 and is a rotationally invariant D_1 space. A rough statement of the theorem is given below and a more explicit description will be given at Theorem 3.

Theorem 1. *For any $k \geq 1$ and any set $\{0, p_1, \dots, p_k\} \in \mathbb{R}^3$ of $k + 1$ distinct points, there exists $\epsilon_0 > 0$, such that for any $\epsilon \in (0, \epsilon_0)$ there exists a D_{k+1} ALF metric g_ϵ which is a nonlinear superposition of the Atiyah–Hitchin metric centered at 0 and k copies of Taub-NUT metrics centered at $p_i/\epsilon, 1 \leq i \leq k$.*

There has been a lot of progress in the analysis and construction of ALF spaces [Min09, Min10, CC19, BM11]. The A_k type ALF has been well understood as a multi-Taub-NUT space via gluing construction [Min11, GH78]. And D_k spaces have been constructed via twistor methods [Hit84] and desingularization [BM11, Auv18, Dan93]. The construction in [SS21] gives an explicit gluing constructions of D_k type ALF based on Sen’s proposal [Sen97], and the construction follows the physical interpretation of modeling electronically charged particles like protons and electrons [AMS12]. There are also works by Cherkis, Kapustin, Hitchin, Roček et.al [CH05, CK98, CK99, LRc88, IRc96] describing ALF spaces as moduli space of monopoles with prescribed singularities. The construction in [SS21] gave D_k ALF when the k Taub-NUT singularities are well separated from the origin, while the ones constructed in this paper can be viewed as the limit of these D_k spaces when one Taub-NUT singularity approaches zero.

It can be potentially further generalized, as suggested in [SS21], to cases of composing a D_j ALF space with various A_{ν_i-1} ALF to generate other $D_{k'}$ spaces, where $k' = j + \sum \nu_i$, which would give other limiting cases when more TN singularities travel to zero or to each other. Since in this paper we are mainly interested in using the property of Atiyah–Hitchin space, we will only work on one generalization where the center piece is exactly given by AH.

Our construction follows the same procedure and the analysis is done following [SS21], as the change of the center piece from \mathcal{M}_0^2 to AH does not change the asymptotic behavior. The mapping properties of such metrics can be analyzed using the fiber boundary calculus, or ϕ -calculus [MM98, Vai01, HHM04], which captures the asymptotic metric structure of ALF spaces.

As a consequence of the construction, we obtain a family of D_k spaces that, with properly chosen marked points, degenerate to the Atiyah–Hitchin space as the Gromov Hausdorff limit. One special feature of the AH space is that it contains a stably minimal sphere that is not holomorphic with any of the complex structures in the hyperKähler rotation [MW93]. With this construction we can obtain such special sphere in all nearby D_k space.

Theorem 2 (same as Theorem 4). *For any $k > 1$, there is a family of D_k ALF metrics $\{g_\epsilon\}_{\epsilon \in (0, \epsilon)}$, such that each of them contains a minimal sphere that is not holomorphic with respect to any of the compatible complex structure.*

In [Fos19] Foscolo constructed a collapsing sequence of K3 surfaces using ALF spaces as part of building blocks. As a corollary he used Atiyah–Hitchin space as the building block to show the existence of minimal non-holomorphic sphere in some K3 surface near the degeneration limit. Now with more ALF spaces constructed above with non-holomorphic spheres, we extend this result of Foscolo and show that there is a neighborhood in the moduli space of K3 surfaces such that each of these has a special sphere.

Corollary 1 (same as Corollary 3). *There is an open neighborhood in the moduli space of hyperKähler metrics on K3, such that each metric in this neighborhood contains a stably minimal sphere that is not holomorphic with respect to any compatible complex structures.*

This corollary is connected to one interpretation of the Weak Gravity Conjecture [AHMNV07], which predicts the existence of “non-BPS” particles. In mathematics such particles can be interpreted as non-holomorphic submanifolds in Calabi–Yau manifolds in the large complex limit [HLM⁺20, DLMS20, LSVY23]. The corollary above can be seen as one of the examples.

The paper is organized in the following order: Section 2 describes the two building blocks, the multi-Taub-NUT space and the Atiyah-Hitchin metric, then gives the gluing construction and perturbative argument to find the hyperKähler metric. Section 3 proves the existence of non-holomorphic spheres in some D_k ALF spaces and K3 surfaces.

Acknowledgement: The author is partially supported by NSF DMS-2305363

2. THE GLUING CONSTRUCTION OF D_k ALF SPACES

The gluing construction is a modification from [SS21], and we describe the construction below. The basic idea is to first construct an

adiabatic Gibbons–Hawking Ansatz with symmetric Taub–NUT singularities which give the correct D_k asymptotic topology at infinity after the quotient of an involution action. After that we glue in a double branched cover of Atiyah Hitchin metric in the center. This produces an approximate hyperKähler metric after involution. Finally we solve for the actual hyperKähler metric by perturbation of hyperKähler triples. The set up is similar to [SS21] where the center piece is a double branched cover of \mathcal{M}_2^0 . We also need to modify the coefficients in the adiabatic Gibbons–Hawking ansatz to be compatible with AH metric. On the other hand, the asymptotic behavior of \mathcal{M}_2^0 and Atiyah Hitchin metric are similar since AH is the double cover of \mathcal{M}_2^0 , the analysis used for perturbing the approximate triple to get the exact solution can be applied directly.

Now we describe the two pieces needed for gluing

2.1. The multi-Taub-NUT space M_ϵ . Fix $\delta \in (0, 1/2)$, and let $\epsilon \in (0, \delta^2)$ which will go to 0 in the limit later. Fix $p_1, \dots, p_k \in \mathbb{R}^3$ sufficiently large, we consider a family of 4-manifolds M_ϵ , each of which is a singular circle fibration over $\{x \in \mathbb{R}^3 : |x| > 2\epsilon\}$. We consider the following adiabatic family of Gibbon–Hawking ansatz on M_ϵ

$$(1) \quad g_\epsilon = h_\epsilon \frac{|dx|^2}{\epsilon^2} + h_\epsilon^{-1} \alpha^2, \quad \alpha = *_\epsilon dh_\epsilon$$

where α is a connection 1-form of the circle bundle, and $*_\epsilon$ is the Hodge star with respect to the flat metric $\frac{|dx|^2}{\epsilon^2}$ on the base $\mathbb{R}_{|x|>2\epsilon}^3$; the singular harmonic function h_ϵ on $\mathbb{R}_{|x|>2\epsilon}^3$ is given by

$$(2) \quad h_\epsilon = 1 - \frac{\epsilon}{|x|} + \sum_{v=1}^k \left(\frac{\epsilon}{2|x + p_v|} + \frac{\epsilon}{2|x - p_v|} \right).$$

We take the singular fibration associated with the Gibbons–Hawking ansatz (1) by

$$(3) \quad \phi : M_\epsilon \rightarrow \mathbb{R}_{|x|>2\epsilon}^3$$

and note the fibration is only singular at $P := \{\pm p_1, \dots, \pm p_2\}$. This ansatz is modified from [SS21, equation (1.4)] by replacing the coefficients of $\frac{1}{|x|}$ from 2ϵ to ϵ which will match the the Atiyah–Hitchin metric in the center later. At $\pm p_i$ this metric has a Taub–NUT type singularity, which is natural since A_k ALF spaces can be viewed as a multi-Taub–NUT space.

The involution action of $\iota : x \mapsto -x$ on \mathbb{R}^3 applies to the fibration. Note that the asymptotic of h_ϵ is given by

$$(4) \quad h_\epsilon = 1 + \frac{\epsilon(2k-2)}{2|x|} + O(\epsilon|x|^{-3}), \quad |x| \gg 1$$

which, after quotient by the involution ι , gives the asymptotic topology of a dihedral D_{k+1} ALF.

We recall one fact about the behavior of g_ϵ when lifted to the compactified space, where the detailed proof was given in [SS21, Proposition 3.4].

Definition 1. A compact manifold with boundary (X, g) has a fiber boundary structure, or ϕ -structure, if there is a fibration $\phi : \partial X \rightarrow Y$ where Y is a compact manifold without boundary and F is the fiber. The space of ϕ -vector fields \mathcal{V}_ϕ is given by b-vector fields (i.e. they are tangent to ∂X) which also satisfy

$$v|_{\partial X} \in C^\infty(\partial X, T(\partial X/Y)),$$

and

$$v(\rho) \in \rho^2 C^\infty(X)$$

where ρ is a boundary defining function for ∂X .

Locally using coordinates $\phi(y_1, \dots, y_b, z_1, \dots, z_f) = (y_1, \dots, y_b)$, the ϕ -vector fields are spanned by

$$\rho^2 \frac{\partial}{\partial \rho}, \rho \frac{\partial}{\partial y_1}, \dots, \rho \frac{\partial}{\partial y_b}, \frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_f}.$$

And a ϕ -metric is a smooth metric on the ϕ -tangent bundle $T_\phi X$, locally given by

$$g_\phi = \frac{d\rho^2}{\rho^4} + \frac{\phi^* g_Y}{\rho^2} + g_F.$$

The fibration $\phi : M_\epsilon \rightarrow \mathbb{R}_{|x|>2\epsilon}^3$ can be lifted to the radially-compactified base $\overline{\mathbb{R}^3} \setminus \{|x| \leq 2\epsilon\}$. The radial compactification of \mathbb{R}^3 is obtained by adding a sphere $\mathbb{S}^2 = \partial \overline{\mathbb{R}^3}$ at infinity. We use the inverted polar coordinates $\rho = 1/|x|, y = \rho x$ near the boundary, and the metric g_ϵ in (1) lifts to a fiber boundary metric of the form

$$g_\epsilon = h_\epsilon \left(\frac{d\rho^2}{\rho^4} + \frac{|dy|^2}{\rho^2} \right) + h_\epsilon^{-1} \bar{\alpha}^2,$$

where h_ϵ and $\bar{\alpha} = *_\epsilon dh_\epsilon$ lift to be smooth on $\overline{\mathbb{R}^3} \setminus \{|x| \leq 2\epsilon\}$ (one can check this by changing variables in (2)).

2.2. The branched cover \widehat{AH}_ϵ . We will glue the (double branched cover) of Atiyah–Hitchin metric into the center. Following the notation in [SS21], we denote by \widehat{AH} the space $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \setminus$ anti-diagonal, or in coordinates:

$$(5) \quad \widehat{AH} = \{(z, w) \in \mathbb{C}^2 \times \mathbb{C}^2 : z_1 \bar{w}_1 + z_2 \bar{w}_2 \neq 0\} / \mathbb{C}^* \times \mathbb{C}^*$$

This space can be identified via the space of X- and Y-ellipsoid (see [SS21, Appendix A] for details). In the region near infinity, one can use coordinates $\{X \in \mathbb{C}^3 \mid \sum_{i=1}^3 X_i^2 = 1\}$. If we write $X = \tilde{x} + i\xi$, $\tilde{x}, \xi \in \mathbb{R}^3$, then they satisfy

$$|\tilde{x}|^2 = 1 + |\xi|^2, \quad \tilde{x} \cdot \xi = 0.$$

There are two involution maps on \widehat{AH} :

$$\begin{aligned} s : (\tilde{x}, \xi) &\mapsto (-\tilde{x}, -\xi) \\ r : (\tilde{x}, \xi) &\mapsto (\tilde{x}, -\xi). \end{aligned}$$

The relation between \widehat{AH} , the Atiyah–Hitchin space AH , and the D_0 monopole moduli space \mathcal{M}_2^0 is given by

$$(6) \quad AH = \widehat{AH}/s, \quad \mathcal{M}_2^0 = AH/r.$$

The relation implies that near infinity \widehat{AH} is a circle bundle $\psi : \widehat{AH} \rightarrow \mathbb{R}^3 \setminus \{|x'| > 1\}$ with the following asymptotic

$$(7) \quad g_{\widehat{AH}} = \left(1 - \frac{1}{|x'|}\right) |dx'|^2 + \left(1 - \frac{1}{|x'|}\right)^{-1} (\alpha')^2 + O(e^{-|x'|})$$

where x' is bijectively related to \tilde{x} in an implicit way. In particular, AH and its branched cover \widehat{AH} are both strongly ALF spaces by [SS21, Definition 3.5] and the asymptotic analysis at infinity can be applied the same way.

To connect with the adiabatic Gibbons–Hawking ansatz constructed before, we consider the following metric

$$(8) \quad g_{\widehat{AH}, \epsilon} = h'_\epsilon |dx'|^2 + (h'_\epsilon)^{-1} (\alpha')^2$$

where

$$h'_\epsilon = 1 - \frac{1}{|x'|} + \sum_{v=1}^k \frac{\epsilon}{|p_v|}, \quad \alpha' = *dh'_\epsilon.$$

We call the associated circle fibration total space \widehat{AH}_ϵ and the fibration

$$(9) \quad \psi : \widehat{AH}_\epsilon \rightarrow \mathbb{R}^3 \setminus \{|x'| > 1/\delta\}.$$

With the same compactification at infinity of \mathbb{R}^3 , we can see that $g_{\widehat{AH}, \epsilon}$ also lifts to be a ϕ -metric for each ϵ .

2.3. Gluing construction. The gluing is done on the connecting neck region

$$\left\{\frac{1}{\delta} < |x'| < \frac{\delta}{\epsilon}\right\} \subset \widehat{AH}_\epsilon$$

and

$$\left\{\frac{\epsilon}{\delta} < |x| < \delta\right\} \subset M_\epsilon$$

we use the identification map $x' = \kappa(x) = x/\epsilon$, $\kappa^*(\alpha_0) = \alpha'$ to identify g_ϵ and $g_{\widehat{AH}_\epsilon}$, and note that under this map h_ϵ and $h_{\epsilon'}$ are identified up to $O(\epsilon^3)$.

However in order to apply the analysis, we need to modify the space. The total space is glued from two parts: \mathcal{W}_0 corresponding to the Atiyah-Hitchin metrics in the center, and \mathcal{W}_1 corresponding to the multi-Taub-NUT singularities, where \mathcal{W}_0 and \mathcal{W}_1 are essentially M_ϵ and \widehat{AH}_ϵ with the additional steps of compactification at infinity and singularity resolution at $\epsilon = 0$. The construction of \mathcal{W}_1 and \mathcal{W}_0 are almost the same as in [SS21], while some modification is needed.

The construction is first done on the double cover $\widetilde{\mathcal{W}}_i$, $i = 0, 1$, and the final space \mathcal{W} is the quotient by the involution ι . First we construct $\widetilde{\mathcal{W}}_0$, which is defined as

$$(10) \quad \widetilde{\mathcal{W}}_0 = \{(z', \epsilon') \in \widehat{AH}_\epsilon \times [0, \epsilon_0) : \frac{1}{\delta} < |\psi(z')| < \frac{\delta}{\epsilon'}\}.$$

The construction of $\widetilde{\mathcal{W}}_1$ is the same as in [SS21, Section 4.1], obtained by pulling back the product of interval $[0, \delta^2)_\epsilon \times M_\epsilon$ to the resolved base \widetilde{B} . We briefly explain the construction of \widetilde{B} here. Define the blow up

$$(11) \quad \widetilde{B} = [\mathbb{R}_x^3 \times [0, \delta^2)_\epsilon; \{(0, 0), \pm p_i \times \{0\}, i = 1, \dots, k\}] \cap \{|x| > \epsilon/\delta\}$$

which introduced $2k + 1$ front faces. We then pull back the circle fibration $\phi : M_\epsilon \rightarrow \mathbb{R}^3 \setminus \{|x| \leq \epsilon/\delta\}$ to \widetilde{B} , such that when restricting to each $\epsilon > 0$ it is the same as the original fibration $\phi : M_\epsilon \rightarrow \mathbb{R}^3 \setminus \{\frac{\epsilon}{\delta} < |x| \leq \epsilon/\delta\}$. On the other hand, when restricting to $\epsilon = 0$ it lifts to a circle bundle over the blown up space $[\mathbb{R}^3; \{0, \pm p_i\}]$.

We construct the space gluing $\widetilde{\mathcal{W}}_1$ into $\widetilde{\mathcal{W}}_0$, identifying the torus neighborhood using the lift of the map κ described earlier, i.e. we identify the neighborhood

$$\{(z', \epsilon') : 0 \leq \epsilon' \leq \epsilon_0, \delta^{-1} < |\psi(z')| < \delta\epsilon^{-1}\} \subset \widetilde{\mathcal{W}}_0$$

and

$$\{(z, \epsilon) : 0 \leq \epsilon \leq \epsilon_0, \epsilon\delta^{-1} \leq |\phi(z)| \leq \delta\} \subset \widetilde{\mathcal{W}}_1$$

via the lift of the map $\phi(z) = x := \psi(z')\epsilon, \epsilon = \epsilon'$, and we denote this map by κ . The total space is defined as

$$(12) \quad \widetilde{\mathcal{W}} := \widetilde{\mathcal{W}}_1 \coprod_{\kappa} \widetilde{\mathcal{W}}_2, \quad \mathcal{W} := \widetilde{\mathcal{W}}/\iota.$$

Here \mathcal{W} is a manifold with corners, where the boundary faces include: I_∞ which corresponds to the fibration over the boundary of the compactified \mathbb{R}^3 of \mathcal{W}_1 ; X_0 that is the lift of the fibration to the boundary face created by blowing up $(0, 0)$; $X_i, i = 1, \dots, k$, that are the lift of the fibration to the boundary faces created by blowing up $\{\pm p_i, 0\}$; X_{ad} the remaining part of $\epsilon = 0$. Here is a picture of the total space:




FIGURE 1. An illustration of the resolved total space \mathcal{W} obtained from gluing \mathcal{W}_0 and \mathcal{W}_1 , with boundary faces and associated boundary defining functions on $\pi^{-1}(0)$

There is a projection map $\pi : \mathcal{W} \rightarrow [0, \epsilon_0)_\epsilon$, such that for each nonzero ϵ the fiber is an approximate ALF metric, while for $\epsilon = 0$ the fiber is a manifold with corners. We also note that ϵ is the total boundary defining function, i.e. if we define σ_ν to be the boundary defining function of $X_\nu, \nu = 0, 1, \dots, k$, and ρ to be the boundary defining function of X_{ad} , then the product of all boundary defining functions on the front face is

$$(13) \quad \epsilon = \rho \sigma_0 \dots \sigma_k.$$

2.4. Solving for the hyperKähler triples. By the gluing construction discussed above, we arrived at a family of approximate metric, each living on the fiber $\pi^{-1}(\epsilon)$ of \mathcal{W} . We will follow the set up in [SS21] to perturb these metrics and solve for the exact hyperKähler triples on each fiber. Recall that every hyperKähler metric on a 4-manifold gives a hyperKähler triple $\omega = (\omega_1, \omega_2, \omega_3)$ such that each ω_i is a symplectic form and they satisfy the following equation [Don06]

$$(14) \quad 0 = Q(\omega)_{jk} := \omega_j \wedge \omega_k - \frac{1}{3} \left(\sum_{i=1}^3 \omega_i^2 \right) \delta_{jk}$$

Conversely a hyperKähler triple determines a hyperKähler metric [SS21, Theorem 2.3].

For a adiabatic Gibbons-hawking ansatz in (1), the triple is given by the following $\omega_\epsilon = (\omega_1, \omega_2, \omega_3)$ [SS21, (2.21)]

$$(15) \quad \begin{aligned} \omega_1 &= \alpha \wedge \frac{dx_1}{\epsilon} + h_\epsilon \frac{dx_2 \wedge dx_3}{\epsilon^2} \\ \omega_2 &= \alpha \wedge \frac{dx_2}{\epsilon} + h_\epsilon \frac{dx_3 \wedge dx_1}{\epsilon^2} \\ \omega_3 &= \alpha \wedge \frac{dx_3}{\epsilon} + h_\epsilon \frac{dx_1 \wedge dx_2}{\epsilon^2} \end{aligned}$$

which can be written as $\Omega_\epsilon + db$ where $b \in O(|\epsilon|^3)$ where Ω_ϵ is given by replacing h_ϵ by its leading term $1 - \frac{\epsilon}{|x|} + \sum_{v=1}^k \frac{\epsilon}{|p_v|}$. Similarly for the branched cover $\widehat{AH}_{\epsilon'}$ the triple is given by $\omega_{AH, \epsilon'} = \Omega_{\epsilon'} + da$ where $a = O(e^{-|x'|})$ and $\Omega_{\epsilon'}$ is given by

$$(16) \quad \begin{aligned} \omega'_1 &= \alpha' \wedge dx'_1 + h'_\epsilon dx'_2 \wedge dx'_3 \\ \omega'_2 &= \alpha' \wedge dx'_2 + h'_\epsilon dx'_3 \wedge dx'_1 \\ \omega'_3 &= \alpha' \wedge dx'_3 + h'_\epsilon dx'_1 \wedge dx'_2 \end{aligned}$$

We will patch the two triples on each fiber to an initial approximation, then solve the equation and show that the solutions are uniformly controlled up to the $\pi^{-1}(0)$ face. This will give a family of smooth hyperKähler triples hence the corresponding family of hyperKähler metrics. To give a more precise statement, we use the following space:

Definition 2. Let $T_\phi(\mathcal{W}/I)$ denote the subspace of the total ϕ -tangent space on \mathcal{W} that is spanned by ϕ -tangent vectors along the ϵ - fiber direction. Similarly the $T_\phi^*(\mathcal{W}/I)$ and $\Omega_\phi^2(\mathcal{W}/I)$ denote the fiberwise ϕ -cotangent space and (smooth) 2-forms.

The symplectic forms will be in the space of $\Omega_\phi^2(\mathcal{W}/I)$, and the operator Q maps into the space $S_0^2 \mathbb{R}^3 \otimes \Lambda^4 T_\phi^*(\mathcal{W}/I)$. With these notations, now we state the theorem.

Theorem 3. Fix $\{0, p_1, \dots, p_k\}$ a set of distinct $k + 1$ points. Then there exists $\epsilon_0 > 0$, such that on $\pi^{-1}([0, \epsilon_0]) \subset \mathcal{W}$, there exists smooth triple $\omega_{S_\epsilon} = (\omega_1, \omega_2, \omega_3)$ with $\omega_i \in \Omega_\phi^2(\mathcal{W}/I)$ satisfying $Q(\omega) = 0$. The corresponding fiber metric $g_{S_\epsilon, \epsilon}$ satisfies

- (1) $\{g_{S_\epsilon, \epsilon}\}_{\epsilon \in I}$ is a family of smooth fiberwise ϕ -metric on \mathcal{W} ;
- (2) $g_{S_\epsilon, 0}|_{X_0}$ restricts to be the Atiyah–Hitchin metric, and $g_{S_\epsilon, 0}|_{X_i}$ restricts to be the Taub–NUT metric for $i = 1, \dots, k$.

Proof. The proof follows almost the same as in [SS21]. Such a strategy has also been used in [MZ18, MZ19]. We first construct an approximate

solution ζ_0 for the hyperKähler triple by patching the two model metrics together using a cut off function χ that is supported near \mathcal{W}_0 :

$$\omega_\chi := \chi\omega_{AH,\epsilon'} + (1 - \chi)\omega_\epsilon$$

where the two hyperKähler triples are defined in (15) and (16). Note that the harmonic function in $\omega_{AH,\epsilon'}$ and ω_ϵ agree up to a closed form of order ϵ^3 , therefore the initial approximation solves the equation up to the following error

$$Q(\omega_\chi) \in \epsilon^3 C^\infty(\mathcal{W}, S_0^2 \mathbb{R}^3 \otimes \Lambda^4 T_\phi^*(\mathcal{W}/I))$$

which also vanishes to infinite order away from X_ν .

The second step is to solve the equation iteratively to ϵ^∞ error. This step follows from the discussion in [SS21, Section 5], where they solve the error iteratively near the corner of $X_\nu \cap X_{ad}$ on the two faces. Recall that the linearized operator of Q is given by 12 copies of scalar Laplacians [SS21,]. When restricting to the solvability near X_ν , note that the replacement AH is still a strongly ALF metric hence the analysis carries through. The solvability near X_{ad} is the same as before. As a result we can iteratively construct a formal power series solution $\zeta = \omega_\chi + dc$ such that

$$Q(\zeta) \in \dot{C}^\infty(\mathcal{W}, S_0^2 \mathbb{R}^3 \otimes \Lambda^4 T_\phi^*(\mathcal{W}/I))$$

that is, the error vanishes to infinite order on every boundary face.

The third step is to show there is a bounded inverse of the linearized operator on \mathcal{W} . This step again relies on that one can patch inverses on X_ν and X_{ad} . Since the only difference in the structure of X_ν from [SS21] is that X_0 is changed from $\overline{M_0^2}$ by \overline{AH} , the inverse is constructed in the same way as in [SS21, Theorem C.3] for a strongly ALF space. On the other hand, on X_{ad} the inverse is constructed explicitly using Fourier series and Green's function where the only difference is the coefficient in h_ϵ which does not change the proof. As a result Δ_ζ is an invertible operator on the domain constructed in [SS21, Theorem 6.6].

Finally by implicit function theorem we get the exact solution ω_{S_ϵ} with $Q(\omega_{S_\epsilon}) = 0$. The smoothness comes from the iterative series and invertibility of the linearized operator, and the restriction on X_ν follows because AH and TN are the 0-th order terms in the metric expansion. \square

As a corollary, such a family of metrics degenerates into the front face, and (under correct scaling and choice of base points) give Gromov-Hausdorff convergence to AH and TN respectively.

- Corollary 2.** (1) Take $\{z_\epsilon\}$ satisfying $|\phi(z_\epsilon)| < \frac{\epsilon}{\delta}$, then the sequence $(g_{S_{e,\epsilon}}, z_\epsilon)$ converges to $(g_{AH}, 0)$ in the Gromov-Hausdorff limit as $\epsilon \rightarrow 0$;
- (2) Take $\{z'_\epsilon\}$ satisfying $|\phi(z'_\epsilon) - p_i| < \epsilon$, then $(g_{S_{e,\epsilon}}, z'_\epsilon)$ converges to $(g_{TN}, 0)$ in the Gromov-Hausdorff limit.

Remark. Different types of gravitational instantons are expected to relate to each other, and similar results of this type where gravitational instantons degenerate into another type can be seen in [Auv18, Che14, LSZ, Sal24b, Sal24a].

3. NON-HOLOMORPHIC MINIMAL SPHERES

In [MW93] Micallef and Wolfson showed that in the Atiyah–Hitchin space there is a special sphere that is stably minimal and not holomorphic with respect to any complex structures in the hyperähler rotation. Using the gluing result above, we extend this result to D_{k+1} spaces constructed above. As a perturbation result, the existence can be proved for D_{k+1} spaces where the Taub-NUT singularities are sufficiently far from the origin.

Theorem 4. *There exists ϵ_0 , such that for every $\epsilon \in (0, \epsilon_0)$ there exists a minimal sphere in $g_{S_{e,\epsilon}}$ that is non-holomorphic with respect to any of the compatible complex structures.*

Proof. We start with the approximate metric $g_{0,\epsilon}$ constructed via patching the $g_{\widehat{AH},\epsilon}$ and g_ϵ . Since the minimal sphere in $g_{\widehat{AH}}$ is compact, so by shrinking ϵ_0 if necessary, there exists $\Sigma_\epsilon \subset (AH|_{|x'| < \delta/\epsilon}, g_{\widehat{AH},\epsilon}) \subset (\pi^{-1}(\epsilon), g_{0,\epsilon})$ for every $\epsilon \in (0, \epsilon_0)$. Note that this minimal sphere Σ satisfies $[\Sigma] \cdot [\Sigma] = -4$ which is not $2 * \text{genus}(\Sigma) - 2$, hence it cannot be holomorphic in a hyperKähler manifold via adjoint formula. Now we apply White’s implicit function theorem [Whi91] (which does not require the ambient manifold to be compact) to the actual hyperKähler metric $g_{S_{E,\epsilon}}$ perturbed from $g_{0,\epsilon}$, we get a stably minimal non-holomorphic sphere in the D_{k+1} space. \square

Corollary 3. *There exists a neighborhood of K3 such that every K3 has a minimal sphere that is non-holomorphic with respect to any of the complex form.*

Remark. The statement follows from [Fos19, Theorem 7.1] via perturbation, however this construction provides a larger neighborhood and gives more explicit examples compared to the statement in [MW06].

Proof. The proof follows exactly the same as in [Fos19, Theorem 7.1] by considering any D_{k+1} metric constructed above as a building block in

the collapsing sequence. Since the approximate metric g_ϵ constructed using D_k bubbles contains the special sphere, by the same perturbation argument in the previous theorem we obtain the existence of non-holomorphic minimal sphere. \square

REFERENCES

- [AH88] Michael Atiyah and Nigel Hitchin, *The geometry and dynamics of magnetic monopoles*, M. B. Porter Lectures, Princeton University Press, Princeton, NJ, 1988. MR 934202
- [AHMNV07] Nima Arkani-Hamed, Luboš Motl, Alberto Nicolis, and Cumrun Vafa, *The string landscape, black holes and gravity as the weakest force*, J. High Energy Phys. (2007), no. 6, 060, 17. MR 2326591
- [AMS12] M. F. Atiyah, N. S. Manton, and B. J. Schroers, *Geometric models of matter*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **468** (2012), no. 2141, 1252–1279. MR 2910348
- [Auv18] Hugues Auvray, *From ALE to ALF gravitational instantons*, Compos. Math. **154** (2018), no. 6, 1159–1221. MR 3826456
- [BM11] Olivier Biquard and Vincent Minerbe, *A Kummer construction for gravitational instantons*, Comm. Math. Phys. **308** (2011), no. 3, 773–794. MR 2855540
- [CC19] Gao Chen and Xiuxiong Chen, *Gravitational instantons with faster than quadratic curvature decay (II)*, J. Reine Angew. Math. **756** (2019), 259–284. MR 4026454
- [CC21] ———, *Gravitational instantons with faster than quadratic curvature decay. I*, Acta Math. **227** (2021), no. 2, 263–307. MR 4366415
- [CH05] Sergey A. Cherkis and Nigel J. Hitchin, *Gravitational instantons of type D_k* , Comm. Math. Phys. **260** (2005), no. 2, 299–317. MR 2177322
- [Che14] Sergey A. Cherkis, *Phases of five-dimensional theories, monopole walls, and melting crystals*, Journal of High Energy Physics **2014** (2014), no. 6, 1–38.
- [CK98] Sergey A. Cherkis and Anton Kapustin, *D_k gravitational instantons and Nahm equations*, Adv. Theor. Math. Phys. **2** (1998), no. 6, 1287–1306. MR 1693628
- [CK99] ———, *Singular monopoles and gravitational instantons*, Comm. Math. Phys. **203** (1999), no. 3, 713–728. MR 1700937
- [CVZ20] Gao Chen, Jeff Viaclovsky, and Ruobing Zhang, *Collapsing Ricci-flat metrics on elliptic $K3$ surfaces*, Comm. Anal. Geom. **28** (2020), no. 8, 2019–2133. MR 4201795
- [Dan93] Andrew S. Dancer, *Dihedral singularities and gravitational instantons*, J. Geom. Phys. **12** (1993), no. 2, 77–91. MR 1231230
- [DLMS20] Mehmet Demirtas, Cody Long, Liam McAllister, and Mike Stillman, *Minimal surfaces and weak gravity*, J. High Energy Phys. (2020), no. 3, 021, 23. MR 4090116
- [Don06] S. K. Donaldson, *Two-forms on four-manifolds and elliptic equations*, Inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 153–172. MR 2313334

- [Fos19] Lorenzo Foscolo, *ALF gravitational instantons and collapsing Ricci-flat metrics on the $K3$ surface*, J. Differential Geom. **112** (2019), no. 1, 79–120. MR 3948228
- [GH78] Gary W Gibbons and Stephen W Hawking, *Gravitational multi-instantons*, Physics Letters B **78** (1978), no. 4, 430–432.
- [GW00] Mark Gross and P. M. H. Wilson, *Large complex structure limits of $K3$ surfaces*, J. Differential Geom. **55** (2000), no. 3, 475–546. MR 1863732
- [HHM04] Tamás Hausel, Eugenie Hunsicker, and Rafe Mazzeo, *Hodge cohomology of gravitational instantons*, Duke Math. J. **122** (2004), no. 3, 485–548. MR 2057017
- [Hit84] N. J. Hitchin, *Twistor construction of Einstein metrics*, Global Riemannian geometry (Durham, 1983), Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1984, pp. 115–125. MR 757213
- [HLM⁺20] Ben Heidenreich, Cody Long, Liam McAllister, Tom Rudelius, and John Stout, *Instanton resummation and the weak gravity conjecture*, J. High Energy Phys. (2020), no. 11, 166, 41. MR 4204112
- [HSVZ22] Hans-Joachim Hein, Song Sun, Jeff Viaclovsky, and Ruobing Zhang, *Nilpotent structures and collapsing Ricci-flat metrics on the $K3$ surface*, J. Amer. Math. Soc. **35** (2022), no. 1, 123–209. MR 4322391
- [IRc96] Ivan T. Ivanov and M. Roček, *Supersymmetric σ -models, twistors, and the Atiyah-Hitchin metric*, Comm. Math. Phys. **182** (1996), no. 2, 291–302. MR 1447294
- [LRc88] U. Lindström and M. Roček, *New hyper-Kähler metrics and new supermultiplets*, Comm. Math. Phys. **115** (1988), no. 1, 21–29. MR 929144
- [LSVY23] Cody Long, Artan Sheshmani, Cumrun Vafa, and Shing-Tung Yau, *Non-holomorphic cycles and non-BPS black branes*, Comm. Math. Phys. **399** (2023), no. 3, 1991–2043. MR 4580537
- [LSZ] Yu-Shen Lin, Sidharth Soundararajan, and Xuwen Zhu, *Non-collapsing degeneration of gravitational instantons*, In preparation.
- [Min09] Vincent Minerbe, *A mass for ALF manifolds*, Comm. Math. Phys. **289** (2009), no. 3, 925–955. MR 2511656
- [Min10] ———, *On the asymptotic geometry of gravitational instantons*, Ann. Sci. Éc. Norm. Supér. (4) **43** (2010), no. 6, 883–924. MR 2778451
- [Min11] ———, *Rigidity for multi-Taub-NUT metrics*, J. Reine Angew. Math. **656** (2011), 47–58. MR 2818855
- [MM98] Rafe Mazzeo and Richard B. Melrose, *Pseudodifferential operators on manifolds with fibred boundaries*, vol. 2, 1998, Mikio Sato: a great Japanese mathematician of the twentieth century, pp. 833–866. MR 1734130
- [MW93] Mario J. Micallef and Jon G. Wolfson, *The second variation of area of minimal surfaces in four-manifolds*, Math. Ann. **295** (1993), no. 2, 245–267. MR 1202392
- [MW06] M. Micallef and J. Wolfson, *Area minimizers in a $K3$ surface and holomorphicity*, Geom. Funct. Anal. **16** (2006), no. 2, 437–452. MR 2231469

- [MZ18] Richard Melrose and Xuwen Zhu, *Resolution of the canonical fiber metrics for a Lefschetz fibration*, J. Differential Geom. **108** (2018), no. 2, 295–317. MR 3763069
- [MZ19] ———, *Boundary behaviour of Weil-Petersson and fibre metrics for Riemann moduli spaces*, Int. Math. Res. Not. IMRN (2019), no. 16, 5012–5065. MR 4001023
- [Sal24a] Willem Adriaan Salm, *Construction of gravitational instantons with non-maximal volume growth via codimension-1 collapse*, 2024.
- [Sal24b] ———, *Elliptic analysis on collapsing gravitational instantons modelled using the gibbons-hawking ansatz*, 2024.
- [Sen97] Ashoke Sen, *A note on enhanced gauge symmetries in M- and string theory*, J. High Energy Phys. (1997), no. 9, Paper 1, 10. MR 1608180
- [SS21] B. J. Schroers and M. A. Singer, *D_k gravitational instantons as superpositions of Atiyah-Hitchin and Taub-NUT geometries*, Q. J. Math. **72** (2021), no. 1-2, 277–337. MR 4271388
- [SZ22] Song Sun and Ruobing Zhang, *Collapsing geometry of hyperkähler 4-manifolds and applications*, 2022.
- [Vai01] Boris Vaillant, *Index- and spectral theory for manifolds with generalized fibred cusps*, Bonner Mathematische Schriften [Bonn Mathematical Publications], vol. 344, Universität Bonn, Mathematisches Institut, Bonn, 2001, Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 2001. MR 1933455
- [Whi91] Brian White, *The space of minimal submanifolds for varying Riemannian metrics*, Indiana Univ. Math. J. **40** (1991), no. 1, 161–200. MR 1101226

NORTHEASTERN UNIVERSITY, BOSTON, MA 02115

Email address: x.zhu@northeastern.edu