

Electroweak Primordial Magnetic Blackhole: Cosmic Production and Physical Implication

Y. M. Cho,^{1,2,*} Sang-Woo Kim,^{3,†} and Seung Hun Oh^{4,‡}

¹*School of Physics and Astronomy, Seoul National University, Seoul 08826, Korea*

²*Center for Quantum Spacetime, Sogang University, Seoul 04107, Korea*

³*Department of Liberal Arts, Konkuk University, Seoul 143701, Korea*

⁴*Department of Liberal Arts, Korea Polytechnic University, Siheung 15073, Korea*

The electroweak monopole, when coupled to gravity, turns to the Reissner-Nordstrom type primordial magnetic blackhole whose mass is bounded below, with the lower bound $M_P\sqrt{\alpha}$. This changes the overall picture of the monopole production mechanism in the early universe drastically and has deep implications in cosmology. In particular, this enhances the possibility that the electroweak monopoles turned to the primordial magnetic blackholes could become the seed of stellar objects and galaxies, and account for the dark matter of the universe. Moreover, this tells that we have a new type of primordial blackhole different from the popular primordial blackhole in cosmology, the electroweak primordial magnetic blackhole based on a totally different production mechanism. We discuss the physical implications of the electroweak primordial magnetic blackhole.

Keywords: cosmological production of electroweak monopole, Ginzburg temperature, remnant electroweak monopole density, Cho-Maison monopole as the electroweak primordial magnetic black hole, evolution of the electroweak primordial magnetic blackholes, electroweak primordial magnetic black hole as seed of stellar objects and galaxies, source of intergalactic magnetic field and ultra high energy cosmic rays, electroweak primordial magnetic black hole as dark matter

I. INTRODUCTION

Since Dirac proposed the magnetic monopole generalizing the Maxwell's theory, the monopole has become an obsession in physics, experimentally as well as theoretically [1, 2]. After the Dirac monopole we have had the Wu-Yang monopole, the 't Hooft-Polyakov monopole, and the grand unification monopole [3–5]. But the electroweak monopole (also known as the Cho-Maison monopole) stands out as the most realistic monopole that could actually exist in nature and be detected [6–12]. This is because it exists in the standard model. So, if the standard model is correct as we believe, this monopole must exist.

Indeed the Dirac monopole in electrodynamics should transform to the electroweak monopole after the unification of the electromagnetic and weak interactions, and the Wu-Yang monopole in QCD is supposed to make the monopole condensation to confine the color. Moreover, the 'tHooft-Polyakov monopole exists only in a hypothetical theory, and the grand unification monopole which could have been amply produced at the grand unification scale in the early universe probably has become completely irrelevant at present universe after the inflation.

This makes the experimental confirmation of the electroweak monopole one of the most urgent issues in the

standard model after the discovery of the Higgs particle. In fact the detection of this monopole, not the Higgs particle, should be regarded as the final test of the standard model. For this reason the newest MoEDAL detector at LHC is actively searching for the monopole [13–16].

To detect at LHC, we need to keep in mind some basic facts about the monopole [6]. First, the magnetic charge of the electroweak monopole is not $2\pi/e$ but $4\pi/e$, twice that of the Dirac monopole. This is because the period of the electromagnetic U(1) subgroup of the standard model becomes 4π , not 2π . This comes from the fact the electromagnetic U(1) comes (partly) from the U(1) subgroup of SU(2). Second, the mass of the monopole is estimated to be of the order of several TeV, or roughly M_W/α . This is because the mass basically comes from the same Higgs mechanism which makes the W boson massive, except that here the coupling is magnetic (i.e., $4\pi/e$). This makes the monopole mass $1/\alpha$ times heavier than the W boson mass, of the order of 11 TeV. Third, in spite of this, the size of the monopole is set by the W boson mass. This is because the monopole solution has the weak boson dressing which shows that the size is fixed by the W boson masses. Finally, it exists within (not beyond) the standard model as the electroweak generalization of the Dirac monopole, which can be viewed as a hybrid between Dirac and 't Hooft-Polyakov monopoles.

Because of these unique characteristics of the electroweak monopole MoEDAL could detect the monopole without much difficulty, if LHC could produce it. However, the 14 TeV LHC may have no chance to produce the monopole if the mass becomes larger than 7 TeV [12].

* ymcho0416@gmail.com

† sangwoo7616@gmail.com

‡ shoh.physics@gmail.com

In this case we must try to detect the remnant monopoles at present universe produced in the early universe.

To detect the remnant monopoles, we need to know how the monopoles are produced in the early universe and how many of them are left over in the present universe. In the literature there have been discussions on the cosmological production of monopoles, but most of them have been on the grand unification monopole [17–20]. The general consensus is that the grand unification monopoles would have overclosed the universe without the inflation, but the inflation might have completely diluted them in such a way that they could have no visible impact on the present universe [21].

For the electroweak monopole, we have a totally different situation. A recent study showed that the electroweak monopole amply produced during the electroweak phase transition does not alter the standard cosmology in any significant way, but could play important roles in cosmology [22]. They could become the seed of the stellar objects and galaxies and thus play an important role in the formation of the large scale structure of the universe, and could even account for the dark matter of the universe. This was based on the observation that, as the only absolutely stable topological elementary particle which has a huge mass in the early universe, it could generate the density perturbation and evolve to the primordial magnetic blackholes (PMBHs). Moreover, the electroweak monopoles could generate the intergalactic magnetic field, and explain the ultra high energy cosmic rays [23].

Actually, the electroweak monopoles can turn to the primordial magnetic blackholes without any density perturbation. It has been well known that the monopoles, when coupled to gravity, automatically become the Reissner-Nordstrom (RN) type magnetic blackholes [24–28]. This means that the electroweak monopoles produced in the early universe automatically become the PMBHs. This makes the above proposal more credible. On the other hand, this necessitates us to re-analyse the electroweak monopole production in the early universe more carefully.

The purpose of this paper is two-fold. First, to study the electroweak monopole production in the early universe in more detail and discuss how the electroweak monopoles turn to the PMBHs when coupled to gravity. Second, to discuss physical implications of the PMBHs in cosmology. We show that the electroweak PMBHs enhance the possibility that they become the seed of the stellar objects and galaxies and thus play important roles in the formation of the large scale structure of the universe. Moreover, they enhance the possibility for the magnetic blackholes to become the dark matter of the universe greatly.

In doing so we also discuss serious shortcomings of the popular monopole production mechanism in the early universe. It has been generally believed that the monopole production in the early universe takes place

during the phase transition, and that the monopole production mechanism critically depends on the type of the phase transition. In the first order phase transition the monopole production is thought to take place by the vacuum bubble collision, which makes the production probability exponentially decreasing. But in the second order phase transition the monopoles are supposed to be produced by the Kibble-Zurek mechanism without the exponential suppression. This view, however, neglects the fact that the monopoles are produced by the change of topology made by the thermal fluctuation of the Higgs vacuum, which starts at the phase transition but continues till the universe cools down to the Ginzburg temperature even after the phase transition.

This means that the monopoles are produced not just during the phase transition but for quite a long time after the phase transition, as long as the Ginzburg temperature is considerably lower than the critical temperature. This necessitates a serious modification of the monopole production mechanism in the early universe. In fact this tells that what is important for the monopole production in the early universe is not the type of the phase transition but the Ginzburg temperature.

Moreover, it has generally been believed that in the second order phase transition the initial monopole density is determined by the correlation length set by the Higgs mass. We find that this view also has a problem. This is because the monopole mass is roughly $1/\alpha$ times bigger than the W boson mass, while the monopole size is fixed by the W boson mass. This implies that the correct correlation length for the monopole production should be more likely $1/\alpha^{1/3}$ times the correlation length set by the Higgs mass. This could reduce the initial monopole density by the factor α .

But a most important shortcoming is that the monopole production in the early universe has completely neglected the fact that, when coupled to gravity, the monopole automatically transforms to magnetically charged RN type blackhole. This means that the monopoles produced in the early universe are destined to play important roles as PMBHs, so that we can not separate the monopoles and PMBHs in the early universe when we have the monopoles. And this point has not been properly appreciated so far.

To understand this, remember that the monopole mass is expected to be of M_W/α , or of 11 TeV. But after it turns to the PMBH the mass can be anywhere between 10^{20} GeV for the extremal blackholes and the infinity in principle for non-extremal ones. This means that the monopole mass changes at least by the factor 10^{16} when they become the blackholes. Moreover, the PMBHs and monopoles have totally different cosmic evolution. Most of the monopoles and antimonopoles produced around $T_i \simeq 102$ GeV are immediately annihilated, and this annihilation continues long time till the temperature reaches $T_f \simeq 50$ MeV.

But for the PMBHs this annihilation do not take place, so that they evolve adiabatically with the Hubble expansion. Moreover, they can not evaporate completely, because they carry the conserved magnetic charge. This means that they have a totally different impacts in cosmology. In particular, the PMBHs increase the possibility for the monopole to become the seed of the stellar objects and galaxies, and the dark matter of the universe greatly.

It must be emphasized that this PMBH is different from the well known primordial blackhole (PBH) proposed by Zeldovich and Novikov, which is supposed to be produced by the statistical density perturbation in the very early universe close to the Planck time [29–31]. Recently the PBH has become a fascinating subject in cosmology because it could account for the dark matter of the universe [32–34]. Moreover, it has been asserted that some of these PBHs could also carry color charge [35]. But it is not clear that the PBH proposed by Zeldovich and Novikov could account for the dark matter [36].

In contrast, our PMBH does not come from the density perturbation, but comes from the gravitational interaction of the electroweak monopole in the standard model during the electroweak phase transition. So the production mechanism and the production time are totally different. And we can not avoid this as far as the standard model is correct. More importantly, our PMBH could account for the dark matter as we will see in the following. And some of our PMBHs could also carry color charge, because they were formed before the QCD color confinement set in.

The paper is organized as follows. In Section II we briefly review the electroweak phase transition which produces the monopole. In Section III we review the cosmic production of the electroweak monopole. In Section IV we discuss the evolution of the electroweak monopoles and the remnant monopole density. In section V we discuss how the Cho-Maison monopole turns to the PMBH when coupled to gravity in the early universe. In Section VI we discuss cosmic production and evolution of the PMBHs. In Section VII we discuss the density of the remnant electroweak PMBHs at present universe and study the plausibility that they become the seed of the stellar objects and galaxies, and account for the dark matter. Finally in Section VIII we discuss the physical implications of the electroweak PMBH.

II. ELECTROWEAK PHASE TRANSITION AND GINZBURG TEMPERATURE: A REVIEW

To discuss the monopole production in the electroweak phase transition, we start from the temperature dependent effective action of the standard model [22, 37–




FIG. 1. The effective potential (1) at T_1 , T_c , T_2 . Here the unit of V_{eff} is chosen to be $V_0 = (\lambda/8)\rho_0^4 = 1$.

39]

$$\begin{aligned}
 V_{eff}(\rho) &= V_0(\rho) - \frac{C_1}{12\pi}\rho^3 T + \frac{C_2}{2}\rho^2 T^2 - \frac{\pi^2}{90}g_*(T)T^4, \\
 V_0(\rho) &= \frac{\lambda}{8}(\rho^2 - \rho_0^2)^2, \\
 C_1 &= \frac{6M_W^3 + 3M_Z^3}{\rho_0^3} \simeq 0.36, \\
 C_2 &= \frac{4M_W^2 + 2M_Z^2 + M_H^2 + 4m_t^2}{8\rho_0^2} \simeq 0.36, \\
 g_*(T) &= \Sigma_B g_B(T) + \Sigma_F \frac{7}{8}g_F(T), \quad (1)
 \end{aligned}$$

where V_0 (with $\lambda \simeq 0.26$ and $\rho_0 \simeq 254.6$ GeV) is the zero-temperature potential, $g_*(T)$ is the total number of distinct helicity states of the particles with mass smaller than T , C_1 and C_2 are the contributions from the ρ dependent part of the masses of gauge bosons, Higgs field, and fermions, M_W , M_Z , M_H , and m_t are the W-boson, Z-boson, Higgs boson, and the top quark masses.

It has three local extrema at

$$\begin{aligned}
 \rho_s &= 0, \\
 \rho_{\pm}(T) &= \left[\frac{C_1}{4\pi\lambda} \pm \sqrt{\left(\frac{C_1}{4\pi\lambda}\right)^2 + \left(\frac{\rho_0}{T}\right)^2 - \frac{2C_2}{\lambda}} \right] T. \quad (2)
 \end{aligned}$$

The $\rho_s = 0$ represents the Higgs vacuum of the symmetric (unbroken) phase, the $\rho_-(T)$ represents the local maximum, and the $\rho_+(T)$ represents the local minimum Higgs vacuum of the broken phase. But the two extrema ρ_{\pm} appear only when T becomes smaller than T_1

$$T_1 = \frac{4\pi\lambda}{\sqrt{32\pi^2\lambda C_2 - C_1^2}} \rho_0 \simeq 146.13 \text{ GeV}. \quad (3)$$

So above this temperature only $\rho_s = 0$ becomes the true vacuum of the effective potential, and the electroweak symmetry remains unbroken.

At $T = T_1$ we have

$$\rho_- = \rho_+ = (C_1/4\pi\lambda) T_1 \simeq 16.11 \text{ GeV}, \quad (4)$$




FIG. 2. The macroscopic view of the effective potential (1). The effective potentials at T_1 , T_c , T_2 are shown in dotted lines, and the effective potentials at the Ginzburg temperature T_G and $T = 0$ are shown in red and black curves. Notice that V_{eff} is almost indistinguishable at T_1 , T_c , and T_2 .

but as temperature cools down below T_1 we have two local minima at ρ_s and ρ_+ with $V_{eff}(0) < V_{eff}(\rho_+)$, until T reaches the critical temperature T_c where $V_{eff}(0)$ becomes equal to $V_{eff}(\rho_+)$,

$$T_c = \sqrt{\frac{18}{36\pi^2\lambda C_2 - C_1^2}} \pi\lambda\rho_0 \simeq 146.09 \text{ GeV},$$

$$\rho_+(T_c) = \frac{C_1}{3\pi\lambda} T_c \simeq 21.47 \text{ GeV}. \quad (5)$$

So $\rho_s = 0$ remains the minimum of the effective potential for $T > T_c$.

Below this critical temperature ρ_+ becomes the true minimum of the effective potential, but $\rho_s = 0$ remains a local (unstable) minimum till the temperature reaches T_2 . At $T = T_c$ the new vacuum bubbles start to nucleate at $\rho = \rho_+$, which takes over the unstable vacuum $\rho_s = 0$ completely at $T = T_2$,

$$T_2 = \sqrt{\frac{\lambda}{2C_2}} \rho_0 \simeq 145.82 \text{ GeV},$$

$$\rho_+(T_2) = \frac{C_1}{2\pi\lambda} T_2 \simeq 32.15 \text{ GeV}. \quad (6)$$

From this point ρ_+ becomes the only (true) minimum, which approaches to the well-known Higgs vacuum ρ_0 at zero temperature. This tells that the electroweak phase transition is of the first order.

However, the energy barrier is extremely small,

$$\left. \frac{V_{eff}(\rho_-) - V_{eff}(\rho_+)}{V_{eff}(\rho_+)} \right|_{T_c} \simeq 3.4 \times 10^{-6}. \quad (7)$$

Moreover, the barrier lasts only for short period since the temperature difference between T_1 and T_c is very small, $\delta = (T_1 - T_c)/T_c \simeq 0.0003$. So for all practical purpose we could neglect this barrier and treat the electroweak phase transition as a second order phase transition. The microscopic view of the effective potential (1) near T_c is




FIG. 3. The determination of the Ginzburg temperature in the electroweak phase transition. Here the red and blue curve represents $\xi^3\Delta F$ and the black line represents T .

shown in Fig. 1, and the macroscopic view of the effective potential at different temperatures is shown in Fig. 2 in comparison.

It has generally been believed that the monopole production in the early universe critically depends on the type of the phase transition. In the first order phase transition, the vacuum bubble collisions in the unstable vacuum are supposed to create the monopoles through the quantum tunneling to the stable vacuum during the phase transition, so that the monopole production is exponentially suppressed by the vacuum tunneling [18]. This, of course, is totally different from the monopole production mechanism in the second order phase transition, where the monopoles are created through the Kibble-Zurek mechanism without any exponential suppression [17, 20].

But as we have emphasized, the monopole production should involve a change of topology which takes place through the thermal fluctuation of the Higgs vacuum from non-vanishing to vanishing values. And this thermal fluctuation continues as far as [40]

$$\xi^3\Delta F \leq T, \quad \Delta F(T) = V(\rho_s) - V(\rho_+), \quad (8)$$

where $\xi(T)$ is the correlation length of the Higgs field given by $\xi = 1/M_H$ and $\Delta F(T)$ is the difference in free energy density between two phases. And the temperature at which this fluctuation stops is given by the Ginzburg temperature T_G , which is defined by the condition $\xi^3\Delta F = T$ [22].

In the electroweak phase transition we can find the Ginzburg temperature graphically from (1) and (8). This is shown in Fig. 3. From this we have [22]

$$T_G \simeq 57.6 \text{ GeV}, \quad \rho_+(T_G) \simeq 232.4 \text{ GeV}. \quad (9)$$

So we can say that the monopole formation takes place between $T_2 \simeq 153.0 \text{ GeV}$ and $T_G \simeq 57.6 \text{ GeV}$, or roughly




FIG. 4. The temperature dependent Higgs and W-boson masses. The blue and red curves represent the Higgs and W-boson masses.

around T_i ,

$$T_i = \frac{T_c + T_G}{2} \simeq 102.09 \text{ GeV},$$

$$\rho_+(T_i) \simeq 187.83 \text{ GeV}. \quad (10)$$

The effective potential (1) at the Ginzburg temperature is shown in Fig. 2 in red curve.

This observation tells that the popular monopole production mechanism in the early universe has a potentially serious problem. Consider the first order phase transition. When T_G becomes lower than T_2 , we have the monopole production even after T_2 without any exponential suppression. In this case the monopole production in the first order phase transition becomes qualitatively the same as in the second order phase transition.

This tells that the exponential suppression of the monopole production in the first order phase transition is only half of the full story which could be totally misleading. *In particular, this shows that what is important in the monopole production in the early universe is not the type of the phase transition, but the Ginzburg temperature.* As far as the Ginzburg temperature becomes lower than T_2 , the monopole production in the first and second order phase transitions is qualitatively the same.

The effective potential (1) gives us two important parameters of the electroweak phase transition, the temperature dependent Higgs mass \bar{M}_H which determines the correlation length $\xi = 1/\bar{M}_H$

$$\bar{M}_H^2 = \left. \frac{d^2 V_{eff}}{d\rho^2} \right|_{\rho_{min}}$$

$$= \begin{cases} \frac{[(T/T_2)^2 - 1]}{2} M_H^2, & T > T_c, \\ \frac{[(\rho_+/\rho_0)^2 + 1 - (T/T_2)^2]}{2} M_H^2, & T \leq T_c, \end{cases} \quad (11)$$

and the W boson mass \bar{M}_W which determines the

	T	$\rho_+(T)$	$\bar{M}_H(T)$	$\bar{M}_W(T)$	$\bar{M}_m(T)$
T_1	146.74	16.3	5.9	0	0
T_c	146.70	21.7	5.5	7.1	971.7
T_2	146.42	32.5	11.7	10.6	1454.8
T_i	102.09	188.4	92.9	61.5	8428.2
T_G	57.49	232.9	116.9	76.0	10419.6
0	0.00	246.2	125.3	80.4	11014.5

TABLE I. The values of ρ_+ , \bar{M}_H , \bar{M}_W , and the expected monopole mass $\bar{M}_m = \bar{M}_W/\alpha$ at various temperatures. All numbers are in GeV.

monopole mass $M_m \simeq \bar{M}_W/\alpha$,

$$\bar{M}_W^2 = \begin{cases} 0, & T > T_c, \\ \frac{g^2}{4} \rho_+^2, & T \leq T_c. \end{cases} \quad (12)$$

The temperature dependent Higgs and W-boson masses are shown in Fig. 4.

Notice that \bar{M}_H acquires its minimum value 5.5 GeV at $T = T_c$ and approaches to the zero temperature value 125.2 GeV as the universe cools down. Moreover, we have $\bar{M}_H(T_G) \simeq 121.7$ GeV. On the other hand, the W boson starts massless at high temperature (before the symmetry breaking), but becomes massive toward the value 7.1 GeV at T_c . Moreover, we have $\bar{M}_W(T_G) \simeq 76.0$ GeV. This implies that the infant monopole masses at T_c and T_i are around 0.97 TeV and 8.4 TeV, assuming $\bar{M}_m \simeq \bar{M}_W/\alpha$. Notice that \bar{M}_m at T_G already becomes almost the adolescent value. In Table I we show the masses of the Higgs, W boson and monopole at various temperatures.

We can translate the monopole production in time scale, converting temperature to time. In the radiation dominant era the age of the universe t is given by [41]

$$t = \left(\frac{90}{32\pi G g_*(T)} \right)^{1/2} \frac{1}{T^2}. \quad (13)$$

From this we have, (with $g_* \simeq 106.75$ including $\gamma, \nu, g, e, \mu, \pi, u, d, s, c, b, \tau, W, Z, H$) [41],

$$t = 0.09 \frac{M_P}{T^2} \simeq 7 \times 10^{-7} \left(\frac{\text{GeV}}{T} \right)^2 \text{ sec}. \quad (14)$$

So we can say that the electroweak monopole production starts from 3.3×10^{-11} sec to 2.1×10^{-10} sec after the big bang for the period of 17.7×10^{-11} sec, or around 6.7×10^{-11} sec after the big bang in average.

We can also estimate how many times the thermal fluctuations take place between T_c and T_G . From the uncertainty principle and (10) we can estimate the time Δt for one fluctuation,

$$\Delta t \simeq \frac{1}{\Delta E} \simeq 3.34 \times 10^{-27} \text{ sec}. \quad (15)$$

From this we have

$$N_f \simeq \frac{\bar{t}}{\Delta t} \simeq 3.1 \times 10^{16}. \quad (16)$$

This assures that we have (more than) enough fluctuations of the Higgs vacuum to produce the monopoles.

III. COSMIC PRODUCTION OF ELECTROWEAK MONOPOLE: INITIAL DENSITY

The electroweak monopole production in the early universe and its cosmological implications have been discussed before, and it has been argued that the electroweak monopole could have deep impact on cosmology, without altering the standard cosmology in any significant way. In particular, it has been proposed that the electroweak monopoles could generate the density perturbation and evolve to primordial blackholes, and could become the seed of the stellar objects and galaxies, account for the dark matter of the universe, could generate the intergalactic magnetic field [22].

In this section we review the electroweak monopole production in the early universe. According to the Kibble's original estimate the initial electroweak monopole density is given by [17]

$$n_i \simeq \frac{g_P}{d_H^3(T_c)}, \quad (17)$$

where $d_H(T_c)$ is the horizon distance at T_c and g_P is the probability that the monopole topology is actually realized in one correlation volume, which has generally been assumed to be around 10 % in the literature [37, 38]. Since $d_H = 2t$ in the radiation dominant era we have

$$d_H(T_c) = 2t(T_c) \simeq \frac{2.6 \times 10^{15}}{T_c} \simeq 0.35 \text{ cm},$$

$$n_i \simeq \left(\frac{T_c}{2.6 \times 10^{15}} \right)^3 \times g_p \simeq 23 \times g_p \text{ cm}^{-3}. \quad (18)$$

As we have emphasized, however, this estimate (based on the assumption that the monopole is produced at T_c) is not realistic. Since the monopoles are produced not just during but even after the phase transition till the universe cools down to the Ginzburg temperature, we have to replace d_H in (17) by the mean value of two correlation lengths at T_c and T_G [22],

$$\xi_i = \frac{1}{\bar{M}_H(T_i)} \simeq 2.1 \times 10^{-16} \text{ cm}. \quad (19)$$

Notice that this is near the electroweak scale ($1/M_W \simeq 2.5 \times 10^{-16} \text{ cm}$). From this we can estimate the initial monopole density,

$$n_i \simeq \frac{g_P}{\xi_i^3} = g_p \times \bar{M}_H(T_i)^3$$

$$\simeq 1.04 \times 10^{47} \times g_p \text{ cm}^{-3}. \quad (20)$$

This should be compared with (18). Obviously this huge difference comes from the difference between d_H and ξ_i .

However, this estimate may also have a defect. Since the monopole size is fixed by the electroweak scale while the mass becomes $1/\alpha$ times bigger than the W boson mass, there appears the possibility that the energy within one correlation volume may not become enough to create one monopole. So we have to make sure that the radiation energy in one correlation volume is no less than the monopole mass, and should require

$$E = \rho(T_i) \times \xi_i^3 \geq \frac{\bar{M}_W(T_i)}{\alpha} \simeq 8.4 \text{ TeV}, \quad (21)$$

where $\rho(T_i)$ is the energy density of the universe at T_i . Now, with

$$\rho(T_i) = \frac{\pi^2}{30} g_* T_i^4 \simeq 34.8 T_i^4, \quad (22)$$

we can calculate the energy within one correlation volume at T_i ,

$$E = \rho(T_i) \times \xi_i^3 \simeq \frac{\pi^2}{30} g_* \left(\frac{T_i}{\bar{M}_H(T_i)} \right)^3 T_i$$

$$\simeq 4.73 \text{ TeV}. \quad (23)$$

This is problematic, because this tells that the energy condition (21) is not satisfied.

A logical way to cure this problem is to make the correlation length ξ by $(1/\alpha)^{1/3}$ times bigger. So we introduce a new correlation length $\bar{\xi}$ by

$$\bar{\xi}_i = \left(\frac{1}{\alpha} \right)^{1/3} \times \frac{1}{\bar{M}_H(T_i)} \simeq 1.08 \times 10^{-15} \text{ cm}. \quad (24)$$

With this we have the new initial monopole density

$$n_i \simeq \frac{g_P}{\bar{\xi}_i^3} = g_p \times \alpha \bar{M}_H(T_i)^3$$

$$\simeq 7.6 \times 10^{44} \times g_p \text{ cm}^{-3}. \quad (25)$$

This should be compared with (20).

IV. EVOLUTION OF ELECTROWEAK MONOPOLE AND REMNANT MONOPOLE DENSITY

The initial monopole density changes as the universe evolves. There are two processes which changes the initial monopole density, the Hubble expansion and the annihilation of monopole-antimonopole pairs, and the evolution of the monopole density n is governed by the Boltzmann equation [17, 18]

$$\frac{dn}{dt} + 3Hn = -\sigma n^2. \quad (26)$$




FIG. 5. The relevant scales, ξ in purple, l_m in blue, and r_{capt} in red, against T . They are normalized by the correlation length ξ_i at T_i . Here we set $M_m(T_i) = 8.4$ TeV.

where H and σ are the Hubble expansion parameter and the monopole-antimonopole annihilation cross section.

The annihilation of the monopoles is controlled by two things, the thermal Brownian motion (random walk) of the monopole in hot thermal bath and the magnetic attraction between monopole and anti-monopole. After the creation the monopoles diffuse in hot thermal plasma by the Brownian motion with the mean free path l_m given by

$$l_m = v_t t_m \simeq \frac{1}{BT} \sqrt{\frac{M_m}{T}}, \quad B = \frac{1}{T} \sum_i k_i \sigma_i, \quad (27)$$

where $v_t \simeq \sqrt{T/M_m}$ and t_m are the thermal velocity and the mean free time of the monopoles, k_i and σ_i are the number density and the cross section of the i -th relativistic charged particles and the sum is the sum over all spin states [18]. With

$$k_i \simeq \frac{3\zeta(3)}{4\pi^2} T^3, \quad \sigma_i \simeq \left(\frac{q_i}{e}\right)^2 \frac{1}{T^2}, \quad (28)$$

we have

$$B \simeq \frac{3\zeta(3)}{4\pi^2} \sum_i \left(\frac{q_i}{e}\right)^2 \simeq 0.09 \times \sum_i \left(\frac{q_i}{e}\right)^2, \quad (29)$$

where q_i is the electric charge of the i -th particle and ζ (with $\zeta(3) = 1.202\dots$) is the Riemann zeta function. Since the charged particles in the plasma are the leptons and quarks, we may put $B \simeq 3$. From this we have $v_t \simeq 0.1$ and $l_m \simeq 5.8 \times 10^{-16}$ cm around T_i .

Against the thermal random walk of the monopoles, the Coulombic attraction between monopole and anti-monopole makes them drift towards each other. The drift velocity v_d of the monopole at a distance r from the anti-monopole in the non-relativistic approximation is given by [42]

$$v_d \simeq \frac{1}{\alpha} \times \frac{1}{BT^2 r^2}. \quad (30)$$

Notice that we have $v_d \simeq 7.4$ with $r = \bar{\xi}_i$. This is unrealistic, which is due to the non-relativistic approximation. But this does tell that $v_t \ll v_d$, which shows that the Coulombic magnetic attraction is much stronger than the thermal diffusion. In Fig. 5 we plot the relevant scales ξ , l_m , and r_{capt} , against T for comparison. This clearly shows that the capture radius is much bigger than the mean free length and correlation length in a wide range of T . This tells that (25) is an overestimation.

Now, assuming the mean distance r between the monopole and anti-monopole is $r \simeq n^{-1/3}$, we can express the capture time by

$$t_{\text{capt}} \simeq \frac{r}{v_d} \simeq \alpha \times \frac{BT^2}{n}. \quad (31)$$

From this we have the monopole-antimonopole annihilation cross section

$$\sigma \simeq \frac{1}{t_{\text{capt}} n} = \frac{1}{\alpha BT^2}. \quad (32)$$

So, with

$$H = \frac{\dot{R}}{R} = \frac{T^2}{CM_P}, \quad C = \left(\frac{45}{4\pi^3 g_*}\right)^{1/2} \simeq 3.74, \quad (33)$$

we can express the Boltzmann equation in term of $\tau = M_m/T$

$$\frac{d}{d\tau} \left(\frac{n_m}{T^3}\right) = -\frac{CM_P}{\alpha BM_m} \left(\frac{n_m}{T^3}\right)^2 = -\frac{\sigma T^3}{\tau H} \left(\frac{n_m}{T^3}\right)^2. \quad (34)$$

Solving this we have [22]

$$n = \frac{T^3}{A(M_m/T - M_m/T_i) + B}, \quad A = \frac{CM_P}{\alpha BM_m}, \quad B = \frac{T_i^3}{n_i}, \quad (35)$$

where M_m is treated as a constant. But in reality it depends on time, so that it should be understood as a mean value.

Notice that when $\tau_i \ll \tau$, only the first term in the denominator becomes important. So the monopole density approaches to

$$n \rightarrow \alpha \times \frac{BT^4}{CM_P}, \quad (36)$$

regardless of the initial condition [18]. The diffusive capture process is effective only when $l_m < r_{\text{capt}}$, which determines the temperature T_f below which the monopole-antimonopole annihilation ceases,

$$T_f \simeq \alpha^2 \times \frac{M_m}{B^2} \simeq 49.7 \text{ MeV}. \quad (37)$$

This is below the muon decoupling temperature, which tells that the annihilation continues very long time. The




FIG. 6. The evolution of the monopole density n against $\tau = M_m/T$ with $M_m = 8.4$ TeV.

evolution of the monopole density n against M_m/T is shown in Fig. 6, where we have put $M_m = M_m(T_i) \simeq 8.4$ TeV.

From this we can estimate the remnant monopole density in the present universe. The monopole density after the annihilation around T_f becomes (with $M_m \simeq 11$ TeV)

$$n_f \simeq \alpha^3 \times \frac{M_m}{BCM_P} T_f^3 \simeq 4.95 \times 10^{14} \text{ cm}^{-3}. \quad (38)$$

where now we have put $M_m = M_m(0) \simeq 11$ TeV since the temperature has cooled down very much. Obviously this is much lower than the initial monopole density given by (25). The number of monopole within the co-moving volume is conserved thereafter. But they still interact with the electron pairs in the hot plasma before the decoupling around $T_d \simeq 0.5$ MeV, when the electron pairs disappear and the interaction rate becomes less than the Hubble expansion rate.

Assuming that the expansion is adiabatic, the current number density and the energy density of the monopole is given by

$$n_0 = \frac{g_{s,0}}{g_{s,f}} \left(\frac{T_0}{T_f}\right)^3 n_f \simeq \alpha^3 \frac{g_{s,0}}{g_{s,f}} \times \frac{M_m}{BCM_P} T_0^3, \quad (39)$$

where g_s is the effective number of degrees of freedom in entropy and $T_0 = 2.73$ K = 2.35×10^{-13} GeV is the temperature of the universe today. With $g_{s,f} \simeq 10.75$ and $g_{s,0} \simeq 3.9$, we have

$$n_0 \simeq 1.1 \times 10^{-23} T_0^3 \simeq 1.86 \times 10^{-20} \text{ cm}^{-3}. \quad (40)$$

This means that there are roughly 6.6×10^{66} monopoles in the observable universe (with the radius 4.4×10^{28} cm), or roughly 2.04×10^7 monopoles per volume of the earth in the present universe. This implies that there are enough electroweak monopoles left over in the present universe that we could detect.

With (40) the density parameter of monopole at

present universe can be written as

$$\begin{aligned} \rho_{m,0} &= n_0 M_m \simeq 2.04 \times 10^{-7} \text{ eV cm}^{-3}, \\ \Omega_m h^2 &= \frac{\rho_{m,0}}{\rho_{c,0}} h^2 \simeq 4.3 \times 10^{-8}, \end{aligned} \quad (41)$$

where $\rho_{c,0} = 3H^2/8\pi G \simeq 1.05 h^2 \times 10^{-5} \text{ GeV cm}^{-3} \simeq 0.48 \times 10^{-5} \text{ GeV cm}^{-3}$ is the critical density of present universe and $h \simeq 0.678$ is the scaled Hubble constant in the unit $H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$. This is about 0.38×10^{-13} of the baryon number density $n_b \simeq 2.5 \times 10^{-7} \text{ cm}^{-3}$, which assures that the electroweak monopole cannot be a dark matter candidate.

Actually, however, the free streaming remnant monopole density could be much less than the above estimate [22]. There are good reasons for this. First, most of them could turn to the PMBHs and become the seed of the stellar objects and galaxies. So many of them might have been buried in stellar objects and galactic centers. The recent observations that there is a strong radial magnetic field near the galactic center could be the experimental evidences of this [43].

Second, the monopoles have a very short penetration length in the matter because they have strong magnetic interaction. In fact a relativistic electroweak monopole can travel only a few meters (less than 10 m) in Aluminium before they are trapped [44]. This means that most of the monopoles left over which did not become the PMBHs could have been trapped and filtered out by the stellar objects, when they collide with them. This strongly implies that the actual free streaming remnant monopole density could be much less than (41). In fact, the Parker bound on the free streaming monopole density implies that the remnant monopole density could be 10^{-4} times less than (41) [22, 45]. Unfortunately it is difficult to estimate how much of them are free streaming at present universe.

Notice that, since the decoupling temperature of the electroweak monopole is much less than the monopole mass, the free streaming monopoles just after the decoupling start as completely non-relativistic. But eventually they are accelerated by the intergalactic magnetic field and become extremely relativistic. So they can acquire the energy which exceeds the Greisin-Zatsepin-Kuzmin (GZK) energy limit, and thus could become identified as the ultra high energy cosmic rays [23, 46, 47].

We can estimate the number of the monopoles which arrive on earth per year. Suppose the density of the free streaming monopoles at present universe per km^3 is \bar{n}_0 . Let the radius of the earth be r_0 and consider a point located at the distance r ($r > r_0$) from the center of the earth. Since the solid angle of the earth viewed from this point is given by

$$\Omega(r) = 2\pi(r^2 - r_0^2) \left(1 - \frac{\sqrt{r^2 - r_0^2}}{r}\right), \quad (42)$$

the number n_{ES} of the monopoles coming from the sphere of radius r_1 ($r_1 > r_0$) from the center of the earth to the earth surface is given by

$$\begin{aligned} n_{ES} &= \bar{n}_0 \int_{r_0}^{r_1} r^2 \frac{\Omega(r)}{4\pi(r^2 - r_0^2)} \sin\theta dr d\theta d\varphi \\ &= \frac{2\pi\bar{n}_0}{3} \left[r_1^3 - r_0^3 - (r_1^2 - r_0^2)^{3/2} \right] \\ &= \frac{2}{3} \left[L^3 + 3L^2 r_0 \left(1 + \frac{r_0}{L}\right) - L^3 \left(1 + \frac{2r_0}{L}\right)^{3/2} \right] \\ &\simeq \pi\bar{n}_0 L r_0^2, \quad (r_0 \ll L), \end{aligned} \quad (43)$$

where we have put $L = r_1 - r_0$. This means that the number of the monopoles arriving on earth per year (with 1 year $\simeq 3.1558 \times 10^7$ sec and $r_0 \simeq 6400$ km) per km^2 is given by

$$\begin{aligned} n_{ES}(1 \text{ year}/km^2) &\simeq \bar{n}_0 \times \frac{3.1558 \times 3}{4 \times (6.4)^2} \times 10^8 \\ &\simeq 5.78 \bar{n}_0 \times 10^6. \end{aligned} \quad (44)$$

So, with (40) we have $n_{ES}/\text{year } km^2 \simeq 10.75$. But we have to keep in mind that this estimate could be lower than this, since the density of the free streaming monopoles at present universe could be much lower than (40). In this case the above estimate becomes consistent with the ultra high energy cosmic ray detection rate [23, 46]

Notice that the monopoles could create a huge density perturbation in the matter dominant era and thus become PMBHs [22]. Indeed, with the monopole energy density ρ_m and the energy density of the universe at the radiation-matter equality time ρ_e given by

$$\begin{aligned} \rho_m &\simeq 11 \text{ TeV} \times \frac{3}{4\pi} M_W^3 \simeq 1.78 \times 10^{50} \text{ GeV } cm^{-3}, \\ \rho_e &\simeq 5.9 \times 10^6 \text{ GeV } cm^{-3}, \end{aligned} \quad (45)$$

we have the monopole density perturbation

$$\frac{\delta\rho}{\rho} = \frac{\rho_m - \rho_e}{\rho_e} \simeq 0.3 \times 10^{44}. \quad (46)$$

This strongly implies that the monopoles could turn to PMBHs [22].

V. CHO-MAISON MONOPOLE VERSUS PRIMORDIAL REISSNER-NORDSTROM MAGNETIC BLACKHOLE

Actually, for the Cho-Maison monopole to become a magnetic blackhole, we do not need the density perturbation. When coupled to gravity, it automatically turns to a magnetic blackhole [24–28]. To show this, we start from

(the bosonic sector of) the Einstein-Weinberg-Salam Lagrangian

$$\begin{aligned} \mathcal{L}_{EWS} &= \frac{\sqrt{-g}}{16\pi G} \left\{ R - |\mathcal{D}_\mu\phi|^2 - \frac{\lambda}{2} \left(\phi^\dagger\phi - \frac{\mu^2}{\lambda} \right)^2 \right. \\ &\quad \left. - \frac{1}{4} \vec{F}_{\mu\nu}^2 - \frac{1}{4} G_{\mu\nu}^2 \right\}, \\ \mathcal{D}_\mu\phi &= (\partial_\mu - i\frac{g}{2}\vec{\tau} \cdot \vec{A}_\mu - i\frac{g'}{2}B_\mu)\phi, \end{aligned} \quad (47)$$

where R is the scalar curvature, ϕ is the Higgs doublet, $\vec{F}_{\mu\nu}$ and $G_{\mu\nu}$ are the gauge field strengths of $SU(2)$ and $U(1)_Y$ with the potentials \vec{A}_μ and B_μ , g and g' are the corresponding coupling constants.

To proceed we choose the following spherically symmetric monopole ansatz in the spherical coordinates (t, r, θ, φ) ,

$$\begin{aligned} \phi &= \frac{i}{\sqrt{2}}\rho(r) \begin{pmatrix} \sin(\theta/2) e^{-i\varphi} \\ -\cos(\theta/2) \end{pmatrix}, \\ \vec{A}_\mu &= \frac{1}{g}(f(r) - 1) \hat{r} \times \partial_\mu \hat{r}, \\ B_\mu &= -\frac{1}{g'}(1 - \cos\theta)\partial_\mu\varphi. \end{aligned} \quad (48)$$

In terms of the physical fields the ansatz can be written as [6, 8]

$$\begin{aligned} \rho &= \rho(r), \\ A_\mu^{(em)} &= -\frac{1}{e}(1 - \cos\theta)\partial_\mu\varphi, \\ W_\mu &= \frac{i}{g} \frac{f(r)}{\sqrt{2}} e^{i\varphi} (\partial_\mu\theta + i \sin\theta \partial_\mu\varphi), \\ Z_\mu &= 0. \end{aligned} \quad (49)$$

This clearly shows that the ansatz is for the electroweak monopole dressed by the W boson.

Now, adopting the static spherically symmetric space-time metric

$$\begin{aligned} ds^2 &= -N^2(r)A(r)dt^2 + \frac{dr^2}{A(r)} \\ &\quad + r^2(d^2\theta + \sin^2\theta d\varphi^2), \end{aligned} \quad (50)$$

we find that the Einstein-Weinberg-Salam action reduces to

$$\begin{aligned} S &= - \int N \left[\frac{r\dot{A} + A - 1}{8\pi G} + AK + U \right] dr, \\ K &= \frac{\dot{f}^2}{g^2} + \frac{r^2}{2}\dot{\rho}^2, \\ U &= \frac{(1 - f^2)^2}{2g^2r^2} + \frac{1}{4}f^2\rho^2 + \frac{\lambda}{8}r^2(\rho^2 - \rho_0^2)^2 \\ &\quad + \frac{1}{2g'^2r^2}. \end{aligned} \quad (51)$$

From this we have the following equations of motion

$$\begin{aligned}\frac{\dot{N}}{N} &= 8\pi G \frac{K}{r}, \\ \dot{A} + \frac{A-1}{r} &= -\frac{8\pi G}{r}(AK+U), \\ A\ddot{\rho} + \left(\dot{A} + \frac{\dot{N}}{N}A + \frac{2A}{r}\right)\dot{\rho} - \frac{f^2}{2r^2}\rho &= \frac{\lambda}{2}(\rho^2 - \rho_0^2)\rho, \\ A\ddot{f} + \left(\dot{A} + \frac{\dot{N}}{N}A\right)\dot{f} + \frac{1-f^2}{r^2}f &= \frac{1}{4}g^2\rho^2f.\end{aligned}\quad (52)$$

Notice that, when the gravitation is switched off (i.e., when $G = 0$ and $A = N = 1$) this describes the non-gravitating Cho-Maison monopole which has the Coulombic monopole singularity at the origin [8].

When the gravity is included the above equation has two types of solutions, the globally defined gravitating Cho-Maison monopole solution and the RN type magnetic black hole solutions [22, 24–28]. Consider the gravitating Cho-Maison monopole first. For this we may adopt the boundary condition

$$\begin{aligned}A(0) &= 1, \quad f(0) = 1, \quad \rho(0) = 0, \\ A(\infty) &= 1, \quad f(\infty) = 0, \quad \rho(\infty) = \rho_0.\end{aligned}\quad (53)$$

and obtain the following expansions for the solution near $r = 0$,

$$\begin{aligned}f(r) &= 1 - f_1x^2 + \dots, \\ \rho(r) &= h\rho_0x^\delta + \dots, \quad \delta = \frac{\sqrt{3}-1}{2}, \\ A(r) &= 1 - \frac{16\pi}{e^2}\sin^2\theta_W\left(\frac{M_W}{M_P}\right)^2\frac{\delta^2}{\sqrt{3}}h^2x^{2\delta} + \dots,\end{aligned}\quad (54)$$

where f_1 and h are constants, $x = M_W r$, and $M_P \simeq 1.22 \times 10^{19}$ GeV is the Planck mass. This tells that a gravitating Cho-Maison monopole solution which has the weak boson dressing for $0 < r < \infty$ exists. Notice, however, that the factor $(M_W/M_P)^2$ in A is extremely small, which makes the metric almost flat.

This tells that the gravitational modification is extremely small, almost non-existent. Only when the W boson mass M_W becomes unrealistically large, of the order of the Planck mass, the gravitational modification becomes explicit here. From this we can conclude that the gravitating Cho-Maison monopole is almost identical to the non-gravitating Cho-Maison monopole. In particular, the gravity has practically no influence on the mass of the monopole.

What is more relevant for us here is the second type, the RN type blackhole solutions carrying the magnetic charge $4\pi/e$ [22, 24–28]. To discuss this, we first switch off the weak bosons and let $f = 0$ and $\rho = \rho_0$. With this we can easily solve (52) and find

$$\begin{aligned}N &= 1, \\ A(r) &= 1 - \frac{2GM}{r} + \frac{4\pi G}{e^2}\frac{1}{r^2},\end{aligned}\quad (55)$$

where M is the ADM mass of the blackhole. This has the outer horizon

$$r_H = r_+ = MG + \sqrt{M^2G^2 - 4\pi G/e^2},\quad (56)$$

with

$$\begin{aligned}M &\geq M_{eRN} = \frac{M_P}{\sqrt{\alpha}} \simeq 11.7 \times M_P \\ &\simeq 1.43 \times 10^{20} \text{ GeV},\end{aligned}\quad (57)$$

where M_{eRN} is the extremal (minimal) mass of the RN blackhole. From this we have

$$M_{RN} = \frac{r_H}{2G} + \frac{2\pi}{e^2 r_H},\quad (58)$$

so that the horizon of the extremal RN blackhole is fixed by

$$r_H^{eRN} = \frac{L_P}{\sqrt{\alpha}} \simeq 1.87 \times 10^{-32} \text{ cm},\quad (59)$$

where $L_P \simeq 1.6 \times 10^{-33}$ cm is the Planck length.

This tells that the naked Cho-Maison monopole without the weak boson dressing, when coupled to gravity, becomes the RN blackhole which carries the magnetic charge $4\pi/e$, which can have any mass bounded below by the extremal mass $M_P/\sqrt{\alpha}$ [24, 25]. So the gravity sets the minimum mass of the naked Cho-Maison monopole of the order of ten times the Planck mass, making it an extremal RN blackhole.

Moreover, with non-trivial f and ρ , the above RN monopole can be generalized to the modified RN blackhole which has the weak boson hair [27, 28]. To see this we choose the boundary condition

$$\begin{aligned}A(r_H) &= 0, \quad f(r_H) = f_H, \quad \rho(r_H) = \rho_H, \\ 0 &\leq f_H \leq 1, \quad 0 \leq \rho_H \leq \rho_0, \\ A(\infty) &= 1, \quad f(\infty) = 0, \quad \rho(\infty) = \rho_0.\end{aligned}\quad (60)$$

With this we have the following equation from (52),

$$\begin{aligned}\dot{A}_H &= \frac{1}{r_H}\left(1 - 8\pi G U_H\right), \\ \dot{A}_H \dot{\rho}_H - \frac{f_H^2}{2r_H^2}\rho_H &= \frac{\lambda}{2}(\rho_H^2 - \rho_0^2)\rho_H, \\ \dot{A}_H \dot{f}_H + \frac{1-f_H^2}{r_H^2}f_H &= \frac{1}{4}g^2\rho_H^2f_H.\end{aligned}\quad (61)$$

Now, assuming $\dot{A}_H \geq 0$, we have from the first equation

$$1 \geq 8\pi G \left(\frac{\lambda}{8}\rho_0^4 r_H^2 + \frac{1}{2g^2 r_H^2}\right) > \frac{4\pi G}{g^2 r_H^2},\quad (62)$$

or

$$r_H > \frac{\sqrt{4\pi G}}{g} = \cos\theta_W \frac{L_P}{\sqrt{\alpha}} \simeq 1.37 \times 10^{-32} \text{ cm}.\quad (63)$$




FIG. 7. The modified extremal Reissner-Nordstrom type magnetic blackhole in the Einstein-Weinberg-Salam theory which has the W boson f (blue) and Higgs field ρ (red) dressing. Notice that with $N(r) \simeq 1$, $A(r)$ (green) represents the metric. The unit in the x-axis is cm .

This tells that r_H is smaller than r_H^{eRN} by the factor $\cos \theta_W$.

To proceed, we also assume that f and ρ are monotonically decreasing and increasing functions. With this we have the following constraint on f_H and ρ_H from the last two equations of (61)

$$\begin{aligned} \lambda \rho_0^2 r_H^2 \left(1 - \frac{\rho_H^2}{\rho_0^2}\right) &< f_H^2 < 1 - \frac{g^2}{4} \rho_H^2 r_H^2, \\ \left(\rho_0^2 - \frac{f_H^2}{\lambda r_H^2}\right) &< \rho_H^2 < \frac{4}{g^2 r_H^2} (1 - f_H^2) \end{aligned} \quad (64)$$

From this we have

$$\left(1 - \frac{g^2}{4\lambda}\right) f_H^2 < 1 - \frac{g^2}{4} \rho_0^2 r_H^2. \quad (65)$$

But the left-hand side is positive, since $g = e/\sin \theta_W \simeq 0.65$ and $\lambda \simeq 0.26$ in the standard model. So we have

$$r_H \leq \frac{1}{M_W} \simeq 2.5 \times 10^{-16} \text{ cm}. \quad (66)$$

This means that the maximum horizon of the modified RN blackhole is set by the size of the Cho-Maison monopole. This is very important, because this tells that the blackhole is inside of the Cho-Maison monopole. Another outcome is that for the Cho-Maison monopole we have $f(0) = 1$ and $\rho(0) = 0$, but the boundary condition $f_H = 1$ and $\rho_H = 0$ is consistent with (52) only when $r_H \simeq 0.64/M_W$. The solution of the extremal RN blackhole with the weak boson hair is shown in Fig. 7.

To find the mass of the blackhole, notice that with

(52) we have

$$\begin{aligned} A(r) &= 1 - \frac{8\pi G}{r} \left[e^{P(r)} \int_{r_H}^r (K + U) e^{-P(r')} dr' + \frac{r_H}{2G} \right], \\ P(r) &= 8\pi G \int_r^\infty \frac{K}{r'} dr', \quad (r_H \leq r), \end{aligned} \quad (67)$$

So, with (60) the ADM mass of the modified RN blackhole is given by

$$M_{mRN} = 4\pi \int_{r_H}^\infty (K + U) e^{-P(r)} dr + \frac{r_H}{2G}, \quad (68)$$

which assures the positivity of the ADM mass. With this we can estimate the mass of the blackhole. When $r_H \ll 1/M_W$, we can switch off the gravity with flat metric, and calculate the leading contribution of the integral (68), and have

$$\begin{aligned} M_{mRN} &\simeq 0.75 \times \frac{4\pi \rho_0}{g} + \frac{2\pi}{g^2 r_H} + \frac{r_H}{2G} \\ &\geq \cos \theta_W \frac{M_P}{\sqrt{\alpha}} + 1.5 \times \sin^2 \theta_W \frac{M_W}{\alpha} \\ &\simeq 10.3 M_P + 47.3 M_W \simeq (1.3 \times 10^{17} + 3.8) \text{ TeV} \\ &\simeq 2.24 \times 10^{-4} g. \end{aligned} \quad (69)$$

This tells that the minimum mass of the modified Reissner-Nordstrom blackhole is set by the Planck mass, and that the contribution of the weak boson dressing to the mass is tiny (of 10^{-17}), only 3.8 TeV. This is very interesting, because this 3.8 TeV is almost identical to the predicted Cho-Maison monopole mass 3.96 TeV based on the scaling argument (without gravity) [8, 22].

On the other hand, when $r_H \simeq 1/M_W$, we have

$$\begin{aligned} M_{mRN} &\leq M_{mRN}^{max} = \frac{M_P}{2M_W} M_P + O(M_W) \\ &\simeq 9.3 \times 10^{32} \text{ TeV} \simeq 1.67 \times 10^{12} g. \end{aligned} \quad (70)$$

which sets the maximum mass of the blackhole. So the mass of the modified RN blackhole made of the Cho-Maison monopole can be anywhere between 10 and 10^{36} times the Planck mass. It is interesting that the weak boson dressing with the gravity could set the range of the mass of the modified RN blackhole. Without the weak boson dressing the mass of the RN blackhole has only the lower bound.

VI. COSMIC PRODUCTION AND EVOLUTION OF ELECTROWEAK PRIMORDIAL MAGNETIC BLACKHOLE

Obviously the cosmic production and evolution of the electroweak monopoles do not apply to the above electroweak PMBHs. This is because the mass becomes totally different. As we have seen, the mass range of

the modified RN type PMBH could be anywhere between $M_P/\sqrt{\alpha}$ (the extremal case) and $(M_P/2M_W) M_P$ (the non-extremal case). Moreover, the mass of the RN PMBH in principle has no upper limit. Now we discuss how this could change the initial density and evolution of the electroweak PMBHs.

Consider the extremal case first, whose mass and size is set by $1/\sqrt{\alpha}$ times the Planck scale. In this case the energy condition (21) should change to

$$E = \frac{\pi^2}{30} g_* T_i^4 \times \xi_{ebh}^3 \geq \frac{M_P}{\sqrt{\alpha}}, \quad (71)$$

where ξ_{ebh} is the size (the correlation length) of the extremal RN magnetic blackhole. From this we have

$$\begin{aligned} \xi_{ebh} &\simeq \left(\frac{30}{\pi^2 g_* \sqrt{\alpha}} \frac{M_P^4}{T_i^4} \right)^{1/3} \frac{1}{M_P} \simeq \frac{0.55 \times 10^{23}}{M_P} \\ &\simeq 0.89 \times 10^{-10} \text{ cm} \end{aligned} \quad (72)$$

This is 0.8×10^5 bigger than the monopole correlation length $\bar{\xi}$ shown in (24) which determines the monopole density. Notice that for the non-extremal modified RN PMBH with mass $(M_P/2M_W) M_P$, we have $\xi_{mbh} \geq 0.86 \times 10^{-4} \text{ cm}$.

From this we have the initial number density N_i of the extremal RN PMBH

$$\begin{aligned} N_i^{ebh} &\simeq \frac{g_P}{\xi_{ebh}^3} \simeq g_P \times 6.0 \times 10^{-69} M_P^3 \\ &\simeq 1.4 \times 10^{30} \times g_p \text{ cm}^{-3}. \end{aligned} \quad (73)$$

This is smaller than the initial monopole density (25) by the factor 1.8×10^{-15} . Similarly for the non-extremal modified RN PMBH with mass $(M_P/2M_W) M_P$, we have $N_i^{mbh} \simeq 1.57 \times 10^{12} \times g_p \text{ cm}^{-3}$. This shows that there is a huge difference between the initial monopole density and magnetic blackhole density in the early universe. Again, this difference originates from the mass difference between the monopole and the magnetic blackhole.

It must be clear that these PMBHs are formed around the same time as the electroweak monopoles do, between T_c and T_G , or around $T_i \simeq 102.09 \text{ GeV}$. In time scale they are formed between $3.3 \times 10^{-11} \text{ sec}$ to $2.1 \times 10^{-10} \text{ sec}$ after the big bang for the period of $17.7 \times 10^{-11} \text{ sec}$, or around $6.7 \times 10^{-11} \text{ sec}$ after the big bang in average.

As we have pointed out, the mass of the RN PMBH theoretically has no upper limit, but obviously the number density of the PMBH depends on the mass. So one may ask what is the number density of RN PMBH with mass M . To answer this notice that in this case the energy condition (71) for the mass M RN PMBH changes to

$$E = \frac{\pi^2}{30} g_* T_i^4 \times (\xi_i^{rnbh})^3 \geq M, \quad (74)$$

so that we have

$$\xi_i^{rnbh} \simeq \left(\frac{30}{\pi^2 g_*} \frac{M^4}{T_i^4} \right)^{1/3} \frac{1}{M}. \quad (75)$$

However, since ξ_i^{rnbh} can not be bigger than the particle horizon in cosmology, we have

$$\begin{aligned} \left(\frac{30}{\pi^2 g_*} \frac{M^4}{T_i^4} \right)^{1/3} \frac{1}{M} &\leq d_H(T_i) \\ &\simeq 2 \left(\frac{90}{32\pi G g_*(T_i)} \right)^{1/2} \frac{1}{T_i}. \end{aligned} \quad (76)$$

From this we have the maximum average mass of the RN PMBH,

$$\begin{aligned} M &\leq M_{mmbh} = 0.5 \times 10^{32} M_P \\ &\simeq 1.1 \times 10^{27} g \simeq 0.55 \times 10^{-6} M_\odot, \end{aligned} \quad (77)$$

which is bigger than the maximum mass of the modified RN BH by the factor 10^{15} . In this case we have

$$r_H \simeq 1.0 \times 10^{32} L_P \simeq 0.16 \text{ cm}. \quad (78)$$

With this, the initial density of the RN PMBH of the maximum average mass M_{mmbh} at T_i is given by

$$\begin{aligned} N_i^{mmbh} &\simeq \frac{g_P}{(\xi_i^{rnbh})^3} \simeq \frac{g_P}{d_H(T_i)^3} \\ &\simeq 1.3 \times 10^{-2} \times g_p \text{ cm}^{-3}. \end{aligned} \quad (79)$$

This should be compared with (73).

It should be mentioned, however, that theoretically the maximum particle horizon for the RN PMBH mass is given (not by $d_H(T_i)$ but) by $d_H(T_G)$. This sets the upper limit of the RN PMBH mass M in cosmology

$$\begin{aligned} M &\leq \bar{M} = 1.57 \times 10^{32} M_P \\ &\simeq 3.46 \times 10^{27} g \simeq 1.73 \times 10^{-6} M_\odot. \end{aligned} \quad (80)$$

With this, the initial density \bar{N}_i of the RN PMBH of the mass \bar{M} at T_i is given by

$$\begin{aligned} \bar{N}_i &\simeq \frac{g_P}{(\xi_i^{rnbh})^3} \simeq \frac{g_P}{d_H(T_G)^3} \\ &\simeq 4.14 \times 10^{-4} \times g_p \text{ cm}^{-3}. \end{aligned} \quad (81)$$

This should be compared with (79). As we will see, this has deep implications in cosmology.

The PBH has become important in cosmology because there is a possibility that the PBH could account for the dark matter of the universe. It has been claimed that the PBH with the mass range $10^{17} g$ to $10^{22} g$ produced between 10^{-21} sec and 10^{-16} sec after the big bang (when the temperature becomes around 10^5 to 10^7 GeV) could account for all dark matter of the universe today [32–34]. Moreover, recently it has been suggested that some of the PBH could carry net color charge, when

the PBH is formed before the QCD confinement sets in around $T \simeq \Lambda_{QCD} \simeq 0.17$ GeV [35]. This has made the PBH more interesting object, and NASA is planning to hunt for such PBH with the new Roman Space Telescope [36]. So one might wonder if our PMBH could also account for the dark matter, and could carry the color flux.

We could think of two possible candidates for such PBHs in cosmology, the well known PBH proposed by Zeldovich and Novikov produced by the statistical density perturbation in the very early universe close to the Planck time [29–31], and our electroweak PMBH produced by the gravitational interaction of the electroweak monopole between 3.3×10^{-11} sec to 2.1×10^{-10} sec after the big bang. So the production mechanism and production period of our PMBH are totally different. And our PMBH carries the conserved magnetic charge, so that it can not evaporate. Moreover, with the mass range $M_P/\sqrt{\alpha}$ to $1.57 \times 10^{32} M_P$, our electroweak PMBH could become a real candidate for the dark matter of the universe. To discuss if our PMBH could really account for the dark matter, we have to discuss the cosmic evolution of the PMBH.

The evolution of the PMBH is totally different from the evolution of the electroweak monopole. Here the annihilation of two magnetic blackholes which have opposite magnetic charges does not play any important role for two reasons. First, the correlation length of the blackhole ξ_{bh} is roughly 10^5 times bigger than that of the monopole, which means that the average distance between the monopole-antimonopole blackhole pairs is much bigger than the capture radius. Second, the mass of the magnetic blackholes is at least 10^{18} times bigger than the monopole. So the magnetic attraction between the blackhole pairs is not strong enough to make the annihilation. Moreover, even if they do, the two blackholes do not disappear. They merge to a magnetically neutral blackhole. This means that they evolve adiabatically with the Hubble expansion, with $N \propto 1/R^3$.

A new aspect of the evolution of the PMBHs is the accretion and evaporation. Zeldovich and Novikov have suggested that their PBHs could have too much accretion from the nearby matters and become supermassive blackholes at present universe, which could be in conflict with the standard big bang cosmology [29–31]. So we need to check if this could also apply to our electroweak PMBH.

Assuming that the accretion is a quasi-stationary process, one could express the rate of the accretion of the primordial blackhole during the radiation dominant era by [29–31]

$$\begin{aligned} \frac{dM}{dt} &= \rho(t) \times 4\pi r_H^2 = \frac{2\pi^3}{15} g_*(t) T(t)^4 \times r_H^2 \\ &= \frac{3\pi^2}{8G} \times \frac{r_H^2}{t^2}. \end{aligned} \quad (82)$$

This, with (13), (22), and (56), is written by

$$\begin{aligned} \frac{dx}{dt} &= \frac{3\pi^2}{4M_P} \left(1 - \frac{2\pi}{e^2} \frac{1}{x^2} + \sqrt{1 - \frac{4\pi}{e^2} \frac{1}{x^2}} \right) \times \frac{x^2}{t^2} \\ &\simeq \frac{3\pi^2}{2M_P} \times \frac{x^2}{t^2}, \end{aligned} \quad (83)$$

where $x = M/M_P$. Notice that the approximation in the second line follows from the fact that for our primordial blackhole x is expected to be very large ($x^2 \geq 1/\alpha$). Integrating this from the initial time $t_i \simeq 6.7 \times 10^{-11}$ sec to the radiation-matter equilibrium time t_e given by [41]

$$t_e = 1.4 \times 10^3 (\Omega_0 h^2)^{-2} \text{ years} \simeq 2.1 \times 10^{11} \text{ sec}, \quad (84)$$

we have the mass M_e of the accreted blackhole at t_e (with $t_i \ll t_e$),

$$\begin{aligned} M_e - M_i &= \frac{3\pi^2}{2M_P^2} \times \frac{M_e M_i}{t_e t_i} \times (t_e - t_i) \\ &\simeq \frac{3\pi^2}{2M_P^2} \times \frac{M_e M_i}{t_i}. \end{aligned} \quad (85)$$

So, for the extremal RN blackhole we have

$$M_e - M_i = \frac{3\pi^2}{2\sqrt{\alpha}} \frac{M_e}{M_P} \times \frac{1}{t_i} \simeq 4.4 \times 10^{-38} M_e, \quad (86)$$

which means that the extremal blackhole has almost no accretion. Similarly for the modified RN blackhole with mass $(M_P/2M_W)M_P$ and horizon size $1/M_W$ at t_i , the accretion becomes negligible. This is basically because the size of the extremal blackhole is very small compared to the particle horizon. Indeed, for the modified RN blackhole with horizon $1/M_W \simeq 2.5 \times 10^{-16}$ cm, the particle horizon d_H at t_i is given by $d_H(t_i) \simeq 4$ cm, bigger than the blackhole horizon by the factor 10^{16} .

For the maximum average mass of RN PMBH given by (77), however, we have

$$M_e - M_i \simeq \frac{3\pi^2}{2} \times \frac{M_i M_e}{M_P^2 t_i} \simeq 0.6 M_e, \quad (87)$$

or

$$M_e \simeq 2.5 M_i \simeq 1.4 \times 10^{-6} M_\odot. \quad (88)$$

In this case the PMBH does have a significant accretion, but does not become a supermassive blackhole. From this one might conclude that the electroweak PMBHs do not grow to a supermassive blackhole.

Indeed, for (85) to be valid, we must have

$$\begin{aligned} M_i &< \frac{2M_P t_i}{3\pi^2} M_P \simeq 0.84 \times 10^{32} M_P \\ &\simeq 1.83 \times 10^{27} g. \end{aligned} \quad (89)$$

Moreover, for the accretion to be effective, M_i has to be very close to this value. Clearly this requirement is not

easy to satisfy. This implies that our PMBHs are not likely to end up with supermassive blackholes at present universe.

On the other hand it must be clear from (89) that, for our electroweak PNBH to become a supermassive blackhole, M_i must be very close to $1.83 \times 10^{27} g$. And the upper limit of the RN PMBH mass given by (80) tells that some of our RN PMBH could meet with this condition. This is remarkable, because this tells that we can not completely exclude the possibility that our PMBHs could indeed grow to become supermassive blackholes.

Now, we consider the evaporation of the PMBHs. Hawking has argued that a PBH of mass M has temperature of $10^{-6} (M_\odot/M) K$, so that blackholes with mass as large as $10^{15} g$ would have radiated away all their mass by now [30]. Fortunately this does not happen to our electroweak PMBHs, because they have the conserved magnetic charge which can not be evaporated. So the evaporation can only make them extremal.

This suggests that our electroweak PMBHs survive to the present universe in a wide range of blackholes, from the extremal RN type magnetic blackholes to supermassive magnetic blackholes. This strongly implies that the electroweak PMBHs could indeed become the seed of the stellar objects and galaxies, and could even account for the dark matter of the universe.

VII. REMNANT ELECTROWEAK PRIMORDIAL MAGNETIC BLACKHOLE

To test the plausibility that the electroweak PMBHs could account for the dark matter of the universe, we have to know the density of the remnant electroweak PMBHs. As we have pointed out, the electroweak PMBHs (unlike the monopoles) have almost no annihilation after they are formed, so that the evolution equation of the PMBH becomes simple,

$$\frac{dN}{dt} + 3HN = 0, \quad (90)$$

where N is the number density of the PMBHs. In this case the number density of the PMBH at present universe is given by

$$N_0 = \frac{g_{s,0}}{g_{s,i}} \left(\frac{T_0}{T_i}\right)^3 N_i \simeq 0.45 \times 10^{-45} N_i \quad (91)$$

So, for the RN PMBHs with maximum average mass $M_{mmbh} = 0.5 \times 10^{32} M_P$ with the initial density given by (79), we have $N_0 \simeq 0.59 \times 10^{-47} g_P cm^{-3}$. With this we have the energy density of the maximum average mass

RN PMBH at T_0 ,

$$\begin{aligned} \rho_{mmbh}(T_0) &= N_0 \times 0.5 \times 10^{32} M_P \\ &\simeq 0.36 \times 10^4 \text{ GeV } cm^{-3}, \\ \Omega_{mmbh} &= \frac{\rho_{bh,0}}{\rho_{c,0}} \simeq 0.74 \times 10^9. \end{aligned} \quad (92)$$

This is too much, which implies that the PMBHs would overclose the universe. Certainly this is not acceptable.

A simple way to avoid this difficulty is to assume that the PMBHs have the Hawking radiation during the matter dominant era and reduce the mass. To test this idea, let us consider the maximum average mass PMBHs and suppose they (after the radiation dominant era) undergo the evaporation which reduces the mass to M_{mmbh0} . In this case the condition that the maximum average mass PMBHs not to overclose the universe is given by

$$\begin{aligned} \rho_{mmbh}(T_0) &= N_0 \times M_{mmbh0} \leq \rho_{c,0} \\ &\simeq 0.48 \times 10^{-5} \text{ GeV } cm^{-3}, \end{aligned} \quad (93)$$

or

$$\begin{aligned} M_{mmbh0} &\leq 0.81 \times 10^{42} g_P^{-1} \text{ GeV} \\ &\simeq 0.67 \times 10^{23} M_P. \end{aligned} \quad (94)$$

So, the maximum average mass PMBHs should evaporate and reduce the mass from $0.5 \times 10^{32} M_P$ to less than $0.67 \times 10^{23} M_P$, not to overclose the universe. This of course is less than the maximum average mass M_{mmbh} , but much bigger than the extremal blackhole mass $M_{ebh} = M_P/\sqrt{\alpha}$. Certainly this seems possible.

In fact, if we assume that the maximum average mass PMBHs become the extremal RN PMBHs after the evaporation, we have

$$\begin{aligned} \rho_{mmbh}(T_0) &= N_0 \times \frac{M_P}{\sqrt{\alpha}} \simeq 8.43 \times 10^{-28} \text{ GeV } cm^{-3}, \\ \Omega_{mmbh} &= \frac{\rho_{bh,0}}{\rho_{c,0}} \simeq 1.76 \times 10^{-22}. \end{aligned} \quad (95)$$

In this case the remnant PMBHs has almost no contribution to the energy density at present universe. This strongly implies that the evaporation of the PMBHs during the matter dominant era could allow them not to overclose the universe. This is nice.

Actually we could find the condition on the initial density of the PMBH at T_i for the PMBHs not to overclose the universe. Suppose the mass of the PMBHs at present universe is M_0 in average. Then, from (91) we have

$$\begin{aligned} \rho_{bh}(T_0) &= N_0 \times M_0 = 0.45 \times 10^{-45} N_i \times M_0 \\ &\leq \rho_{c,0} \simeq 0.48 \times 10^{-5} \text{ GeV } cm^{-3}. \end{aligned} \quad (96)$$

With this we have

$$N_i \leq \frac{1.07 \times 10^{40} \text{ GeV}}{M_0} cm^{-3}. \quad (97)$$

So, assuming that the present PMBHs are extremal (i.e., assuming $M_0 = M_P/\sqrt{\alpha}$), we have

$$N_i \leq 7.5 \times 10^{19} \text{ cm}^{-3}. \quad (98)$$

From this we can deduce

$$\xi_i^{ebh} \geq 0.24 g_P^{1/3} \times 10^{-6} \text{ cm}, \quad (99)$$

which implies that the initial mass M_i^{ebh} at T_i could be around

$$\begin{aligned} M_i^{ebh} &\geq \frac{\pi^2}{30} g_* T_i^4 \times (\xi_i^{ebh})^3 \simeq 6.57 \times 10^{30} \text{ GeV} \\ &\simeq 5.39 \times 10^{21} M_P. \end{aligned} \quad (100)$$

Unfortunately, at the moment we have no way to predict the average mass M_0 of the remnant PMBHs at present universe.

The above discussion implies that Ω_{mbh} of the PMBHs could have the value similar to that of the dark matter, and could account for the dark matter of the universe. And this could happen within the framework of the standard cosmology. This is a very interesting possibility. But, of course, this is a proposal (a possibility) which has yet to be verified. Ideally, we should be able to predict the initial and final density of the electroweak PMBHs, and predict the average mass of the PMBHs. Unfortunately we could not do this in this work. Instead, we provided the plausibility argument how this could happen with the electroweak PMBHs. What is comforting is that there seems no obvious obstacle which can forbid this.

VIII. PHYSICAL IMPLICATIONS

In this paper we have compared the cosmic production of the electroweak monopoles and the electroweak PMBHs, and discussed the cosmological implications of the electroweak PMBHs. In particular, we have studied the possibility that the electroweak PMBHs become the seed of the stellar objects and galaxies, and account for the dark matter of the universe. The justification of this study is that, if the standard model is correct, we can not avoid the electroweak PMBH in cosmology. This is because the electroweak monopole automatically turns to the RN type PMBH when coupled to the gravity.

The cosmological implications of the electroweak monopole has been studied before, and it has been suggested that the electroweak monopoles could generate the density perturbation and turns to the PMBHs, and thus become the seed of the stellar objects and galaxies [22]. In this paper we have re-analyzed the cosmic production of the electroweak monopole in the early universe to argue that the popular monopole production mechanisms,

the vacuum bubble collisions in the first order phase transition and the Kibble-Zurek mechanism in the second order phase transition, may have serious defects. This is because the monopoles are created by the thermal fluctuations of the Higgs vacuum which could continue long after the phase transition till the temperature cools down to the Ginzburg temperature.

This suggests that in both first and second order phase transitions the major monopole production could take place after the phase transition. In this case the main monopole production in the first order phase transition could take place after the phase transition after the vacuum tunneling, as far as the Ginzburg temperature is considerably lower than the critical temperature. This means that what is important in the monopole production in cosmology is not the type of the phase transition but what is the Ginzburg temperature.

We have also argued that another popular wisdom, the assumption that the initial monopole density is determined by the correlation length fixed by the Higgs mass, may have a shortcoming. This is because the mass of the monopole is given by $1/\alpha$ times the W boson mass but the size of the monopole is fixed by the W boson mass, so that the energy within this correlation volume may not be enough to create the monopole. To cure this defect we have suggested to adopt a new correlation length $1/\alpha^{1/3}$ times bigger than the correlation length fixed by the Higgs mass.

Our study tells that the cosmic evolution of the electroweak PMBH is totally different from that of the electroweak monopole. There are many reasons for this. First, the electroweak PMBH has a huge mass with a wide range of uncertainty at the initial stage, from $M_P/\sqrt{\alpha}$ (for the extremal PMBH) to $1.57 \times 10^{32} M_P \simeq 1.73 \times 10^{-6} M_\odot$, compared to the monopole mass fixed by 11 TeV. This changes the initial density of the PMBHs greatly. Second, unlike the monopoles, the PMBHs have virtually no annihilation at the initial stage, basically because the monopole-antimonopole capture radius becomes much bigger than the correlation length which determines the initial monopole density.

Instead, what is important for the PMBHs is the accretion and evaporation. We have shown that our PMBHs could have significant accretion which in principle could make some of PMBHs super massive blackholes under certain conditions. As for the evaporation, the electroweak PMBHs can not evaporate completely, because they carry the conserved magnetic charge. So the evaporation could only make them the extremal RN blackhole. On the other hand the evaporation could play important roles in cosmology. All in all, we have argued that the electroweak PMBHs could become the seed of the stellar objects and galaxies, and could even account for the dark matter of the universe without conflicting with the standard cosmology.

Unfortunately, we have provided only the plausibility

argument how this could happen in this paper. One reason why we could not prove this is the huge uncertainty in the mass range of the PMBH. Another reason is that, in principle we have two possible solutions, the PMBH and the gravitationally modified electroweak monopole which is almost identical to the electroweak monopole, when the electroweak monopole is coupled to gravity. So we do not know how many electroweak PMBHs we could have at the initial stage of the monopole formation in cosmology. These are the big uncertainties that we have to remove before we could make a precise prediction.

But a most important point of our study is that in cosmology we have a new PBH, the electroweak PMBH, which is totally different from the popular PBH proposed by Zeldovich and Novikov. This PBH is a prediction based on the density perturbation, but our PMBH is an inevitable outcome of the standard model. So, if the standard model is correct, we have to deal with this. Moreover, our study in this paper strongly suggests that,

with the mass range $M_P/\sqrt{\alpha}$ to $1.57 \times 10^{32} M_P$, the electroweak PMBH could actually account for the dark matter, and could even carry the color flux.

This brings us interesting questions. Do we really need two primordial blackholes in cosmology? Which could be more realistic? Could they transform to each other merging together or absorbing the other? Can our PMBH could also carry the color charge? If so, how? And obviously we can ask many more questions. Certainly we need more research work to answer these questions.

ACKNOWLEDGEMENT

This work is supported in part by the National Research Foundation of Korea funded by the Ministry of Science and Technology (Grant 2022-R1A2C1006999), Center for Quantum Spacetime, Sogang University, and Department of Liberal Arts, Korea Polytechnic University, Korea.

-
- [1] P.A.M. Dirac, Proc. Roy. Soc. London, **A133**, 60 (1931); Phys. Rev. **74**, 817 (1948).
 - [2] B. Cabrera, Phys. Rev. Lett. **48**, 1378 (1982).
 - [3] T.T. Wu and C.N. Yang, in *Properties of Matter under Unusual Conditions*, edited by H. Mark and S. Fernbach (Interscience, New York) 1969; Phys. Rev. **D12**, 3845 (1975); Y.M. Cho, Phys. Rev. Lett. **44**, 1115 (1980); Phys. Lett. **B115**, 125 (1982).
 - [4] G. 't Hooft, Nucl. Phys. **B79**, 276 (1974); A.M. Polyakov, JETP Lett. **20**, 194 (1974); B. Julia and A. Zee, Phys. Rev. **D11**, 2227 (1975); M. Prasad and C. Sommerfield, Phys. Rev. Lett. **35**, 760 (1975).
 - [5] C. Dokos and T. Tomaras, Phys. Rev. **D21**, 2940 (1980).
 - [6] Y.M. Cho and D. Maison, Phys. Lett. **B391**, 360 (1997).
 - [7] Yisong Yang, Proc. Roy. Soc. London, **A454**, 155 (1998); Yisong Yang, *Solitons in Field Theory and Nonlinear Analysis* (Springer Monographs in Mathematics), p. 322 (Springer-Verlag) 2001.
 - [8] Kyoungtae Kimm, J.H. Yoon, and Y.M. Cho, Eur. Phys. J. **C75**, 67 (2015).
 - [9] J. Ellis, N.E. Mavromatos, and T. You, Phys. Lett **B756**, 29, (2016).
 - [10] F. Blaschke and P. Benes, Prog. Theor. Exp. Phys. **073B03** (2018).
 - [11] Pengming Zhang, Liping Zou, and Y.M. Cho, Euro. Phys. J. **C80**, 280 (2020).
 - [12] P. Benes, F. Blaschke, and Y.M. Cho, arXiv:2024.10747[hep-ph], to be published.
 - [13] B. Acharya et al. (MoEDAL Collaboration), Phys. Rev. Lett. **118**, 061801 (2017).
 - [14] B. Acharya et al. (MoEDAL Collaboration), Phys. Rev. Lett. **123**, 021802 (2019).
 - [15] B. Acharya et al. (MoEDAL Collaboration), Nature **602**, 63 (2022).
 - [16] G. Aad. et al. (ATLAS Collaboration), Phys. Rev. Lett. **124**, 031802 (2020).
 - [17] T.W.B. Kibble, J. Phys. **A9** 1387 (1976).
 - [18] J.P. Preskill, Phys. Rev. Lett. **43**, 1365 (1979).
 - [19] A.H. Guth and E.J. Weinberg, Nucl. Phys. **B212**, 321 (1983).
 - [20] W.H. Zurek, Phys. Rep. **276** 177 (1996).
 - [21] A.H. Guth, Phys. Rev. **D23**, 347 (1981); A.D. Linde, Phys. Lett. **B108** 389 (1982).
 - [22] Y.M. Cho, Phil. Trans. R. Soc. **A377**, 0038 (2019).
 - [23] Y.M. Cho and Franklin H. Cho, Phys. Lett **B851**, 138598, (2024).
 - [24] F. Bais and R. Russel, Phys. Rev. **D11**, 2692 (1975).
 - [25] Y.M. Cho and P.G.O. Freund, Phys. Rev. **D12**, 1588 (1975).
 - [26] Y.M. Cho, Kyoungtae Kimm, J.H. Yoon, Phys. Lett. **B761**, 203 (2016).
 - [27] Yang Bai and Mrunal Korwar, JHEP **04**, 119 (2021).
 - [28] D. Zhu, K.M. Wong and G.Q. Wong, Commun. Theor. Phys. **76**, 035201 (2024).
 - [29] Y.B. Zeldovich and I.D. Novikov, Sov. Astro. AJ **10**, 602 (1967).
 - [30] S. Hawking, Nature **248**, 30 (1974).
 - [31] B. Carr and S. Hawking, Mon. Not. R. astr. Soc. **168**, 399 (1974).
 - [32] B. Carr and F. Kuhnel, Annu. Rev. Nucl. Part. Sci. **70**, 355 (2020).
 - [33] A.M. Green and B.J. Kavanagh, J. Phys. **G48**, 043001 (2021).
 - [34] B. Carr, S. Clesse, J. Garcia-Bellido, M. Hwkins, and F. Kuhnel, Phys. Rep. **1054**, 1 (2024).
 - [35] E. Alonso-Monsalve and D. Kaiser, Phys. Rev. Lett. **132**, 230402 (2024).
 - [36] W. DeRocco, E. Frangipane, N. Hamer, S. Profumo, N. Smyth, Phys. Rev. **D109**, 023013 (2024).
 - [37] D.A. Krizhnits and A.D. Linde, Phys. Lett. **B42**, 471 (1972); C. Bernard, Phys. Rev. **D9**, 3312 (1974).
 - [38] G. Anderson and L. Hall, Phys. Rev. **D45**, 2685 (1992); M. Dine, R. Leigh, P. Huet, A. Linde, and D. Linde, Phys. Rev. **D46**, 550 (1992).
 - [39] S. Arunasalam and A. Kobahidze, Euro. Phys. J. **C77**, 444 (2017); S. Arunasalam, D. Collision, and A. Kobah-

- hidze, hep-ph/1810.10696 (2018).
- [40] V. Ginzburg, Sov. Phys. Solid State **2**, 1824 (1960).
- [41] See, e.g., E. Kolb and M. Turner, *The Early Universe* (Addition-Wesley Publishing Co.) 1990.
- [42] A. Vilenkin and E. Shellard, *Cosmic Strings and other Topological Defects* (Cambridge University Press) 1994.
- [43] R. Eatough et al., Nature **591**, 391 (2013); M. Zamaninasab, E. Clausen-Brown, T. Savolainen, A. Tchekhovskoy, Nature **510**, 126 (2014).
- [44] S. Cecchini, L. Patrizii, Z. Sahnoun, G. Sirri, and V. Togo, arXiv: hep-ph/1606.01220 (2016).
- [45] E.N. Parker, Astrophys. J. **160**, 383 (1970).
- [46] Telescope Array Collaboraltion, Science **382**, 903 (2023).
- [47] K. Greisen, Phys. Rev. Lett. **16**, 748 (1996); G. Zatsepin, V. Kuzmin, Pisma Zh. Eksp. Teor. Fiz. **4**, 114 (1996).