

Dwellers in the Deep: Biological Consequences of Dark Oxygen

Manasvi Lingam,^{1,2,3*} Amedeo Balbi⁴ and Madhur Tiwari¹

¹Department of Aerospace, Physics and Space Sciences, Florida
Institute of Technology, Melbourne, FL 32901, USA

²Department of Chemistry and Chemical Engineering, Florida
Institute of Technology, Melbourne, FL 32901, USA

³Department of Physics, The University of Texas at Austin,
Austin, TX 78712, USA

⁴Dipartimento di Fisica, Università di Roma “Tor Vergata”, 00133
Roma, Italy

*To whom correspondence should be addressed;

E-mail: mlingam@fit.edu

Keywords: dark oxygen | deep oceans | aerobic respiration | complex life |
abiogenesis

Abstract: The striking recent putative detection of “dark oxygen”
(dark O₂) sources on the abyssal ocean floor in the Pacific at ~ 4 km
depth raises the intriguing scenario that complex (i.e., animal-like)
life could exist in underwater environments sans oxygenic photosyn-

thesis. In this work, we thus explore the possible (astro)biological implications of this discovery. From the available data, we roughly estimate the concentration of dissolved O_2 and the corresponding O_2 partial pressure, as well as the flux of O_2 production, associated with dark oxygen sources. Based on these values, we infer that organisms limited by internal diffusion may reach maximal sizes of ~ 0.1 – 1 mm in habitats with dark O_2 , while those with circulatory systems might achieve sizes of ~ 0.1 – 10 cm. Optimistically, the estimated dark oxygen flux can potentially support biomass densities up to ~ 3 – 30 g m $^{-2}$, perhaps surpassing typical reported densities at similar depths in global deep-sea surveys. Finally, we outline how oceanic settings with dark O_2 may facilitate the origin(s) of life via the emergence of electrotrophy. Our findings indicate that complex life fueled by dark oxygen is plausibly capable of inhabiting submarine environments devoid of photosynthesis on Earth, conceivably extending likewise to extraterrestrial locations such as icy worlds with subsurface oceans (e.g., Enceladus and Europa), which are likely common throughout the Universe.

1 Introduction and Motivation

The non-monotonic increase of molecular oxygen (O_2) across Earth's geological history exerted a panoply of profound consequences ranging from the formation of the ozone layer (which partially screens ultraviolet radiation) and new minerals to the creation of novel ecological niches and metabolisms [Lane, 2002, Canfield, 2014, Lyons et al., 2014, Knoll, 2015, Fischer et al., 2016, Knoll and Nowak, 2017, Judson, 2017, Lingam and Balbi, 2024]. Aerobic metabolic reactions are conventionally documented to yield free energy that is roughly an order of mag-

nitude greater than their anaerobic counterparts for a fixed amount of food [McCollom, 2007, Koch and Britton, 2008]. On the other hand, molecular oxygen can readily abstract electrons from certain biomolecules and enable the formation of reactive oxygen species, thus posing toxicity issues and engendering an “oxygen paradox” [Davies, 1995, Halliwell and Gutteridge, 2015].

Drawing partly on the above energy-rich nature of aerobic metabolisms, a variety of classic [Nursall, 1959, Cloud Jr., 1968, Runnegar, 1982, Knoll, 1985] and modern [Catling et al., 2005, Knoll and Sperling, 2014, Knoll, 2015, Lyons et al., 2021, Mills et al., 2023] studies have advanced and/or elaborated upon the hypothesis that O_2 is a prerequisite of sorts for the evolution of complex, motile, macroscopic, multicellular life – or “complex” life for short [cf. Knoll, 2011]; we caution, however, that a causal link continues to be debated and analyzed [Cole et al., 2020, Bozdogan et al., 2021, Sperling et al., 2022]. Despite this ambiguity, it is still tenable that O_2 was beneficial for the advent of such complex life [Knoll and Sperling, 2014, Lingam and Loeb, 2021, Mills et al., 2023], which is implicitly treated in the paper as being near-synonymous with metazoans (i.e., animals), since the latter evince the prior attributes and are potent ecosystem engineers [Jones et al., 1994, Butterfield, 2011].

If we now turn our attention to the provenance of Earth’s oxygenated atmosphere and oceans, there are a bevy of factors at play [Knoll, 2015, Catling and Kasting, 2017], but the root cause is the emergence of oxygenic photosynthesis [Canfield, 2014, Lyons et al., 2014, Fischer et al., 2016, Lyons et al., 2021], which synthesizes molecular oxygen as a metabolic product. In this respect, it should be appreciated that only a small fraction of Earth’s oceans are amenable to (oxygenic) photosynthesis. The euphotic zone depth or the compensation depth – both of which crudely encapsulate the spatial extent wherein photosynthesis (chiefly) takes place – happen to be $\lesssim 100$ m [Sarmiento and Gruber, 2006,

Lee et al., 2007, Middelburg, 2019]; evidently much shallower than the average ocean depth of 3.7 km. Moreover, on temperate rocky planets orbiting M-dwarfs, the most widespread type of stars in the Cosmos, these depths could decline to only a few meters [Ritchie et al., 2018, Lingam and Loeb, 2020].

Therefore, significant portions of Earth’s habitats – as well as presumably Earth-like planets in the habitable zones of other stars [Kasting et al., 1993, Kopparapu et al., 2013] – do not seem conducive to photosynthesis. The situation is exacerbated if we consider worlds that may have habitable conditions in subsurface oceans under substantial icy shells: notable examples in our solar system are Europa and Enceladus [Nimmo and Pappalardo, 2016, Hendrix et al., 2019]. This group of worlds are not anticipated to support photosynthesis in their oceans, even if they could otherwise be habitable [Gaidos et al., 1999, Chyba and Hand, 2001]. This issue is pertinent because such worlds might be among the most common class of habitable worlds in the Milky Way, namely, up to ~ 3 orders of magnitude more prevalent than rocky planets in the habitable zone [Mojzsis, 2021, Lingam et al., 2023].

On the basis of the preceding paragraphs, it would appear that a multitude of habitable worlds and environments in the Universe do not allow for the production of O_2 via oxygenic photosynthesis. The ensuing scarcity of O_2 would, *prima facie*, potentially diminish the prospects for complex life. However, this reasoning is rendered invalid if a novel source of O_2 not dependent on photosynthesis could be identified. In this context, the remarkable putative discovery of “*dark oxygen*” generation at the abyssal seafloor in the (~ 4 km deep) Clarion–Clipperton Zone of the Pacific Ocean [Sweetman et al., 2024] stands out as a conceivable avenue in this regard, perhaps by way of polymetallic nodules.

Thus, by taking inspiration from this ostensibly striking result, we explore possible ramifications of the existence of dark oxygen for the genesis and evolu-

tion of complex life in Earth’s deep oceans and on other worlds, paralleling the “follow the oxygen” strategy espoused in Ward et al. [2019].

2 Prospects for complex life

In this section, we delve into the prospects for complex life and generic biomass sustained by dark oxygen at the seafloor.

2.1 Constraints on dark oxygen availability and biomass

To assess the consequences of dark oxygen production (DOP) for complex life, we begin with sketching some tentative findings inferred from the empirical data of Sweetman et al. [2024, pg. 737], which will be utilized subsequently.

- To loosely estimate the concentration of dissolved O_2 (denoted by \mathcal{C}) furnished exclusively by means of DOP, we may subtract the final steady-state value of 201–819 μM from the initial (ambient) O_2 abundance of 185.2 μM that could likely be supplied by external sources (e.g., current Earth’s oxygenated oceans) – as a consistency check, this ambient concentration is similar to deep-sea measurements of O_2 in the Pacific ocean [Smith et al., 2022] – which thereupon yields $\mathcal{C} \approx 15.8\text{--}633.8 \mu\text{M}$.
- Next, we investigate the following cognate scenario. In lieu of the actual DOP, if we envision an oxygen source/reservoir (e.g., atmosphere) in contact with a water body, as on Earth during its history, we can then ask: what is the O_2 partial pressure (labeled by \mathcal{P}) that would essentially translate to an effective concentration of \mathcal{C} (stemming from DOP)? To answer this question, we harness Henry’s Law for dilute aqueous solutions widely

used in physical chemistry [Atkins and De Paula, 2017, Chapter 4D]:

$$\mathcal{C} \approx k_H \mathcal{P}, \quad (1)$$

where $k_H(T)$ is Henry’s Law “constant” for (dilute) aqueous solutions. In reality, $k_H(T)$ is a function of the ambient temperature T , canonically modeled through the Van’t Hoff equation for simplicity [Sander, 1999, Section 2.5]. For the case of dissolved O_2 , we adopt [Sander, 2023]:¹

$$k_H(T) \approx 1300 \mu\text{M bar}^{-1} \exp\left(\frac{1500}{T} - \frac{1500}{298.15 \text{ K}}\right). \quad (2)$$

On substituting $T = 274.75 \text{ K}$, akin to the Clarion–Clipperton Zone, in Eq. (2), we obtain $k_H \approx 2 \times 10^3 \mu\text{M bar}^{-1}$; the external total pressure has a relatively modest effect on the solubility (e.g., Hofmann et al. 2011, Cole et al. 2020 and Averill and Eldredge 2020, Chapter 13.4), and is therefore not taken into account. After plugging k_H in Eq. (1) along with the prior range for \mathcal{C} , we end up with $\mathcal{P} \approx 7.9\text{--}318 \text{ mbar}$, the latter of which is tantamount to $\mathcal{P} \approx 3.7\text{--}150\% \text{ PAL}$ (viz., present atmospheric level).

- Measurements of abyssal dark oxygen suggest that the local flux (denoted by \mathcal{F}) attached to DOP is perhaps $\mathcal{F} \approx 1.7\text{--}18 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$.

Thus, as per the above inferences and calculations, the lower bound of the potential O_2 abundance arising from DOP is estimated to be $\mathcal{C}_{\min} \approx 15.8 \mu\text{M}$, or equivalently $\mathcal{P}_{\min} \approx 7.9 \text{ mbar} \approx 3.7\% \text{ PAL}$ for the scenario of the oxygen source/reservoir interfacing with water. In some instances, we will employ these conservative choices for dark O_2 , although we shall mostly avail ourselves of the entire O_2 range(s).

¹<https://henrys-law.org/henry/casrn/7782-44-7>

Before proceeding onward to carrying out a theoretical analysis of organismal size, we highlight a couple of important features from Earth’s biota. A variety of initial analyses indicated that O_2 levels of $\sim 1\text{--}10\%$ PAL may be sufficient for the evolution of early metazoans [Berkner and Marshall, 1965, Cloud, 1976, Runnegar, 1991], while select publications subsequently revised this threshold downwards to $\gtrsim 0.1\%$ PAL [Sperling et al., 2013b]. On comparing this range of proposed values for the desired O_2 partial pressure with \mathcal{P}_{\min} , this range apparently overlaps with the latter, hinting that animal-like organisms could possibly exist in environs hosting DOP.

From an empirical standpoint, a plethora of *Polychaete* species are documented in oxygen-scarce benthic regions with O_2 concentrations of $< 10 \mu\text{M}$ (e.g., Levin et al. 1991 and Sperling et al. 2013a, Table 1), which are distinctly lower than \mathcal{C}_{\min} . Likewise, controlled laboratory experiments conducted on the demosponge *Halichondria panicea* have shown that this animal can survive at oxygen levels of $\lesssim 15 \mu\text{M}$, or 0.5–4% PAL to be accurate [Mills et al., 2014], which are respectively compatible with \mathcal{C}_{\min} and \mathcal{P}_{\min} . Hence, even on empirical grounds, the O_2 abundances attributable to DOP may suffice to sustain metazoan-type organisms. On the other hand, according to modern data, carnivory might be rendered challenging at O_2 concentrations of $\lesssim 15 \mu\text{M}$ [Sperling et al., 2013a, pg. 13448], which is close to \mathcal{C}_{\min} and should be duly borne in mind.

Another key datum that might be partly adduced in favor of harboring animal-like lifeforms (in abyssal settings with dark oxygen) is that at least one known metazoan, the parasite *Henneguya salminicola* of the family *Myxobolidae* [Yahalomi et al., 2020], lacks mitochondrial genomes and the capacity for aerobic respiration. Although a rather extreme outlier – in the sense of being a parasitic and minuscule animal (sub-mm in size) – this example nevertheless

illustrates that animals do not necessarily require O_2 for their survival.

Finally, let us contemplate the idealized setup where we assume that DOP operates continuously to maintain the flux \mathcal{F} , and that all the associated O_2 is fully consumed by metazoan-like organisms with no other sinks active. In this setup, we may estimate an upper bound on the biomass area density σ_{\max} on the seafloor by resorting to the following order-of-magnitude approach:

$$\sigma_{\max} \sim \frac{\mathcal{F}}{\mathcal{B}}, \quad (3)$$

where \mathcal{B} is the mass-specific aerobic metabolic rate (expressed in terms of the wet tissue mass), chosen to be $\mathcal{B} \sim 0.54 \text{ cm}^3 O_2 \text{ g}^{-1} \text{ h}^{-1}$ from Sperling et al. [2013b] for potential early (or “primitive”) metazoans. We will now invoke the molar volume – a quantity of relevance for performing unit conversion [Hofmann et al., 2011, pg. 1213] – which yields $\mathcal{B} \sim 0.58 \text{ mmol } O_2 \text{ g}^{-1} \text{ d}^{-1}$.

We caution that the mass-specific metabolic rate exhibits variation across and within taxa, and as a function of body mass of temperature – the extent is not precisely settled (contrast Glazier 2005, Clarke 2017, Glazier 2022 with Makarieva et al. 2008, Hoehler et al. 2023) – owing to which the results involving \mathcal{B} should be viewed as broadly heuristic estimates. At the same time, however, the salient quantities all scale as \mathcal{B}^α ; wherein we have $|\alpha| \leq 1$, thereby reflecting a modest dependence on \mathcal{B} , and revealing that our conclusions ought not be particularly sensitive to the magnitude of \mathcal{B} . This opportune facet should be borne in mind henceforth.

After substituting the preceding values for \mathcal{B} and \mathcal{F} in Eq. (3), we arrive at $\sigma_{\max} \sim 2.9\text{--}31 \text{ g m}^{-2}$. This optimistic range (i.e., likely embodying the greatest viable biomass density) compares favorably with the observed macrofaunal (wet) biomass density of order $0.01\text{--}1 \text{ g m}^{-2}$ at an ocean depth of $\sim 4 \text{ km}$ in a global survey [Rex et al., 2006, Figure 3], but specialized habitats such as some

submarine canyons (e.g., Kaikoura Canyon) comprise an unusually high (wet) biomass density on the order of 1000 g m^{-2} [De Leo et al., 2010, pg. 2787].

Lastly, an interesting property of abyssal regions of the Clarion–Clipperton Zone is that they contain high megafaunal number densities of order 1 m^{-2} [Amon et al., 2016, Simon-Lledó et al., 2019, Durden et al., 2021], which could be mapped to a biomass density of $\sim 1\text{--}10 \text{ g m}^{-2}$ for fiducial cm-sized metazoans [Simon-Lledó et al., 2019, Table 1]. This value is loosely similar to σ_{\max} , with the proviso that the latter is an optimistic biomass estimate spanning all types of organisms.

2.2 Theoretical diffusion and circulation size limits

It is well-established that unicellular prokaryote-like organisms may not typically exceed a maximal size imposed by the diffusive uptake of ambient substrates [Denny, 1993, Kiørboe, 2008]. In fact, this bottleneck can restrict the sizes attainable by the biggest prokaryotes [Koch, 1996, Schulz and Jørgensen, 2001], albeit other factors also influence the upper-size thresholds of organisms [e.g., Andersen et al., 2016], as elaborated hereafter.

We obtain a rough upper bound for the organismal radius (R_{\max}) and size ($L_{\max} \equiv 2R_{\max}$) by balancing the supply rate of O_2 through diffusion with the metabolic rate of the spheroidal organism, corresponding to left-hand side (LHS) and right-hand side (RHS) of the following equation:

$$4\pi R_{\max} DC \sim \frac{4\pi}{3} \rho \mathcal{B} R_{\max}^3, \quad (4)$$

where the LHS constitutes a standard biophysics textbook treatment [Berg, 1993, Chapter 2], and the RHS (viz., the whole-organism metabolic rate) is constructed by multiplying the organismal mass with the mass-specific aerobic metabolic rate \mathcal{B} . In the above equation, D signifies the diffusion coefficient,

and $\rho \approx 1 \text{ g cm}^{-3}$ is the mass density of the organism [Milo and Phillips, 2016, pg. 72]. On simplifying the above equation, we end up with

$$L_{\max} \sim \sqrt{\frac{12DC}{\rho\mathcal{B}}}, \quad (5)$$

for the upper-size bound of a spheroidal organism that acquires O_2 via diffusive uptake from the ambient environment.

At normal temperature and pressure (NTP), the diffusion coefficient is $2 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ for O_2 [Han and Bartels, 1996]. Next, we specify $T \approx 274.75 \text{ K}$ and total pressure of $\sim 400 \text{ bar}$ in the Clarion–Clipperton Zone, enabling us to compute the dynamic viscosity η from Schmelzer et al. [2005, Table 1]. Upon combining this data with the Stokes–Einstein equation – namely, $D \propto T/\eta$ [Milo and Phillips, 2016, pg. 212] – we arrive at $D \approx 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ in this region. On substituting the preceding values in Eq. (5), we obtain $L_{\max} \sim 0.17\text{--}1.1 \text{ mm}$. Hence, we conclude that organisms solely reliant on diffusive uptake of dark O_2 might reach sizes up to 0.1–1 mm.

For the upcoming two parts pertaining to maximal size thresholds of idealized representations of multicellular organisms [Alexander, 1971], it is worth recognizing that the original models employed – which are known to satisfactorily reproduce properties of (modern) Earth’s biota – were formulated as a function of the partial O_2 pressure. Thus, given the usage of the effective O_2 partial pressure (\mathcal{P}) as a proxy, the ensuing results should be carefully interpreted because of possible subtle difference(s) between the dissolved O_2 concentration \mathcal{C} at the abyssal seafloor and the scenario delineated previously vis-à-vis determining \mathcal{P} . Furthermore, as some parameters are not precisely quantified for early metazoans and analogous organisms, we rely on fiducial values in such cases.

In the first group, organisms composed of simple aggregates of cells may

be reliant on the diffusion of O_2 to all cells in their interior for supporting the organism's metabolic needs [Harvey, 1928]. For a spheroidal lifeform, this constraint yields a upper-size limit of [Catling et al., 2005, pg. 433]:

$$L_{\max} \sim \sqrt{\frac{6\mathcal{K}}{\rho\mathcal{B}} (\mathcal{P} - \mathcal{P}_{\text{crit}})}, \quad (6)$$

where \mathcal{K} is the O_2 permeability within the lifeform, and $\mathcal{P}_{\text{crit}}$ is the minimal (internal) oxygen level required for organismal functioning. An extra factor of ρ appears in the denominator because we work with mass-specific metabolic rate \mathcal{B} , and not its volume-specific counterpart. We adopt $\mathcal{K} \sim 2.24 \times 10^{-3} \text{ cm}^2 \text{ bar}^{-1} \text{ h}^{-1}$ and $\mathcal{P}_{\text{crit}} \sim 6.67 \times 10^{-4} \text{ bar}$ from Sperling et al. [2013b]. On plugging these values along with \mathcal{P} , \mathcal{B} , and ρ from earlier, we obtain $L_{\max} \sim 0.13\text{--}0.89 \text{ mm}$, implying that the largest organisms constrained by internal diffusion may attain sizes of order 0.1–1 mm in habitats with DOP.

In the second category, the putative spheroidal organisms are presumed to host a circulation system that can distribute O_2 throughout their bodies, instead of relying on diffusion. For such lifeforms, the upper bound for the organismal size is expressible as [Catling et al., 2005, pg. 424]:

$$L_{\max} \sim \frac{3\mathcal{K}}{\rho\mathcal{B}\ell} (\mathcal{P} - \mathcal{P}_b), \quad (7)$$

where \mathcal{P}_b is the average dissolved O_2 concentration in the blood, which is modeled as $\mathcal{P}/4$ [Alexander, 1971], and ℓ is the mean distance between the organism's surface and circulation system (i.e., the thickness of the epidermal layer). As with Eq. (6), an extra factor of ρ is included in the denominator for conversion purposes. After setting $\mathcal{B} \sim 0.1 \text{ cm}^3 \text{ O}_2 \text{ g}^{-1} \text{ h}^{-1}$ for bigger invertebrate-like organisms with lower mass-specific metabolic rates [Payne et al., 2011] and $\ell \sim 2 \times 10^{-3} \text{ cm}$ [Sperling et al., 2013b] in Eq. (7),

we end up with $L_{\max} \sim 0.2\text{--}8$ cm. Hence, DOP might possess the capacity, in principle, to sustain “megafauna”, to wit, lifeforms exhibiting sizes of $\sim 1\text{--}10$ cm [Rex et al., 2006, pg. 3].

3 Implications for abiogenesis

The chief purpose of this paper was to analyze the consequences of DOP for complex life. We will, however, delve briefly into a potential feature identified in Sweetman et al. [2024]: the authors conjectured that the source of dark oxygen may be seawater electrolysis engendered by polymetallic nodules, which seem to have a potential difference reaching $\Delta\Phi_0 \approx 0.95$ V on their surfaces. We tackle the possible ramifications of this feature for the origin(s) of life, a vast and rapidly expanding field of research [Luisi, 2016, Deamer, 2019, Preiner et al., 2020, Sasselov et al., 2020, Lingam and Loeb, 2021].

In 1961, Peter Mitchell elucidated the mechanism(s) of chemiosmosis [Mitchell, 1961], which led to the 1978 Nobel Prize in Chemistry [Mitchell, 1979]. This ubiquitous biochemical phenomenon involves the flow of charged particles across a semipermeable membrane to synthesize adenosine triphosphate (ATP) [Nicholls and Ferguson, 2013]. The proton motive force (PMF), denoted by Δp (units of V), underpinning chemiosmosis is given by [Nicholls and Ferguson, 2013, pg. 44]:

$$\Delta p = \Delta\phi - \frac{2.3RT}{F}\Delta(pH), \quad (8)$$

where $\Delta\phi$ is the transmembrane electric potential difference, and $\Delta(pH) \equiv pH_{\text{in}} - pH_{\text{out}}$ is the pH difference between the interior and exterior of the cell. Note that R represents the gas constant, and F labels the Faraday constant. From inspecting Eq. (8), it is regulated by multiple variables (ranging from organismal to environmental), owing to which a single value cannot be assigned

to the PMF; with that said, a fiducial choice of $\Delta p \sim 0.2 \pm 0.1$ V is compatible with experiments and modeling for sundry species [e.g., Rao et al., 2008, Soga et al., 2011, Nicholls and Ferguson, 2013, Milo and Phillips, 2016].

Taking inspiration from chemiosmosis and the PMF, the famous alkaline hydrothermal vent (AHV) hypothesis for the origin of life (i.e., abiogenesis) postulates that life emerged in a submarine AHV [Martin et al., 2008, Russell et al., 2014, Sojo et al., 2016, Russell, 2021, 2023] by harnessing the naturally occurring redox and proton gradients in these sites [Russell and Hall, 1997], which may have translated to a total electrochemical gradient equivalent to ~ 1 V as per theoretical analyses and laboratory experiments [Russell et al., 2014, Barge et al., 2015, Sojo et al., 2016]. These steep gradients are conjectured to have facilitated carbon fixation, prebiotic synthesis, and abiogenesis [Russell, 2021, Schwander et al., 2023].

If we compare the reported magnitude of $\Delta\Phi_0$, associated with the nodules, against the above values of the PMF and the putative electrochemical gradient at AHVs, it is evident that the former matches and/or surpasses the latter duo. Therefore, this aspect raises the tantalizing notion that the surfaces of polymetallic nodules might also permit the instantiation of electrochemistry conducive to prebiotic chemistry. However, one crucial divergence is that cells as well as AHVs – the latter via thin inorganic layers composed of minerals (e.g., iron oxyhydroxides) [Russell, 2018] – evince sharp gradients operational across membranes, whereas these nodules, with typical sizes of $\lesssim 1\text{--}10$ cm [Hein et al., 2020], are not anticipated to have those distinctive properties.

To circumvent the preceding caveat, we underscore that electrical currents have been detected in a variety of submarine hydrothermal vents [Yamamoto et al., 2013, 2017], which can be attributed to a sustained redox potential difference of around 0.5 V between hydrothermal mineral deposits and the surrounding

seawater. This current drives the avenue of electrotrophy, whereby electrons serve as the energy source (i.e., electron donor) for microbial metabolisms [Lovley, 2022]. Experiments mimicking the electrical current in hydrothermal vent settings, in conjunction with *in situ* analyses, have shown that microbes (e.g., from genus *Thiomicrothrix*) may survive and grow by means of electrotrophy [Pillot et al., 2021, Yamamoto et al., 2023]. Drawing on these strands, Pillot et al. [2023] proposed that a prototype of electrotrophy might have sparked abiogenesis, and adumbrated a series of conceivable intermediate steps.

With an electric potential of $\lesssim \Delta\Phi_0$ measured at the surfaces of polymetallic nodules, an electron flow and current is rendered credible, which may in turn pave the way for the emergence of life by way of (proto-)electrotrophy. Moreover, these nodules contain a plethora of metals (e.g., manganese) and minerals (e.g., iron oxyhydroxides) [Hein et al., 2020], with strong connections to the origin(s) of life [Sahai et al., 2016, Russell, 2018, Muchowska et al., 2020], thence perhaps boosting (directly or otherwise) the significance of polymetallic nodules for abiogenesis.

While the exposition in this section hints at these sites (viz., polymetallic nodules in the abyssal seafloor) comprising intriguing candidates for the nurseries of abiogenesis, further theoretical and empirical investigations of the surface potential difference and the material composition of the nodules, *inter alia*, will be important for gauging their viability; a research program paralleling the modeling and experiments delineated in Russell [2021, Sections 9 & 12] is arguably warranted.

4 Discussion and conclusions

Motivated by the striking ostensible discovery of dark oxygen (O_2), we analyzed the possible ramifications of dark O_2 for “complex” life and the origin(s) of life on Earth and beyond; our major surmises are summarized as follows:

1. The maximal sizes attainable by idealized unicellular or multicellular organisms (i.e., constrained by internal or external diffusion processes) for the estimated concentrations of dark O_2 may be $\sim 0.1\text{--}1$ mm.
2. In contrast, the upper-size bounds of organisms with internal circulation systems for the distribution of oxygen could range between ~ 0.1 cm to ~ 10 cm, with the latter threshold falling under the umbrella of “megafauna”.
3. Under optimistic circumstances, the biomass densities might reach as high as $\sim 3\text{--}30$ g m $^{-2}$, in principle exceeding the reported macrofaunal densities at depths of ~ 4 km in global deep-sea surveys, but perhaps of the same order as the inferred metazoan densities in the abyssal locales of the Clarion–Clipperton Zone.
4. Polymetallic nodules on the seafloor with a surface potential difference of $\lesssim 1$ V are tentatively considered the provenance of dark oxygen production (DOP). We sketched how this ensuing voltage may conceivably aid the emergence of life by enabling prebiotic synthesis to transpire through the pathway of electrotrophy, where electrons function as the energy source.

In the paper, we have chronicled the limitations and/or unknowns of our overall approach (which has a heuristic slant emphasizing simplicity), where appropriate.

With regard to the limitations, we have often computed theoretical upper bounds that are not necessarily the characteristic values, and not incorporated

factors whose impacts are not clearly resolved (e.g., role of salinity in modulating O_2 supply and organismal size). Likewise, it was implicitly posited that DOP and the O_2 concentration do not change substantially in time (albeit they can exhibit spatial heterogeneity). This facet patently merits follow-up *in situ* explorations to ascertain the spatiotemporal nature of both dark O_2 sources and sinks. As for unknowns, while certain environmental parameters (e.g., temperature and pressure) are reasonably attested, the biological variables should be construed as extrapolations based on available knowledge.

Notwithstanding these caveats, most of our findings are endowed with the key advantage that they are consistent with empirical data (e.g., O_2 requirements for metazoans outlined previously). The capacity for possibly harboring macrofauna or megafauna (mentioned above) fueled by dark O_2 is roughly compatible with the high biodiversity of such metazoans (e.g., *Xenophyophorea* and *Ophiuroidea*) in the Clarion–Clipperton Zone, including nodule-rich regions [Amon et al., 2016, Gooday et al., 2017, Christodoulou et al., 2020, Rabone et al., 2023, Uhlenkott et al., 2023], although this feature does not translate to a causal link. In addition, laboratory experiments entailing the reproduction of poly-metallic nodules and DOP may help assess whether organisms and/or biomass densities outside/within the preceding ranges are viable.

Furthermore, as many of the results do not explicitly take the root cause of DOP into account, they could be adapted to study lifeforms and ecosystems supported by alternative channels of DOP. A couple of tenable routes worth highlighting in this realm are as follows:

- The production of oxidants on the surface (e.g., powered by radiolysis) and their delivery to the ocean can effectively input O_2 to the latter even sans photosynthesis [Chyba and Hand, 2001]. This mechanism is presumably active on icy moons like Europa [Lingam and Balbi, 2024, Chapter 10],

whose ~ 100 km deep subsurface ocean is estimated to accrue O_2 at supply rates of $\sim 3 \times 10^8 - 3 \times 10^{11} \text{ mol yr}^{-1}$ [Greenberg, 2010, Vance et al., 2016].

- Microbial dismutation (e.g., chlorite dismutation) could generate dark O_2 . This metabolic phenomenon has been recently documented in groundwater samples extracted from multiple aquifers in Canada, where the mean dissolved O_2 concentration was determined to be $\sim 16.3 \mu\text{M}$ [Ruff et al., 2023], which is nearly equal to the minimal O_2 abundance (\mathcal{C}_{\min}) employed in our calculations.

To round off our preliminary venture into this eclectic subject, we reiterate our prefatory statement that marine habitable settings implausible for photosynthesis, especially on icy worlds with subsurface oceans, are likely widespread in the Universe. Therefore, if DOP is feasible and commonplace on this class of worlds – whether via seawater electrolysis or the prior two routes – then our analysis may broadly encapsulate the profound consequences of dark oxygen for the prevalence of abiogenesis, complex multicellularity, and perhaps even technological intelligence [Lingam and Loeb, 2021] in the Cosmos.

Acknowledgments

TBA.

References

- R. M. Alexander. *Size and Shape*. Number 29 in Studies in Biology. London: Edward Arnold, 1971.
- D. J. Amon, A. F. Ziegler, T. G. Dahlgren, A. G. Glover, A. Goineau, A. J. Goday, H. Wiklund, and C. R. Smith. Insights into the abundance and diversity

- of abyssal megafauna in a polymetallic-nodule region in the eastern Clarion-Clipperton Zone. *Sci. Rep.*, 6:30492, July 2016. doi: 10.1038/srep30492.
- K. H. Andersen, T. Berge, R. J. Gonçaves, M. Hartvig, J. Heuschele, S. Hylander, N. S. Jacobsen, C. Lindemann, E. A. Martens, A. B. Neuheimer, K. Olsson, A. Palacz, A. E. F. Prowe, J. Sainmont, S. J. Traving, A. W. Visser, N. Wadhwa, and T. Kiørboe. Characteristic Sizes of Life in the Oceans, from Bacteria to Whales. *Annu. Rev. Mar. Sci.*, 8:217–241, Jan. 2016. doi: 10.1146/annurev-marine-122414-034144.
- P. Atkins and J. De Paula. *Elements of Physical Chemistry*. Oxford: Oxford University Press, 2017.
- B. Averill and P. Eldredge. *General Chemistry: Principles, Patterns, and Applications*. Boston: FlatWorld, 2.1 edition, 2020.
- L. M. Barge, Y. Abedian, M. J. Russell, I. J. Doloboff, J. H. E. Cartwright, R. D. Kidd, and I. Kanik. From Chemical Gardens to Fuel Cells: Generation of Electrical Potential and Current Across Self-Assembling Iron Mineral Membranes. *Angew. Chem.*, 127(28):8302–8305, July 2015. doi: 10.1002/ange.201501663.
- H. C. Berg. *Random Walks in Biology*. Princeton: Princeton University Press, 1993.
- L. V. Berkner and L. C. Marshall. On the Origin and Rise of Oxygen Concentration in the Earth’s Atmosphere. *J. Atmos. Sci.*, 22(3):225–261, May 1965. doi: 10.1175/1520-0469(1965)022<0225:OTOARO>2.0.CO;2.
- G. O. Bozdog, E. Libby, R. Pineau, C. T. Reinhard, and W. C. Ratcliff. Oxygen suppression of macroscopic multicellularity. *Nat. Commun.*, 12:2838, Jan. 2021. doi: 10.1038/s41467-021-23104-0.

- N. J. Butterfield. Animals and the invention of the Phanerozoic Earth system. *Trends Ecol. Evol.*, 26(2):81–87, 2011. doi: 10.1016/j.tree.2010.11.012.
- D. E. Canfield. *Oxygen: A Four Billion Year History*. Science Essentials. Princeton: Princeton University Press, 2014.
- D. C. Catling and J. F. Kasting. *Atmospheric Evolution on Inhabited and Lifeless Worlds*. Cambridge: Cambridge University Press, 2017.
- D. C. Catling, C. R. Glein, K. J. Zahnle, and C. P. McKay. Why O₂ Is Required by Complex Life on Habitable Planets and the Concept of Planetary “Oxygenation Time”. *Astrobiology*, 5(3):415–438, June 2005. doi: 10.1089/ast.2005.5.415.
- M. Christodoulou, T. O’Hara, A. F. Hugall, S. Khodami, C. F. Rodrigues, A. Hilario, A. Vink, and P. Martinez Arbizu. Unexpected high abyssal ophiuroid diversity in polymetallic nodule fields of the northeast Pacific Ocean and implications for conservation. *Biogeosciences*, 17(7):1845–1876, Apr. 2020. doi: 10.5194/bg-17-1845-2020.
- C. F. Chyba and K. P. Hand. Life Without Photosynthesis. *Science*, 292(5524): 2026–2027, 2001. doi: 10.1126/science.1060081.
- A. Clarke. *Principles of Thermal Ecology: Temperature, Energy and Life*. Oxford: Oxford University Press, 2017.
- P. Cloud. Beginnings of biospheric evolution and their biogeochemical consequences. *Paleobiology*, 2(4):351–387, Jan. 1976. doi: 10.1017/S009483730000498X.
- P. E. Cloud Jr. Atmospheric and Hydrospheric Evolution on the Primitive Earth. *Science*, 160(3829):729–736, May 1968. doi: 10.1126/science.160.3829.729.

- D. B. Cole, D. B. Mills, D. H. Erwin, E. A. Sperling, S. M. Porter, C. T. Reinhard, and N. J. Planavsky. On the co-evolution of surface oxygen levels and animals. *Geobiology*, 18(3):260–281, May 2020. doi: 10.1111/gbi.12382.
- K. J. A. Davies. Oxidative stress: the paradox of aerobic life. *Biochem. Soc. Symp.*, 61:1–31, 1995. doi: 10.1042/bss0610001.
- F. C. De Leo, C. R. Smith, A. A. Rowden, D. A. Bowden, and M. R. Clark. Submarine canyons: hotspots of benthic biomass and productivity in the deep sea. *Proc. R. Soc. B*, 277(1695):2783–2792, 2010. doi: 10.1098/rspb.2010.0462.
- D. W. Deamer. *Assembling Life: How Can Life Begin on Earth and Other Habitable Planets?* Oxford: Oxford University Press, 2019.
- M. Denny. *Air and Water: The Biology and Physics of Life’s Media*. Princeton: Princeton University Press, 1993.
- J. M. Durden, M. Putts, S. Bingo, A. B. Leitner, J. C. Drazen, A. J. Gooday, D. O. B. Jones, A. K. Sweetman, T. W. Washburn, and C. R. Smith. Megafaunal Ecology of the Western Clarion Clipperton Zone. *Front. Mar. Sci.*, 8:671062, 2021. doi: 10.3389/fmars.2021.671062.
- W. W. Fischer, J. Hemp, and J. E. Johnson. Evolution of Oxygenic Photosynthesis. *Annu. Rev. Earth Planet. Sci.*, 44:647–683, June 2016. doi: 10.1146/annurev-earth-060313-054810.
- E. J. Gaidos, K. H. Nealson, and J. L. Kirschvink. Life in Ice-Covered Oceans. *Science*, 284(5420):1631–1633, 1999. doi: 10.1126/science.284.5420.1631.
- D. S. Glazier. Beyond the ‘3/4-power law’: variation in the intra- and inter-specific scaling of metabolic rate in animals. *Biol. Rev.*, 80(4):611–662, 2005. doi: 10.1017/S1464793105006834.

- D. S. Glazier. Variable metabolic scaling breaks the law: from ‘Newtonian’ to ‘Darwinian’ approaches. *Proc. R. Soc. B*, 289(1985):20221605, 2022. doi: 10.1098/rspb.2022.1605.
- A. J. Gooday, M. Holzmann, C. Cauille, A. Goineau, O. Kamenskaya, A. A. T. Weber, and J. Pawlowski. Giant protists (xenophyophores, Foraminifera) are exceptionally diverse in parts of the abyssal eastern Pacific licensed for polymetallic nodule exploration. *Biol. Conserv.*, 207:106–116, Mar. 2017. doi: 10.1016/j.biocon.2017.01.006.
- R. Greenberg. Transport Rates of Radiolytic Substances into Europa’s Ocean: Implications for the Potential Origin and Maintenance of Life. *Astrobiology*, 10(3):275–283, Apr. 2010. doi: 10.1089/ast.2009.0386.
- B. Halliwell and J. M. C. Gutteridge. *Free Radicals in Biology and Medicine*. Oxford: Oxford University Press, 5th edition, 2015.
- P. Han and D. M. Bartels. Temperature Dependence of Oxygen Diffusion in H₂O and D₂O. *J. Phys. Chem.*, 100(13):5597–5602, 1996. doi: 10.1021/jp952903y.
- E. N. Harvey. The oxygen consumption of luminous bacteria. *J. Gen. Physiol.*, 11(5):469–475, 1928. doi: 10.1085/jgp.11.5.469.
- J. R. Hein, A. Koschinsky, and T. Kuhn. Deep-ocean polymetallic nodules as a resource for critical materials. *Nat. Rev. Earth Environ.*, 1(3):158–169, Feb. 2020. doi: 10.1038/s43017-020-0027-0.
- A. R. Hendrix, T. A. Hurford, L. M. Barge, M. T. Bland, J. S. Bowman, W. Brinckerhoff, B. J. Buratti, M. L. Cable, J. Castillo-Rogez, G. C. Collins, S. Diniega, C. R. German, A. G. Hayes, T. Hoehler, S. Hosseini, C. J. A. Howett, A. S. McEwen, C. D. Neish, M. Neveu, T. A. Nordheim, G. W. Patterson, D. A. Patthoff, C. Phillips, A. Rhoden, B. E. Schmidt, K. N. Singer,

- J. M. Soderblom, and S. D. Vance. The NASA Roadmap to Ocean Worlds. *Astrobiology*, 19(1):1–27, Jan. 2019. doi: 10.1089/ast.2018.1955.
- T. M. Hoehler, D. J. Mankel, P. R. Girguis, T. M. McCollom, N. Y. Kiang, and B. B. Jørgensen. The metabolic rate of the biosphere and its components. *Proc. Natl. Acad. Sci.*, 120(25):e2303764120, June 2023. doi: 10.1073/pnas.2303764120.
- A. F. Hofmann, E. T. Peltzer, P. M. Walz, and P. G. Brewer. Hypoxia by degrees: Establishing definitions for a changing ocean. *Deep Sea Res. I: Oceanogr. Res.*, 58(12):1212–1226, Dec. 2011. doi: 10.1016/j.dsr.2011.09.004.
- C. G. Jones, J. H. Lawton, and M. Shachak. Organisms as Ecosystem Engineers. *Oikos*, 69(3):373–386, Apr. 1994. doi: 10.2307/3545850.
- O. P. Judson. The energy expansions of evolution. *Nat. Ecol. Evol.*, 1(6):0138, 2017. doi: 10.1038/s41559-017-0138.
- J. F. Kasting, D. P. Whitmire, and R. T. Reynolds. Habitable Zones around Main Sequence Stars. *Icarus*, 101(1):108–128, Jan. 1993. doi: 10.1006/icar.1993.1010.
- T. Kjørboe. *A Mechanistic Approach to Plankton Ecology*. Princeton: Princeton University Press, 2008.
- A. H. Knoll. The precambrian evolution of terrestrial life. In M. D. Papagiannis, editor, *The Search for Extraterrestrial Life: Recent Developments*, volume 112 of *IAU Symposium*, pages 201–211. Dordrecht: Springer, Jan. 1985. doi: 10.1007/978-94-009-5462-5_29.
- A. H. Knoll. The Multiple Origins of Complex Multicellularity. *Annu. Rev. Earth Planet. Sci.*, 39:217–239, May 2011. doi: 10.1146/annurev.earth.031208.100209.

- A. H. Knoll. *Life on a Young Planet: The First Three Billion Years of Evolution on Earth*. Princeton Science Library. Princeton University Press, 2nd edition, 2015.
- A. H. Knoll and M. A. Nowak. The timetable of evolution. *Sci. Adv.*, 3(5): e1603076, May 2017. doi: 10.1126/sciadv.1603076.
- A. H. Knoll and E. A. Sperling. Oxygen and animals in Earth history. *Proc. Natl. Acad. Sci.*, 111(11):3907–3908, Mar. 2014. doi: 10.1073/pnas.1401745111.
- A. L. Koch. What size should a bacterium be? A question of scale. *Annu. Rev. Microbiol.*, 50(1):317–348, 1996. doi: 10.1146/annurev.micro.50.1.317.
- L. G. Koch and S. L. Britton. Aerobic metabolism underlies complexity and capacity. *J. Physiol.*, 586(1):83–95, 2008. doi: 10.1089/ast.2006.0119.
- R. K. Kopparapu, R. Ramirez, J. F. Kasting, V. Eymet, T. D. Robinson, S. Mahadevan, R. C. Terrien, S. Domagal-Goldman, V. Meadows, and R. Deshpande. Habitable Zones around Main-sequence Stars: New Estimates. *Astrophys. J.*, 765(2):131, Mar. 2013. doi: 10.1088/0004-637X/765/2/131.
- N. Lane. *Oxygen: The Molecule that Made the World*. Oxford: Oxford University Press, 2002.
- Z. Lee, A. Weidemann, J. Kindle, R. Arnone, K. L. Carder, and C. Davis. Euphotic zone depth: Its derivation and implication to ocean-color remote sensing. *J. Geophys. Res. Oceans*, 112(C3):C03009, Mar 2007. doi: 10.1029/2006JC003802.
- L. A. Levin, C. L. Huggett, and K. F. Wishner. Control of deep-sea benthic community structure by oxygen and organic-matter gradients in the eastern Pacific Ocean. *J. Mar. Res.*, 49(4):763–800, 1991.

- M. Lingam and A. Balbi. *From Stars to Life: A Quantitative Approach to Astrobiology*. Cambridge: Cambridge University Press, 2024. URL <https://cambridge.org/9781009411219>.
- M. Lingam and A. Loeb. Constraints on Aquatic Photosynthesis for Terrestrial Planets around Other Stars. *Astrophys. J. Lett.*, 889(1):L15, Jan. 2020. doi: 10.3847/2041-8213/ab6a14.
- M. Lingam and A. Loeb. *Life in the Cosmos: From Biosignatures to Technosignatures*. Cambridge: Harvard University Press, 2021.
- M. Lingam, A. Balbi, and S. M. Mahajan. A Bayesian Analysis of Technological Intelligence in Land and Oceans. *Astrophys. J.*, 945(1):23, Mar. 2023. doi: 10.3847/1538-4357/acb6fa.
- D. R. Lovley. Electrotrophy: Other microbial species, iron, and electrodes as electron donors for microbial respirations. *Bioresour. Technol.*, 345:126553, Feb. 2022. doi: 10.1016/j.biortech.2021.126553.
- P. L. Luisi. *The Emergence of Life: From Chemical Origins to Synthetic Biology*. Cambridge: Cambridge University Press, 2nd edition, 2016.
- T. W. Lyons, C. T. Reinhard, and N. J. Planavsky. The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488):307–315, Feb. 2014. doi: 10.1038/nature13068.
- T. W. Lyons, C. W. Diamond, N. J. Planavsky, C. T. Reinhard, and C. Li. Oxygenation, Life, and the Planetary System during Earth's Middle History: An Overview. *Astrobiology*, 21(8):906–923, Aug. 2021. doi: 10.1089/ast.2020.2418.
- A. M. Makarieva, V. G. Gorshkov, B.-L. Li, S. L. Chown, P. B. Reich, and V. M. Gavrillov. Mean mass-specific metabolic rates are strikingly similar

- across life's major domains: evidence for life's metabolic optimum. *Proc. Natl. Acad. Sci.*, 105(44):16994–16999, 2008. doi: 10.1073/pnas.0802148105.
- W. Martin, J. Baross, D. Kelley, and M. J. Russell. Hydrothermal vents and the origin of life. *Nat. Rev. Microbiol.*, 6(11):805–814, 2008. doi: 10.1038/nrmicro1991.
- T. M. McCollom. Geochemical Constraints on Sources of Metabolic Energy for Chemolithoautotrophy in Ultramafic-Hosted Deep-Sea Hydrothermal Systems. *Astrobiology*, 7(6):933–950, Dec. 2007. doi: 10.1089/ast.2006.0119.
- J. J. Middelburg. *Marine Carbon Biogeochemistry: A Primer for Earth System Scientists*. Springer Briefs in Earth System Sciences. Cham: Springer, 2019. doi: 10.1007/978-3-030-10822-9.
- B. J. W. Mills, A. J. Krause, I. Jarvis, and B. D. Cramer. Evolution of Atmospheric O₂ Through the Phanerozoic, Revisited. *Annual Review of Earth and Planetary Sciences*, 51:253–276, May 2023. doi: 10.1146/annurev-earth-032320-095425.
- D. B. Mills, L. M. Ward, C. Jones, B. Sweeten, M. Forth, A. H. Treusch, and D. E. Canfield. Oxygen requirements of the earliest animals. *Proc. Natl. Acad. Sci.*, 111(11):4168–4172, Mar. 2014. doi: 10.1073/pnas.1400547111.
- R. Milo and R. Phillips. *Cell Biology by the Numbers*. New York: Garland Science, 2016.
- P. Mitchell. Coupling of Phosphorylation to Electron and Hydrogen Transfer by a Chemi-Osmotic type of Mechanism. *Nature*, 191(4784):144–148, July 1961. doi: 10.1038/191144a0.
- P. Mitchell. Keilin's Respiratory Chain Concept and Its Chemiosmotic Con-

- sequences. *Science*, 206(4423):1148–1159, Dec. 1979. doi: 10.1126/science.388618.
- S. J. Mojzsis. Habitable potentials. *Nat. Astron.*, 5:1083–1085, Nov. 2021. doi: 10.1038/s41550-021-01529-3.
- K. B. Muchowska, S. J. Varma, and J. Moran. Nonenzymatic Metabolic Reactions and Life’s Origins. *Chem. Rev.*, 120(15):7708–7744, 2020. doi: 10.1021/acs.chemrev.0c00191.
- D. G. Nicholls and S. J. Ferguson. *Bioenergetics*. London: Academic Press, 4th edition, 2013.
- F. Nimmo and R. T. Pappalardo. Ocean worlds in the outer solar system. *J. Geophys. Res. Planets*, 121(8):1378–1399, Aug. 2016. doi: 10.1002/2016JE005081.
- J. R. Nursall. Oxygen as a Prerequisite to the Origin of the Metazoa. *Nature*, 183(4669):1170–1172, Apr. 1959. doi: 10.1038/1831170b0.
- J. L. Payne, C. R. McClain, A. G. Boyer, J. H. Brown, S. Finnegan, M. Kowalewski, R. A. Krause Jr., S. K. Lyons, D. W. McShea, P. M. Novack-Gottshall, F. A. Smith, P. Spaeth, J. A. Stempien, and S. C. Wang. The evolutionary consequences of oxygenic photosynthesis: a body size perspective. *Photosynth. Res.*, 107(1):37–57, Jan. 2011. doi: 10.1007/s11120-010-9593-1.
- G. Pillot, O. Amin Ali, S. Davidson, L. Shintu, A. Godfroy, Y. Combet-Blanc, P. Bonin, and P.-P. Liebgott. Identification of enriched hyperthermophilic microbial communities from a deep-sea hydrothermal vent chimney under electrolithoautotrophic culture conditions. *Sci. Rep.*, 11:14782, July 2021. doi: 10.1038/s41598-021-94135-2.
- G. Pillot, Ó. Santiago, S. Kerzenmacher, and P.-P. Liebgott. Spark of Life: Role

- of Electrotrophy in the Emergence of Life. *Life*, 13(2):356, Jan. 2023. doi: 10.3390/life13020356.
- M. Preiner, S. Asche, S. Becker, H. C. Betts, A. Boniface, E. Camprubi, K. Chandru, V. Erastova, S. G. Garg, N. Khawaja, G. Kostyrka, R. Machné, G. Moggioli, K. B. Muchowska, S. Neukirchen, B. Peter, E. Pichlhöfer, Á. Radványi, D. Rossetto, A. Salditt, N. M. Schmelling, F. L. Sousa, F. D. K. Tria, D. Vörös, and J. C. Xavier. The Future of Origin of Life Research: Bridging Decades-Old Divisions. *Life*, 10(3):20, Feb. 2020. doi: 10.3390/life10030020.
- M. Rabone, J. H. Wiethase, E. Simon-Lledó, A. M. Emery, D. O. B. Jones, T. G. Dahlgren, G. Bribiesca-Contreras, H. Wiklund, T. Horton, and A. G. Glover. How many metazoan species live in the world’s largest mineral exploration region? *Curr. Biol.*, 33(12):2383–2396, 2023. doi: 10.1016/j.cub.2023.04.052.
- S. P. S. Rao, S. Alonso, L. Rand, T. Dick, and K. Pethe. The protonmotive force is required for maintaining ATP homeostasis and viability of hypoxic, nonreplicating *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci.*, 105(33): 11945–11950, Aug. 2008. doi: 10.1073/pnas.0711697105.
- M. A. Rex, R. J. Etter, J. S. Morris, J. Crouse, C. R. McClain, N. A. Johnson, C. T. Stuart, J. W. Deming, R. Thies, and R. Avery. Global bathymetric patterns of standing stock and body size in the deep-sea benthos. *Mar. Ecol. Prog. Ser.*, 317:1–8, July 2006. doi: 10.3354/meps317001.
- R. J. Ritchie, A. W. D. Larkum, and I. Ribas. Could photosynthesis function on Proxima Centauri b? *Int. J. Astrobiol.*, 17(2):147–176, Apr 2018. doi: 10.1017/S1473550417000167.
- S. E. Ruff, P. Humez, I. H. de Angelis, M. Diao, M. Nightingale, S. Cho, L. Connors, O. O. Kuloyo, A. Seltzer, S. Bowman, S. D. Wankel, C. N. McClain,

- B. Mayer, and M. Strous. Hydrogen and dark oxygen drive microbial productivity in diverse groundwater ecosystems. *Nat. Commun.*, 14:3194, June 2023. doi: 10.1038/s41467-023-38523-4.
- B. Runnegar. The Cambrian explosion: Animals or fossils? *J. Geol. Soc. Australia*, 29(3):395–411, Oct. 1982. doi: 10.1080/00167618208729222.
- B. Runnegar. Precambrian oxygen levels estimated from the biochemistry and physiology of early eukaryotes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 97(1-2):97–111, Dec. 1991. doi: 10.1016/0031-0182(91)90186-U.
- M. Russell. Green Rust: The Simple Organizing ‘Seed’ of All Life? *Life*, 8(3):35, Aug. 2018. doi: 10.3390/life8030035.
- M. J. Russell. The “Water Problem” (sic), the Illusory Pond and Life’s Submarine Emergence—A Review. *Life*, 11(5):429, 2021. doi: 10.3390/life11050429.
- M. J. Russell. A self-sustaining serpentinization mega-engine feeds the fougérite nanoengines implicated in the emergence of guided metabolism. *Front. Microbiol.*, 14:1145915, 2023. doi: 10.3389/fmicb.2023.1145915.
- M. J. Russell and A. J. Hall. The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *J. Geol. Soc.*, 154(3):377–402, May 1997. doi: 10.1144/gsjgs.154.3.0377.
- M. J. Russell, L. M. Barge, R. Bhartia, D. Bocanegra, P. J. Bracher, E. Branscomb, R. Kidd, S. McGlynn, D. H. Meier, W. Nitschke, T. Shibuya, S. Vance, L. White, and I. Kanik. The Drive to Life on Wet and Icy Worlds. *Astrobiology*, 14(4):308–343, Apr. 2014. doi: 10.1089/ast.2013.1110.
- N. Sahai, H. Kaddour, and P. Dalai. The Transition from Geochemistry to Biogeochemistry. *Elements*, 12(6):389–394, Dec. 2016. doi: 10.2113/gselements.12.6.389.

- R. Sander. Modeling Atmospheric Chemistry: Interactions between Gas-Phase Species and Liquid Cloud/Aerosol Particles. *Surv. Geophys.*, 20(1):1–31, Jan. 1999. doi: 10.1023/A:1006501706704.
- R. Sander. Compilation of Henry’s law constants (version 5.0.0) for water as solvent. *Atmos. Chem. Phys.*, 23(19):10901–12440, Oct. 2023. doi: 10.5194/acp-23-10901-2023.
- J. L. Sarmiento and N. Gruber. *Ocean Biogeochemical Dynamics*. Princeton University Press, 2006.
- D. D. Sasselov, J. P. Grotzinger, and J. D. Sutherland. The origin of life as a planetary phenomenon. *Sci. Adv.*, 6(6):eaax3419, Feb. 2020. doi: 10.1126/sciadv.aax3419.
- J. W. P. Schmelzer, E. D. Zanotto, and V. M. Fokin. Pressure dependence of viscosity. *J. Chem. Phys.*, 122(7):074511–074511, Feb. 2005. doi: 10.1063/1.1851510.
- H. N. Schulz and B. B. Jørgensen. Big Bacteria. *Annu. Rev. Microbiol.*, 55(1): 105–137, 2001. doi: 10.1146/annurev.micro.55.1.105.
- L. Schwander, M. Brabender, N. Mrnjavac, J. L. E. Wimmer, M. Preiner, and W. F. Martin. Serpentinization as the source of energy, electrons, organics, catalysts, nutrients and pH gradients for the origin of LUCA and life. *Front. Microbiol.*, 14:1257597, 2023. doi: 10.3389/fmicb.2023.1257597.
- E. Simon-Lledó, B. J. Bett, V. A. I. Huvenne, T. Schoening, N. M. A. Benoist, R. M. Jeffreys, J. M. Durden, and D. O. B. Jones. Megafaunal variation in the abyssal landscape of the Clarion Clipperton Zone. *Prog. Oceanogr.*, 170: 119–133, Jan. 2019. doi: 10.1016/j.pocean.2018.11.003.

- K. L. Smith, M. Messié, T. P. Connolly, and C. L. Huffard. Decadal Time-Series Depletion of Dissolved Oxygen at Abyssal Depths in the Northeast Pacific. *Geophys. Res. Lett.*, 49(24):e2022GL101018, Dec. 2022. doi: 10.1029/2022GL101018.
- N. Soga, K. Kinoshita Jr., M. Yoshida, and T. Suzuki. Efficient ATP synthesis by thermophilic *Bacillus* F_oF₁-ATP synthase. *FEBS J.*, 278(15):2647–2654, 2011. doi: 10.1111/j.1742-4658.2011.08191.x.
- V. Sojo, B. Herschy, A. Whicher, E. Camprubí, and N. Lane. The Origin of Life in Alkaline Hydrothermal Vents. *Astrobiology*, 16(2):181–197, Feb. 2016. doi: 10.1089/ast.2015.1406.
- E. A. Sperling, C. A. Frieder, A. V. Raman, P. R. Girguis, L. A. Levin, and A. H. Knoll. Oxygen, ecology, and the Cambrian radiation of animals. *Proc. Natl. Acad. Sci.*, 110(33):13446–13451, Aug. 2013a. doi: 10.1073/pnas.1312778110.
- E. A. Sperling, G. P. Halverson, A. H. Knoll, F. A. Macdonald, and D. T. Johnston. A basin redox transect at the dawn of animal life. *Earth Planet. Sci. Lett.*, 371:143–155, June 2013b. doi: 10.1016/j.epsl.2013.04.003.
- E. A. Sperling, T. H. Boag, M. I. Duncan, C. R. Endriga, J. A. Marquez, D. B. Mills, P. M. Monarrez, J. A. Sclafani, R. G. Stockey, and J. L. Payne. Breathless through Time: Oxygen and Animals across Earth’s History. *Biol. Bull.*, 243(2):184–206, 2022. doi: 10.1086/721754.
- A. K. Sweetman, A. J. Smith, D. S. W. de Jonge, T. Hahn, P. Schroedl, M. Silverstein, C. Andrade, R. L. Edwards, A. J. M. Lough, C. Woulds, W. B. Homoky, A. Koschinsky, S. Fuchs, T. Kuhn, F. Geiger, and J. J. Marlow. Evidence of dark oxygen production at the abyssal seafloor. *Nat. Geosci.*, 17: 737–739, 2024. doi: 10.1038/s41561-024-01480-8.

- K. Uhlenkott, K. Meyn, A. Vink, and P. Martínez Arbizu. A review of megafauna diversity and abundance in an exploration area for polymetallic nodules in the eastern part of the Clarion Clipperton Fracture Zone (North East Pacific), and implications for potential future deep-sea mining in this area. *Mar. Biodivers.*, 53(2):22, Apr. 2023. doi: 10.1007/s12526-022-01326-9.
- S. D. Vance, K. P. Hand, and R. T. Pappalardo. Geophysical controls of chemical disequilibria in Europa. *Geophys. Res. Lett.*, 43(10):4871–4879, May 2016. doi: 10.1002/2016GL068547.
- L. M. Ward, V. Stamenković, K. Hand, and W. W. Fischer. Follow the Oxygen: Comparative Histories of Planetary Oxygenation and Opportunities for Aerobic Life. *Astrobiology*, 19(6):811–824, June 2019. doi: 10.1089/ast.2017.1779.
- D. Yahalomi, S. D. Atkinson, M. Neuhof, E. S. Chang, H. Philippe, P. Cartwright, J. L. Bartholomew, and D. Huchon. A cnidarian parasite of salmon (Myxozoa: Henneguya) lacks a mitochondrial genome. *Proc. Natl. Acad. Sci.*, 117(10):5358–5363, Mar. 2020. doi: 10.1073/pnas.1909907117.
- M. Yamamoto, R. Nakamura, K. Oguri, S. Kawagucci, K. Suzuki, K. Hashimoto, and K. Takai. Generation of Electricity and Illumination by an Environmental Fuel Cell in Deep-Sea Hydrothermal Vents. *Angew. Chem.*, 125(41):10958–10961, Oct. 2013. doi: 10.1002/ange.201302704.
- M. Yamamoto, R. Nakamura, T. Kasaya, H. Kumagai, K. Suzuki, and K. Takai. Spontaneous and Widespread Electricity Generation in Natural Deep-Sea Hydrothermal Fields. *Angew. Chem.*, 129(21):5819–5822, May 2017. doi: 10.1002/ange.201701768.
- M. Yamamoto, Y. Takaki, H. Kashima, M. Tsuda, A. Tanizaki, R. Nakamura, and K. Takai. In situ electrosynthetic bacterial growth using electricity gen-

erated by a deep-sea hydrothermal vent. *ISME J.*, 17(1):12–20, Jan. 2023.
doi: 10.1038/s41396-022-01316-6.