

Equivalent Characterizations of the Aubin Property for Nonlinear Semidefinite Programming*

Liang Chen[†] Ruoning Chen[‡] Defeng Sun[§] Liping Zhang[¶]

August 15 2024

Abstract

In this paper, we study the Aubin property of the Karush-Kuhn-Tucker solution mapping for the nonlinear semidefinite programming (NLSDP) problem at a locally optimal solution. In the literature, it is known that the Aubin property implies the constraint nondegeneracy by Fusek [SIAM J. Optim. 23 (2013), pp. 1041-1061] and the second-order sufficient condition by Ding et al. [SIAM J. Optim. 27 (2017), pp. 67-90]. Based on the Mordukhovich criterion, here we further prove that the strong second-order sufficient condition is also necessary for the Aubin property to hold. Consequently, several equivalent conditions including the strong regularity are established for NLSDP's Aubin property. Together with the recent progress made by Chen et al. on the equivalence between the Aubin property and the strong regularity for nonlinear second-order cone programming [arXiv:2406.13798v1 (2024)], this paper constitutes a significant step forward in characterizing the Aubin property for general non-polyhedral C^2 -cone reducible constrained optimization problems.

Keywords: Nonlinear semidefinite programming, Aubin property, Strong second-order sufficient condition, Constraint nondegeneracy, Strong regularity

MSCcodes: 49J53, 90C22, 90C31, 90C46

1 Introduction

Consider the constrained optimization problem

$$\min_{x \in \mathcal{X}} f(x) \quad \text{s.t.} \quad G(x) \in \mathcal{K}, \quad (1.1)$$

where \mathcal{X} and \mathcal{Y} are two finite-dimensional real Hilbert spaces each endowed with an inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\| \cdot \|$, $f : \mathcal{X} \rightarrow \mathbb{R}$ and $G : \mathcal{X} \rightarrow \mathcal{Y}$ are twice continuously differentiable functions, and $\mathcal{K} \subseteq \mathcal{Y}$ is a closed convex set. The Lagrangian function of (1.1) is defined by

$$\mathcal{L}(x, y) := f(x) + \langle y, G(x) \rangle, \quad (x; y) \in \mathcal{X} \times \mathcal{Y}, \quad (1.2)$$

*This work was funded by the National Key R & D Program of China (No. 2021YFA001300), the National Natural Science Foundation of China (Nos. 12271150 and 12171271), the Hunan Provincial Natural Science Foundation of China (No. 2023JJ10001), the Science and Technology Innovation Program of Hunan Province (No. 2022RC1190), the RGC Senior Research Fellow Scheme (No. SRFS2223-5S02), and the GRF Projects (Nos. 15307822 and 15307523).

[†]School of Mathematics, Hunan University, Changsha, China (chl@hnu.edu.cn).

[‡]Department of Mathematical Sciences, Tsinghua University, Beijing, China; and Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong (crn22@mails.tsinghua.edu.cn, ruoning.chen@polyu.edu.hk).

[§]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong (defeng.sun@polyu.edu.hk).

[¶]Department of Mathematical Sciences, Tsinghua University, Beijing, China (lipingzhang@tsinghua.edu.cn).

and the first-order optimality condition of (1.1) is characterized by the Karush-Kuhn-Tucker (KKT) system

$$0 = \nabla_x \mathcal{L}(x, y) \quad \text{and} \quad y \in \mathcal{N}_{\mathcal{K}}(G(x)), \quad (1.3)$$

where $\nabla_x \mathcal{L}(x, y)$ denotes the adjoint of $\mathcal{L}'_x(x, y)$, the partial derivative of \mathcal{L} with respect to $x \in \mathcal{X}$, and $\mathcal{N}_{\mathcal{K}}$ denotes the normal cone of \mathcal{K} in convex analysis [37]. For any solution $(\bar{x}; \bar{y}) \in \mathcal{X} \times \mathcal{Y}$ of the KKT system (1.3), we say that \bar{x} is a stationary point of (1.1) and \bar{y} is a (Lagrange) multiplier at \bar{x} . The set of all multipliers at \bar{x} is denoted by $\mathcal{M}(\bar{x})$.

The canonically perturbed version [6, Section 5.1.3] of (1.1) is given by

$$\min_{x \in \mathcal{X}} f(x) - \langle a, x \rangle \quad \text{s.t.} \quad G(x) - b \in \mathcal{K}, \quad (1.4)$$

where $a \in \mathcal{X}$ and $b \in \mathcal{Y}$ are the perturbation parameters. With \mathcal{L} being the Lagrangian function defined by (1.2), the KKT system of (1.4), as a perturbed KKT system of (1.1), is given by

$$a = \nabla_x \mathcal{L}(x, y) \quad \text{and} \quad y \in \mathcal{N}_{\mathcal{K}}(G(x) - b). \quad (1.5)$$

Then, one can associate (1.5) with the solution mapping

$$\mathbb{S}_{\text{KKT}}(a, b) := \{(x; y) \mid a = \nabla_x \mathcal{L}(x, y), y \in \mathcal{N}_{\mathcal{K}}(G(x) - b)\}. \quad (1.6)$$

As a core research topic in perturbation analysis of optimization problems, how the solution set $\mathbb{S}_{\text{KKT}}(a, b)$ of (1.5) varies along with (a, b) around the origin has been studied for a long time [6, 13, 14, 19, 22, 32, 40]. In a landmark paper of Robinson [34], the definition of strong regularity was introduced to extend the inverse and implicit function theorems to generalized equations. Note that the KKT system (1.3) can be equivalently reformulated to the generalized equation

$$0 \in \Phi(x, y) := \begin{pmatrix} \nabla_x \mathcal{L}(x, y) \\ -G(x) \end{pmatrix} + \begin{pmatrix} 0 \\ \mathcal{N}_{\mathcal{K}}^{-1}(y) \end{pmatrix}. \quad (1.7)$$

Then, we say that $(\bar{x}; \bar{y})$ is a strongly regular solution of generalized equation (1.7) (or the KKT system (1.3)) if the inverse of the set-valued mapping

$$\bar{\Phi}(x, y) := \begin{pmatrix} \nabla_{xx}^2 \mathcal{L}(\bar{x}, \bar{y})(x - \bar{x}) + \nabla G(\bar{x})(y - \bar{y}) \\ -G(\bar{x}) - G'(\bar{x})(x - \bar{x}) \end{pmatrix} + \begin{pmatrix} \mathcal{N}_{\mathcal{X}}(x) \\ \mathcal{N}_{\mathcal{K}}^{-1}(y) \end{pmatrix}$$

has a Lipschitz continuous single-valued localization around $(0; 0) \in \mathcal{X} \times \mathcal{Y}$ for $(\bar{x}; \bar{y})$, where $\nabla_{xx}^2 \mathcal{L}$ is the partial Hessian of \mathcal{L} with respect to x . According to [11], such a strong regularity condition is equivalent to the condition that the solution mapping \mathbb{S}_{KKT} has a Lipschitz continuous single-valued localization around $((0; 0); (\bar{x}; \bar{y}))$, or equivalently, the mapping Φ in (1.7) is strongly metrically regular at $(\bar{x}; \bar{y})$ for $(0; 0)$. Another significant yet less restrictive concept in studying the variation of \mathbb{S}_{KKT} with respect to the perturbation is the Aubin property (c.f. [40, Section 9F] or [13, Section 3E] for a systematic introduction), which was originally called the ‘‘pseudo-Lipschitzian’’ property by Aubin [1]. The Aubin property of \mathbb{S}_{KKT} holds at $(0; 0) \in \mathcal{X} \times \mathcal{Y}$ for $(\bar{x}; \bar{y}) \in \mathbb{S}_{\text{KKT}}(0, 0)$ if there exist a constant $\kappa > 0$ and open neighborhoods \mathcal{U} of $(0; 0)$ and \mathcal{V} of $(\bar{x}; \bar{y})$ such that

$$\mathbb{S}_{\text{KKT}}(a', b') \cap \mathcal{V} \subseteq \mathbb{S}_{\text{KKT}}(a, b) + \kappa \|(a', b') - (a, b)\| \mathbb{B}_{\mathcal{X} \times \mathcal{Y}} \quad \forall (a, b), (a', b') \in \mathcal{U},$$

where $\mathbb{B}_{\mathcal{X} \times \mathcal{Y}}$ denotes the closed unit ball in $\mathcal{X} \times \mathcal{Y}$ centered at the origin. It is easy to see from definition that the Aubin property of \mathbb{S}_{KKT} holds at $(0; 0)$ for $(\bar{x}; \bar{y})$ if $(\bar{x}; \bar{y})$ is a strongly regular solution to (1.3). Moreover, such an Aubin property is equivalent to the metric regularity or the linear openness of Φ at (\bar{x}, \bar{y}) (e.g. [40, Theorem 9.43]).

Since the solution mapping \mathbb{S}_{KKT} is implicitly defined, verifying both the strong regularity and the Aubin property from their definitions is generally unachievable. Consequently, equivalent characterizations for the two conditions have evolved into a central topic in optimization theory and variational analysis. For conventional nonlinear programming with \mathcal{K} in (1.1) being a convex polyhedral cone, such equivalent characterizations have been well established for about

three decades. Specifically, Robinson [34] defined the strong second-order sufficient condition (SSOSC) for the nonlinear programming problem. Moreover, at a stationary point \bar{x} satisfying the linear independence constraint qualification (LICQ), he also showed in [34, Theorem 4.1] that the KKT system is strongly regular at $(\bar{x}; \bar{y}) \in \mathbb{S}_{\text{KKT}}$ if the SSOSC holds at $(\bar{x}; \bar{y})$. At the same time, Kojima [25] introduced the concept of strong stability to stationary points of the nonlinear programming problem, and showed in [25, Theorems 6.4 & 6.5] that for a locally optimal solution satisfying the LICQ, it is strongly stable if and only if the SSOSC holds. It was later revealed by Jongen et al. [20, Section 3] that the strong regularity and the strong stability are equivalent for stationary points of nonlinear programming where the LICQ holds. Furthermore, the strong regularity was characterized by Kummer [26] based on a generalized “implicit function theorem” on nonsmooth equations, and one may refer to [21] and [24] for a related approach, especially [24, Theorem 4.3] for a survey of equivalent characterizations. By combining the results of [34], [25], and [24], one has that at a locally optimal solution of the nonlinear programming problem, the strong regularity is equivalent to the condition that both the SSOSC and the LICQ hold (c.f. [5, Remark 4.11]). Such a result is also available in [5, Theorem 4.10] and [6, Proposition 5.38]. In addition to these equivalent characterizations of the strong regularity, a surprising result of Dontchev and Rockafellar [12, Theorems 1, 4, & 5] for the nonlinear programming problem is that the Aubin property of \mathbb{S}_{KKT} holds at $(0; 0)$ for $(\bar{x}; \bar{y}) \in \mathbb{S}_{\text{KKT}}$ with \bar{x} being a locally optimal solution can imply that the strong regularity of the KKT system (1.3) at $(\bar{x}; \bar{y})$. Consequently, a comprehensive class of equivalent characterizations of the Aubin property for the nonlinear programming problem was achieved.

When the set \mathcal{K} in (1.1) is no longer a polyhedral set, characterizing the two concepts becomes much more involved. A more realistic setting is that \mathcal{K} is C^2 -cone reducible (c.f. [6, Definition 3.135]), which is practical enough for encompassing many important classes of optimization problems, including the nonlinear programming, the nonlinear second-order cone programming (NLSOCP), and the nonlinear semidefinite programming (NLSDP) [41]. In this setting, the perturbation analysis of problem (1.1) has been extensively studied [2, 3, 4, 10, 42, 43], and characterizing the strong regularity via second-order optimality conditions has been recognized as a prominent topic. Although deriving a result akin to [6, Proposition 5.38] for problem (1.1) in its general form has not been achieved, achievements have been made for the most representative classes of problems with significant importance in the form of (1.1).

According to [6, Theorem 5.24], the strong regularity of the KKT system (1.3) at a solution $(\bar{x}; \bar{y})$ implies that the constraint nondegeneracy condition holds at \bar{x} (or \bar{x} is nondegenerate, c.f. (2.13)). With the help of this result, Bonans and Ramírez [4] established a counterpart of [6, Proposition 5.38] for the NLSOCP problem. For the NLSDP problem, Sun [42] defined the SSOSC by introducing an approximation set, and finally obtained a collection of equivalent characterizations of the strong regularity condition, including the SSOSC accompanied by the constraint nondegeneracy. These results properly extend the characterizations of the strong regularity from the conventional nonlinear programming to problem (1.1) with \mathcal{K} being a non-polyhedral set. Nevertheless, for both the NLSOCP and the NLSDP problems, obtaining such an extension for characterizing the Aubin property has been an open question for a long time.

The first step in addressing this issue was achieved by Outrata and Ramírez [31] (and the erratum by Opazo, Outrata, and Ramírez [30]). They proved that, for a nondegenerate locally optimal solution \bar{x} of the NLSOCP problem (i.e., (1.1) with \mathcal{K} being the Cartesian product of second-order cones), the Aubin property of a solution mapping (akin to (3.1)) at $0 \in \mathcal{X}$ for \bar{x} can imply the SSOSC, but under an assumption regarding the strict complementarity. Recently, Chen et al. [7] finalized this conclusion by removing these assumptions, and showed that, with $\bar{y} \in \mathcal{M}(\bar{x})$, the Aubin property of \mathbb{S}_{KKT} at $(0; 0)$ for $(\bar{x}; \bar{y})$ is equivalent to the strong regularity of $(\bar{x}; \bar{y})$ to the KKT system, constituting a counterpart of the seminal result of Dontchev and Rockafellar [12] for the NLSOCP. As a result, the extension of the characterizations for the Aubin property is realized from the nonlinear programming problem to the NLSOCP problem.

In this paper, we study the equivalent characterizations of the Aubin property for the NLSDP problem

$$\min_{x \in \mathcal{X}} f(x) \quad \text{s.t.} \quad h(x) = 0, \quad g_j(x) \in \mathcal{S}_+^{p_j}, \quad j = 1, \dots, J, \quad (1.8)$$

where $h : \mathcal{X} \rightarrow \mathbb{R}^m$ and $g_j : \mathcal{X} \rightarrow \mathcal{S}_+^{p_j}$, $j = 1, \dots, J$ are twice continuously differentiable functions, \mathcal{S}^{p_j} is the linear space of $p_j \times p_j$ real symmetric matrices endowed with the inner product $\langle A, B \rangle := \text{trace}(AB)$ for $A, B \in \mathcal{S}^{p_j}$, where $\text{trace}(\cdot)$ denotes the sum of the diagonal elements, and $\mathcal{S}_+^{p_j}$ is the closed convex cone of all positive semidefinite matrices in \mathcal{S}^{p_j} . When $J = 1$ in (1.8), the strong regularity of the corresponding KKT system is equivalent to the SSOSC together with the constraint nondegeneracy [42, Theorem 4.1]. For the case that $J > 1$, it is easy to see from the analysis in [42] that such an equivalence still holds. Moreover, in the case that $J = 1$, Fusek [15] showed that the Aubin property at $(0; 0)$ for a solution $(\bar{x}; \bar{y})$ can imply the constraint nondegeneracy at \bar{x} . Such a result was later extended by Klatte and Kummer [23] to the general case of problem (1.1).

The main contribution of this paper is that, at a locally optimal solution \bar{x} of (1.8) with $\bar{y} \in \mathcal{M}(\bar{x})$, we prove that the Aubin property of \mathbb{S}_{KKT} at $(0; 0)$ for $(\bar{x}; \bar{y})$ implies the SSOSC. We achieve this by designing an auxiliary optimization problem and fully exploiting its properties, and using the Mordukhovich criterion for characterizing the Aubin property [27]. We should emphasize that the key tools we used in this paper are essentially different from those in [7] for the nonlinear second-order cone programming. Specifically, the main progress in [7] is based on a lemma of alternative choices on cones, and the fact that the spectral factorization associated with second-order cones admits only two “eigenvalues” is indispensable. This approach can be used here when $\max_j \{p_j\} \leq 3$, but does not apply to the general cases. Consequently, it is not hard to see that the tools in [7] are far from sufficient for deriving a counterpart of its main result in the setting of the NLSDP problem (1.8). Based on the equivalence between the Aubin property and the strong regularity established in this paper, we finally obtain a comprehensive set of equivalent characterizations of the Aubin property for the NLSDP problem (1.8).

The remaining parts of this paper are organized as follows. Section 2 introduces the notation, the definitions, and the preliminary results used throughout this paper. In Section 3, we study the implications of the Aubin property of the solution mapping $\mathbb{S}_{\text{KKT}}(a, b)$, especially the SSOSC. The equivalent characterizations of the Aubin property are given in Section 4. We conclude this paper in Section 5.

2 Notation and preliminaries

Let \mathcal{E} and \mathcal{F} be two finite-dimensional real Hilbert spaces each endowed with an inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\|\cdot\|$. Given a set of vectors $\{z_1, \dots, z_r\} \subset \mathcal{E}$, we use $\text{span}\{z_1, \dots, z_r\}$ to denote the linear subspace it spans. The notation $(\cdot; \cdot)$ signifies the arrangement of two vectors or linear operators symbolically in a column-wise manner. The inner product on $\mathcal{E} \times \mathcal{F}$ is defined by $\langle (z_1; z'_1), (z_2; z'_2) \rangle := \langle z_1, z_2 \rangle + \langle z'_1, z'_2 \rangle$ for all $(z_1; z_2), (z'_1; z'_2) \in \mathcal{E} \times \mathcal{F}$, and the norm $\|\cdot\|$ on $\mathcal{E} \times \mathcal{F}$ is induced by this inner product. For a vector $z \in \mathcal{E}$ (or a subspace $\mathcal{E}_0 \subseteq \mathcal{E}$), z^\perp (or \mathcal{E}_0^\perp) denotes its orthogonal complement in \mathcal{E} . Given a cone $C \subseteq \mathcal{E}$, $C^\circ := \{v \in \mathcal{E} \mid \langle v, z \rangle \leq 0 \forall z \in C\}$ is the polar cone of C . For a linear operator $\mathcal{A} : \mathcal{E} \rightarrow \mathcal{F}$, we use \mathcal{A}^* , $\text{rge}(\mathcal{A})$ and $\ker(\mathcal{A})$ to denote its adjoint, range space, and null space, respectively. Note that $\text{rge}(\mathcal{A}) = (\ker(\mathcal{A}^*))^\perp$. For a continuously differentiable function $\psi : \mathcal{E} \rightarrow \mathcal{F}$, we use $\psi'(z)$ to denote the Fréchet derivative or the Jacobian of ψ at $z \in \mathcal{E}$, and define $\nabla\psi(z) := (\psi'(z))^*$.

Given a matrix $A \in \mathbb{R}^{l \times q}$, we use A_{ik} to denote the entry at the i -th row and the k -th column of A , and use A_k to denote the k -th column of A . The transpose of A is denoted by A^\top . When $l = q$, we use A^{-1} to denote the inverse of A if it is nonsingular. Given a subset $S \subseteq \{1, \dots, q\}$, we use $|S|$ to denote its cardinality and use A_S to denote the sub-matrix of A by eliminating all the columns that are not indexed by S from A . For the given index sets $I \subseteq \{1, \dots, l\}$ and $S \subseteq \{1, \dots, q\}$, we use A_{IS} to denote the $|I| \times |S|$ sub-matrix of A by removing all the rows and columns not in I and S , respectively. Given two matrices $A, B \in \mathbb{R}^{l \times q}$, $A \circ B$ denotes their Hadamard product. The inner product of A, B is defined by $\langle A, B \rangle = \text{trace}(AB)$, and $\|A\| = \sqrt{\langle A, A \rangle}$ is the Frobenius norm. We use E to denote an all-ones matrix, whose dimension will be specified from the context. For a matrix $A \in \mathcal{S}^p$, we use $A \succ 0$ (or $A \succeq 0$) to say that A is positive definite (or positive semidefinite). Moreover, $A \prec 0$ (or $A \preceq 0$) means that $-A \succ 0$ (or $-A \succeq 0$).

Given a set $C \subseteq \mathcal{E}$, we use $\text{lin}(C)$ to denote the largest linear subspace contained in C (the lineality space of C), $\text{aff}(C)$ to denote the smallest linear subspace that contains C (the affine hull of C), and $\text{cl}(C)$ to denote the closure of C . The paratingent cone of C at \bar{z} is defined by

$$\mathcal{T}_C^P(\bar{z}) := \limsup_{z \xrightarrow{C} \bar{z}, t \searrow 0} \frac{C - z}{t},$$

where “lim sup” is the outer limit in Painlevé-Kuratowski convergence for subsets, and $z \xrightarrow{C} \bar{z}$ means that $z \rightarrow \bar{z}$ with $z \in C$. The regular (Fréchet) normal cone of C at $\bar{z} \in C$ is defined by

$$\widehat{\mathcal{N}}_C(\bar{z}) := \{v \in \mathcal{E} \mid \langle v, z - \bar{z} \rangle \leq o(\|z - \bar{z}\|) \ \forall z \in C\}$$

and the limiting (Mordukhovich) normal cone of C at $\bar{z} \in C$ is defined by

$$\mathcal{N}_C(\bar{z}) := \limsup_{z \xrightarrow{C} \bar{z}} \widehat{\mathcal{N}}_C(z).$$

When C is a closed convex set, one has $\widehat{\mathcal{N}}_C(\bar{z})$ coincides with $\mathcal{N}_C(\bar{z})$, and both of them are simply called the normal cone of C at \bar{z} (in convex analysis [37]), i.e.,

$$\mathcal{N}_C(\bar{z}) = \begin{cases} \{v \in \mathcal{E} \mid \langle v, z - \bar{z} \rangle \leq 0 \ \forall z \in \mathcal{E}\}, & \text{if } \bar{z} \in C, \\ \emptyset, & \text{otherwise.} \end{cases}$$

When C is a closed convex set, the tangent cone $\mathcal{T}_C(\bar{z})$ of C at $\bar{z} \in C$ can be defined by $\mathcal{T}_C(\bar{z}) := (\mathcal{N}_C(\bar{z}))^\circ$ [40, Example 6.24].

Given a function $\psi : \mathcal{E} \rightarrow (-\infty, \infty]$ with $\bar{z} \in \mathcal{E}$ such that $\psi(\bar{z})$ is finite, according to [40, Theorem 8.9] the (limiting) subdifferential of ψ at \bar{z} can be defined as

$$\partial\psi(\bar{z}) := \{v \mid (v, -1) \in \mathcal{N}_{\text{epi}\psi}(\bar{z}; \psi(\bar{z}))\}, \quad (2.1)$$

where $\text{epi}\psi := \{(z; t) \mid t \geq \psi(z)\}$ is the epigraph of ψ .

Given a set-valued mapping $\Psi : \mathcal{E} \rightrightarrows \mathcal{F}$, we use $\text{gph}\Psi \in \mathcal{E} \times \mathcal{F}$ to denote the the graph of Ψ , i.e.,

$$\text{gph}\Psi := \{(z; w) \in \mathcal{E} \times \mathcal{F} \mid w \in \Psi(z)\}.$$

The strict graphical (paringent) derivative of Ψ at $(\bar{z}; \bar{w}) \in \text{gph}\Psi$ is defined by

$$\mathcal{D}_*\Psi(\bar{z}, \bar{w})(u) = \{v \in \mathcal{F} \mid (u; v) \in \mathcal{T}_{\text{gph}\Psi}^P(\bar{z}; \bar{w})\}, \quad u \in \mathcal{E}. \quad (2.2)$$

Meanwhile, the limiting coderivative of Ψ at $(\bar{z}; \bar{w}) \in \text{gph}\Psi$ is defined by

$$\mathcal{D}^*\Psi(\bar{z}, \bar{w})(v) := \{u \in \mathcal{E} \mid (u; -v) \in \mathcal{N}_{\text{gph}\Psi}(\bar{z}; \bar{w})\}, \quad v \in \mathcal{F}. \quad (2.3)$$

The set-valued mapping $\Psi : \mathcal{E} \rightrightarrows \mathcal{F}$ is said to have the Aubin property at $\bar{z} \in \mathcal{E}$ for $\bar{w} \in \Psi(\bar{z})$ if there exist a constant $\kappa > 0$ and open neighborhoods \mathcal{U} of \bar{z} and \mathcal{V} of \bar{w} such that

$$\Psi(z) \cap \mathcal{V} \subset \Psi(z') + \kappa\|z - z'\|\mathbb{B}_{\mathcal{F}} \quad \forall z, z' \in \mathcal{U},$$

where $\mathbb{B}_{\mathcal{F}}$ is the closed unit ball in \mathcal{F} . Moreover, it has been perfectly characterized by the Mordukhovich criterion [27] that, under the assumption that $\text{gph}\Psi$ is locally closed around $(\bar{z}; \bar{w}) \in \text{gph}\Psi$, the Aubin property of Ψ holds at \bar{z} for \bar{w} if and only if $\mathcal{D}^*\Psi(\bar{z}, \bar{w})(0) = \{0\}$.

2.1 Coderivative related to positive semidefinite cone

This subsection briefly introduces the explicit formula of the limiting coderivative of the normal cone mapping to \mathcal{S}_+^p , which was calculated by Ding et al. [9]. Let $A \in \mathcal{S}^p$ be given. It admits an eigenvalue decomposition in the form of

$$A = P\Lambda P^\top = P \begin{pmatrix} \Lambda_{\alpha\alpha} & & \\ & 0_{\beta\beta} & \\ & & \Lambda_{\gamma\gamma} \end{pmatrix} P^\top \quad \text{with } P = (P_\alpha, P_\beta, P_\gamma), \quad (2.4)$$

where $P \in \mathbb{R}^{p \times p}$ is an orthogonal matrix and Λ is a diagonal matrix with non-increasing diagonal elements such that $\Lambda_{\alpha\alpha} \succ 0$ and $\Lambda_{\gamma\gamma} \prec 0$. Here α, β and γ are three sets of indices with the cardinalities $|\alpha|$, $|\beta|$ and $|\gamma|$. One has $P_\alpha \in \mathbb{R}^{p \times |\alpha|}$, $P_\beta \in \mathbb{R}^{p \times |\beta|}$ and $P_\gamma \in \mathbb{R}^{p \times |\gamma|}$. We use $\Pi_{\mathcal{S}_+^p}(A)$ to denote the metric projection of A to the cone \mathcal{S}_+^p (under the Frobenius norm). For convenience, we denote $A_+ := \Pi_{\mathcal{S}_+^p}(A)$ and $A_- := A - A_+ \in -\mathcal{S}_+^p$. Then, it is obvious that $|\alpha|$ and $|\gamma|$ are the ranks of A_+ and A_- , respectively. Recall that \mathcal{S}_+^p is a closed convex cone, and one has

$$\mathcal{T}_{\mathcal{S}_+^p}(A_+) = \{Z \in \mathcal{S}^p \mid P_{\beta \cup \gamma}^\top Z P_{\beta \cup \gamma} \succeq 0\}.$$

Consequently, it holds that

$$\text{lin}(\mathcal{T}_{\mathcal{S}_+^p}(A_+)) = \{Z \in \mathcal{S}^p \mid P_\beta^\top Z P_\beta = 0, P_\beta^\top Z P_\gamma = 0, P_\gamma^\top Z P_\gamma = 0\}.$$

Denote the set of all partitions of the index set β by \mathcal{P}_β . Let $\mathbb{R}_{\geq}^{|\beta|}$ be the set of all the vectors in $\mathbb{R}^{|\beta|}$ whose components are arranged in non-increasing order, i.e.,

$$\mathbb{R}_{\geq}^{|\beta|} := \{z \in \mathbb{R}^{|\beta|} \mid z_1 \geq z_2 \geq \dots \geq z_{|\beta|}\}.$$

For any $z \in \mathbb{R}_{\geq}^{|\beta|}$, one can define $D(z) \in \mathbb{R}^{|\beta| \times |\beta|}$ as the matrix whose elements $(D(z))_{ik}$, $i, k \in \{1, \dots, |\beta|\}$, are given by

$$(D(z))_{ik} := \begin{cases} \frac{\max\{z_i, 0\} - \max\{z_k, 0\}}{z_i - z_k} \in [0, 1], & \text{if } z_i \neq z_k, \\ 1, & \text{if } z_i = z_k > 0, \\ 0, & \text{if } z_i = z_k \leq 0. \end{cases}$$

Define the set

$$\Upsilon_{|\beta|} := \left\{ \bar{Z} \in \mathcal{S}^{|\beta|} \mid \bar{Z} = \lim_{k \rightarrow \infty} D(z^k), z^k \rightarrow 0, z^k \in \mathbb{R}_{\geq}^{|\beta|} \right\} \subseteq \mathcal{S}^{|\beta|}. \quad (2.5)$$

Let $\Xi_1 \in \Upsilon_{|\beta|}$. Then, there exists a partition $\pi(\beta) := (\beta_+, \beta_0, \beta_-) \in \mathcal{P}_\beta$ such that

$$\Xi_1 = \begin{pmatrix} E_{\beta_+ \beta_+} & E_{\beta_+ \beta_0} & (\Xi_1)_{\beta_+ \beta_-} \\ E_{\beta_+ \beta_0}^\top & 0 & 0 \\ (\Xi_1)_{\beta_+ \beta_-} & 0 & 0 \end{pmatrix}, \quad (2.6)$$

where each element of $(\Xi_1)_{\beta_+ \beta_-}$ belongs to the interval $[0, 1]$. Moreover, based on Ξ_1 one can define the matrix

$$\Xi_2 := \begin{pmatrix} 0 & 0 & E_{\beta_+ \beta_-} - (\Xi_1)_{\beta_+ \beta_-} \\ 0 & 0 & E_{\beta_0 \beta_-} \\ (E_{\beta_+ \beta_-} - (\Xi_1)_{\beta_+ \beta_-})^\top & E_{\beta_0 \beta_-}^\top & E_{\beta_- \beta_-} \end{pmatrix}. \quad (2.7)$$

Building up on the above definitions, the coderivative of $\mathcal{N}_{\mathcal{S}_+^p}$ can be explicitly given in the following result.

Lemma 2.1. [9, Theorem 3.1 & Proposition 3.3] *Suppose that $A \in \mathcal{S}^p$ has the eigenvalue decomposition in (2.4). Then, $U \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^p}(A_+, A_-)(V)$ if and only if*

$$U = P \begin{pmatrix} 0 & 0 & \tilde{U}_{\alpha\gamma} \\ 0 & \tilde{U}_{\beta\beta} & \tilde{U}_{\beta\gamma} \\ \tilde{U}_{\gamma\alpha} & \tilde{U}_{\gamma\beta} & \tilde{U}_{\gamma\gamma} \end{pmatrix} P^\top \quad \text{and} \quad V = P \begin{pmatrix} \tilde{V}_{\alpha\alpha} & \tilde{V}_{\alpha\beta} & \tilde{V}_{\alpha\gamma} \\ \tilde{V}_{\beta\alpha} & \tilde{V}_{\beta\beta} & 0 \\ \tilde{V}_{\gamma\alpha} & 0 & 0 \end{pmatrix} P^\top \quad (2.8)$$

with

$$\tilde{U}_{\beta\beta} \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta|}}(0, 0)(\tilde{V}_{\beta\beta}) \quad \text{and} \quad \Sigma_{\alpha\gamma} \circ \tilde{U}_{\alpha\gamma} - (E_{\alpha\gamma} - \Sigma_{\alpha\gamma}) \circ \tilde{V}_{\alpha\gamma} = 0, \quad (2.9)$$

where $\tilde{U} := P^\top U P \in \mathcal{S}^p$, $\tilde{V} := P^\top V P \in \mathcal{S}^p$, and $\Sigma \in \mathbb{R}^{p \times p}$ is the matrix defined by

$$\Sigma_{ik} := \frac{\max\{\lambda_i, 0\} - \max\{\lambda_k, 0\}}{\lambda_i - \lambda_k}, \quad i, k \in \{1, \dots, p\}.$$

In addition, $\tilde{U}_{\beta\beta} \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta|}}(0, 0)(\tilde{V}_{\beta\beta})$ holds if and only if there exist a matrix $\Xi_1 \in \Upsilon_{|\beta|}$ in (2.5) and an orthogonal matrix $O \in \mathbb{R}^{|\beta| \times |\beta|}$ such that

$$\Xi_1 \circ O^\top \tilde{U}_{\beta\beta} O = \Xi_2 \circ O^\top \tilde{V}_{\beta\beta} O, \quad O_{\beta_0}^\top \tilde{U}_{\beta\beta} O_{\beta_0} \preceq 0, \quad \text{and} \quad O_{\beta_0}^\top \tilde{V}_{\beta\beta} O_{\beta_0} \preceq 0, \quad (2.10)$$

where $(\beta_+, \beta_0, \beta_-) \in \mathcal{P}(\beta)$ is a partition such that Ξ_1 takes the form of (2.6) and Ξ_2 is given by (2.7).

We make the following remark to Lemma 2.1.

Remark 2.1. For the case that $A = 0$ in Lemma 2.1, one has $|\beta| = p$. Thus, by taking P as the identity matrix, one can get $\tilde{V}_{\beta\beta} = V$. One can further take $\beta_+ = \beta$ and $\beta_- = \beta_0 = \emptyset$ to get a partition of β . In this case, one has from (2.6) and (2.7) that $\Xi_1 = E_{\beta\beta}$ and $\Xi_2 = 0 \in \mathcal{S}^{|\beta|}$. Then, one has that (2.10) holds with $\tilde{U}_{\beta\beta} := 0 \in \mathcal{S}^{|\beta|}$ for any orthogonal matrix $O \in \mathbb{R}^{|\beta| \times |\beta|}$. Consequently, one has $0 \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^p}(0, 0)(V)$ for any $V \in \mathcal{S}^p$.

2.2 Second-order sufficient conditions

In this subsection, we briefly introduce the second-order (sufficient) optimality conditions of the NLSDP (1.8), which can be viewed as an extension of the discussions in [42].

Let \bar{x} be a stationary point of the NLSDP (1.8) and $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$ be a multiplier at \bar{x} , where $\bar{\zeta} \in \mathbb{R}^m$ and $\bar{\Gamma}_j \in \mathcal{S}^{p_j}$ for all $j = 1, \dots, J$. From the KKT system (1.3) we know that $g_j(\bar{x}) \in \mathcal{S}_+^{p_j}$ and $\bar{\Gamma}_j \in \mathcal{N}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x}))$. For convenience, define $g(\bar{x}) := (g_1(\bar{x}); \dots; g_J(\bar{x}))$ and $A_j := g_j(\bar{x}) + \bar{\Gamma}_j$, $j = 1, \dots, J$. Then one has $(A_j)_+ = g_j(\bar{x})$ and $(A_j)_- = \bar{\Gamma}_j$. Moreover, one can write the eigenvalue decomposition of A_j as in (2.4) with $P_j = ((P_j)_{\alpha_j}, (P_j)_{\beta_j}, (P_j)_{\gamma_j})$ being the corresponding orthogonal matrix and Λ^j being the corresponding diagonal matrix of eigenvalues. Then, the critical cone of (1.8) at \bar{x} can be explicitly given as

$$\mathcal{C}(\bar{x}) = \{d \mid h'(\bar{x})d = 0, g'_j(\bar{x})d \in \mathcal{C}(A_j; \mathcal{S}_+^{p_j}), j = 1, \dots, J\}, \quad (2.11)$$

where for any $j = 1, \dots, J$,

$$\begin{aligned} \mathcal{C}(A_j; \mathcal{S}_+^{p_j}) &:= \mathcal{T}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x})) \cap \bar{\Gamma}_j^\perp \\ &= \{Z \in \mathcal{S}^{p_j} \mid (P_j)_{\beta_j}^\top Z (P_j)_{\beta_j} \succeq 0, (P_j)_{\beta_j}^\top Z (P_j)_{\gamma_j} = 0, (P_j)_{\gamma_j}^\top Z (P_j)_{\gamma_j} = 0\}. \end{aligned} \quad (2.12)$$

According to [6, Section 4.6.1], for a feasible point $x \in \mathcal{X}$ of (1.1), it is called nondegenerate [35], or the constraint nondegeneracy [36] holds at x , if

$$G'(x)\mathcal{X} + \text{lin}(\mathcal{T}_\mathcal{K}(G(x))) = \mathcal{Y}, \quad (2.13)$$

The constraint nondegeneracy condition (2.13) at \bar{x} can be written as

$$\begin{pmatrix} h'(\bar{x}) \\ g'(\bar{x}) \end{pmatrix} \mathcal{X} + \left(\prod_{j=1}^J \text{lin}(\mathcal{T}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x}))) \right) = \begin{pmatrix} \mathbb{R}^m \\ \prod_{j=1}^J \mathcal{S}^{p_j} \end{pmatrix}. \quad (2.14)$$

Note that (2.14) implies that $\mathcal{M}(\bar{x})$ is a singleton, i.e., \bar{y} is the unique multiplier at \bar{x} . Meanwhile, by following the proof of [42, Proposition 3.1] one can get that

$$\begin{aligned} \text{aff}(\mathcal{C}(\bar{x})) &= \{d \in \mathcal{X} \mid h'(\bar{x})d = 0, g'_j(\bar{x})d \in \text{aff}(\mathcal{C}(A_j; \mathcal{S}_+^{p_j})), j = 1, \dots, J\} \\ &= \left\{ d \in \mathcal{X} \mid \begin{array}{l} h'(\bar{x})d = 0, (P_j)_{\beta_j}^\top (g'_j(\bar{x})d) (P_j)_{\gamma_j} = 0, \\ (P_j)_{\gamma_j}^\top (g'_j(\bar{x})d) (P_j)_{\gamma_j} = 0, j = 1, \dots, J \end{array} \right\}. \end{aligned} \quad (2.15)$$

As a consequence, the second-order sufficient condition and the SSOSC for the NLSDP problem (1.8) can be defined as follows.

Definition 2.1. Let \bar{x} be a stationary point of (1.8) with $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$. Let $\mathcal{Q} : \mathcal{X} \rightarrow \mathcal{X}$ be the self-adjoint linear operator defined by

$$\langle d, \mathcal{Q}d \rangle = \langle d, \nabla_{xx}^2 \mathcal{L}(\bar{x}, \bar{y})d \rangle - 2 \sum_{j=1}^J \langle \bar{\Gamma}_j, (g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \rangle \quad \forall d \in \mathcal{X}, \quad (2.16)$$

where $(g_j(\bar{x}))^\dagger$ denotes the Moore–Penrose pseudoinverse of $g_j(\bar{x})$ for all $j = 1, \dots, J$. We say that the second-order sufficient condition holds at $(\bar{x}; \bar{y})$ if

$$\langle d, \mathcal{Q}d \rangle > 0 \quad \forall d \in \mathcal{C}(\bar{x}) \setminus \{0\}. \quad (2.17)$$

Moreover, we say that the SSOSC holds at $(\bar{x}; \bar{y})$ if

$$\langle d, \mathcal{Q}d \rangle > 0 \quad \forall d \in \text{aff}(\mathcal{C}(\bar{x})) \setminus \{0\}. \quad (2.18)$$

Remark 2.2. In definition 2.1, the second-order sufficient condition (2.17) follows from [3], while the SSOSC is a straightforward extension of [42, Definition 3.2] from the case that $J = 1$ to the general setting.

2.3 Technical lemmas

In this part, we provide three technical lemmas for our discussions. The first one is about the polar cones.

Lemma 2.2 ([40, Corollary 11.25(d)]). Let $\mathcal{H} : \mathcal{E} \rightarrow \mathcal{F}$ be a linear operator, and $K \subset \mathcal{F}$ be a nonempty closed convex cone. Then the set $C := \{z \in \mathcal{E} \mid \mathcal{A}z \in K\}$ is also a closed convex cone and one has

$$C^\circ = \text{cl} \{ \mathcal{A}^*v \mid v \in K^\circ \},$$

where the closure operation is superfluous if 0 is in the interior of $\{K - \text{rge}\mathcal{H}\}$.

The following lemma gives a variational characterization of self-adjoint positive definite operators. One can also see [18, Proposition 3.1] for a more straightforward proof based on the Moreau decomposition.

Lemma 2.3 ([17, Theorem 3.6]). Let $\mathcal{H} : \mathcal{E} \rightarrow \mathcal{E}$ be an invertible self-adjoint linear operator and C be a closed convex cone in \mathcal{E} . Then

$$\langle z, \mathcal{H}z \rangle > 0 \quad \forall z \in \mathcal{E} \quad \Leftrightarrow \quad \begin{cases} \langle z, \mathcal{H}z \rangle > 0 \quad \forall z \in C \setminus \{0\} \text{ and} \\ \langle z, \mathcal{H}^{-1}z \rangle > 0 \quad \forall z \in C^\circ \setminus \{0\}. \end{cases}$$

The following lemma discusses Robinson's constraint qualification [33], via its equivalent reformulation [6, Proposition 2.97], of a constraint system.

Lemma 2.4. Let $\mathcal{H} : \mathcal{E} \rightarrow \mathcal{F}$ be a linear operator, and $K \subseteq \mathcal{F}$ be a nonempty closed convex cone. Then, Robinson's constraint qualification holds at any feasible point

$$(\bar{z}; \bar{v}) \in \left\{ (z; v) \in \mathcal{E} \times \mathcal{F} \mid z - \mathcal{H}^*v = 0, \frac{1}{2}(\|v\|^2 - 1) = 0, v \in K \right\},$$

in the sense that for any given $(\Delta_z; \delta; \Delta_v) \in \mathcal{E} \times \mathbb{R} \times \mathcal{F}$, one can find $z \in \mathcal{E}$, $v \in \mathcal{F}$ and $u \in \mathcal{T}_K(\bar{v})$ such that

$$\begin{cases} z - \mathcal{H}^*v = \Delta_z, & (2.19a) \\ \langle \bar{v}, v \rangle = \delta, & (2.19b) \\ v + u = \Delta_v. & (2.19c) \end{cases}$$

Proof. Note that $\mathcal{T}_K(\bar{v})$ is a closed convex cone. From [40, Example 6.24] we know that $\mathcal{N}_K(\bar{v}) = (\mathcal{T}_K(\bar{v}))^\circ$. Then by Moreau's decomposition theorem [37, Theorem 31.5] we know that one can uniquely decompose Δ_v by

$$\Delta_v = \Delta'_v + \Delta''_v, \quad \Delta'_v \in \mathcal{T}_K(\bar{v}), \quad \Delta''_v \in \mathcal{N}_K(\bar{v}), \quad \langle \Delta'_v, \Delta''_v \rangle = 0.$$

One can take $v_0 := \Delta''_v \in \mathcal{N}_K(\bar{v})$ and $u_0 = \Delta'_v \in \mathcal{T}_K(\bar{v})$, so that $v_0 + u_0 = \Delta_v$. Moreover, it is easy to see that

$$\lambda \bar{v} \in \mathcal{T}_K(\bar{v}) \quad \text{with} \quad \lambda := \langle \bar{v}, v_0 \rangle - \delta.$$

Then, by letting $v := v_0 - \lambda \bar{v}$, one has $u := u_0 + \lambda \bar{v} \in \mathcal{T}_K(\bar{z})$ and $z := \Delta_z + \mathcal{H}^*v$. Consequently, one has (2.19a) and (2.19c) hold. Moreover, since $\|\bar{v}\| = 1$, it holds that $\langle \bar{v}, v \rangle = \langle \bar{v}, v_0 - \lambda \bar{v} \rangle = \langle \bar{v}, v_0 \rangle - \lambda = \delta$, so that (2.19b) holds. This completes the proof. \square

3 Implications of the Aubin property for NLSDP

In this section, we study the implications of the Aubin property of the solution mapping \mathbb{S}_{KKT} in (1.6) for the NLSDP problem (1.8). Throughout this section, we set $G(x) := (h(x); g(x)) = (h(x); g_1(x); \dots; g_J(x))$, $\mathcal{Y} := \mathbb{R}^m \times \prod_{j=1}^J \mathcal{S}^{p_j}$ and $\mathcal{K} := \{0 \in \mathbb{R}^m\} \times \prod_{j=1}^J \mathcal{S}_+^{p_j}$ in (1.1), and make no distinction between (1.1) and (1.8).

3.1 A reduction method for NLSDP

This part exploits the Aubin property of the solution mapping \mathbb{S}_{KKT} in (1.6). By using a reduction method, we can get more accessible results from the original formula of the Mordukhovich criterion.

Note that if \mathbb{S}_{KKT} in (1.6) has the Aubin property at $(0; 0) \in \mathcal{X} \times \mathcal{Y}$ for $(\bar{x}; \bar{y})$, the mapping \mathbb{S}_{GE} , defined by

$$\mathbb{S}_{\text{GE}}(a) := \{x \in \mathcal{X} \mid a \in \nabla f(x) + \nabla G(x) \mathcal{N}_{\mathcal{K}}(G(x))\}, \quad (3.1)$$

also has the Aubin property at $0 \in \mathcal{X}$ for \bar{x} . Moreover, according to the Mordukhovich criterion [27], the latter Aubin property holds if and only if $\mathcal{D}^* \mathbb{S}_{\text{GE}}(0, \bar{x})(0) = \{0\}$. Since \mathcal{K} is C^2 -cone reducible at every $y \in \mathcal{K}$, one can utilize the second-order chain rule developed in [31, Theorem 7], as a generalization of [29, Theorem 3.4], to get the following result.

Lemma 3.1. *Let \bar{x} be a locally optimal solution of problem (1.8) with $\bar{y} \in \mathcal{M}(\bar{x})$. Suppose that \bar{x} is nondegenerate, i.e., (2.14) holds. Then one has*

$$\mathcal{D}^* \mathbb{S}_{\text{GE}}(0, \bar{x})(0) = \{-d \mid 0 \in \nabla_{xx}^2 \mathcal{L}(\bar{x}, \bar{y})d + \nabla G(\bar{x}) \mathcal{D}^* \mathcal{N}_{\mathcal{K}}(G(\bar{x}), \bar{y})(G'(\bar{x})d)\}.$$

Proof. The proof follows from [31, Theorem 20], in which \mathcal{K} was assumed to be a second-order cone, but it still holds when \mathcal{K} is a closed convex set. \square

Next, we apply Lemmas 2.1 and 3.1 to the NLSDP problem (1.8) to obtain the following result.

Proposition 3.1. *Let \bar{x} be a nondegenerate locally optimal solution of problem (1.8) such that $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$. For $j \in \{1, \dots, J\}$, define $A_j := g_j(\bar{x}) + \bar{\Gamma}_j$ and write its eigenvalue decomposition $A_j = P_j \Lambda^j P_j^\top$ as in (2.4) with $P_j = ((P_j)_{\alpha_j}, (P_j)_{\beta_j}, (P_j)_{\gamma_j})$ being the corresponding orthogonal matrix and Λ^j being the corresponding diagonal matrix of eigenvalues. Then, the mapping \mathbb{S}_{GE} defined by (3.1) has the Aubin property at 0 for \bar{x} if and only if*

$$\begin{aligned} \mathcal{Q}d \notin \text{rge}(\nabla h(\bar{x})) - \left\{ \sum_{j=1}^J \nabla g_j(\bar{x}) (P_j)_{\beta_j \cup \gamma_j} \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} (P_j)_{\beta_j \cup \gamma_j}^\top \right. \\ \left. \begin{array}{l} \tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) ((P_j)_{\beta_j}^\top (g'_j(\bar{x})d) (P_j)_{\beta_j}) \\ \forall \tilde{U}_{\beta_j \gamma_j}^j \in \mathbb{R}^{|\beta_j| \times |\gamma_j|}, \tilde{U}_{\gamma_j \gamma_j}^j \in \mathcal{S}^{|\gamma_j|}, j = 1, \dots, J \end{array} \right\} \quad \forall d \in \text{aff}(\mathcal{C}(\bar{x})) \setminus \{0\}, \end{aligned} \quad (3.2)$$

where \mathcal{Q} is the linear operator defined in (2.16).

Proof. From Lemma 3.1 we know that the mapping \mathbb{S}_{GE} defined by (3.1) has the Aubin property at 0 for \bar{x} if and only if

$$\begin{aligned} \nabla_{xx}^2 \mathcal{L}(\bar{x}, \bar{y})d \notin & -\nabla h(\bar{x})\mathcal{D}^*\mathcal{N}_{\{0\}}(h(\bar{x}), \bar{\zeta})(h'(\bar{x})d) \\ & - \sum_{j=1}^J \nabla g_j(\bar{x})\mathcal{D}^*\mathcal{N}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x}), \bar{\Gamma}_j)(g'_j(\bar{x})d) \quad \forall 0 \neq d \in \mathcal{X}. \end{aligned} \quad (3.3)$$

On the one hand, since $h(\bar{x}) = 0$, it is easy to see from the definition of the coderivative in (2.3) that

$$\mathcal{D}^*\mathcal{N}_{\{0\}}(h(\bar{x}), \bar{\zeta})(h'(\bar{x})d) = \begin{cases} \mathbb{R}^m, & \text{if } h'(\bar{x})d = 0, \\ \emptyset, & \text{otherwise.} \end{cases}$$

On the other hand, for all $j = 1, \dots, J$, by applying Lemma 2.1 to $A_j = g_j(\bar{x}) + \bar{\Gamma}_j$ and $V^j := g'_j(\bar{x})d$ we know that $U^j \in D^*\mathcal{N}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x}), \bar{\Gamma}_j)(g'_j(\bar{x})d)$ holds if and only if $U := U^j$ and $V := g'_j(\bar{x})d$ can be expressed as in (2.8) with $\alpha := \alpha_j$, $\beta := \beta_j$ and $\gamma := \gamma_j$ such that (2.9) holds. Since \bar{x} is nondegenerate, from the formulation of $\text{aff}(\mathcal{C}(\bar{x}))$ given in (2.15) one can see that (3.3) holds if and only if

$$\nabla_{xx}^2 \mathcal{L}(\bar{x}, \bar{y})d \notin \text{rge}(\nabla h(\bar{x})) - \sum_{j=1}^J \nabla g_j(\bar{x})\mathbb{U}_d^j \quad \forall d \in \text{aff}(\mathcal{C}(\bar{x})) \setminus \{0\}, \quad (3.4)$$

where with $\tilde{V}_{\alpha_j \gamma_j}^j := (P_j)_{\alpha_j}^\top (g'_j(\bar{x})d)(P_j)_{\gamma_j}$ and $\tilde{V}_{\beta_j \beta_j}^j := (P_j)_{\beta_j}^\top (g'_j(\bar{x})d)(P_j)_{\beta_j}$, the set \mathbb{U}_d^j is defined by

$$\mathbb{U}_d^j := \left\{ P_j \begin{pmatrix} 0 & 0 & \tilde{U}_{\alpha_j \gamma_j}^j \\ 0 & \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\alpha_j \gamma_j}^j)^\top & (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} P_j^\top \right. \\ \left. \begin{array}{l} \tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^*\mathcal{N}_{\mathcal{S}_+^{p_j}}(0, 0)(\tilde{V}_{\beta_j \beta_j}^j), \\ \Sigma_{\alpha_j \gamma_j}^j \circ \tilde{U}_{\alpha_j \gamma_j}^j = (E_{\alpha_j \gamma_j} - \Sigma_{\alpha_j \gamma_j}^j) \circ \tilde{V}_{\alpha_j \gamma_j}^j, \quad j = 1, \dots, J \end{array} \right\},$$

in which the matrix Σ^j is defined by

$$\Sigma_{ik}^j := \frac{\max\{\lambda_i^j, 0\} - \max\{\lambda_k^j, 0\}}{\lambda_i^j - \lambda_k^j}, \quad i, k \in \{1, \dots, p_j\}, \quad j = 1, \dots, J, \quad (3.5)$$

where each λ_i^j is the i -th diagonal element of Λ^j . Note that for any $d \in \mathcal{X}$ one has $g'_j(\bar{x})d \in \mathcal{S}^{p_j}$ and $(g_j(\bar{x}))^\dagger \in \mathcal{S}^{p_j}$. Therefore, it holds that

$$\begin{aligned} & 2\langle \bar{\Gamma}_j, (g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \rangle \\ &= \langle \bar{\Gamma}_j (g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger, g'_j(\bar{x})d \rangle + \langle (g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \bar{\Gamma}_j, g'_j(\bar{x})d \rangle, \\ &= \langle \nabla g_j(\bar{x})(\bar{\Gamma}_j (g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger + (g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \bar{\Gamma}_j), d \rangle \quad \forall j = 1, \dots, J. \end{aligned} \quad (3.6)$$

Moreover, for any $d \in \mathcal{X}$ and $j \in \{1, \dots, J\}$, one can get from the eigenvalue decomposition $A_j = P_j \Lambda^j P_j^\top$ that

$$\begin{aligned} (g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \bar{\Gamma}_j &= P_j \begin{pmatrix} (\Lambda_{\alpha_j \alpha_j}^j)^{-1} & & \\ & 0 & \\ & & 0 \end{pmatrix} P_j^\top (g'_j(\bar{x})d) P_j \begin{pmatrix} 0 & & \\ & 0 & \\ & & \Lambda_{\gamma_j \gamma_j}^j \end{pmatrix} P_j^\top \\ &= P_j \begin{pmatrix} 0 & 0 & (\Lambda_{\alpha_j \alpha_j}^j)^{-1} \tilde{V}_{\alpha_j \gamma_j}^j \Lambda_{\gamma_j \gamma_j}^j \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} P_j^\top. \end{aligned} \quad (3.7)$$

Similarly, one also has that

$$\bar{\Gamma}_j(g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger = P_j \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \Lambda_{\gamma_j\gamma_j}^j (\tilde{V}_{\alpha_j\gamma_j}^j)^\top (\Lambda_{\alpha_j\alpha_j}^j)^{-1} & 0 & 0 \end{pmatrix} P_j^\top. \quad (3.8)$$

Then, by putting (3.6), (3.7) and (3.8) together, one can see that for any $d \in \mathcal{X}$ and any $j = 1, \dots, J$, it holds that

$$\begin{aligned} & 2\langle \bar{\Gamma}_j, (g'_j(\bar{x})d)(g_j(\bar{x}))^\dagger (g'_j(\bar{x})d) \rangle \\ &= \left\langle \nabla g_j(\bar{x}) P_j \begin{pmatrix} 0 & 0 & (\Lambda_{\alpha_j\alpha_j}^j)^{-1} \tilde{V}_{\alpha_j\gamma_j}^j \Lambda_{\gamma_j\gamma_j}^j \\ 0 & 0 & 0 \\ \Lambda_{\gamma_j\gamma_j}^j (\tilde{V}_{\alpha_j\gamma_j}^j)^\top (\Lambda_{\alpha_j\alpha_j}^j)^{-1} & 0 & 0 \end{pmatrix} P_j^\top, d \right\rangle. \end{aligned} \quad (3.9)$$

Note that for any $U^j \in \mathbf{U}_d^j$, $j = 1, \dots, J$, it holds that

$$U^j = P_j \begin{pmatrix} 0 & 0 & \tilde{U}_{\alpha_j\gamma_j}^j \\ 0 & \tilde{U}_{\beta_j\beta_j}^j & \tilde{U}_{\beta_j\gamma_j}^j \\ (\tilde{U}_{\alpha_j\gamma_j}^j)^\top & (\tilde{U}_{\beta_j\beta_j}^j)^\top & \tilde{U}_{\gamma_j\gamma_j}^j \end{pmatrix} P_j^\top$$

and $\Sigma_{\alpha_j\gamma_j}^j \circ \tilde{U}_{\alpha_j\gamma_j}^j = (E_{\alpha_j\gamma_j} - \Sigma_{\alpha_j\gamma_j}) \circ \tilde{V}_{\alpha_j\gamma_j}^j$. Then, from the definition of $\Sigma_{\alpha_j\gamma_j}$ in (3.5) one can see that for any $i \in \alpha_j$ and $k \in \gamma_j$ it holds that

$$\frac{\lambda_i^j}{\lambda_i^j - \lambda_k^j} \tilde{U}_{ik}^j + \frac{\lambda_k^j}{\lambda_i^j - \lambda_k^j} \tilde{V}_{ik}^j = 0.$$

Consequently, one has that $\lambda_i^j \tilde{U}_{ik}^j + \lambda_k^j \tilde{V}_{ik}^j = 0$. Therefore, it holds that

$$(\tilde{U}^j)_{\alpha_j\gamma_j} + (\Lambda_{\alpha_j\alpha_j}^j)^{-1} \tilde{V}_{\alpha_j\gamma_j}^j \Lambda_{\gamma_j\gamma_j}^j = 0,$$

which, together with (3.9) and (2.16), implies that (3.2) holds if and only if (3.4) holds. This completes the proof. \square

When the assumptions in Proposition 3.1 hold, one can define the linear operator $\mathcal{B} : \mathcal{X} \rightarrow \mathbb{R}^m \times \prod_{j=1}^J (\mathbb{R}^{|\beta_j| \times |\gamma_j|} \times \mathcal{S}^{|\gamma_j|})$ by

$$\mathcal{B}d := \begin{pmatrix} h'(\bar{x})d; \\ (P_{\beta_1}^\top [g'_1(\bar{x})d] P_{\gamma_1}; P_{\gamma_1}^\top [g'_1(\bar{x})d] P_{\gamma_1}) \\ \vdots \\ (P_{\beta_J}^\top [g'_J(\bar{x})d] P_{\gamma_J}; P_{\gamma_J}^\top [g'_J(\bar{x})d] P_{\gamma_J}) \end{pmatrix}, \quad d \in \mathcal{X}. \quad (3.10)$$

From (2.15) one can see that $\ker(\mathcal{B}) = \text{aff}(\mathcal{C}(\bar{x}))$, which is a finite-dimensional subspace of \mathcal{X} . Let r be the dimension of $\ker(\mathcal{B})$. Then, one can find a collection of linearly independent vectors $\omega_1, \dots, \omega_r \in \mathcal{X}$ such that $\text{span}\{\omega_1, \dots, \omega_r\} = \ker(\mathcal{B})$. Moreover, one can define the linear operator $\mathcal{W} : \mathbb{R}^r \rightarrow \ker(\mathcal{B})$ via

$$\mathcal{W}\nu = \sum_{i=1}^r \nu_i \omega_i, \quad \nu = (\nu_1; \dots; \nu_r) \in \mathbb{R}^r. \quad (3.11)$$

Consequently, one has $\text{rge}(\mathcal{W}) = \ker(\mathcal{B})$. Based on the definition of \mathcal{W} in (3.11) one can get the following result from Proposition 3.1.

Proposition 3.2. *Let \bar{x} be a nondegenerate locally optimal solution of problem (1.8) such that $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$. For $j \in \{1, \dots, J\}$, define $A_j := g_j(\bar{x}) + \bar{\Gamma}_j$ and write its eigenvalue decomposition $A_j = P_j \Lambda^j P_j^\top$ as in (2.4) with $P_j = ((P_j)_{\alpha_j}, (P_j)_{\beta_j}, (P_j)_{\gamma_j})$ being the*

corresponding orthogonal matrix and Λ^j being the corresponding diagonal matrix of eigenvalues. Let \mathcal{Q} be the linear operator defined by (2.16) and \mathcal{W} be the linear operator defined by (3.11). Then, the mapping \mathbb{S}_{GE} defined by (3.1) has the Aubin property at 0 for \bar{x} if and only if

$$\mathcal{W}^* \mathcal{Q} \mathcal{W} \nu \notin \left\{ \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top \right. \\ \left. \mid \tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0)((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \nu)(P_j)_{\beta_j}) \right\} \quad \forall \nu \in \mathbb{R}^r \setminus \{0\}. \quad (3.12)$$

Proof. From proposition 3.1 we know that the the mapping \mathbb{S}_{GE} defined by (3.1) for the NLSDP problem (1.8) has the Aubin property at 0 for \bar{x} if and only if (3.2) holds. Note that \mathcal{W} in (3.11) is well-defined and $\text{rge}(\mathcal{W}) = \text{aff}(\mathcal{C}(\bar{x}))$. Therefore, such an Aubin property of \mathbb{S}_{GE} holds if and only if

$$\mathcal{Q} \mathcal{W} \nu \notin \text{rge}(\nabla h(\bar{x})) - \left\{ \sum_{j=1}^J \nabla g_j(\bar{x})(P_j)_{\beta_j \cup \gamma_j} \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} (P_j)_{\beta_j \cup \gamma_j}^\top \right. \\ \left. \mid \begin{array}{l} \tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0)((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \nu)(P_j)_{\beta_j}) \\ \forall \tilde{U}_{\beta_j \gamma_j}^j \in \mathbb{R}^{|\beta_j| \times |\gamma_j|}, \tilde{U}_{\gamma_j \gamma_j}^j \in \mathcal{S}^{|\gamma_j|}, j = 1, \dots, J \end{array} \right\} \quad \forall \nu \in \mathbb{R}^r \setminus \{0\}. \quad (3.13)$$

Consequently, it is sufficient to prove that (3.13) and (3.12) are equivalent.

First, suppose that (3.12) does not hold, i.e., there exists a nonzero vector $\bar{\nu} \in \mathbb{R}^r$ such that

$$\mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\nu} = - \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top \quad (3.14)$$

with

$$\tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0)((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\nu})(P_j)_{\beta_j}), \quad j = 1, \dots, J.$$

Denote $\bar{\mu} := \mathcal{Q} \mathcal{W} \bar{\nu}$. From (3.14) one has that

$$\bar{\mu} + \sum_{j=1}^J \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top \in \ker(\mathcal{W}^*). \quad (3.15)$$

Recall that the linear operator \mathcal{B} in (3.10) is well-defined. Moreover, for any $d \in \mathcal{X}$, $\xi \in \mathbb{R}^m$, $U_{\beta_j \gamma_j}^j \in \mathbb{R}^{|\beta_j| \times |\gamma_j|}$, $U_{\gamma_j \gamma_j}^j \in \mathcal{S}^{|\gamma_j|}$, $j = 1, \dots, J$, it holds that

$$\begin{aligned} & \left\langle \mathcal{B}^* \left(\xi; (U_{\beta_j \gamma_j}^1; U_{\gamma_j \gamma_j}^1); \dots; (U_{\beta_j \gamma_j}^J; U_{\gamma_j \gamma_j}^J) \right), d \right\rangle \\ &= \left\langle \left(\xi; (U_{\beta_j \gamma_j}^1; U_{\gamma_j \gamma_j}^1); \dots; (U_{\beta_j \gamma_j}^J; U_{\gamma_j \gamma_j}^J) \right), \mathcal{B} d \right\rangle \\ &= \langle \xi, h'(\bar{x}) d \rangle \\ & \quad + \sum_{j=1}^J \langle U_{\beta_j \gamma_j}^j, (P_j)_{\beta_j}^\top [g'_j(\bar{x}) d] (P_j)_{\gamma_j} \rangle + \sum_{j=1}^J \langle U_{\gamma_j \gamma_j}^j, (P_j)_{\gamma_j}^\top [g'_j(\bar{x}) d] (P_j)_{\gamma_j} \rangle \\ &= \langle \xi, h'(\bar{x}) d \rangle \\ & \quad + \sum_{j=1}^J \left\langle (P_j)_{\beta_j} U_{\beta_j \gamma_j}^j (P_j)_{\gamma_j}^\top, g'_j(\bar{x}) d \right\rangle + \sum_{j=1}^J \left\langle (P_j)_{\gamma_j} U_{\gamma_j \gamma_j}^j (P_j)_{\gamma_j}^\top, g'_j(\bar{x}) d \right\rangle \\ &= \left\langle \nabla h(\bar{x}) \xi + \sum_{j=1}^J \nabla g(\bar{x})(P_j)_{\beta_j} U_{\beta_j \gamma_j}^j (P_j)_{\gamma_j}^\top + \sum_{j=1}^J \nabla g(\bar{x})(P_j)_{\gamma_j} U_{\gamma_j \gamma_j}^j (P_j)_{\gamma_j}^\top, d \right\rangle. \end{aligned} \quad (3.16)$$

Note that $\ker(\mathcal{W}^*) = \text{rge}(\mathcal{B}^*) = (\ker(\mathcal{B}))^\perp$. Then by combining (3.15) and (3.16) one can get that

$$\begin{aligned} \bar{\mu} \in \text{rge}(\nabla h(\bar{x})) - \left\{ \sum_{j=1}^J \nabla g_j(\bar{x})(P_j)_{\beta_j \cup \gamma_j} \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} (P_j)_{\beta_j \cup \gamma_j}^\top \right. \\ \left. \begin{array}{l} \tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) ((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\nu}) (P_j)_{\beta_j}) \\ \forall \tilde{U}_{\beta_j \gamma_j}^j \in \mathbb{R}^{|\beta_j| \times |\gamma_j|}, \tilde{U}_{\gamma_j \gamma_j}^j \in \mathcal{S}^{|\gamma_j|}, j = 1, \dots, J \end{array} \right\}, \end{aligned}$$

which contradicts to (3.13). Therefore, (3.13) implies (3.12).

Next, suppose that (3.13) does not hold, i.e., there exist two vectors $\bar{\nu} \in \mathbb{R}^r$ and $\xi \in \mathbb{R}^m$, the matrices

$$\tilde{U}_{\beta_j \gamma_j}^j \in \mathbb{R}^{|\beta_j| \times |\gamma_j|}, \quad \tilde{U}_{\gamma_j \gamma_j}^j \in \mathcal{S}^{|\gamma_j|}, \quad j = 1, \dots, J,$$

and the matrices

$$\tilde{U}_{\beta_j \beta_j}^j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) ((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\nu}) (P_j)_{\beta_j}), \quad j = 1, \dots, J,$$

such that

$$\mathcal{Q} \mathcal{W} \bar{\nu} = \nabla h(\bar{x}) \xi - \sum_{j=1}^J \nabla g_j(\bar{x})(P_j)_{\beta_j \cup \gamma_j} \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} (P_j)_{\beta_j \cup \gamma_j}^\top.$$

Note that for any vector $\nu \in \mathbb{R}^r$, it holds that $\mathcal{W} \nu \in \ker(\mathcal{B}) = \text{aff}(\mathcal{C}(\bar{x}))$. Then by using (2.15) one has for all $\nu \in \mathbb{R}^r$ it holds that

$$\begin{aligned} & \langle \nu, \mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\nu} \rangle \\ &= \langle \nu, \mathcal{W}^* \nabla h(\bar{x}) \xi \rangle \\ & \quad - \sum_{j=1}^J \left\langle \nu, \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j \cup \gamma_j} \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} (P_j)_{\beta_j \cup \gamma_j}^\top \right\rangle \\ &= \langle h'(\bar{x}) \mathcal{W} \nu, \xi \rangle \\ & \quad - \sum_{j=1}^J \left\langle (P_j)_{\beta_j \cup \gamma_j}^\top (g'_j(\bar{x}) \mathcal{W} \nu) (P_j)_{\beta_j \cup \gamma_j}, \begin{pmatrix} \tilde{U}_{\beta_j \beta_j}^j & \tilde{U}_{\beta_j \gamma_j}^j \\ (\tilde{U}_{\beta_j \gamma_j}^j)^\top & \tilde{U}_{\gamma_j \gamma_j}^j \end{pmatrix} \right\rangle \\ &= - \sum_{j=1}^J \left\langle (P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \nu) (P_j)_{\beta_j}, \tilde{U}_{\beta_j \beta_j}^j \right\rangle \\ &= - \sum_{j=1}^J \left\langle \nu, \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top \right\rangle, \end{aligned}$$

which means that $\mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\nu} = - \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top$. Therefore, (3.12) also implies (3.13). Consequently, (3.12) is equivalent to (3.13), which completes the proof. \square

We can readily get the following result based on Proposition 3.2.

Corollary 3.1. *Suppose that the assumptions in Proposition 3.2 hold. Then the linear operator $\mathcal{W}^* \mathcal{Q} \mathcal{W}$ is a nonsingular matrix in \mathcal{S}^r .*

Proof. Note that $\mathcal{W}^* \mathcal{Q} \mathcal{W} \in \mathcal{S}^r$ holds by definition. Suppose on the contrary that there exists a nonzero vector $\bar{\nu} \in \mathbb{R}^r$ such that $\mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\nu} = 0$. According to the discussions in Remark 2.1 one can set

$$\tilde{U}_{\beta_j \beta_j}^j := 0 \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) ((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\nu}) (P_j)_{\beta_j}), \quad j = 1, \dots, J.$$

As a result, one has $\mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\nu} = 0 = - \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \tilde{U}_{\beta_j \beta_j}^j (P_j)_{\beta_j}^\top$, which contradicts (3.12). Consequently, $\mathcal{W}^* \mathcal{Q} \mathcal{W}$ is nonsingular. \square

3.2 Aubin property implies SSOSC

The following lemma comes from [10, Corollary 25].

Lemma 3.2. *Let \bar{x} be a locally optimal solution of problem (1.8) with $\bar{y} \in \mathcal{M}(\bar{x})$. Suppose that the Aubin property of \mathbb{S}_{KKT} in (1.6) holds at $(0,0)$ for (\bar{x}, \bar{y}) . Then, \bar{x} is nondegenerate, i.e., (2.14) holds, and the second-order sufficient condition (2.17) in Definition 2.1 holds with \mathcal{Q} being defined in (2.16).*

Here, we provide a remarkable improvement to Lemma 3.2.

Proposition 3.3. *Let \bar{x} be a locally optimal solution to the NLSDP problem (1.8) and $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$ be a multiplier at \bar{x} . Suppose that the solution mapping \mathbb{S}_{KKT} in (1.6) has the Aubin property at $(0,0)$ for (\bar{x}, \bar{y}) . Then, the SSOSC (2.18) in Definition 2.1 holds at $(\bar{x}; \bar{y})$ with \mathcal{Q} being defined in (2.16).*

Proof. From Lemma 3.2 we know that the constraint nondegeneracy (2.14) holds, implying that $\mathcal{M}(\bar{x})$ is a singleton, and that the second-order sufficient condition (2.17) holds. For all $j \in \{1, \dots, J\}$, define $A_j := g_j(\bar{x}) + \bar{\Gamma}_j$ and write its eigenvalue decomposition as in (2.4) with $P_j = ((P_j)_{\alpha_j}, (P_j)_{\beta_j}, (P_j)_{\gamma_j})$ being the corresponding orthogonal matrix and Λ^j being the corresponding diagonal matrix of eigenvalues. Then the linear operators \mathcal{B} in (3.10) and \mathcal{W} in (3.11) are well-defined. Consequently, (3.12) in Proposition 3.2 holds.

Define $\mathcal{A} : \mathbb{R}^r \rightarrow \mathcal{S}^{|\beta_1|} \times \dots \times \mathcal{S}^{|\beta_J|}$ by $\mathcal{A}\nu := (\mathcal{A}_1\nu; \dots; \mathcal{A}_J\nu)$, where the linear operators $\mathcal{A}_j : \mathbb{R}^r \rightarrow \mathcal{S}^{|\beta_j|}$, $j = 1, \dots, J$, are defined by

$$\mathcal{A}_j\nu := (P_j)_{\beta_j}^\top (g'_j(\bar{x})\mathcal{W}\nu)(P_j)_{\beta_j}, \quad \nu \in \mathbb{R}^r. \quad (3.17)$$

Then, one can define the closed convex cone $\Omega \subseteq \mathbb{R}^r$ by

$$\Omega := \{\nu \in \mathbb{R}^r \mid \mathcal{A}_j\nu \succeq 0, \quad j = 1, \dots, J\}.$$

Since (2.14) holds, for any given $Y_j \in \mathcal{S}^{p_j}$, $j = 1, \dots, J$, one can always find a vector $d \in \mathcal{X}$ and the matrices

$$\begin{aligned} Z_j &\in \text{lin}(\mathcal{T}_{\mathcal{S}_+^{p_j}}(g_j(\bar{x}))) \\ &= \{Z_j \in \mathcal{S}^{p_j} \mid (P_j)_{\beta_j \cup \gamma_j}^\top Z_j (P_j)_{\beta_j \cup \gamma_j} = 0\}, \quad j = 1, \dots, J \end{aligned}$$

such that

$$\begin{cases} h'(\bar{x})d = 0, \\ g'_j(\bar{x})d + Z_j = P_j \begin{pmatrix} (Y_j)_{\alpha_j \alpha_j} & (Y_j)_{\alpha_j \beta_j} & (Y_j)_{\alpha_j \gamma_j} \\ (Y_j)_{\beta_j \alpha_j} & (Y_j)_{\beta_j \beta_j} & 0 \\ (Y_j)_{\gamma_j \alpha_j} & 0 & 0 \end{pmatrix} P_j^\top, \quad j = 1, \dots, J. \end{cases}$$

Consequently, it holds that

$$\begin{aligned} &P_j^\top (g'_j(\bar{x})d) P_j + \begin{pmatrix} (P_j)_{\alpha_j}^\top Z_j (P_j)_{\alpha_j} & (P_j)_{\alpha_j}^\top Z_j (P_j)_{\beta_j} & (P_j)_{\alpha_j}^\top Z_j (P_j)_{\gamma_j} \\ (P_j)_{\beta_j}^\top Z_j (P_j)_{\alpha_j} & 0 & 0 \\ (P_j)_{\gamma_j}^\top Z_j (P_j)_{\alpha_j} & 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} (Y_j)_{\alpha_j \alpha_j} & (Y_j)_{\alpha_j \beta_j} & (Y_j)_{\alpha_j \gamma_j} \\ (Y_j)_{\beta_j \alpha_j} & (Y_j)_{\beta_j \beta_j} & 0 \\ (Y_j)_{\gamma_j \alpha_j} & 0 & 0 \end{pmatrix}, \quad j = 1, \dots, J. \end{aligned}$$

Then by (2.15) we know that $d \in \text{aff}(C(\bar{x}))$. Recall that $\text{rge}(\mathcal{W}) = \ker(\mathcal{B}) = \text{aff}(C(\bar{x}))$. Then there exists a vector $\nu \in \mathbb{R}^r$ such that $d = \mathcal{W}\nu$. Since $(Y_j)_{\beta_j \beta_j}$ can be any matrix in $\mathcal{S}^{|\beta_j|}$, one can see that the linear operator \mathcal{A} defined by (3.17) is surjective. Recall from (2.11) and (2.12) that

$$C(\bar{x}) = \left\{ d \in \mathcal{X} \mid \begin{cases} h'(\bar{x})d = 0, (P_j)_{\beta_j}^\top (g'_j(\bar{x})d)(P_j)_{\beta_j} \succeq 0, \\ (P_j)_{\beta_j}^\top (g'_j(\bar{x})d)(P_j)_{\gamma_j} = 0, \\ (P_j)_{\gamma_j}^\top (g'_j(\bar{x})d)(P_j)_{\gamma_j} = 0, \quad j = 1, \dots, J \end{cases} \right\} \subseteq \text{rge}(\mathcal{W}).$$

Then the second-order sufficient condition (2.17) can be equivalently recast as

$$\langle \nu, \mathcal{W}^* \mathcal{Q} \mathcal{W} \nu \rangle > 0 \quad \forall \nu \in \Omega \setminus \{0\}. \quad (3.18)$$

Also, the SSOSC (2.18) can be equivalently written as

$$\langle \nu, \mathcal{W}^* \mathcal{Q} \mathcal{W} \nu \rangle > 0 \quad \forall \nu \in \mathbb{R}^r \setminus \{0\}. \quad (3.19)$$

Moreover, since \mathcal{A} is surjective, one can get from Lemma 2.2 that

$$\begin{aligned} \Omega^\circ &= \{ \mathcal{A}^*(\Theta_1; \dots; \Theta_J) \mid \Theta_j \in -\mathcal{S}_+^{|\beta_j|}, j = 1, \dots, J \} \\ &= \left\{ \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \Theta_j (P_j)_{\beta_j}^\top \mid \Theta_j \preceq 0, j = 1, \dots, J \right\}. \end{aligned}$$

Note that by Corollary 3.1, the matrix $\mathcal{W}^* \mathcal{Q} \mathcal{W}$ is nonsingular. Then, by using Lemma 2.3 and (3.18) we know that (3.19) holds if and only if

$$\langle \eta, (\mathcal{W}^* \mathcal{Q} \mathcal{W})^{-1} \eta \rangle > 0 \quad \forall \eta \in \Omega^\circ \setminus \{0\}. \quad (3.20)$$

In the following, we shall show that (3.20) holds.

Suppose on the contrary that (3.20) does not hold. Consider the following auxiliary optimization problem

$$\begin{aligned} \min_{\eta \in \mathbb{R}^r, \Theta_j \in \mathcal{S}^{|\beta_j|}} \quad & \frac{1}{2} \langle \eta, (\mathcal{W}^* \mathcal{Q} \mathcal{W})^{-1} \eta \rangle \\ \text{s.t.} \quad & \begin{cases} \eta - \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \Theta_j (P_j)_{\beta_j}^\top = 0, \\ \frac{1}{2} (\sum_{j=1}^J \|\Theta_j\|^2 - 1) = 0, \\ \Theta_j \preceq 0, j = 1, \dots, J. \end{cases} \end{aligned} \quad (3.21)$$

Since the feasible set of (3.21) is compact, the minimum of the objective function can be attained at a certain solution $(\bar{\eta}; \bar{\Theta}_1; \dots; \bar{\Theta}_J)$. Moreover, it is easy to see that the optimal value of (3.21) is not positive since (3.20) does not hold. By applying Lemma 2.4 with $\mathcal{H} = \mathcal{A}$, $K = -(\mathcal{S}_+^{|\beta_1|} \times \dots \times \mathcal{S}_+^{|\beta_J|})$, $\mathcal{E} = \mathbb{R}^r$ and $\mathcal{F} = \mathcal{S}^{|\beta_1|} \times \dots \times \mathcal{S}^{|\beta_J|}$, one has that Robinson's constraint qualification holds for the constraint in (3.21) at $(\bar{\eta}; \bar{\Theta}_1; \dots; \bar{\Theta}_J)$. Consequently, we know from [6, Theorem 3.9] that there exists a Lagrange multiplier $(\bar{\rho}; \bar{\tau}; \bar{\Delta}_1; \dots; \bar{\Delta}_J) \in \mathbb{R}^r \times \mathbb{R} \times \mathcal{S}^{|\beta_1|} \times \dots \times \mathcal{S}^{|\beta_J|}$ at $(\bar{\eta}; \bar{\Theta}_1; \dots; \bar{\Theta}_J)$ such that the following KKT system of (3.21) holds:

$$\begin{cases} (\mathcal{W}^* \mathcal{Q} \mathcal{W})^{-1} \bar{\eta} + \bar{\rho} = 0, \\ -(P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\rho})(P_j)_{\beta_j} + \bar{\tau} \bar{\Theta}_j + \bar{\Delta}_j = 0, \\ \bar{\eta} - \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \bar{\Theta}_j (P_j)_{\beta_j}^\top = 0, \\ \frac{1}{2} (\sum_{j=1}^J \|\bar{\Theta}_j\|^2 - 1) = 0, \\ \bar{\Theta}_j \preceq 0, \bar{\Delta}_j \succeq 0, \langle \bar{\Delta}_j, \bar{\Theta}_j \rangle = 0, j = 1, \dots, J. \end{cases} \quad (3.22)$$

Note that

$$\mathcal{W}^* \mathcal{Q} \mathcal{W} \bar{\rho} = -\bar{\eta} = -\sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \bar{\Theta}_j (P_j)_{\beta_j}^\top. \quad (3.23)$$

Moreover, from (3.22) one can also see that $\sum_{j=1}^J \|\bar{\Theta}_j\|^2 = 1$, so that

$$\begin{aligned} \bar{\tau} &= \langle (\bar{\Delta}_1; \dots; \bar{\Delta}_J) + \bar{\tau}(\bar{\Theta}_1; \dots; \bar{\Theta}_J), (\bar{\Theta}_1; \dots; \bar{\Theta}_J) \rangle \\ &= \sum_{j=1}^J \left\langle (P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\rho})(P_j)_{\beta_j}, \bar{\Theta}_j \right\rangle \\ &= \left\langle \bar{\rho}, \sum_{j=1}^J \mathcal{W}^* \nabla g_j(\bar{x})(P_j)_{\beta_j} \bar{\Theta}_j (P_j)_{\beta_j}^\top \right\rangle = \langle \bar{\rho}, \bar{\eta} \rangle = -\langle \bar{\eta}, (\mathcal{W}^* \mathcal{Q} \mathcal{W})^{-1} \bar{\eta} \rangle \geq 0, \end{aligned}$$

where the last inequality holds because the optimal value of (3.21) is not positive. Note that one has the eigenvalue decomposition that

$$\bar{\Theta}_j + \bar{\Delta}_j = \bar{O}_j \begin{pmatrix} (\bar{\Lambda}_j)_{\bar{\alpha}_j \bar{\alpha}_j} & 0 & 0 \\ 0 & 0_{\bar{\beta}_j \bar{\beta}_j} & 0 \\ 0 & 0 & (\bar{\Lambda}_j)_{\bar{\gamma}_j \bar{\gamma}_j} \end{pmatrix} \bar{O}_j^\top, \quad j = 1, \dots, J,$$

where \bar{O}_j is an orthogonal matrix. Moreover, from the last line of (3.22) one has that

$$\bar{\Theta}_j = \bar{O}_j \begin{pmatrix} 0_{\bar{\alpha}_j \bar{\alpha}_j} & 0 & 0 \\ 0 & 0_{\bar{\beta}_j \bar{\beta}_j} & 0 \\ 0 & 0 & (\bar{\Lambda}_j)_{\bar{\gamma}_j \bar{\gamma}_j} \end{pmatrix} \bar{O}_j^\top, \quad j = 1, \dots, J$$

and

$$\bar{\Delta}_j + \bar{\tau} \bar{\Theta}_j = \bar{O}_j \begin{pmatrix} (\bar{\Lambda}_j)_{\bar{\alpha}_j \bar{\alpha}_j} & 0 & 0 \\ 0 & 0_{\bar{\beta}_j \bar{\beta}_j} & 0 \\ 0 & 0 & \bar{\tau} (\bar{\Lambda}_j)_{\bar{\gamma}_j \bar{\gamma}_j} \end{pmatrix} \bar{O}_j^\top, \quad j = 1, \dots, J.$$

For arbitrarily given $j \in \{1, \dots, J\}$, one can take $\beta_+ = \bar{\alpha}_j$, $\beta_0 = \bar{\beta}_j \cup \bar{\gamma}_j$, and $\beta_- = \emptyset$ in (2.6) and (2.7) to obtain

$$\Xi_1 = \begin{pmatrix} E_{\bar{\alpha}_j \bar{\alpha}_j} & E_{\bar{\alpha}_j \bar{\beta}_j} & E_{\bar{\alpha}_j \bar{\gamma}_j} \\ E_{\bar{\beta}_j \bar{\alpha}_j} & 0 & 0 \\ E_{\bar{\gamma}_j \bar{\alpha}_j} & 0 & 0 \end{pmatrix} \quad \text{and} \quad \Xi_2 = 0 \in \mathcal{S}^{|\beta_j|}.$$

Consequently, it holds that

$$\begin{cases} \Xi_1 \circ \bar{O}^\top \bar{\Theta}_j \bar{O} = 0 = \Xi_2 \circ \bar{O}^\top (\bar{\Delta}_j + \bar{\tau} \bar{\Theta}_j) \bar{O}, \\ \bar{O}_{\beta_0}^\top \bar{\Theta}_j \bar{O}_{\beta_0} \preceq 0, \\ \bar{O}_{\beta_0}^\top (\bar{\Delta}_j + \bar{\tau} \bar{\Theta}_j) \bar{O}_{\beta_0} \preceq 0, \end{cases}$$

which, together with Lemma 2.1 and (3.22), implies that

$$\bar{\Theta}_j \in \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) (\bar{\Delta}_j + \bar{\tau} \bar{\Theta}_j) = \mathcal{D}^* \mathcal{N}_{\mathcal{S}_+^{|\beta_j|}}(0, 0) ((P_j)_{\beta_j}^\top (g'_j(\bar{x}) \mathcal{W} \bar{\rho}) (P_j)_{\beta_j}).$$

Note that such an inclusion holds simultaneously for all $j = 1, \dots, J$. Thus, this inclusion, together with (3.23), makes a contradiction to (3.12) in Proposition 3.2 (with $\nu = \bar{\rho}$ and $\tilde{U}_{\beta_j \beta_j}^j = \bar{\Theta}_j$ for all $j = 1, \dots, J$). As a consequence, we know that (3.20) is valid. Hence, one has (3.19) holds, which leads to the SSOSC (2.18) in Definition 2.1. This completes the proof. \square

4 Equivalent characterizations of the Aubin property

This section establishes the equivalent characterizations of the Aubin property of \mathbb{S}_{KKT} (1.6) at $(0; 0)$ for $(\bar{x}; \bar{y})$ with \bar{x} being a locally optimal solution to the NLSDP problem (1.8) and $\bar{y} \in \mathcal{M}(\bar{x})$.

We first review some related concepts in variational analysis. As was mentioned in Section 1, the Aubin property is related to the strong metric regularity [16, Definition 2.5]. Specifically, for a set-valued mapping $\Psi : \mathcal{E} \rightrightarrows \mathcal{F}$, one has Ψ is strongly metrically regular at $(\bar{z}; \bar{w}) \in \text{gph} \Psi$ if Ψ^{-1} has the Aubin property at \bar{w} for \bar{z} , and there exist neighborhoods \mathcal{U} of \bar{z} and \mathcal{V} of \bar{w} , such that $\Psi^{-1}(w) \cap \mathcal{U}$ is a singleton for all $w \in \mathcal{V}$. The following result provides a criterion for characterizing the strong metric regularity.

Lemma 4.1. [16, Theorem 2.7] *Ψ is strongly metrically regular at $(\bar{z}; \bar{w}) \in \text{gph}(\Psi)$ if and only if for all $w \in \mathcal{F}$ and $z \in \mathcal{E}$, one has*

$$0 \in \mathcal{D}^* \Psi(\bar{z}, \bar{w})(w) \Rightarrow w = 0 \quad \text{and} \quad 0 \in \mathcal{D}_* \Psi(\bar{z}, \bar{w})(z) \Rightarrow z = 0,$$

where \mathcal{D}_* refers to the strict graphical derivative defined in (2.2).

Recall that the KKT system (1.3) can be equivalently written as the nonsmooth equation

$$F(x, y) := \begin{pmatrix} \nabla_x \mathcal{L}(x, y) \\ -G(x) + \Pi_{\mathcal{K}}(G(x) + y) \end{pmatrix} = 0. \quad (4.1)$$

Since F is locally Lipschitz continuous around $(\bar{x}; \bar{y})$, it is almost everywhere differentiable in a neighborhood \mathcal{V} of $(\bar{x}; \bar{y})$ by Rademacher's theorem [40, Theorem 9.60]. We use $D_F \subseteq \mathcal{V}$ to denote the set of points at which F is differentiable. The Bouligand subdifferential of F at $(\bar{x}; \bar{y})$ is defined by

$$\partial_B F(\bar{x}, \bar{y}) := \{v \in \mathcal{X} \times \mathcal{Y} \mid \exists (x_k; y_k) \xrightarrow{D_F} (\bar{x}; \bar{y}) \text{ with } F'(x_k, y_k) \rightarrow v\}.$$

The Clarke generalized jacobian of F at $(\bar{x}; \bar{y})$ is defined by

$$\bar{\partial} F(\bar{x}, \bar{y}) := \text{conv}(\partial_B F(\bar{x}, \bar{y})),$$

i.e., the convex hull of $\partial_B F(\bar{x}, \bar{y})$.

The perturbed KKT system (1.4) corresponds to a two-parametric optimization problem

$$\min_{x \in \mathcal{X}, a \in \mathcal{X}, b \in \mathcal{Y}} \phi(x, b) - \langle a, x \rangle, \quad (4.2)$$

where

$$\phi(x, b) := f(x) + \delta_{\mathcal{K}}(G(x) - b) \quad (4.3)$$

with $\delta_{\mathcal{K}}(\cdot)$ being the indicator function of \mathcal{K} in convex analysis [37]. When $a = 0 \in \mathcal{X}$ and $b = 0 \in \mathcal{Y}$, one has (4.2) is exactly (1.1). Given $\iota > 0$ and $(\bar{x}; \bar{b}) \in \mathcal{X} \times \mathcal{Y}$ such that $\phi(\bar{x}, \bar{b})$ is finite, one can define

$$\begin{cases} M_\iota(a, b) := \arg \min_{x \in \mathcal{X}} \{\phi(x, b) - \langle a, x \rangle \mid \|x - \bar{x}\| \leq \iota\}, \\ m_\iota(a, b) := \inf_{x \in \mathcal{X}} \{\phi(x, b) - \langle a, x \rangle \mid \|x - \bar{x}\| \leq \iota\}. \end{cases}$$

We say the point \bar{x} is a Lipschitzian fully stable local minimizer [28, Definition 3.2] of (4.2) at $(\bar{a}; \bar{b})$ if there exist numbers $\kappa > 0, \iota > 0$ and a neighborhood \mathcal{V} of $(\bar{a}; \bar{b})$ such that the mapping $M_\iota(a, b)$ is single-valued on \mathcal{V} with $M_\iota(\bar{a}, \bar{b}) = \bar{x}$ satisfying

$$\|M_\iota(a_1, b_1) - M_\iota(a_2, b_2)\| \leq \kappa \|(a_1; b_1) - (a_2; b_2)\| \quad \forall (a_1; b_1), (a_2; b_2) \in \mathcal{V},$$

and that the function $m_\iota(a, b)$ is also Lipschitz continuous on \mathcal{V} . Different from \mathbb{S}_{KKT} , the mapping $M_\iota(a, b)$ focuses mainly on locally optimal solutions.

Recently, Rockafellar [38, 39] introduced the strong variational sufficient condition for local optimality, and provided several characterizations of this abstract property. With ϕ being defined in (4.3), one can reformulate (1.1) to

$$\min_{x \in \mathcal{X}, b \in \mathcal{Y}} \phi(x, b) \quad \text{s.t.} \quad b = 0.$$

Define the function $\phi_\ell(x, b) := \phi(x, b) + \frac{\ell}{2} \|b\|^2$. The following definition comes from [39, Section 2].

Definition 4.1. *The (strong) variational sufficient condition for local optimality in (1.1) holds with respect to a solution $(\bar{x}; \bar{y})$ to the KKT system (1.3) if there exists $\ell > 0$ such that ϕ_ℓ is (strongly) variationally convex with respect to the pair $((\bar{x}; 0), (0; \bar{y})) \in \text{gph}(\partial \phi_\ell)$, i.e., there exist open convex neighborhoods \mathcal{U} of $(\bar{x}; 0)$ and \mathcal{V} of $(0; \bar{y})$, and a closed proper (strongly) convex function $\varrho \leq \phi_\ell$ on \mathcal{U} such that*

$$(\mathcal{U} \times \mathcal{V}) \cap \text{gph}(\partial \varrho) = (\mathcal{U} \times \mathcal{V}) \cap \text{gph}(\partial \phi_\ell),$$

and $\varrho(x, b) = \phi_\ell(x, b)$ holds for $((x; b); (a; y))$ belonging to this common set. Here $\partial \varrho$ is the subdifferential in convex analysis and $\partial \phi_\ell$ is the limiting subdifferential defined in (2.1).

Remark 4.1. For a special case ($J = 1$ and $m = 0$) of the NLSDP problem (1.8), the strong variational sufficient condition (with respect to a solution $(\bar{x}; \bar{y})$ to the KKT system (1.3)) and the SSOSC (2.18) (for the same $(\bar{x}; \bar{y})$) were proved to be equivalent in [43, Theorem 3.3]. This equivalence can be extended to the NLSDP problem (1.8) in its general form with ease by directly following their proof.

Based on the above definitions, we can establish the following result regarding the equivalent characterizations of the Aubin property for the solution mapping \mathbb{S}_{KKT} in (1.6) of the NLSDP problem (1.8).

Theorem 4.1. Let \bar{x} be a locally optimal solution to the NLSDP problem (1.8) and $\bar{y} = (\bar{\zeta}; \bar{\Gamma}_1; \dots; \bar{\Gamma}_J) \in \mathcal{M}(\bar{x})$ be a multiplier at \bar{x} . Then, the following statements are equivalent:

- (i) The solution mapping \mathbb{S}_{KKT} in (1.6) has the Aubin property at $(0; 0)$ for $(\bar{x}; \bar{y})$.
- (ii) The strong second-order sufficient (2.18) holds at $(\bar{x}; \bar{y})$ and \bar{x} is nondegenerate, i.e., (2.14) holds.
- (iii) Any element of the Clarke generalized Jacobian $\bar{\partial}F(\bar{x}, \bar{y})$ is nonsingular, where the function F is defined in (4.1).
- (iv) The KKT point $(\bar{x}; \bar{y})$ is a strongly regular solution to the generalized equation (1.7) (or the KKT system (1.5)).
- (v) The mapping Φ in (1.7) is strongly metrically regular at $(\bar{x}; \bar{y})$ for $(0; 0)$.
- (vi) For any $w \in \mathcal{X} \times \mathcal{Y}$ with $0 \in \mathcal{D}^*\Phi(\bar{x}, \bar{y})(w)$, one has $w = 0$.
- (vii) The strong variational sufficient condition in Definition 4.1 holds with respect to $(\bar{x}; \bar{y})$, and \bar{x} is nondegenerate, i.e., (2.14) holds.
- (viii) \bar{x} is a Lipschitzian fully stable local minimizer of (4.2), and \bar{x} is nondegenerate, i.e., (2.14) holds.

Proof. One has (i) \Rightarrow (ii) from Proposition 3.3 and [23, Theorem 1]. By simply repeating the proof of [42, Proposition 3.2] one can get (ii) \Rightarrow (iii) \Rightarrow (iv). It follows by [34, Corollary 2.2] and the definition that (iv) \Rightarrow (v) holds. From Lemma 4.1 and the Mordukhovich criterion for the Aubin property [40, Theorem 9.40] one has (v) \Rightarrow (vi) \Rightarrow (i). According to Remark 4.1 we know that (ii) \Leftrightarrow (vii). One also has from [28, Theorem 5.6] that (iv) \Leftrightarrow (viii). This completes the proof. \square

Remark 4.2. Note that the eight equivalent conditions listed in Theorem 4.1 are not exhaustive. For example, one additional equivalent condition, according to [42, Remark 3.1], is that F is a locally Lipschitz homeomorphism [14, Definition 2.1.9] near $(\bar{x}; \bar{y})$. Furthermore, by [42, Theorem 4.1], another condition is that \bar{x} is nondegenerate and strongly stable [6, Definition 5.33]. For more details and equivalent conditions, one may refer to [42] and the references therein.

Remark 4.3. According to the equivalence between the two conditions (v) and (vi) in Theorem 4.1, one can see from Lemma 4.1 that (vi) is a sufficient condition for

$$0 \in \mathcal{D}_*\Phi(\bar{x}, \bar{y})(z) \Rightarrow z = 0 \quad \forall z \in \mathcal{X} \times \mathcal{Y}.$$

Remark 4.4. It should be emphasized that, for (1.1) with \mathcal{K} being an arbitrary C^2 -cone reducible set, it is still unknown if the strong regularity of the KKT system (1.7) is equivalent to the constraint nondegeneracy (2.13) combined with a certain second-order optimality condition. The currently known cases include the nonlinear programming [12], the NLSOCP [4], the NLSDP problem [42] and a composite matrix programming regarding matrix eigenvectors [8].

5 Conclusions

In this paper, we prove that at a locally optimal solution to the nonlinear semidefinite programming problem (1.8), the Aubin property of \mathbb{S}_{KKT} (1.6) is equivalent to the strong second-order sufficient condition plus the constraint nondegeneracy. This enables us to derive a series of equivalent characterizations of the Aubin property, which includes the strong regularity of the Karush-Kuhn-Tucker system (1.3). As a byproduct, for nonlinear semidefinite programming this paper answers the open question posed in [10, Section 5] if the Aubin property can be characterized by an exact form of a certain second-order optimality condition together with the constraint nondegeneracy. It is noted that our analysis for nonlinear semidefinite programming (and also for nonlinear second-order cone programming in [7]) relies on the explicit formulas of the coderivative for the underlying normal cone mapping. Currently, it is not clear to us how to extend these results to the generic non-polyhedral C^2 -cone reducible constrained optimization problems. We leave this as our future research topic.

References

- [1] Aubin, J.-P.: Lipschitz behavior of solutions to convex minimization problems. *Math. Oper. Res.* **9**, 87–111 (1984)
- [2] Bonnans, J.F., Cominetti, R., Shapiro. A.: Sensitivity analysis of optimization problems under second order regularity constraints. *Math. Oper. Res.* **23**, 803–832 (1998)
- [3] Bonnans, J.F., Cominetti, R., Shapiro. A.: Second order optimality conditions based on parabolic second order tangent sets. *SIAM J. Optim.* **9**, 466–492 (1999)
- [4] Bonnans, J.F., Ramírez, H.: Perturbation analysis of second-order cone programming problems. *Math. Program.* **104**(2), 205–227 (2005)
- [5] Bonnans, J.F., Sulem A.: Pseudopower expansion of solutions of generalized equations and constrained optimization problems. *Math. Program.* **70**, 123–148 (1995)
- [6] Bonnans, J.F., Shapiro, A.: *Perturbation Analysis of Optimization Problems*. Springer, New York (2000)
- [7] Chen, L., Chen, R.N., Sun, D.F., Zhu, J.Y.: Aubin property and strong regularity are equivalent for NLSOCP (2024). [arXiv:2406.13798v1](https://arxiv.org/abs/2406.13798v1)
- [8] Cui, Y., Ding, C.: Composite matrix optimization: strong regularity, constraint nondegeneracy and beyond (2019). [arXiv:1907.13253v2](https://arxiv.org/abs/1907.13253v2)
- [9] Ding, C., Sun D.F., Ye, J.J.: First order optimality conditions for mathematical programs with semidefinite cone complementarity constraints. *Math. Program.* **147**, 539–579 (2014)
- [10] Ding, C., Sun, D.F., Zhang, L.W.: Characterization of the robust isolated calmness for a class of conic programming problems. *SIAM J. Optim.* **27**, 67–90 (2017)
- [11] Dontchev, A.L.: Implicit function theorems for generalized equations. *Math. Program.* **70**, 91–106 (1995)
- [12] Dontchev, A.L., Rockafellar, R.T.: Characterization of strong regularity for variational inequalities over polyhedral convex sets. *SIAM J. Optim.* **6**, 1087–1105 (1996)
- [13] Dontchev, A.L., Rockafellar, R.T.: *Implicit Functions and Solution Mappings*, 2nd edn. Springer, New York (2014)
- [14] Facchinei, F., Pang, J.-S.: *Finite-Dimensional Variational Inequalities and Complementarity Problems: Volume I*. Springer, New York (2003)
- [15] Fusek, P.: On metric regularity for weakly almost piecewise smooth functions and some applications in nonlinear semidefinite programming. *SIAM J. Optim.* **23**, 1041–1061 (2013)

- [16] Gfrerer, H., Outrata, J.V.: On (local) analysis of multifunctions via subspaces contained in graphs of generalized derivatives. *J. Math. Anal. Appl.* **508**, 125895 (2022)
- [17] Han, S.-P., Mangasarian, O.L.: Conjugate cone characterization of positive definite and semidefinite matrices. *Linear Algebra Appl.* **56**, 89–103 (1984)
- [18] Hiriart-Urruty, J.-B., Malick, J.: A fresh variational-analysis look at the positive semidefinite matrices world. *J. Optim. Theory Appl.* **153**, 551–577 (2012)
- [19] Hoheisel, T., Kanzow, C.: Stationary conditions for mathematical programs with vanishing constraints using weak constraint qualifications. *J. Math. Anal. Appl.* **337**, 292–310 (2008)
- [20] Jongen, H.T., Möbert, T., Rückmann, J., Tammer, K.: On inertia and Schur complement in optimization. *Linear Algebra Appl.* **95**, 97–109 (1987)
- [21] Jongen, H.T., Klatte, D., Tammer, K.: Implicit functions and sensitivity of stationary points. *Math. Program.* **19**, 123–138 (1990)
- [22] Klatte, D., Kummer, B.: *Nonsmooth Equations in Optimization: Regularity, Calculus, Methods and Applications*. Kluwer Academic Publishers, Boston (2002)
- [23] Klatte, D., Kummer, B.: Aubin property and uniqueness of solutions in cone constrained optimization. *Math. Methods Oper. Res.* **77**, 291–304 (2013)
- [24] Klatte, D., Tammer, K.: Strong stability of stationary solutions and Karush-Kuhn-Tucker points in nonlinear optimization. *Ann. Oper. Res.* **27**, 285–308 (1990)
- [25] Kojima, M.: Strongly stable stationary solutions in nonlinear programs. In: Robinson, S.M. (ed.) *Analysis and Computation of Fixed Points*, pp. 93–138. Academic Press, New York (1980)
- [26] Kummer, B.: Lipschitzian inverse functions directional derivatives and application in $C^{1,1}$ optimization. *J. Optim. Theory Appl.* **70**, 561–582 (1991)
- [27] Mordukhovich, B.S.: Stability theory for parametric generalized equations and variational inequalities via nonsmooth analysis. *Trans. Amer. Math. Soc.* **343**, 609–657 (1994)
- [28] Mordukhovich, B.S., Nghia, T.T.A., Rockafellar, R.T.: Full stability in finite-dimensional optimization. *Math. Oper. Res.* **40**, 226–252 (2015)
- [29] Mordukhovich, B.S., Outrata, J.V.: On second-order subdifferentials and their applications. *SIAM J. Optim.* **12**, 139–169 (2001)
- [30] Opazo, F., Outrata, J.V., Ramírez, H.: Erratum: on the Aubin property of critical points to perturbed second-order cone programs. *SIAM J. Optim.* **27**, 2143–2151 (2017)
- [31] Outrata, J.V., Ramírez, H.: On the Aubin property of critical points to perturbed second-order cone programs. *SIAM J. Optim.* **21**, 798–823 (2011)
- [32] Pang, J.-S., Sun, D.F., Sun, J.: Semismooth homeomorphisms and strong stability of semidefinite and Lorentz complementarity problems. *Math. Oper. Res.* **28**, 39–63 (2003)
- [33] Robinson, S. M.: First order conditions for general nonlinear optimization. *SIAM J. Appl. Math.* **30**, 597–607 (1976)
- [34] Robinson, S.M.: Strongly regular generalized equations. *Math. Oper. Res.* **5**, 43–62 (1980)
- [35] Robinson, S. M.: Local structure of feasible sets in nonlinear programming, Part II: Nondegeneracy. *Math. Program. Study* **22**, 217–230 (1984)
- [36] Robinson, S.M.: Constraint nondegeneracy in variational analysis. *Math. Oper. Res.* **28**, 201–232 (2003)
- [37] Rockafellar, R.T.: *Convex Analysis*. Princeton University Press, Princeton, New Jersey (1970)

- [38] Rockafellar, R.T.: Progressive decoupling of linkages in monotone variational inequalities and convex optimization. In Hojo, M., Hoshino, M., Takahashi W. (eds.) Proceedings of the 10th International Conference on Nonlinear Analysis and Convex Analysis (Chitose, Japan, 2017), pp. 271–291. Yokohama Publishers (2019)
- [39] Rockafellar, R.T.: Augmented Lagrangians and hidden convexity in sufficient conditions for local optimality. *Math. Program.* **198**, 159–194 (2023)
- [40] Rockafellar, R.T., Wets R.J.-B.: *Variational Analysis*. Springer-Verlag, New York (1998)
- [41] Shapiro, A.: Sensitivity analysis of generalized equations. *J. Math. Sci.* **115**, 2554–2565 (2003)
- [42] Sun, D.F.: The strong second order sufficient condition and constraint nondegeneracy in nonlinear semidefinite programming and their implications. *Math. Oper. Res.* **31**, 761–776 (2006)
- [43] Wang, S.W., Ding, C., Zhang, Y.J., Zhao, X.Y.: Strong variational sufficiency for nonlinear semidefinite programming and its implications. *SIAM J. Optim.* **33**, 2988–3011 (2023)