# Blow-up solutions for the steady state of the Keller-Segel system on Riemann surfaces

Zhengni Hu Thomas Bartsch Mohameden Ahmedou

September 4, 2024

#### Abstract

We study the following Neumann boundary problem related to the stationary solutions of the Keller-Segel system, a basic model of chemotaxis phenomena:

$$\begin{cases} -\Delta_g u + \beta u = \lambda \left( \frac{V e^u}{\int_{\Sigma} V e^u dv_g} - \frac{1}{|\Sigma|_g} \right) & \text{in } \mathring{\Sigma} \\ \\ \partial_{\nu_g} u = 0 & \text{on } \partial \Sigma \end{cases},$$

on a compact Riemann surface  $(\Sigma, g)$  with smooth boundary  $\partial \Sigma$ . Here,  $\lambda, \beta \geq 0$  are non-negative parameters, and V is a smooth non-negative function with a finite zero set. For any given integers  $m \geq k \geq 0$ , we establish a sufficient condition on V for the existence of a sequence of blow-up solutions as  $\lambda$  approaches the critical values  $4\pi(m+k)$ . Moreover, the study expands to address the corresponding singular problem.

 $\textbf{Keywords:} \ \textit{Keller-Segel models; Blow-up solutions; Lyapunov-Schmidt reduction}$ 

2020 AMS Subject Classification: 35J57, 58J05

#### 1 Introduction

The Keller-Segel system was first introduced in [23] to show the aggregation of biological species. It is a coupled parabolic system for the concentration of species u(x,t) and chemical released v(x,t) as follows.

(1) 
$$\begin{cases} u_{t} = \Delta u - \chi \nabla(u \nabla v), & x \in \Omega, t > 0 \\ \Gamma v_{t} = \Delta v - \beta_{0} v + \delta u, & x \in \Omega, t > 0 \\ u(x, 0) = u_{0}(x), & \text{for } x \in \Omega \\ v(x, 0) = v_{0}(x), & \text{for } x \in \Omega \\ \frac{\partial u}{\partial \nu_{q}} = \frac{\partial v}{\partial \nu_{q}} = 0, & \text{on } \partial \Omega \end{cases}$$

where  $\Omega \subset \mathbb{R}^N$   $(N \ge 1)$ ,  $\chi$ ,  $\Gamma$ ,  $\beta_0$  and  $\delta$  are positive parameters. The mass of u(x,t) is preserved in (1), i.e.,

$$\int_{\Omega} u(x,t) = \int_{\Omega} u_0(x).$$

Considering the stationary solutions of (1), the problem turns out to be an elliptic system. After a transformation (see [18,21,34], for instance),  $u = Ce^v$  for some constant C. For v, we obtain the following problem with the Neumann boundary condition,

(2) 
$$\begin{cases} -\Delta v + \beta v = \lambda \left( \frac{e^v}{\int_{\Omega} e^v} - \frac{1}{|\Omega|} \right), & x \in \Omega \\ \frac{\partial v}{\partial \nu_g} = 0, & \text{on } \partial \Omega \end{cases},$$

where  $\nu_g$  is the unit outward normal on  $\partial\Omega$ ,  $\beta$  and  $\lambda$  are parameters.

In the one-dimensional case, Schaaf demonstrates the existence of non-trivial solutions using a bifurcation technique, as shown in [30]. For the higher-dimensional case with  $N \geq 3$ , we refer to [1,4,29] and references therein.

This paper specifically concentrates on the case where N=2. We will now delve into the literature on this particular setting.

By Struwe's technique and blow-up analysis, Wang and Wei in [34] obtain non-constant solutions of (2) for  $\beta > \frac{\lambda}{|\Omega|} - \lambda_1$  and  $\lambda \in (4\pi, +\infty) \setminus 4\pi\mathbb{N}_+$ , where  $\lambda_1$  is the first eigenvalue of  $-\Delta$  with the Neumann boundary condition. Independently, Senba and Suzuki obtain the same result in [31]. Battaglia generates their result for  $\lambda \in (0, +\infty) \setminus 4\pi\mathbb{N}_+$  and  $\beta$  with any sign in [3]. He proves the existence of non-constant solutions of (2) with some algebraic conditions involved with  $\beta$ ,  $\lambda$  and eigenvalues  $\{\lambda_i\}$  by the variational method and Moser theory.

However, the blow-up phenomena may occur when  $\lambda$  approaches the critical value  $4\pi N_+$ . Del Pino and Wei in [28] construct positive value bubbling solutions for the Neumann boundary

problem on a bounded domain  $\Omega$  with  $\beta > 0$ 

(3) 
$$\begin{cases} -\Delta u + \beta u = \varepsilon^2 e^u & \text{in } \Omega \\ \partial_{\nu} u = 0 & \text{on } \partial \Omega \end{cases},$$

by the Lyapunov-Schmidt reduction. In particular, the sequence of bubbling solutions blows up at k distinct points  $\xi_1, \dots, \xi_k$  inside the domain  $\Omega$  and m-k distinct points  $\xi_{k+1}, \dots, \xi_m$  on the boundary of  $\Omega$ , i.e., as  $\varepsilon \to 0$ 

$$u_{\varepsilon} \rightharpoonup \sum_{i=1}^{k} 8\pi \delta_{\xi_i} + \sum_{i=k+1}^{m} 4\pi \delta_{\xi_i},$$

where  $\delta_{\xi}$  is the Dirac mass. Subsequently, Del Pino, Pistoia and Vaira in [14] construct solutions of (3) which blow up along the whole boundary  $\partial\Omega$ .

This paper studies the Neumann boundary problem on a compact Riemann surface  $\Sigma$  with a smooth boundary  $\partial \Sigma$ :

(4) 
$$\begin{cases} -\Delta_g u + \beta u = \lambda \left( \frac{V e^u}{\int_{\Sigma} V e^u dv_g} - \frac{1}{|\Sigma|_g} \right) & \text{in } \mathring{\Sigma} \\ \partial_{\nu_g} u = 0 & \text{on } \partial \Sigma \end{cases},$$

where the parameters  $\lambda, \beta \in \mathbb{R}$  and V is a non-negative smooth function with a finite zero set denoted as  $\{q_1, \dots, q_\iota\}$  for some  $\iota \in \mathbb{N}$ ,  $\mathring{\Sigma} := \Sigma \setminus \partial \Sigma$  is the interior of  $\Sigma$ ,  $\Delta_g$  is the Laplace-Beltrami operator,  $dv_g$  is the area element in  $(\Sigma, g)$ ,  $|\Sigma|_g = \int_{\Sigma} dv_g$ , and  $\nu_g$  is the unit outward normal of  $\partial \Sigma$ .

This paper delves into the study of the blow-up solutions of the problem (4). For integers  $m \geq k \geq 0$ , we establish sufficient conditions for blow-up solutions. Moreover, the precise locations of blow-up points are explicitly characterized by the "stable" critical point of a functional  $\mathcal{F}_{k,m}$  in terms of Green's function on  $\Sigma$  and the function V.

The non-linear equation in (2) with  $\beta \equiv 0$  transforms into the well-known mean-field equation. This equation arises in various branches of mathematics and physics, such as statistical mechanics (see [6,7,24]), Abelian Chern-Simons gauge theory (see [5,27,32,35]), and conformal geometry (see [8–10,12,22,33]). When coupled with Dirichlet boundary conditions, in the spirit of the Lyapunov-Schmidt reduction the blow-up solutions of the mean-field equations are well-studied both in Euclidean spaces (refer to [13,16,28] and the references therein) and on Riemann surfaces without boundaries (refer to [2,15,17]). Recently, [20] obtained blow-up solutions with Neumann boundary conditions on Riemann surfaces with boundaries

under nonvanishing of a quantity related to V and Gaussian curvature  $K_g$ . As in these papers, our approach to finding blow-up solutions of (4) is based on variational methods combined with the Lyapunov-Schmidt reduction. In comparison to [20], we relax the condition on the nonvanishing quantity and extend our analysis to the case where  $\beta \neq 0$ .

It is noteworthy that we allow V to be 0 at  $q_i$  for any  $i = 1, \dots, \iota$  where  $\iota \in \mathbb{N}$ . So, it is also possible to establish blow-up solutions for the following singular problem:

(5) 
$$\begin{cases} -\Delta_g \tilde{u} + \beta \tilde{u} = \lambda \left( \frac{\tilde{V}e^{\tilde{u}}}{\int_{\Sigma} \tilde{V}e^{\tilde{u}}dv_g} - \frac{1}{|\Sigma|_g} \right) - 4\pi \sum_{i=1}^{\iota} n_i \left( \delta_{q_i} - \frac{1}{|\Sigma|_g} \right) & \text{in } \mathring{\Sigma} \\ \partial_{\nu_g} \tilde{u} = 0 & \text{on } \partial \Sigma \end{cases}$$

Here,  $\tilde{V}$  is a positive smooth function, and  $n_i \in \mathbb{N}$  for  $i = 1, \dots, \iota$ . Notably, the problem (5) emerges as a specific instance of (4). To elucidate, we define the Green's function through the following equations for any  $\xi \in \Sigma$ :

(6) 
$$\begin{cases} -\Delta_g G^g(x,\xi) + \beta G^g(x,\xi) = \delta_{\xi} - \frac{1}{|\Sigma|_g} & x \in \mathring{\Sigma} \\ \partial_{\nu_g} G^g(x,\xi) = 0 & x \in \partial \Sigma \\ \int_{\Sigma} G^g(x,\xi) d\nu_g(x) = 0 \end{cases}$$

We take  $u(x) = \tilde{u}(x) + 4\pi \sum_{i=1}^{l} n_i G^g(x, q_i)$  and  $V(x) = \tilde{V}(x) e^{-4\pi \sum_{i=1}^{l} n_i G^g(x, q_i)}$ . u satisfies the equations (4) in which V is a non-negative smooth function with the zero set  $\{q_1, \dots, q_t\}$ .

We now present the main results, starting with defining the "stable" critical points set as in [13, 16, 25].

**Definition 1.1.** Let  $F: D \to \mathbb{R}$  be a  $C^1$ -function and K be a compact subset of critical points of F, i.e

$$K\subset\subset\{x\in D:\nabla F(x)=0\}.$$

A critical set K is  $C^1$ -stable if for any closed neighborhood U of K in D, there exists  $\varepsilon > 0$  such that if  $G: D \to \mathbb{R}$  is a  $C^1$ -function with  $|F - G|_{C^1(U)} < \varepsilon$ , then G has at least one critical point in U.

The main theorem asserts the existence of a sequence of blow-up solutions for (4), with these solutions exhibiting blow-up behaviour at the stable critical points of a function  $\mathcal{F}_{k,m}$  (refer to Section 2). The function is defined on  $\mathscr{D} := \mathring{\Sigma}^k \times (\partial \Sigma)^{m-k} \setminus \{\xi = (\xi_1, \dots, \xi_m) : \xi_i = \xi_j \text{ for some } i \neq j \text{ or } V(\xi_i) = 0 \text{ for some } i = 1, \dots, m\}$ . Specifically,  $\mathcal{F}_{k,m} : \mathscr{D} \subset \mathring{\Sigma}^k \times (\partial \Sigma)^{m-k}$ 

$$(\partial \Sigma)^{m-k} \to \mathbb{R},$$

$$\mathcal{F}_{k,m}(\xi_1,\cdots,\xi_m) = \sum_{i=1}^m \varrho^2(\xi_i) H^g(\xi_i,\xi_i) + \sum_{\substack{i,j=1\\i\neq j}}^m \varrho(\xi_i) \varrho(\xi_j) G^g(\xi_i,\xi_j)$$

$$+\sum_{i=1}^{m} 2\varrho(\xi_i) \log V(\xi_i),$$

where  $\varrho(\xi)$  equals  $8\pi$ , if  $\xi \in \overset{\circ}{\Sigma}$  and equals  $4\pi$ , if  $\xi \in \partial \Sigma$ .

**Theorem 1.1.** For any given integers  $0 \le k \le m$ , if  $K \subset \mathcal{D}$  is a  $C^1$ -stable critical point set of  $\mathcal{F}_{k,m}$ , then for any  $n \in \mathbb{N}+$ , there exists  $\lambda_n$  in a small neighborhood of  $\lambda_{k,m}$  and a solution  $u_n$  of the problem (4) with respect to the parameter  $\lambda_n$  such that  $\lambda_n \to \lambda_{k,m}$  as  $n \to +\infty$ . Furthermore, the sequence of solutions  $\{u_n\}$  blows up precisely at points  $\xi_1, \dots, \xi_m$ , such that  $(\xi_1, \dots, \xi_m) \in K$  (up to a subsequence), in a sense that

$$\frac{\lambda_n V e^{u_n}}{\int_{\Sigma} V e^{u_n} dv_g} \to \sum_{i=1}^m \varrho(\xi_i) \delta_{\xi_i}, \text{ as } n \to +\infty$$

which is convergent as measures on  $\Sigma$ .

Remark 1.1. If the zero set of V is empty, the behavior of  $\mathcal{F}_{k,m}$  as  $\xi$  approaches  $\partial \mathscr{D}$  results in its divergence towards  $+\infty$  (as in [20, Lemma 7.3]). This divergence suggests the presence of at least one global minimum point within the interior of  $\mathscr{D}$ . Additionally, a local minimum point is inherently "stable." Consequently, in this scenario, the blow-up solutions must exist. However, when V(q) = 0 for some  $q \in \Sigma$ , a complication arises. As  $\xi$  approaches  $\partial \mathscr{D}$ , there are cases where the sum of the first terms tends to  $+\infty$  while the last term approaches  $-\infty$ , leading to an indeterminate behaviour of  $\mathcal{F}_{k,m}$ .

# 2 Preliminaries

Throughout this paper, we use the terms "sequence" and "subsequence" interchangeably, as the distinction is not crucial for the context of our analysis. The constant denoted by C in our deduction may assume different values across various equations or even within different lines of equations. We denote  $B_r(y) := \{y \in \mathbb{R}^2 : |y| < r\}, A_r(y) := B_{2r}(y) \setminus B_r(y), B_r := B_r(0)$  and  $A_r := A_r(0)$ .

To construct the ansatz for solutions of problem (4), we firstly introduce a family of isothermal coordinates (see [11,15,37], for instance). For any  $\xi \in \mathring{\Sigma}$ , there exists an isothermal

coordinate system  $(U(\xi), y_{\xi})$  such that  $y_{\xi}$  maps an open neighborhood  $U(\xi)$  around  $\xi$  onto an open disk  $B^{\xi}$  with radius  $2r_{\xi}$  in which  $g = \sum_{i=1}^{2} e^{\hat{\varphi}_{\xi}(y_{\xi}(x))} dx^{i} \otimes dx^{i}$ . W.l.o.g., we can assume that  $y_{\xi}(\xi) = (0,0)$  and  $\overline{U(\xi)} \subset \mathring{\Sigma}$ . For  $\xi \in \partial \Sigma$  there exists an isothermal coordinate system  $(U(\xi), y_{\xi})$  around  $\xi$  such that the image of  $y_{\xi}$  is a half disk  $B^{\xi} := \{y = (y_{1}, y_{2}) \in \mathbb{R}^{2} : |y| < 2r_{\xi}, y_{2} \geq 0\}$  with a radius  $2r_{\xi}$  and  $y_{\xi}(U(\xi) \cap \partial \Sigma) = \{y = (y_{1}, y_{2}) \in \mathbb{R}^{2} : |y| < 2r_{\xi}, y_{2} = 0\}$  with  $g = \sum_{i=1}^{2} e^{\hat{\varphi}_{\xi}(y_{\xi}(x))} dx^{i} \otimes dx^{i}$ . W.l.o.g., we can assume that  $y_{\xi}(\xi) = (0,0)$ . Here  $\hat{\varphi}_{\xi} : B^{\xi} \to \mathbb{R}$  is related to  $K_{g}$ , the Gaussian curvature of  $\Sigma$ , by the equation

(7) 
$$-\Delta \hat{\varphi}_{\xi}(y) = 2K_g(y_{\xi}^{-1}(y))e^{\hat{\varphi}_{\xi}(y)} \quad \text{for all } y \in B^{\xi}.$$

For  $\xi \in \Sigma$  and  $0 < r \le 2r_{\xi}$  we set

$$B_r^{\xi} := B^{\xi} \cap \{ y \in \mathbb{R}^2 : |y| < r \} \text{ and } U_r(\xi) := y_{\xi}^{-1}(B_r^{\xi}).$$

Both  $y_{\xi}$  and  $\hat{\varphi}_{\xi}$  are assumed to depend smoothly on  $\xi$  as in [15]. Additionally,  $\hat{\varphi}_{\xi}$  satisfies  $\hat{\varphi}_{\xi}(0) = 0$  and  $\nabla \hat{\varphi}_{\xi}(0) = 0$ . Specifically, as in [37] the Neumann boundary conditions preserved by the isothermal coordinates as for any  $\xi \in \partial \Sigma$  and  $x \in y_{\xi}^{-1}(B^{\xi} \cap \partial \mathbb{R}^{2}_{+})$ , we have

(8) 
$$(y_{\xi})_* (\nu_g(x)) = -e^{-\frac{\hat{\varphi}_{\xi}(y)}{2}} \frac{\partial}{\partial y_2} \Big|_{y=y_{\xi}(x)}.$$

Let  $\chi$  be a radial cut-off function in  $C^{\infty}(\mathbb{R},[0,1])$  such that  $\chi(s)=\begin{cases} 1, & \text{if } |s|\leq 1\\ 0, & \text{if } |s|\geq 2 \end{cases}$ . And we denote that  $\chi_{\xi}(x)=\chi\left(\frac{4|y_{\xi}(x)|}{r_{\xi}}\right)$  and  $\varphi_{\xi}(x)=\hat{\varphi}_{\xi}(y_{\xi}(x))$ . Let the function

$$\Gamma_{\xi}^{g}(x) = \Gamma^{g}(x,\xi) = \begin{cases} \frac{1}{2\pi} \chi_{\xi}(x) \log \frac{1}{|y_{\xi}(x)|} & \text{if } \xi \in \mathring{\Sigma} \\ \frac{1}{\pi} \chi_{\xi}(x) \log \frac{1}{|y_{\xi}(x)|} & \text{if } \xi \in \partial \Sigma \end{cases}.$$

Decomposing the Green's function  $G^g(x,\xi) = \Gamma_{\xi}^g(x) + H_{\xi}^g(x)$ , we have the function  $H_{\xi}^g(x) := H^g(x,\xi)$  that solves the following equations:

(9) 
$$\begin{cases} -\Delta_g H_{\xi}^g + \beta H_{\xi}^g &= -\beta \frac{4}{\varrho(\xi)} \chi_{\xi} \log \frac{1}{|y_{\xi}|} + \frac{4}{\varrho(\xi)} (\Delta_g \chi_{\xi}) \log \frac{1}{|y_{\xi}|} \\ &+ \frac{8}{\varrho(\xi)} \langle \nabla \chi_{\xi}, \nabla \log \frac{1}{|y_{\xi}|} \rangle_g - \frac{1}{|\Sigma|_g}, & \text{in } \mathring{\Sigma} \\ \partial_{\nu_g} H_{\xi}^g &= -\frac{4}{\varrho(\xi)} (\partial_{\nu_g} \chi_{\xi}) \log \frac{1}{|y_{\xi}|} - \frac{4}{\varrho(\xi)} \chi_{\xi} \partial_{\nu_g} \log \frac{1}{|y_{\xi}|}, & \text{on } \partial \Sigma \end{cases}$$

$$\int_{\Sigma} H_{\xi}^g dv_g &= -\frac{4}{\varrho(\xi)} \int_{\Sigma} \chi_{\xi} \log \frac{1}{|y_{\xi}|} dv_g$$

By the regularity of elliptic equations (see [26]), there is a unique smooth solution  $H^g(x,\xi)$  that solves (9) in  $C^{1,\alpha}(\Sigma)$ .  $H^g(x,\xi)$  is the regular part of  $G^g(x,\xi)$  and  $H^g(\xi,\xi)$  is also called

Robin's function on  $\Sigma$ . It is easy to check that  $H^g(\xi, \xi)$  is independent of the choice of the cut-off function  $\chi$  and the local chart. Denote

$$\Delta = \{\xi := (\xi_1, \dots, \xi_m) \in \mathring{\Sigma}^k \times (\partial \Sigma)^{m-k} : \xi_i = \xi_j \text{ for some } i \neq j\},$$
 and  $Q := \{\xi := (\xi_1, \dots, \xi_m) \in \mathring{\Sigma}^k \times (\partial \Sigma)^{m-k} : V(\xi_i) = 0 \text{ for some } i = 1, \dots, m\}.$  Let 
$$\mathscr{D} = \mathring{\Sigma}^k \times (\partial \Sigma)^{m-k} \setminus (\Delta \cup Q).$$

For  $\delta > 0$ , we consider

$$M_{\delta}: = \{ \xi \in \mathscr{D} : d_g(\xi_i, \partial \Sigma) \ge \delta \text{ for } i = 1, \dots, k;$$
  
$$d_g(\xi_i, \xi_i) \ge \delta \text{ for any } i \ne j; V(\xi_i) \ge \delta \text{ for } i = 1, \dots, m \},$$

a compact subset  $\mathring{\Sigma}^k \times (\partial \Sigma)^{m-k}$ , where  $d_g(\cdot, \cdot) : \Sigma \times \Sigma \to \mathbb{R}$  is the geodesic distance with respect to metric g and  $d_g(p, \partial \Sigma) := \inf_{q \in \partial \Sigma} d_g(p, q)$  for any  $p \in \mathring{\Sigma}$ . We are now in a position to introduce the function  $\mathcal{F}_{k,m} : M_{\delta} \to \mathbb{R}$ ,

$$\xi \mapsto \sum_{i=1}^{m} 2\varrho(\xi_i) \log V(\xi_i) + \sum_{i=1}^{m} \varrho(\xi_i)^2 H^g(\xi_i, \xi_i) + \sum_{i,j=1}^{m} \varrho(\xi_i) \varrho(\xi_j) G^g(\xi_i, \xi_j).$$

$$i, j = 1$$

$$i \neq j$$

We observe that for any  $\alpha \in (0,1)$ ,  $G^g(x,\xi) \in C^{\infty}(\Sigma \setminus \{\xi\})$  and  $H^g(x,\xi)$  is  $C^{1,\alpha}(\Sigma)$ , too. Thus  $\mathcal{F}_{k,m}$  is  $C^{1,\alpha}(M_{\delta})$  for any fixed  $\delta > 0$ .

Let  $u_n$  be a sequence of solutions of (4) with respect to the parameter  $\lambda$  as  $\lambda \to \lambda_{k,m}$ .

To study the blow-up solutions of (4), we consider the weak solution of the following problem in the space  $\overline{\mathbf{H}}^1 := \left\{ u \in H^1(\Sigma) : \int_{\Sigma} u dv_g = 0 \right\}$ ,

(10) 
$$\begin{cases} (-\Delta_g + \beta)u = \rho^2 V e^u - \overline{\rho^2 V e^u} & \text{in } \mathring{\Sigma} \\ \partial_{\nu_g} u = 0 & \text{on } \partial \Sigma \end{cases}$$

such that  $\rho^2 V e^u \rightharpoonup \sum_{i=1}^m \varrho(\xi_i) \delta_{\xi_i}$ , weakly convergent in a sense of measure in  $\Sigma$  as  $\rho \to 0$ , for some  $\xi = (\xi_1, \dots, \xi_m) \in \mathscr{D}$ . If we take  $\lambda = \rho^2 \int_{\Sigma} V e^u dv_g$ , the weak solutions of (10) must be the weak solutions of (4). So we try to construct a sequence of blow-up solutions of (10) as  $\rho \to 0$  and then pass back to the original problem (4) as  $\lambda \to \lambda_{k,m}$ .

It is well known that  $u_{\tau,\eta}(y) = \log \frac{8\tau^2}{(\tau^2\rho^2 + |y-\eta|^2)^2}$  for  $(\tau,\eta) \in (0,\infty) \times \mathbb{R}^2$  are all the solutions of the Liouville-type equations,

$$\begin{cases} -\Delta u = \rho^2 e^u \text{ in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^u < \infty. \end{cases}$$

Our goal is to construct approximate solutions of (10) applying the pull-back of  $u_{\tau,\eta}$  to  $\Sigma$  by isothermal coordinates and selecting appropriate values for  $\tau$  and  $\xi$ . Define

$$U_{\tau,\xi}(x) = u_{\tau,0}(y_{\xi}(x)) = \log \frac{8\tau^2}{(\tau^2 \rho^2 + |y_{\xi}(x)|^2)}, \text{ for all } x \in y_{\xi}^{-1}(B^{\xi}).$$

For any function  $f \in L^1(\Sigma)$ , we denote its mean over  $\Sigma$  as  $\overline{f} = \frac{1}{|\Sigma|g} \int_{\Sigma} f dv_g$ . Then, we introduce a projection operator P, which is used to project  $U_{\tau,\xi}$  into the space  $\overline{H}^1$ . The projected function  $PU_{\tau,\xi}$  is defined as the solution to the problem:

(11) 
$$\begin{cases} (-\Delta_g + \beta)PU_{\tau,\xi}(x) = \rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} - \overline{\rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}}}, & x \in \mathring{\Sigma}, \\ \partial_{\nu_g} PU_{\tau,\xi} = 0, & x \in \partial \Sigma, \\ \int_{\Sigma} PU_{\tau,\xi} dv_g = 0, & x \in \partial \Sigma, \end{cases}$$

For  $\beta \neq 0$ , the last condition of zero integral of  $PU_{\tau,\xi}$  over  $\Sigma$  can be inferred from the preceding equations via the divergence theorem. However, it explicitly addresses the case when  $\beta = 0$ , ensuring the solution criteria for all  $\beta \geq 0$ . The solution of (11) is unique within  $\overline{H}^1$  and  $PU_{\tau,\xi}$  in  $C^{\infty}(\Sigma)$  as per regularity theory (see [26]), ensuring that  $PU_{\tau,\xi}$  is well-defined.

Let

$$\psi_{\tau,\eta}^{0}(y) = \frac{\partial}{\partial_{\tau}} u_{\tau,\eta}(x) = \frac{2}{\tau} \frac{|y - \eta|^{2} - \tau^{2} \rho^{2}}{|y - \eta|^{2} + \tau^{2} \rho^{2}},$$

and

$$\psi_{\tau,\eta}^{j}(y) = \frac{\partial}{\partial \eta_{j}} u_{\tau,\eta}(x) = 4 \frac{y_{j} - \eta_{j}}{\tau^{2} \rho^{2} + |y - \eta|^{2}},$$

for j=1,2. It is observed that the derivatives above satisfy the equation:  $-\Delta \psi = \rho^2 e^{u_{\tau,\eta}} \psi$  in  $\mathbb{R}^2$ , where  $\psi = \psi^j_{\tau,\eta}$ , for j=0,1,2. The function  $\Psi^j_{\tau,\xi}$  is then defined as the pull-back of  $\psi^j_{\tau,0}$  under the isothermal coordinate  $y_{\xi}$ , i.e.,  $\Psi^j_{\tau,\xi}(x) = \psi^j_{\tau,0}(y_{\xi}(x))$ , for any  $x \in y^{-1}_{\xi}(B^{\xi}_{2r_0})$ . Let  $P\Psi^j_{\tau,\xi}$  be a projection into  $\overline{\mathbb{H}}^1$  of  $\Psi^j_{\tau,\xi}$ , for  $\xi \in \Sigma$  and  $j=0,1,..,J_i$ , where  $J_i$  equals 2 if  $1 \leq i \leq k$  and equals 1 if  $k+1 \leq i \leq m$ .  $P\Psi^j_{\tau,\xi}$  is defined as the solution of

(12) 
$$\begin{cases} (-\Delta_g + \beta)P\Psi^j_{\tau,\xi} = \rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^j_{\tau,\xi} - \overline{\rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^j_{\tau,\xi}}, & x \in \mathring{\Sigma}, \\ \partial_{\nu_g} P\Psi^j_{\tau,\xi} = 0, & x \in \partial \Sigma, \\ \int_{\Sigma} P\Psi^j_{\tau,\xi} = 0. & \end{cases}$$

By the regularity theory, as referenced in [26], the solution to problem (12) is unique and smooth on  $\Sigma$ . Hence,  $P\Psi_{\tau,\xi}^j$  is well-defined and lies in the space  $C^{\infty}(\Sigma)$ .

For any configuration  $\xi = (\xi_1, \dots, \xi_m) \in M_\delta$ , we can establish an isothermal chart around  $y_{\xi_i}$  for each point  $\xi_i$  for  $i = 1, 2, \dots, m$ . Given the compactness of  $\Sigma$ , it is possible to select a uniform radius  $r_{\xi_i} > 0$  for any  $\xi \in M_\delta$ , denoted as  $4r_0$ . This radius is sufficiently small

and depends only on  $\delta$  and  $\partial \Sigma$ . Moreover, we ensure that  $y_{\xi_i}^{-1}(B_{2r_0}^{\xi_i}) \cap y_{\xi_j}^{-1}(B_{2r_0}^{\xi_j}) = \emptyset$  for any  $i, j = 1, 2, \dots, m$  with  $i \neq j$ . For any  $i = 1, \dots, m$ , we define the scaling parameter  $\tau_i$  as:

(13) 
$$\tau_i(\xi) = \sqrt{\frac{1}{8}V(\xi_i)e^{\varrho(\xi_i)H(\xi_i,\xi_i) + \sum_{h \neq i} \varrho(\xi_h)G(\xi_i,\xi_h)}}.$$

For simplicity, we denote that  $U_i = U_{\tau_i(\xi),\xi_i}$ ,  $\chi_i = \chi_{\xi_i}$ ,  $\varphi_i := \varphi_{\xi_i}$ ,  $\hat{\varphi}_i = \hat{\varphi}_{\xi_i}$  and  $\tau_i = \tau_i(\xi)$ . The formulation of the scaling parameter  $\tau_i$  is chosen for technical considerations.

We consider the manifold for given integers  $m \ge k \ge 0$  and a positive constant  $\rho > 0$ ,

$$\mathcal{M}_{\rho}^{k,m} := \left\{ \sum_{i=1}^{m} PU_i : \xi_i \in \mathring{\Sigma} \text{ for } i = 1, 2, \cdots, k \text{ and } \xi_i \in \partial \Sigma \text{ for } i = k+1, \cdots, m \right\}.$$

The functions in manifold  $\mathcal{M}_{\rho}^{k,m}$  serve as approximate solutions of the problem (10). We then denote the projected function for any  $i = 1, 2, \dots, m$  and  $j = 0, 1, \dots, J_i$  as

$$P\Psi_i^j := P\Psi_{\tau_i(\xi),\xi_i}^j.$$

These projected functions generate a subspace of  $\overline{\mathbf{H}}^1$ ,  $\{P\Psi_i^j: i=1,\cdots,m,j=1,\cdots,J_i\}$  denoted as  $K_{\xi}$ . Furthermore, we introduce an inner product for the space  $\overline{\mathbf{H}}^1$  as follows:

$$\langle \psi, \phi \rangle := \int_{\Sigma} \langle \nabla \psi, \nabla \phi \rangle_g dv_g + \beta \int_{\Sigma} \psi \phi dv_g \text{ for any } \psi, \phi \in \overline{\mathbf{H}}^1.$$

The orthogonal complement concerning this inner product, denoted as  $K_{\xi}^{\perp}$ , is then:

$$K_{\xi}^{\perp} = \left\{ \phi \in \overline{\mathbf{H}}^1 : \langle \phi, f \rangle = 0 \text{ for all } f \in K_{\xi} \right\}.$$

We also introduce  $\Pi_{\xi}: \overline{\mathbb{H}}^1 \to K_{\xi}$  and  $\Pi_{\xi}^{\perp}: \overline{\mathbb{H}}^1 \to K_{\xi}^{\perp}$  as the orthogonal projections onto  $K_{\xi}$  and  $K_{\xi}^{\perp}$ , respectively. The solution u can decompose into two parts: one part lies on the manifold  $\mathcal{M}_{\rho}^{k,m}$ ; the other part is on  $K_{\xi}^{\perp}$  near the orthogonal space of the tangent space of the manifold  $\mathcal{M}_{\rho}^{k,m}$ , i.e.,  $u = \sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho}$ , where  $\phi_{\xi}^{\rho}$  is the remainder term.

# 3 The Lyapunov-Schmidt reduction

Utilizing the Moser-Trudinger type inequality on compact Riemann surfaces, as in [36], we have

$$\sup_{\int_{\Sigma} |\nabla_g u|^2 dv_g = 1, \int_{\Sigma} u dv_g = 0} \int e^{2\pi u^2} dv_g < +\infty.$$

Since  $\left(\int_{\Sigma} |\nabla u|_g^2 dv_g + \beta \int_{\Sigma} |u|^2 dv_g\right)^{\frac{1}{2}}$  and  $\left(\int_{\Sigma} |\nabla u|_g^2 dv_g\right)^{\frac{1}{2}}$  are equivalent norms in the Hilbert space  $\overline{H}^1$ , it follows that

$$\begin{split} \log \int_{\Sigma} e^{u} dv_{g} &\leq \log \int_{\Sigma} e^{2\pi \frac{u^{2}}{\|u\|^{2}} + \frac{1}{8\pi} \|u\|^{2}} dv_{g} \quad \text{(by Young's Inequality)} \\ &= \frac{1}{8\pi} \int_{\Sigma} |\nabla_{g} u|^{2} dv_{g} + C, \\ &\leq \frac{1}{8\pi C} \langle u, u \rangle + C \text{ for any } u \in \overline{\mathbf{H}}^{1}, \end{split}$$

where C>0 is a constant. Consequently,  $\overline{\mathrm{H}}^1\to L^p(\Sigma), u\mapsto e^u$  is continuous. For any p>1, let  $i_p^*:L^p(\Sigma)\to\overline{\mathrm{H}}^1$  be the adjoint operator corresponding to the immersion  $i:\overline{\mathrm{H}}^1\to L^{\frac{p}{p-1}}$  and  $i^*:\cup_{p>1}L^p(\Sigma)\to\overline{\mathrm{H}}^1$ . For any  $f\in L^p(\Sigma)$ , we define that  $i^*(f):=i^*(f-\bar{f})$ , i.e., for any  $h\in\overline{\mathrm{H}}^1,\ \langle i^*(f),h\rangle=\int_{\Sigma}(f-\bar{f})hdv_g$ .

The problem (10) has the following equivalent form,

(14) 
$$\begin{cases} u = i^*(\rho^2 V e^u) \\ u \in \overline{H}^1 \end{cases}.$$

### 3.1 The linear operator

We consider the linear operator

$$L_{\varepsilon}^{\rho}(\phi) := \Pi_{\varepsilon}^{\perp}(\phi - i^*(\rho^2 V e^{\sum_{i=1}^{m} PU_i}\phi))$$

for any fixed  $\xi \in M_{\delta}$ . The following lemma shows that the linear operator is invertible for fixed  $\rho$ , and the norm of the inverse operator is controlled by  $|\log \rho|$  as  $\rho \to 0$ , which is a key lemma to solve the problem (10).

**Lemma 3.1.** For any  $\delta > 0$ , let  $\xi = (\xi_1, \dots, \xi_m) \in M_{\delta}$ . There exists  $\rho_0 > 0$  and a constant c > 0 such that for any  $\rho \in (0, \rho_0)$  we have

$$||L_{\xi}^{\rho}(\phi)|| \geqslant \frac{c}{|\log \rho|} ||\phi||, \quad \forall \phi \in K_{\xi}^{\perp}.$$

In particular, the operator  $L_{\xi}^{\rho}$  is invertible and  $\left\| \left( L_{\xi}^{\rho} \right)^{-1} \right\| \leqslant |\log \rho|/c$ .

By [16], the proof of Lemma 3.1 is relatively standard, which is given in Appendix D.

Fix  $\sum_{i=1}^{m} PU_i$ , we try to obtain the solution of

$$\Pi_{\xi}^{\perp} \left( \sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho} - i^* (\rho^2 e^{\sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho}}) \right) = 0,$$

for  $\phi_{\xi}^{\rho} \in K_{\xi}^{\perp}$  applying the fixed-point theorem. Then, it is reduced to a finite-dimensional problem.

**Proposition 3.1.** For any  $\delta > 0$ , and  $\xi = (\xi_1, \dots, \xi_m) \in M_\delta$ . For any  $p \in (1, \frac{6}{5})$  there exist  $\rho_0 > 0$  and R > 0 (uniformly in  $\xi$ ) such that for any  $\rho \in (0, \rho_0)$  there is a unique  $\phi_{\xi}^{\rho} \in K_{\xi}^{\perp}$  such that

(15) 
$$\Pi_{\xi}^{\perp} \left[ \sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho} - i^* \left( \rho^2 e^{\sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho}} \right) \right] = 0.$$

and

$$\|\phi_{\xi}^{\rho}\| \le R\rho^{\frac{2-p}{p}} |\log \rho|.$$

*Proof.* Define an operator  $T_{\xi}^{\rho}$  on  $K_{\xi}^{\perp}$  as follows:

$$T_{\xi}^{\rho}(\phi) = \left[ \left( L_{\xi}^{\rho} \right)^{-1} \circ \Pi_{\xi}^{\perp} \circ i^{*} \right] M_{\xi}^{\rho}(\phi)$$

$$M_{\xi}^{\rho}(\phi) = \rho^{2} e^{\sum_{i=1}^{m} PU_{i}} \left[ e^{\phi} - 1 - \phi \right] + \left[ \rho^{2} e^{\sum_{i=1}^{m} PU_{i}} - \rho^{2} \sum_{i=1}^{m} \chi_{i} e^{-\varphi_{\xi_{i}}} e^{U_{i}} \right]$$

We observe that  $i^*(\rho^2 \sum_{i=1}^m \chi_i e^{-\varphi_{\xi_i}} e^{U_i}) = \sum_{i=1}^m PU_i$ . It is obvious that  $\phi$  is a fixed point of  $T_{\xi}^{\rho}$  if and only if  $\phi$  solves (15) on  $K_{\xi}^{\perp}$ .

Claim. There exists  $\rho_0$  and R > 0 such that  $T_{\xi}^{\rho}$  is a contraction map for any  $\rho \in (0, \rho_0)$  and  $|\phi| \leq R\rho^{\frac{2-p}{p}} |\log \rho|$ .

Applying Lemma 3.1, Lemma C.1, Lemma C.2 and the Moser-Trudinger inequality, we obtain

$$\begin{split} \|T_{\xi}^{\rho}(\phi)\| & \leq C|\log\rho|\|i^{*}\circ M_{\xi}^{\rho}(\phi)\| \leq C|\log\rho| \left|M_{\xi}^{\rho}(\phi)\right|_{L^{p}(\Sigma)} \\ & \leq C|\log\rho| \left(\left|\rho^{2}Ve^{\sum_{i=1}^{m}PU_{i}}(e^{\phi}-1-\phi)\right|_{L^{p}(\Sigma)} \right. \\ & + \left|\rho^{2}Ve^{\sum_{i=1}^{m}PU_{i}}-\rho^{2}\sum_{i=1}^{m}\chi_{i}e^{U_{i}}\right|_{L^{p}(\Sigma)} \right) \\ & \leq C|\log\rho| \left(\|\phi\|^{2}e^{c_{2}\|\phi\|^{2}}\rho^{\frac{2-2pr}{pr}}+\rho^{\frac{2-p}{p}}\right), \end{split}$$

where  $c_2 > 0$  is a constant, r > 1 is sufficiently close to 1, and  $p \in (1, \frac{6}{5})$ . We then fix arbitrary  $p \in (1, \frac{6}{5})$  and choose R > 0 large enough such that  $C(1 + e^{c_2}) \le R$ . Next, we select  $\rho_1 > 0$ 

such that  $\max\{R\rho^{\frac{2-2pr}{pr}+\frac{2-p}{p}}|\log\rho|, R\rho^{\frac{2-p}{p}}|\log\rho|\} \le 1$  for all  $\rho \in (0, \rho_1)$ . Consequently, for any  $|\phi| \le R\rho^{\frac{2-p}{p}}|\log\rho|$ , we have  $|T_{\xi}^{\rho}| \le R\rho^{\frac{2-p}{p}}|\log\rho|$ . And similarly, by Lemma C.2 we deduce that

$$\begin{split} \|T_{\xi}^{\rho}(\phi_{1}) - T_{\xi}^{\rho}(\phi_{2})\| & \leq C' |\log \rho| \left| \rho^{2} e^{\sum_{i=1}^{m} PU_{i}} (e^{\phi_{1}} - e^{\phi_{2}} - (\phi_{1} - \phi_{2})) \right|_{L^{p}(\Sigma)} \\ & \leq C' |\log \rho| e^{c_{2}(\sum_{j=1}^{2} \|\phi_{j}\|^{2})} \rho^{\frac{2-2pr}{pr}} (\sum_{j=1}^{2} \|\phi_{j}\|)) \|\phi_{1} - \phi_{2}\| \\ & \leq 2RC' e^{2c_{2}} \rho^{\frac{2-2pr}{pr} + \frac{1+\alpha_{0}-p}{p}} \log^{2} \rho \|\phi_{1} - \phi_{2}\| \leq \frac{1}{2} \|\phi_{1} - \phi_{2}\|, \end{split}$$

uniformly for any  $\rho \in (0, \rho_2)$  and  $\xi \in M_\delta$ , where  $\rho_2 > 0$  is chosen such that

$$\max\{R\rho^{\frac{2-p}{p}}|\log\rho|, 2RC'e^{c_2}\rho^{\frac{2-2pr}{pr}+\frac{2-p}{p}}\log^2\rho\} < \frac{1}{2}$$

for any  $\rho \in (0, \rho_2)$ . We then define  $\rho_0 = \min\{\rho_1, \rho_2\}$ . Thus  $T_{\xi}^{\rho}(\phi)$  is a contraction map on  $\{\phi \in K_{\xi}^{\perp} : \|\phi\| \leq R\rho^{\frac{2-p}{p}} |\log \rho|\}$ . By the contracting-mapping principle, there exists a unique fixed point of  $T_{\xi}^{\rho}$  on  $\{\phi \in K_{\xi}^{\perp} : \|\phi\| \leq R\rho^{\frac{2-p}{p}} |\log \rho|\}$ .

# 4 The reduced functional and its expansion

The associated functional  $E_{\rho}(u)$  of the problem (10) is defined as following,

(16) 
$$E_{\rho}(u) = \frac{1}{2} \int_{\Sigma} (|\nabla u|_g^2 + \beta |u|^2) dv_g - \rho^2 \int_{\Sigma} V e^u dv_g.$$

Assume u has the form  $\sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho}$ , where  $\phi_{\xi}^{\rho}$  is obtained by Proposition 3.1, the reduced functional can be obtained  $\tilde{E}_{\rho}(\xi) := E_{\rho}(\sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho})$  with  $\|\phi_{\xi}^{\rho}\| \leq R\rho^{\frac{2-p}{p}} |\log \rho|$ , i.e.,

$$\widetilde{E}_{\rho}(\xi) := \frac{1}{2} \int_{\Sigma} \left( |\nabla (\sum_{i=1}^{m} PU_{i} + \phi_{\xi}^{\rho})|_{g}^{2} + \beta \left| \sum_{i=1}^{m} PU_{i} + \phi_{\xi}^{\rho} \right|^{2} \right) dv_{g}$$
$$-\rho^{2} \int_{\Sigma} V e^{\sum_{i=1}^{m} PU_{i} + \phi_{\xi}^{\rho}} dv_{g}.$$

And  $\tilde{E}_{\rho}$  has a  $C^1$ -expansion concerning  $\xi$  as the following expansion proposition.

**Proposition 4.1.** As  $\rho \to 0$ ,

$$\widetilde{E}_{\rho}(\xi) = \sum_{i=1}^{m} \varrho(\xi_{i})(3\log 2 - 2\log \rho) - 2\sum_{i=1}^{m} \varrho(\xi_{i}) - \frac{1}{2}\mathcal{F}_{k,m}(\xi) + o(1),$$

 $C^1$ -uniformly in any compact sets of  $\mathscr{D}$ .

*Proof.* Denote  $\phi = \phi_{\xi}^{\rho}$  to simplify the notation. Then

$$\widetilde{E}_{\rho}(\xi) = \frac{1}{2} \left( \sum_{i=1}^{m} \langle PU_i, PU_i \rangle + \sum_{i \neq j} \langle PU_i, PU_j \rangle \right) + \frac{1}{2} \left( \|\phi\|^2 + 2 \sum_{i=1}^{m} \langle PU_i, \phi \rangle \right)$$
$$- \int_{\Sigma} \rho^2 V e^{\sum_{i=1}^{m} PU_i} dv_g - \int_{\Sigma} \rho^2 (V e^{\sum_{i=1}^{m} PU_i + \phi} - V e^{\sum_{i=1}^{m} PU_i}) dv_g.$$

We notice that  $|e^s - 1| \le e^{|s|} |s| (\forall s \in \mathbb{R})$ . By Lemma B.2,

$$\left| \int_{\Sigma} \rho^{2} (Ve^{\sum_{i=1}^{m} PU_{i} + \phi} - Ve^{\sum_{i=1}^{m} PU_{i}}) dv_{g} \right| \leq \left| \int_{\Sigma} \rho^{2} Ve^{\sum_{i=1}^{m} PU_{i}} e^{|\phi|} |\phi| dv_{g} \right|$$

$$\leq \mathcal{O} \left( \rho^{2} \left( \int_{\Sigma} e^{r \sum_{i=1}^{m} PU_{i}} dv_{g} \right)^{1/r} |e^{|\phi|}|_{L^{s}(\Sigma)} |\phi|_{L^{t}(\Sigma)} \right)$$

$$\leq \mathcal{O} \left( \left| \rho^{2} Ve^{\sum_{i=1}^{m} PU_{i}} \right|_{L^{r}(\Sigma)} \|\phi\| \right)$$

$$= o(1)(\rho \to 0),$$

where  $r \in (1,2)$  with  $\frac{1}{s} + \frac{1}{r} + \frac{1}{t} = 1$  and  $\frac{2(1-r)}{r} + \frac{2-p}{p} > 0$ . By Lemma E.1 and Lemma E.2, as  $\rho \to 0$ 

(17) 
$$\widetilde{E}_{\rho}(\xi) = \sum_{i=1}^{m} \varrho(\xi_{i})(3\log 2 - 2\log \rho) - 2\sum_{i=1}^{m} \varrho(\xi_{i}) - \frac{1}{2}\mathcal{F}_{k,m}(\xi) + o(1).$$

By (15),

(18) 
$$\sum_{i=1}^{m} PU_i + \phi - i^* \left( \rho^2 V e^{\sum_{i=1}^{m} PU_i + \phi} \right) = \sum_{s=1}^{m} \sum_{t=1}^{J_t} c_{st}^{\rho} P\Psi_s^t,$$

where  $c_{st}^{\rho}$  are coefficients. We combine (18) with Lemma E.6

(19) 
$$\sum_{i=1}^{m} \sum_{j=1}^{J_i} \left| c_{ij}^{\rho} \right| = \mathcal{O}(\rho^2),$$

via Lemma B.1 and Remark B.2. For the  $C^1$ -expansion, by Lemma B.3 and Lemma E.6,

$$\partial_{(\xi_h)_j} E_{\rho} \left( \sum_{i=1}^m PU_i + \phi \right)$$

$$= \left\langle \sum_{i=1}^m PU_i + \phi - i^* \left( \rho^2 e^{\sum_{i=1}^m PU_i + \phi} \right), \partial_{(\xi_h)_j} PU_h + \sum_{i=1}^m P\Psi_i^0 \partial_{(\xi_h)_j} \tau_i(\xi) + \partial_{(\xi_h)_j} \phi \right\rangle$$

$$= -\frac{1}{2} \frac{\partial \mathcal{F}}{\partial (\xi_h)_j} \left( \xi_1, \dots, \xi_m \right) + \left\langle \sum_{s=1}^m \sum_{t=1}^{J_s} c_{st}^{\rho} P\Psi_s^t, \sum_{i=1}^m P\Psi_i^0 \partial_{(\xi_h)_j} \tau_i(\xi) + \partial_{(\xi_h)_j} \phi \right\rangle + o(1)$$

$$= -\frac{1}{2} \frac{\partial \mathcal{F}}{\partial (\xi_h)_j} \left( \xi_1, \dots, \xi_m \right) + \sum_{s=1}^m \sum_{t=1}^{J_i} c_{st}^{\rho} \left\langle P\Psi_s^t, \sum_{i=1}^m P\Psi_i^0 \partial_{(\xi_h)_j} \tau_i(\xi) + \partial_{(\xi_h)_j} \phi \right\rangle + o(1),$$

for any  $h=1,2,\cdots,m$  and  $j=1,\cdots,J_h$ . Utilizing Lemma B.4, we have

$$|\langle P\Psi_s^t, P\Psi_i^0 \rangle| \le ||P\Psi_s^t|| ||P\Psi_i^0|| = O\left(\frac{1}{\rho}\right).$$

Considering the fact that  $\langle P\Psi_s^t, \phi \rangle = 0$  and  $|\partial_{(\xi_h)_j} P\Psi_s^t| \leq |\partial_{(\xi_h)_j} \Psi_s^t| = \mathcal{O}(\frac{1}{\rho^2})$  through direct calculation, we obtain

$$\begin{split} \langle P\Psi_{s}^{t}, \partial_{(\xi_{h})_{j}} \phi \rangle &= \partial_{(\xi_{h})_{j}} \langle P\Psi_{s}^{t}, \phi \rangle - \langle \partial_{(\xi_{h})_{j}} P\Psi_{s}^{t}, \phi \rangle \\ &\leq O\left(\left\|\phi\right\| \left\| \partial_{(\xi_{h})_{j}} P\Psi_{s}^{t} \right\|\right) = O\left(\frac{\left\|\phi\right\|}{\rho^{2}}\right) \\ &= o\left(\frac{1}{\rho^{2}}\right). \end{split}$$

Consequently, we have

(20) 
$$\left\langle \sum_{s=1}^{m} \sum_{t=1}^{J_i} c_{st}^{\rho} P \Psi_s^t, \sum_{i=1}^{m} \partial_{(\xi_h)_j} \tau_i(\xi) P \Psi_i^0 + \partial_{(\xi_h)_j} \phi \right\rangle = o\left(\frac{1}{\rho^2} \sum_{s=1}^{m} \sum_{t=1}^{J_i} |c_{st}^{\rho}|\right).$$

It follows that

(21) 
$$\partial_{(\xi_h)_j} \tilde{E}_{\rho}(\xi) = -\frac{1}{2} \frac{\partial \mathcal{F}_{k,m}}{\partial (\xi_h)_j} (\xi_1, \cdots, \xi_m) + o\left(\frac{1}{\rho^2} \sum_{s=1}^m \sum_{t=1}^{J_i} |c_{st}^{\rho}|\right).$$

Considering (19), (21) implies that for any  $h=1,\cdots,m$  and  $j=1,\cdots,J_h$ 

$$\partial_{(\xi_h)_j} E_\rho \left( \sum_{i=1}^m PU_i + \phi \right) = -\frac{1}{2} \frac{\partial \mathcal{F}}{\partial (\xi_h)_j} (\xi_1, \cdots, \xi_m) + o(1),$$

as 
$$\rho \to 0$$
.

On the other hand,  $\sum_{i=1}^{m} PU_i + \phi_{\xi}^{\rho}$  is a critical point of  $E_{\rho}(u)$  in  $\overline{\mathbb{H}}^1$ , which is equivalent to  $\xi$  being a critical point of  $\tilde{E}_{\rho}(\xi)$  in  $\mathscr{D}$ .

**Proposition 4.2.** The function  $\sum_{i=1}^{m} PU_{\tau_i(\xi),\xi_i} + \phi_{\xi}^{\rho}$  is a solution of (10) if and only if  $\xi$  is a critical point of the reduced map

$$\xi \to \tilde{E}_{\rho}(\xi) = E_{\rho} \left( \sum_{i=1}^{m} PU_{\tau_{i}(\xi),\xi_{i}} + \phi_{\xi}^{\rho} \right).$$

*Proof.* Denote  $\phi := \phi_{\xi}^{\rho}$  to simplify the notations. Assume that  $\xi$  is a critical point of the reduced map  $\tilde{E}_{\rho}(\xi)$ . Then  $\xi$  satisfies

(22) 
$$\partial_{(\xi_i)_j} \tilde{E}_{\rho}(\xi) = 0,$$

for any  $i = 1, \dots, m$  and  $j = 1, \dots, J_i$ .

By (15) of Proposition 3.1,  $\sum_{h=1}^{m} PU_h + \phi - i^* \left( \rho^2 V e^{\sum_{h=1}^{m} PU_h + \phi} \right) = \sum_{s=1}^{m} \sum_{t=1}^{J_i} c_{st}^{\rho} P\Psi_s^t$ , where  $c_{st}^{\rho}$  are coefficients. Then

(23) 
$$\left\langle \sum_{s=1}^{m} \sum_{t=1}^{J_i} c_{st}^{\rho} P \Psi_s^t, \partial_{(\xi_i)_j} P U_i + \sum_{h=1}^{m} P \Psi_i^0 \partial_{(\xi_i)_j} \tau_h(\xi) + \partial_{(\xi_i)_j} \phi \right\rangle = 0.$$

And by (20) in Proposition 4.1 and (23),

$$\sum_{s=1}^{m} \sum_{t=1}^{J_i} c_{st}^{\rho} \left\langle P \Psi_s^t, \partial_{(\xi_i)_j} P U_i \right\rangle = o\left(\frac{1}{\rho^2} \sum_{s=1}^{m} \sum_{t=1}^{J_i} |c_{st}^{\rho}|\right).$$

By Remark B.2, we conclude that  $c_{ij}^{\rho} = 0$  for any  $i = 1, \dots, m$  and  $j = 1, \dots, J_i$ . Thus

(24) 
$$\sum_{h=1}^{m} PU_h + \phi - i^* \left( \rho^2 e^{\sum_{h=1}^{m} PU_h + \phi} \right) = 0.$$

If  $\sum_{h=1}^{m} PU_h + \phi_{\xi}^{\rho}$  is a weak solution to (10) in  $\overline{H}^1$ , then (24) is verified. Furthermore, the validity of (22) is apparent, leading to the conclusion that  $\xi$  is a critical point of the reduced function  $\tilde{E}_{\rho}(\xi)$ .

# 5 Proof of the Main Result

Now, we are ready to prove the main result.

**Proof of Theorem 1.1** Let K be a stable critical point set of  $\mathcal{F}_{k,m}$ , as stated in Proposition 4.1. For any  $n \in \mathbb{N}_+$ , there exists a sequence  $\rho_n \to 0$  and points  $\xi_{\rho_n} = (\xi_{1,\rho_n}, \cdots, \xi_{m,\rho_n}) \in \mathscr{D}$  such that  $dist(\xi_{\rho_n}, K) < \frac{1}{n}$  and  $\xi_{\rho_n}$  is a critical point of  $\tilde{E}_{\rho_n} : \mathscr{D} \to \mathbb{R}$ . Without loss of generality, assume that  $\xi_{\rho_n} = (\xi_{1,\rho_n}, \cdots, \xi_{m,\rho_n}) \to \xi = (\xi_1, \cdots, \xi_m) \in K$ , as  $n \to +\infty$ . Then define  $u_n = \sum_{i=1}^m PU_{\tau_i(\xi_{\rho_n}),\xi_{i,\rho_n}} + \phi_{\xi_{\rho}}^{\rho_n}$ . According to Proposition 4.2,  $u_{\rho_n}$  solves (10) along  $\rho_n \to 0$ , which means that  $u_n$  solves problem (4) in a weak sense for some  $\lambda_n = \rho_n^2 \int_{\Sigma} V e^{u_n} dv_g$ . Applying Lemma B.2, Lemma C.2 and Lemma E.2,  $\lambda_n = \lambda_{k,m} + o(1)$ , as  $n \to +\infty$ . Claim. For any  $\Psi \in C(\Sigma)$ ,

$$\rho^2 \int_{\Sigma} V e^{u_n} \Psi dv_g \to \sum_{i=1}^m \varrho(\xi_i) \Psi(\xi_i) \quad \text{as } n \to +\infty.$$

In fact, by the inequality  $|e^s - 1| \leq e^{|s|} |s|$  for any  $s \in \mathbb{R}$  and Lemma C.1, we have

$$\rho^{2} \int_{\Sigma} V e^{u_{n}} \Psi dv_{g} = \rho^{2} \int_{\Sigma} V e^{\sum_{i=1}^{m} PU_{i}} \Psi dv_{g} + o(1)$$

$$= \sum_{i=1}^{m} \int_{\Sigma} \rho^{2} \chi_{\xi_{i}} e^{U_{i}} \Psi dv_{g} + o(1)$$

$$= \sum_{i=1}^{m} \varrho(\xi_{i}) \Psi(\xi_{i}) + o(1)(n \to 0).$$

Therefore,  $u_n$  is a sequence of blow-up solutions of (4) along  $\lambda_n \to \lambda_{k,m}$ . The proof is concluded.

#### A

**Lemma A.1.** For any given  $\alpha \in (0,1)$ ,  $\beta \geq 0$ , let  $(\Sigma,g)$  be a compact Riemann surface with boundary in  $C^{2,\alpha}$ -class and let  $f \in C^{\alpha}(\Sigma)$ ,  $h \in C^{1,\alpha}(\Sigma)$  such that:

$$\int_{\Sigma} f dv_g = \int_{\partial \Sigma} h ds_g.$$

Then, there exists a unique solution to the problem.

$$\begin{cases}
-\Delta_g u + \beta u = f & \text{in } \Sigma \\
\partial_{\nu_g} u = h & \text{on } \partial \Sigma
\end{cases}$$

in the space  $\{u \in C^{2,\alpha}(\Sigma): \int_{\Sigma} u dv_g = 0\}$ . Moreover, it has the following Schauder estimate:

$$||u||_{C^{2,\alpha}(\Sigma)} \le C \left( ||f||_{C^{\alpha}(\Sigma)} + ||h||_{C^{1,\alpha}(\Sigma)} \right),$$

where C is a constant.

*Proof.* Combining the isothermal coordinates in [37, Lemma 4] with the Schauder estimates (see [19, Theorem 6.31], for instance), Lemma A.1 can be concluded.

**Lemma A.2.** Let  $(\Sigma, g)$  be a compact Riemann surface with smooth boundary  $\partial \Sigma$ . For any  $\beta \geq 0$ , if  $f \in L^2(\Sigma, g)$  satisfies  $\int_{\Sigma} f dv_g = 0$ , then there exists a unique weak solution of

(25) 
$$\begin{cases}
-\Delta_g u + \beta u = f & in \quad \Sigma \\
\partial_{\nu_g} u = 0 & on \quad \partial \Sigma \\
\int_{\Sigma} u dv_g = 0
\end{cases}$$

i.e., there exists  $u \in \overline{H}^1$  satisfying that

$$\int_{\Sigma} \langle \nabla u, \nabla \varphi \rangle_g dv_g + \beta \int_{\Sigma} u \varphi dv_g = \int_{\Sigma} f \varphi dv_g, \quad \forall \varphi \in \overline{\mathbf{H}}^1.$$

Moreover, for any p > 1 if  $f \in L^p(\Sigma)$ , there exists  $u \in W^{2,p}$  solving (25) with the following  $W^{2,p}$ -estimate:  $||u||_{W^{2,p}(\Sigma)} \le C||f||_{L^p(\Sigma)}$ .

For the Poisson equation with homogeneous Neumann boundary condition, the lemma above was proven in [37, Lemma 5]. And we can deduce (25) it by the same approach.

*Proof.* For the uniqueness, we assume that  $u_1, u_2$  are two weak solutions of (25) in  $\overline{H}^1$ . It follows that

$$\int_{\Sigma} \langle \nabla (u_1 - u_2), \nabla \varphi \rangle_g dv_g + \beta \int_{\Sigma} (u_1 - u_2) \varphi dv_g = 0,$$

for any  $\varphi \in H^1(\Sigma)$ . Then,  $u_1 = u_2$  up to the addition of a constant. Observing that  $\int_{\Sigma} u_1 dv_g = \int_{\Sigma} u_2 dv_g = 0$ , we deduce that  $u_1 \equiv u_2$ .

We will prove the existence by variational methods. Consider the energy functional

$$J(u) = \frac{1}{2} \int_{\Sigma} (|\nabla u|_g^2 + \beta u^2) dv_g - \int_{\Sigma} f u dv_g.$$

Applying the Hölder inequality and the Poincaré inequality,

$$\left| \int_{\Sigma} f u dv_g \right| \le \|f\|_{L^2(\Sigma)} \|u\|_{L^2(\Sigma)} \le C \|f\|_{L^2(\Sigma)} \|\nabla u\|_{L^2(\Sigma)},$$

which yields that J has a lower bound in  $\overline{H}^1$ . Let  $u_n$  be a sequence in  $\overline{H}^1$  such that J attains the minimum value, i.e.,

$$\lim_{n \to +\infty} J(u_n) = \inf_{u \in \overline{\mathbb{H}}^1} J(u).$$

For any  $n \in \mathbb{N}_+$ ,  $J(u_n) \ge \frac{1}{2} ||u_n||^2 - C||f||_{L^2(\Sigma)} ||u_n||$ . Given that  $\inf_{u \in \overline{\mathbb{H}}^1} J(u) \le J(0) = 0$ ,  $u_n$  is uniformly bounded in  $\overline{\mathbb{H}}^1$ . Up to a subsequence, we assume that  $u_n$  converges to some  $u_0 \in \overline{\mathbb{H}}^1$  weakly. By the Rellich–Kondrachov theorem,  $u_n \to u_0$  strongly in  $L^q(\Sigma)$  for any q > 1 and almost everywhere. Fatou's lemma implies that

$$J(u_0) \le \liminf_{n \to +\infty} J(u_n) = \inf_{u \in \overline{H}^1} J(u).$$

Thus,  $u_0$  is a minimizer of J(u) on  $\overline{H}^1$ .

Next, we consider the  $W^{2,p}$ -estimates of the solutions. Employing the isothermal coordinates introduced in Section 2 it is sufficient to prove the  $L^p$ -regularity locally in an open disk

or half-disk in  $\mathbb{R}^2$ . Specifically, in the case of a half-disk, we can extend the problem evenly to a full open disk, considering that  $\partial_{\nu_g} u = 0$  on the boundary. This extension allows for the application of the standard local  $L^p$ -theory, thereby we can establish the  $L^p$ -regularity for the Neumann boundary problem (25) on a compact Riemann surface  $\Sigma$ .

#### $\mathbf{B}$

This section provides detailed proofs of crucial estimates for  $PU_{\tau,\xi}$  for  $\tau \in (0,\infty)$  and  $\xi \in \Sigma$ . For any  $\xi$  located in a compact subset of  $\mathring{\Sigma}$  or  $\partial \Sigma$ , we set  $r_{\xi}$  to be  $4r_0$ , where  $r_0$  is a positive constant.

The following lemma is the asymptotic expansion of  $PU_{\tau,\xi}$  as  $\rho \to 0$ .

**Lemma B.1.** The function  $PU_{\delta,\xi}$  satisfies

$$PU_{\tau,\xi} = \chi_{\xi} \left( U_{\tau,\xi} - \log(8\tau^2) \right) + \varrho(\xi) H^g(x,\xi) + \mathcal{O}(\rho^{1+\alpha_0})(\rho \to 0),$$

for any  $\alpha_0 \in (0,1)$  and the convergent is locally uniform for  $\xi$  in  $\overset{\circ}{\Sigma}$  and  $\partial \Sigma$  and also locally uniform for  $\tau$  in  $(0,+\infty)$ . In particular,

$$PU_{\tau,\xi} = \varrho(\xi)G^g(x,\xi) + \mathcal{O}(\rho^{1+\alpha_0})(\rho \to 0),$$

locally uniformly in  $\Sigma \setminus \{\xi\}$ .

Proof. Let  $\eta_{\tau,\xi}(x) = PU_{\tau,\xi} - \chi_{\xi}(U_{\tau,\xi} - \log 8\tau^2) - \varrho(\xi)H^g(x,\xi)$ . If  $\xi \in \mathring{\Sigma}$ ,

$$\partial_{\nu_g} \eta_{\tau,\xi} = 2\partial_{\nu_g} \chi_{\xi} \log \left( 1 + \frac{\tau^2 \rho^2}{|y_{\xi}(x)|^2} \right) - 2\chi_{\xi} \partial_{\nu_g} \log \left( 1 + \frac{\tau^2 \rho^2}{|y_{\xi}(x)|^2} \right) \equiv 0$$

on  $\partial \Sigma$ . We observe that for any  $x \in \partial \Sigma \cap U(\xi)$ 

$$\partial_{\nu_g} |y_{\xi}(x)|^2 = -e^{-\frac{1}{2}\hat{\varphi}_{\xi}(y)} \left. \frac{\partial}{\partial y_2} |y|^2 \right|_{y=y_{\xi}(x)} = 0.$$

If  $\xi \in \partial \Sigma$ , for any  $x \in \partial \Sigma$ , as  $\rho \to 0$ .

$$\partial_{\nu_g} \eta_{\tau,\xi}(x) = 2(\partial_{\nu_g} \chi_{\xi}) \frac{\tau^2 \rho^2}{|y_{\xi}(x)|^2} - 2\chi_{\xi} \partial_{\nu_g} \log \left( 1 + \frac{\tau^2 \rho^2}{|y_{\xi}(x)|^2} \right) + \mathcal{O}(\rho^4) = \mathcal{O}(\rho^2).$$

Then for any  $\xi \in \Sigma$  we have  $\partial_{\nu_g} \eta_{\tau,\xi} = \mathcal{O}(\rho^2)$ . For any  $A \subset \mathbb{R}^2$ , denote  $aA := \{ay : y \in A\}$ .

$$\begin{split} &\int_{\Sigma} \eta_{\tau,\xi} dv_g = -\int_{\Sigma} \chi_{\xi} (U_{\tau,\xi} - \log(8\tau^2)) + \varrho(x) \Gamma_{\xi}(x) dv_g(x) \\ &= -\int_{\Sigma} \chi_{\xi} \log \frac{|y_{\xi}(x)|^4}{(\tau^2 \rho^2 + |y_{\xi}(x)|^2)^2} dv_g(x) \\ &= 2\int_{B_{r_0}^{\xi}} \log \frac{\tau^2 \rho^2 + |y|^2}{|y|^2} e^{-\hat{\varphi}_{\xi}(y)} dy + 2\int_{B_{2r_0}^{\xi} \backslash B_{r_0}(0)} \chi(|y|) \left(\frac{\tau^2 \rho^2}{|y|^2} + \mathcal{O}(\rho^4)\right) e^{\hat{\varphi}_{\xi}(y)} dy \\ &= 2\tau^2 \rho^2 \int_{\frac{1}{\tau\rho} (B_{r_0}^{\xi} \cap B_{r_0}(0))} \log \left(1 + \frac{1}{|y|^2}\right) e^{-\hat{\varphi}(\tau\rho y)} dy + \mathcal{O}(\rho^2) \\ &= 2\tau^2 \rho^2 (1 + \mathcal{O}(\rho)) \int_{B_{r_0/(\tau\rho)}(0)} \log \left(1 + \frac{1}{|y|^2}\right) dy + \mathcal{O}(\rho^2) \\ &= \mathcal{O}(\rho^2 |\log \rho|), \end{split}$$

where we applied

$$\int_{|y| < \frac{r_0}{\tau \rho}} \log \left( 1 + \frac{1}{|y|^2} \right) dy = 2\pi \int_0^{r_0/(\tau \rho)} \log \left( 1 + \frac{1}{r^2} \right) r dr$$

$$= \pi \int_0^{r_0^2/(\tau \rho)^2} \log \left( 1 + \frac{1}{t} \right) dt$$

$$= \pi \frac{r_0^2}{\tau^2 \rho^2} \log \left( 1 + \frac{\tau^2 \rho^2}{r_0^2} \right) - \pi \int_0^{r_0^2/(\tau \rho)^2} (1 - \frac{1}{1+t}) dt$$

$$= \pi \frac{r_0^2}{\tau^2 \rho^2} \left( 1 + \frac{\tau^2 \rho^2}{r_0^2} + \mathcal{O}(\rho^4) \right) - \pi \frac{r_0^2}{\tau^2 \rho^2} + \pi \log \left( 1 + \frac{r_0^2}{\tau^2 \rho^2} \right)$$

$$= \mathcal{O}(|\log \rho|).$$

For any  $x \in U_{2r_0}(\xi)$ ,  $-\Delta_g U_{\tau,\xi} = e^{-\hat{\varphi}_{\xi}(y)} \Delta u_{\tau,0}|_{y=y_{\xi}(x)} = e^{-\varphi_{\xi}} e^{U_{\tau,\xi}}$ . It follows that

$$(-\Delta_{g} + \beta)\eta_{\tau,\xi} = (-\Delta_{g} + \beta) \left( PU_{\tau,\xi} - \chi_{\xi}(U_{\tau,\xi} - \log 8\tau^{2}) - \varrho(\xi)H_{\xi}^{g} \right)$$

$$= (\Delta_{g}\chi_{\xi}) \log \frac{|y_{\xi}|^{4}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})^{2}} + 2\langle \nabla\chi_{\xi}, \nabla \log \frac{|y_{\xi}|^{4}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})^{2}} \rangle_{g}$$

$$+ \frac{1}{|\Sigma|_{g}} \left( \varrho(\xi) - \int_{\Sigma} \rho^{2}\chi_{\xi}e^{-\varphi_{\xi}}e^{U_{\tau,\xi}}dv_{g} \right) + 2\beta \log \left( 1 + \frac{\tau^{2}\rho^{2}}{|y_{\xi}|^{2}} \right).$$

We observe that  $\Delta_g \chi_{\xi} \equiv 0$  and  $\nabla \chi_{\xi} \equiv 0$  in  $U_{2r_0}(\xi) \setminus U_{r_0}(\xi)$ . For any  $x \in A_{r_0}(\xi)$ , we have

$$U_{\tau,\xi} - \log(8\tau^2) + 4\log|y_{\xi}(x)| = -2\log\left(1 + \frac{\tau^2 \rho^2}{|y_{\xi}(x)|^2}\right) = -2\tau^2 \rho^2 |y_{\xi}(x)|^{-2} + \mathcal{O}(\rho^4)$$

and

$$\nabla \left( U_{\tau,\xi} - \log(8\tau^2) + 4\log|y_{\xi}(x)| \right) = -2\tau^2 \rho^2 \nabla |y_{\xi}(x)|^{-2} + \mathcal{O}(\rho^4).$$

$$\int_{\Sigma} \rho^{2} \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} dv_{g} = \int_{B_{2r_{0}}^{\xi}} \rho^{2} \chi(|y|) \frac{8\tau^{2}}{(\tau^{2}\rho^{2} + |y|^{2})^{2}} dy$$

$$= \int_{B_{r_{0}}^{\xi}} \rho^{2} \chi(|y|) \frac{8\tau^{2}}{(\tau^{2}\rho^{2} + |y|^{2})^{2}} dy + \mathcal{O}(\rho^{2})$$

$$= \varrho(\xi) + \mathcal{O}(\rho^{2}),$$

where we applied the fact that  $\int_{|y|< r} \frac{\tau^2 \rho^2}{(\tau^2 \rho^2 + |y|^2)^2} dy = \pi - \frac{\pi \tau^2 \rho^2}{r^2} + \frac{\pi \tau^4 \rho^4}{(r^2 + \tau^2 \rho^2)r^2}$  for any  $r \ge 0$ . Hence, for any p > 1

$$\|(-\Delta_g + \beta)\eta_{\tau,\xi}\|_p = \mathcal{O}(\rho^2 + \beta\rho^{\frac{2}{p}}).$$

By the regularity theory in Lemma A.1 and Lemma A.2 and, we have

$$\|\eta_{\tau,\xi} - \overline{\eta_{\tau,\xi}}\|_{\infty} \le C\left(\|\partial_{\nu_g}\eta_{\tau,\xi}\|_{L^{\infty}(\partial\Sigma)} + \|(-\Delta_g + \beta)\eta_{\tau,\xi}\|_{L^p(\Sigma)}\right) \le C(\rho^2 + \beta\rho^{\frac{2}{p}}),$$

for p > 1.

We take  $p \in (1,2)$  such that  $\alpha_0 = \frac{2}{p} - 1 > 0$ , then as  $\rho \to 0$ 

$$\eta_{\tau,\xi} = \mathcal{O}(\rho^{1+\alpha_0}),$$

uniformly in  $C(\Sigma)$ .

**Lemma B.2.** If  $p \geq 1$  then  $|\rho^2 \chi_{\xi} e^{U_{\tau,\xi}}|_{L^p(\Sigma)} = \mathcal{O}(\rho^{\frac{2(1-p)}{p}})$  which is uniform for  $\xi$  in  $\Sigma$  and locally uniform for  $\tau$  in  $(0, +\infty)$ .

Proof.

$$\int_{\Sigma} (\rho^{2} \chi_{\xi} e^{U_{\tau,\xi}})^{p} dv_{g} = \int_{B_{2r_{0}}^{\xi}} e^{\hat{\varphi}_{\xi}(y)} \frac{(8\tau^{2}\rho^{2})^{p}}{(\tau^{2}\rho^{2} + |y|^{2})^{p}} dy$$

$$= \int_{B_{2r_{0}}^{\xi}} \frac{(8\tau^{2}\rho^{2})^{p}}{(\tau^{2}\rho^{2} + |y|^{2})^{p}} dy + \int_{B_{2r_{0}}^{\xi}} (e^{\hat{\varphi}_{\xi}(y)} - 1) \frac{(8\tau^{2}\rho^{2})^{p}}{(\tau^{2}\rho^{2} + |y|^{2})^{p}} dy$$

$$= (\tau^{2}\rho^{2})^{1-p} \int_{\frac{1}{\tau\rho} B_{2r_{0}}^{\xi}} (1 + \mathcal{O}(\tau\rho|y|)) \frac{8}{(1+|y|^{2})^{2}} dy$$

$$= \mathcal{O}(\rho^{2(1-p)}).$$

Thus  $|\rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}}|_{L^p(\Sigma)} = \mathcal{O}(\rho^{\frac{2(1-p)}{p}})$  uniformly in  $\xi \in \Sigma$  and  $\tau$  is bounded away from zero.  $\square$ 

Next, we discuss the asymptotic expansions of  $P\Psi_i^j$  as  $\rho \to 0$  analogue to  $PU_{\tau,\xi}$ . For j=1,2 if  $\xi \in \mathring{\Sigma}$  and j=1 if  $\xi \in \partial \Sigma$ , we define  $H^j(x,\xi)$  to be the unique solution of the

following problem

$$\begin{cases} -\Delta_g H^j(x,\xi) = \frac{4}{\varrho(\xi)} \Delta_g \chi_{\xi} \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} + \frac{8}{\varrho(\xi)} \left\langle \nabla \chi_{\xi}, \nabla \left( \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \right) \right\rangle_g & x \in \mathring{\Sigma} \\ \partial_{\nu_g} H^j(x,\xi) = -\frac{4}{\varrho(\xi)} \partial_{\nu_g} \left( \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \right) \chi_{\xi} - \frac{4}{\varrho(\xi)} \frac{y_{\xi}(x)_j}{|y_{\xi}|^2} \partial_{\nu_g} \chi_{\xi} & x \in \partial \Sigma \\ \int_{\Sigma} H^j(x,\xi) dv_g = -\frac{4}{\varrho(\xi)} \int_{\Sigma} \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \chi_{\xi}(x) dv_g \end{cases}$$

**Lemma B.3.** For any  $\alpha_0 \in (0,1)$ ,

$$P\Psi_{\tau,\xi}^{0}(x) = \chi_{\xi}\left(\Psi_{\tau,\xi}^{0}(x) - \frac{2}{\tau}\right) + \mathcal{O}(\rho^{1+\alpha_{0}}) = -4\chi_{\xi}(x)\frac{\tau\rho^{2}}{\tau^{2}\rho^{2} + |y_{\xi}(x)|^{2}} + \mathcal{O}(\rho^{1+\alpha_{0}})$$

in  $C(\Sigma)$  as  $\rho \to 0$ . And  $P\Psi^0_{\tau,\xi}(x) = \mathcal{O}(\rho^{1+\alpha_0})(\rho \to 0)$ , in  $C_{loc}(\Sigma \setminus \{\xi\})$  uniformly for  $\xi$  in any compact subset of  $\mathring{\Sigma}$  or  $\xi \in \partial \Sigma$  and  $\tau$  is bounded away from zero.

For  $\xi \in \mathring{\Sigma}$  with j = 1, 2, or for  $\xi \in \partial \Sigma$  with j = 1,

$$P\Psi_{\tau,\xi}^{j}(x) = \chi_{\xi}(x)\Psi_{\tau,\xi}^{j}(x) + \varrho(\xi)H^{j}(x,\xi) + \mathcal{O}(\rho^{\alpha_{0}})$$

in  $C(\Sigma)$  as  $\rho \to 0$ , where  $H^j(x,\xi)$  is the unique solution of the following problem

$$\begin{cases}
(-\Delta_g + \beta)H^j(x,\xi) &= -\beta \frac{4}{\varrho(\xi)} \chi_{\xi} \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} + \frac{4}{\varrho(\xi)} (\Delta_g \chi_{\xi}) \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \\
+ \frac{8}{\varrho(\xi)} \left\langle \nabla \chi_{\xi}, \nabla \left( \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \right) \right\rangle_g, \quad x \in \mathring{\Sigma} \\
\partial_{\nu_g} H^j(x,\xi) &= -\frac{4}{\varrho(\xi)} \partial_{\nu_g} \left( \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \right) \chi_{\xi} - \frac{4}{\varrho(\xi)} \frac{y_{\xi}(x)_j}{|y_{\xi}|^2} \partial_{\nu_g} \chi_{\xi}, \quad x \in \partial \Sigma \\
\int_{\Sigma} H^j(x,\xi) dv_g &= -\frac{4}{\varrho(\xi)} \int_{\Sigma} \frac{y_{\xi}(x)_j}{|y_{\xi}(x)|^2} \chi_{\xi}(x) dv_g
\end{cases}$$

In addition, the convergences above are uniform for  $\xi$  in any compact subset of  $\overset{\circ}{\Sigma}$  or  $\xi \in \partial \Sigma$  and  $\tau$  bounded away from zero.

*Proof.* We recall that

$$\Psi^0_{\tau,\xi}(x) = \frac{2}{\tau} \frac{|y_{\xi}(x)|^2 - \tau^2 \rho^2}{|y_{\xi}(x)|^2 + \tau^2 \rho^2} = \frac{2}{\tau} \left( 1 - \frac{2\tau^2 \rho^2}{|y_{\xi}(x)|^2 + \tau^2 \rho^2} \right), x \in U_{2r_0}(\xi).$$

Let  $\eta_{\tau,\xi} = P\Psi^0_{\tau,\xi} - \chi_{\xi} \left( \Psi^0_{\tau,\xi} - \frac{2}{\tau} \right)$ . For  $x \in \partial \Sigma$ ,

$$\partial_{\nu_g} \left( \chi_{\xi} \left( \Psi^0_{\tau,\xi}(x) - \frac{2}{\tau} \right) \right) = -\partial_{\nu_g} \chi_{\xi} \frac{4\tau \rho^2}{|y_{\xi}(x)|^2 + \tau^2 \rho^2} + \chi_{\xi} \frac{8\tau \rho^2 |y_{\xi}(x)|^2}{(|y_{\xi}(x)|^2 + \tau^2 \rho^2)^2} \partial_{\nu_g} \log |y_{\xi}(x)|.$$

If  $\xi \in \mathring{\Sigma}$ ,  $\partial_{\nu_g} \eta_{\tau,\xi} \equiv 0$  in  $\partial \Sigma$ ; if  $\xi \in \partial \Sigma$ ,  $\partial_{\nu_g} \eta_{\tau,\xi} = \mathcal{O}(\rho^2)$  on  $\partial \Sigma$ . By direct calculation, we have

$$\int_{\Sigma} \chi_{\xi} \left( \Psi_{\tau,\xi}^{0} - \frac{2}{\tau} \right) dv_{g} = 2 \int_{\Sigma} \chi_{\xi} \frac{\tau \rho^{2}}{|y_{\xi}(x)|^{2} + \tau^{2} \rho^{2}} dv_{g}(x) = 2\tau \rho^{2} \int_{B_{2r_{0}}^{\xi}} \frac{1}{|y|^{2} + \tau^{2} \rho^{2}} e^{\hat{\varphi}_{\xi}(y)} dy 
= 2\tau \rho^{2} \int_{B_{2r_{0}}^{\xi}} \frac{1}{|y|^{2} + \tau^{2} \rho^{2}} dy + 2\tau \rho^{2} \int_{B_{2r_{0}}^{\xi}} \frac{1}{|y|^{2} + \tau^{2} \rho^{2}} (e^{\hat{\varphi}_{\xi}(y)} - 1) dy 
= \mathcal{O}(\rho^{2} \log \rho)$$

and

$$\begin{aligned} & (-\Delta_g + \beta) \eta_{\tau,\xi} = (-\Delta_g + \beta) \left( P \Psi^0_{\tau,\xi} - \chi_{\xi} \left( \Psi^0_{\tau,\xi} - \frac{2}{\tau} \right) \right) \\ &= (\Delta_g \chi_{\xi}) \left( \Psi^0_{\tau,\xi} - \frac{2}{\tau} \right) + 2 \langle \nabla \chi_{\xi}, \nabla \Psi^0_{\tau,\xi} \rangle_g - \rho^2 \chi_{\xi} e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^0_{\tau,\xi} + \beta \chi_{\xi} \frac{4\tau \rho^2}{|y_{\xi}|^2 + \tau^2 \rho^2} \\ &= \beta \chi_{\xi} \frac{4\tau \rho^2}{|y_{\xi}|^2 + \tau^2 \rho^2} + \mathcal{O}(\rho^2), \end{aligned}$$

where we applied the fact for any fixed r > 0,  $\int_{|y| < r} \frac{\tau^2 \rho^2 - |y|^2}{(\tau^2 \rho^2 + |y|^2)^3} = \mathcal{O}(\rho^2)$  as  $\rho \to 0$ . Via the regularity theory in Lemma A.1 and Lemma A.2, for any p > 1, there exists a constant C > 0 such that

(27) 
$$\|\eta_{\tau,\xi} - \overline{\eta_{\tau,\xi}}\|_{\infty} \leq C \left( \|\partial_{\nu_g} \eta_{\tau,\xi}\|_{L^{\infty}(\partial \Sigma)} + \|(-\Delta_g + \beta)\eta_{\tau,\xi}\|_{L^{p}(\Sigma)} \right)$$
$$\leq C(\rho^2 + \beta \rho^{\frac{2}{p}} |\log \rho|^{\frac{1}{p}}).$$

We choose  $p \in (1,2)$  such that  $\alpha_0 < \frac{2}{p} - 1$ , then  $\eta_{\tau,\xi} = \mathcal{O}(\rho^{1+\alpha_0})$ , uniformly in  $C(\Sigma)$ . We recall that

$$\Psi_{\tau,\xi}^{j}(x) = \frac{4y_{\xi}(x)_{j}}{\tau^{2}\rho^{2} + |y_{\xi}(x)|^{2}}, \text{ for any } x \in U_{2r_{0}}(\xi).$$

If  $\xi \in \mathring{\Sigma}$ ,  $\partial_{\nu_q} H^j(x,\xi) = 0$  for any  $x \in \partial \Sigma$ . If  $\xi \in \partial \Sigma$ , for any  $x \in \partial \Sigma$  by direct calculation,

$$\chi_{\xi}(x)\partial_{\nu_g}\left(\frac{y_{\xi}(x)_1}{|y_{\xi}(x)|^2}\right) = 0.$$

Denote  $\partial_{\nu_g} H^j(\xi,\xi) := 0$ , then  $\partial_{\nu_g} H^j(\cdot,\xi) \in C^{\infty}(\partial \Sigma)$ . By Lemma A.1, there is a unique solution to the problem (26) in  $C^{1,\alpha}(\partial \Sigma)$  for any  $\alpha \in (0,1)$ . Let

$$\zeta_{\tau,\xi}(x) = P\Psi_{\tau,\xi}^{j}(x) - \chi_{\xi}(x)\Psi_{\tau,\xi}^{j}(x) - \varrho(\xi)H^{j}(x,\xi).$$

Since  $\int_B \frac{\rho^3 y_j}{(\rho^2 + |y|^2)^3} dy = 0$  for j = 1, 2 and  $B = B_r$  or j = 1 and  $B = B_r \cap \{y_2 \ge 0\}$ , we have the following estimates:

$$\overline{\rho^{2}\chi_{\xi}e^{-\varphi_{\xi}}e^{U_{\tau,\xi}}\Psi_{\tau,\xi}^{j}} = \int_{B_{2r_{0}}^{\xi}} \frac{8\tau^{2}\rho^{2}\chi(|y|)y_{j}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}}dy$$

$$= \int_{B} \frac{8\tau^{2}\rho^{2}y_{j}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}}dy + \mathcal{O}(\rho^{2}) = \mathcal{O}(\rho^{2}),$$

$$(-\Delta_{g} + \beta)\zeta_{\tau,\xi} = -\frac{4\tau^{2}\rho^{2}(y_{\xi})_{j}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})|y_{\xi}|^{2}}\Delta_{g}\chi_{\xi} - 8\tau^{2}\rho^{2}\left\langle\nabla\chi_{\xi}, \nabla\left(\frac{(y_{\xi})_{j}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})|y_{\xi}|^{2}}\right)\right\rangle_{g}$$
$$-\frac{\rho^{2}\chi_{\xi}e^{-\varphi_{\xi}}e^{U_{\tau,\xi}}\Psi_{\tau,\xi}^{j}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})|y_{\xi}|^{2}}$$
$$= 4\beta\chi_{\xi}\frac{\tau^{2}\rho^{2}(y_{\xi})_{j}}{(\tau^{2}\rho^{2} + |y_{\xi}|^{2})|y_{\xi}|^{2}} + \mathcal{O}(\rho^{2})$$

and

$$\int_{\Sigma} \zeta_{\tau,\xi} dv_g = 4 \int_{\Sigma} \chi_{\xi}(x) \frac{\tau^2 \rho^2 y_{\xi}(x)_j}{|y_{\xi}(x)|^2 (\tau^2 \rho^2 + |y_{\xi}(x)|^2)} dv_g(x) = 4 \int_{B_{2r_0}^{\xi}} \chi\left(\frac{|y|}{r_0}\right) e^{\hat{\varphi}_{\xi}(y)} \frac{\tau^2 \rho^2 y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)}.$$

If  $\xi \in \overset{\circ}{\Sigma}$  and j = 1, 2

$$\left| \int_{B_{2r_0}} \chi(|y|) \frac{\tau^2 \rho^2 (e^{\hat{\varphi}_{\xi}(y)} - 1) y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)} dy \right| \le C \int_{|y| < 2r_0} \frac{\tau^2 \rho^2 |D^2 \hat{\varphi}_{\xi}(0) y^2 + o(|y|^3)|}{|y| (\tau^2 \rho^2 + |y|^2)} dy$$

$$= C \int_{|y| < \frac{2r_0}{\tau \rho}} \frac{\tau^3 \rho^3 |y| (1 + \tau \rho o(|y|))}{(1 + |y|^2)} dy \le C 2\pi \int_0^{\frac{2r_0}{\tau \rho}} \frac{\tau^3 \rho^3 r^2}{(1 + r^2)} dr = \mathcal{O}(\rho^2).$$

In this case, we have

$$\begin{split} & \int_{\Sigma} \zeta_{\tau,\xi} dv_g = 4 \int_{B_{2r_0}(0)} \chi\left(\frac{|y|}{r_0}\right) e^{\hat{\varphi}_{\xi}(y)} \frac{\tau^2 \rho^2 y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)} dy \\ = & -4 \int_{B_{2r_0}} \chi\left(\frac{|y|}{r_0}\right) \frac{\tau^2 \rho^2 y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)} dy + \int_{B_{2r_0}} \chi\left(\frac{|y|}{r_0}\right) (e^{\hat{\varphi}_{\xi}(y)} - 1) \frac{\tau^2 \rho^2 y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)} dy = \mathcal{O}(\rho^2). \end{split}$$

For  $\xi \in \partial \Sigma$  and j = 1

$$\begin{aligned} \left| \int_{\Sigma} \zeta_{\tau,\xi} dv_g \right| &= 4 \left| \int_{B_{2r_0}^{\xi}} \chi \left( \frac{|y|}{r_0} \right) e^{\hat{\varphi}_{\xi}(y)} \frac{\tau^2 \rho^2 y_j}{|y|^2 (\tau^2 \rho^2 + |y|^2)} dy \right| \\ &= \mathcal{O}(\rho^2), \end{aligned}$$

then, it follows for  $\xi \in \partial \Sigma$ ,  $\int_{\Sigma} \zeta_{\tau,\xi} dv_g = \mathcal{O}(\rho^2)$ . If  $\xi \in \mathring{\Sigma}$ ,  $\partial_{\nu_g} \zeta_{\tau,\xi}(x) \equiv 0$  for any  $x \in \partial \Sigma$ . If  $x \in \partial \Sigma$ , by calculation, we deduce that

$$\partial_{\nu_{g}} \zeta_{\tau,\xi}(x) = -\partial_{\nu_{g}} \chi_{\xi}(\Psi_{\tau,\xi}^{j} + \varrho(\xi)H^{j}(x,\xi)) - \chi_{\xi} \partial_{\nu_{g}} \left(\Psi_{\tau,\xi}^{j} + \varrho(\xi)H^{j}(x,\xi)\right) \\
= (\partial_{\nu_{g}} \chi_{\xi}) \frac{4\tau^{2} \rho^{2} y_{\xi}(x)_{j}}{(\tau^{2} \rho^{2} + |y_{\xi}(x)|^{2})|y_{\xi}(x)|^{2}} + \chi_{\xi} \partial_{\nu_{g}} \frac{4\tau^{2} \rho^{2} y_{\xi}(x)_{j}}{(\tau^{2} \rho^{2} + |y_{\xi}(x)|^{2})|y_{\xi}(x)|^{2}} \\
= \mathcal{O}(\rho^{2}).$$

Applying the regularity theory in Lemma A.1 and Lemma A.2, for any  $p \in (1,2)$ , we deduce that

$$\|\zeta_{\tau,\xi} - \overline{\zeta_{\tau,\xi}}\|_{\infty} \le C\left(\|\partial_{\nu_g}\zeta_{\tau,\xi}\|_{L^{\infty}(\partial\Sigma)} + \|(-\Delta_g + \beta)\zeta_{\tau,\xi}\|_{L^p(\Sigma)}\right) \le C(\rho^2 + \beta\rho^{\frac{1}{p}}).$$

We take  $p \in (0,1)$  such that  $\alpha_0 = \frac{1}{p}$ . Then as  $\rho \to 0$ , we have

$$\eta_{\tau,\xi} = \mathcal{O}(\rho^{\alpha_0})$$

uniformly in  $C(\Sigma)$ .

Remark B.1.  $\partial_{\tau}PU_{\tau,\xi} = P\Psi^{0}_{\tau,\xi}$  by the uniqueness of the solution to the problem (12). However,  $\partial_{\xi_{j}}PU_{\tau,\xi} \neq P\Psi^{j}_{\tau,\xi}$ . Analog to the proof of Lemma B.3, we can obtain the following expansion, for any  $\alpha_{0} \in (0,1)$ ,

(28) 
$$\partial_{\xi_j} P U_{\tau,\xi} = \partial_{\xi_j} (\chi_{\xi} U_{\tau,\xi}) + \varrho(\xi) \partial_{\xi_j} H_{\xi}^g + \mathcal{O}(\rho^{\alpha_0}) (\rho \to 0),$$

in  $C(\Sigma)$ , which is uniformly convergent for  $\xi$  in any compact subset of  $\mathring{\Sigma}$  or  $\xi \in \partial \Sigma$  and  $\tau$  in any compact subset of  $(0, \infty)$ .

Indeed, we notice that for any  $y \in U_{2r_0}(\xi)$  as  $y \to 0$ 

$$\partial_{\xi_j} |y_{\xi}(x)|^2 \Big|_{x=y_{\varepsilon}^{-1}(y)} = -2y_j + \mathcal{O}(|y|^3).$$

Let  $\zeta_{\tau,\xi}^* = \partial_{\xi_j} PU_{\tau,\xi} - \partial_{\xi_j} (\chi_{\xi} U_{\tau,\xi}) - \varrho(\xi) \partial_{\xi_j} H^g(x,\xi)$ . It is easy to obtain

$$(-\Delta_g + \beta)\zeta_{\tau,\xi}^* = -\beta \partial_{\xi_i} \left( \chi_{\xi} U_{\tau,\xi} + 4\chi_{\xi} \log |y_{\xi}| \right) + \mathcal{O}(\rho^2 |\log \rho|), \quad \text{in } \mathring{\Sigma}$$

$$\int_{\Sigma} \zeta_{\tau,\xi}^* dv_g = \mathcal{O}(\rho^2 |\log \rho|),$$

and  $\partial_{\xi}\zeta_{\tau,\xi}^* = \mathcal{O}(\rho^2)$  on  $\partial\Sigma$ . Applying the regularity theory in Lemma A.1 and Lemma A.2, we have

$$\zeta_{\tau,\xi}^* = \mathcal{O}(\rho^2 |\log \rho| + \beta \rho^{\frac{1}{p}}),$$

for any  $p \in (1,2)$ . We take  $p \in (1,2)$  such that  $\alpha_0 = \frac{1}{p}$ , then we deduce (28).

The following lemma shows asymptotic "orthogonality" properties of  $P\Psi_i^j$ .

**Lemma B.4.** For any  $\alpha_0 \in (0,1)$ , we have as  $\rho \to 0$  for j, i = 0, 1, 2 for  $\xi \in \mathring{\Sigma}$  and i, j = 0, 1 for  $\xi \in \partial \Sigma$ ,

$$\langle P\Psi^{i}_{\tau,\xi}, P\Psi^{j}_{\tau,\xi} \rangle = \begin{cases} \frac{8\varrho(\xi)D_{i}}{\pi\tau^{2}} \delta_{ij} + \mathcal{O}(\rho^{\alpha_{0}}) & \text{when } i \text{ or } j = 0\\ \frac{8\varrho(\xi)D_{i}}{\pi\tau^{2}\rho^{2}} \delta_{ij} + \mathcal{O}(\rho^{\alpha_{0}-1}) & \text{otherwise} \end{cases},$$

and

$$\langle P\Psi^{i}_{\tau^{0},\xi_{0}}, P\Psi^{j}_{\tau^{1},\xi_{1}} \rangle = \begin{cases} \mathcal{O}(\rho^{\alpha_{0}}) & \text{when } i \text{ or } j = 0\\ \mathcal{O}(\rho^{\alpha_{0}-1}) & \text{otherwise} \end{cases},$$

where three different points  $\xi, \xi_0, \xi_1 \in \Sigma$  and uniformly in  $\tau, \tau^0, \tau^1$  are bounded away from zero and the  $\delta_{ij}$  is the Kronecker symbol, and  $D_0 = \int_{\mathbb{R}^2} \frac{1-|y|^2}{(1+|y|^2)^4} dy$ ,  $D_1 = D_2 = \int_{\mathbb{R}^2} \frac{|y|^2}{(1+|y|^2)^4} dy$ .

Proof.

$$\begin{split} \langle P\Psi^i_{\tau,\xi}, P\Psi^j_{\tau,\xi} \rangle &= \int_{\Sigma} \rho^2 \chi_{\xi}(x) e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^i_{\tau,\xi} P\Psi^j_{\tau,\xi} dv_g(x) \\ &= \int_{\Sigma \cap U_{2r_0}(\xi)} + \int_{\Sigma \backslash U_{2r_0}(\xi)} \rho^2 \chi_{\xi}(x) e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^i_{\tau,\xi} P\Psi^j_{\tau,\xi} dv_g(x). \end{split}$$

For i = j = 0, by Lemma B.3,

$$\begin{split} &\int_{\Sigma \cap U_{2r_0}(\xi)} \rho^2 \chi_{\xi}(x) e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^0_{\tau,\xi} P \Psi^0_{\tau,\xi} dv_g(x) \\ &= 16\tau \rho^2 \int_{B_{2r_0}^{\xi}} \chi\left(\frac{|y|}{r_0}\right) \frac{|y|^2 - \tau^2 \rho^2}{(\tau^2 \rho^2 + |y|^2)^3} \left(-\frac{4\tau \rho^2 \chi\left(\frac{|y|}{r_0}\right)}{\tau^2 \rho^2 + |y|^2} + \mathcal{O}(\rho^{1+\alpha_0})\right) dy \\ &= \frac{64}{\tau^2} \int_{\frac{1}{\tau \rho} B_{r_0}^{\xi}} \frac{1 - |y|^2}{(1 + |y|^2)^4} + \mathcal{O}(\rho^{1+\alpha_0}). \end{split}$$

Considering that  $\frac{64}{\tau^2} \int_{\frac{1}{\tau\rho} B_{r_0}^{\xi}} \frac{1-|y|^2}{(1+|y|^2)^4} = \frac{8\varrho(\xi)}{\tau^2\pi} \int_{\mathbb{R}^2} \frac{1-|y|^2}{(1+|y|^2)^4} dy + \mathcal{O}(\rho^2)$ , as  $\rho \to 0$ ,

$$\langle P\Psi^{0}_{\tau,\xi}, P\Psi^{0}_{\tau,\xi} \rangle = \int_{\Sigma} \rho^{2} \chi_{\xi}(x) e^{-\varphi_{\xi}} e^{U_{\tau,\xi}} \Psi^{0}_{\tau,\xi} P\Psi^{0}_{\tau,\xi} dv_{g}(x) = \frac{8\varrho(\xi)D_{0}}{\pi\tau^{2}} + \mathcal{O}(\rho^{1+\alpha_{0}}),$$

where  $D_0 = \int_{\mathbb{R}^2} \frac{1-|y|^2}{(1+|y|^2)^4} dy$ .

Similarly, for j=0 and i=1,2 for  $\xi\in\mathring{\Sigma}$  and i=1 for  $\xi\in\partial\Sigma$ , we have

$$\begin{split} \langle P\Psi_{\tau,\xi}^{i}, P\Psi_{\tau,\xi}^{0} \rangle &= \rho^{2} \int_{\Sigma \cap U_{2r_{0}}(\xi)} \chi_{\xi}(x) e^{-\varphi_{\xi}(x)} e^{U_{\tau,\xi}} \Psi_{\tau,\xi}^{i} P\Psi_{\tau,\xi}^{0} dv_{g}(x) \\ &= 32\tau^{2} \rho^{2} \int_{B_{2r_{0}}^{\xi}} \chi\left(\frac{|y|}{r_{0}}\right) \frac{y_{i}}{(\tau^{2} \rho^{2} + |y|^{2})^{3}} \left(-\frac{4\tau \rho^{2} \chi\left(\frac{|y|}{r_{0}}\right)}{\tau^{2} \rho^{2} + |y|^{2}} + \mathcal{O}(\rho^{1+\alpha_{0}})\right) dy \\ &= \mathcal{O}(\rho^{\alpha_{0}}). \end{split}$$

Applying Lemma B.3, for  $\xi \in \mathring{\Sigma}$  we have

$$\rho^{2} \int_{\Sigma \cap U_{2r_{0}}(\xi)} \chi_{\xi}(x) e^{-\varphi_{\xi}(x)} e^{U_{\tau,\xi}} \Psi_{\tau,\xi}^{i} P \Psi_{\tau,\xi}^{j} dv_{g}(x)$$

$$= 32\tau^{2} \rho^{2} \int_{B_{2r_{0}}^{\xi}} \chi\left(\frac{|y|}{r_{0}}\right) \frac{y_{i}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}} \left(\chi\left(\frac{|y|}{r_{0}}\right) \frac{4y_{j}}{\tau^{2}\rho^{2} + |y|^{2}} + \varrho(\xi) H^{j}(y_{\xi}^{-1}(y), \xi) + \mathcal{O}(\rho^{\alpha_{0}})\right) dy$$

$$= \frac{128}{\tau^{2}\rho^{2}} \int_{\frac{1}{\tau\rho} B_{r_{0}}^{\xi}} \frac{y_{i}y_{j}}{(1 + |y|^{2})^{4}} dy + \int_{B_{r_{0}}^{\xi}} \frac{32\tau^{2}\rho^{2}\varrho(\xi)y_{i}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}} (H^{j}(y_{\xi}^{-1}(y), \xi) - H^{j}(\xi, \xi)) dy$$

$$+ 32\tau^{2}\rho^{2}\varrho(\xi) H^{j}(\xi, \xi) \int_{B_{r_{0}}^{\xi}} \frac{y_{i}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}} dy + \mathcal{O}(\rho^{\alpha_{0}-1})$$

$$= \frac{128}{\tau^{2}\rho^{2}} \int_{\frac{1}{\tau\rho} B_{r_{0}}^{\xi}} \frac{y_{i}y_{j}}{(1 + |y|^{2})^{4}} dy + \mathcal{O}\left(\int_{B_{r_{0}}^{\xi}} \frac{32\tau^{2}\rho^{2}|y|^{2}}{(\tau^{2}\rho^{2} + |y|^{2})^{3}} dy\right) + \mathcal{O}(\rho^{\alpha_{0}-1})$$

$$= \frac{8\varrho(\xi)D_{i}}{\pi\tau^{2}\rho^{2}} \delta_{ij} + \mathcal{O}(\rho^{\alpha_{0}-1})(\rho \to 0),$$

where  $D_i = \int_{\mathbb{R}^2} \frac{|y|^2}{(1+|y|^2)^4} dy$ . For  $\xi \in \partial \Sigma$ , applying Lemma B.3 again,

$$\begin{split} & \rho^2 \int_{\Sigma \cap U_{2r_0}(\xi)} e^{U_{\tau,\xi}} \Psi_{\tau,\xi}^1 P \Psi_{\tau,\xi}^1 dv_g(x) \\ & = \int_{B_{2r_0}^{\xi}} \chi\left(\frac{|y|}{r_0}\right) \frac{32\tau^2 \rho^2 y_1}{(\tau^2 \rho^2 + |y|^2)^3} \left(\frac{4\chi\left(\frac{|y|}{r_0}\right) y_1}{\tau^2 \rho^2 + |y|^2} + \varrho(\xi) H^1(y_{\xi}^{-1}(y), \xi) + \mathcal{O}(\rho^{\alpha_0})\right) \\ & = \frac{128}{\tau^2 \rho^2} \int_{\frac{1}{\tau \rho} B_{r_0}^{\xi}} \frac{y_1^2}{(1 + |y|^2)^4} + \mathcal{O}(\rho^{\alpha_0 - 1}). \end{split}$$

We observe that

$$\left| \frac{128}{\tau^2 \rho^2} \int_{\frac{1}{\tau \rho} B_{r_0}^{\xi}} \frac{y_1^2}{(1+|y|^2)^4} - \frac{128}{\tau^2 \rho^2} \int_{\mathbb{R}^2_+} \frac{y_1^2}{(1+|y|^2)^4} \right| \leq \frac{128}{\tau^2 \rho^2} \int_{\mathbb{R}^2_+ \setminus \frac{1}{\tau \rho} B_{r_0}^{\xi}} \frac{1}{(1+|y|^2)^3} dy \\
\leq \mathcal{O}(\rho^2),$$

and

$$\rho^2 \int_{\Sigma \setminus U_{2r_0}(\xi)} \chi_{\xi}(x) e^{-\varphi_{\xi}(x)} e^{U_{\tau,\xi}} \Psi^i_{\tau,\xi} P \Psi^j_{\tau,\xi} dv_g = \mathcal{O}(\rho^2 \|P\Psi^j_{\tau,\xi}\|) = \mathcal{O}(\rho)(\rho \to 0),$$

for  $i, j \in \{1, 2\}$ , if  $\xi \in \mathring{\Sigma}, i, j = 1, 2$ , and if  $\xi \in \partial \Sigma, i, j = 1$ . Thus we have

$$\langle P\Psi_{\tau,\xi}^i, P\Psi_{\tau,\xi}^j \rangle = \frac{8\varrho(\xi)D_i}{\pi\tau^2\rho^2}\delta_{ij} + \mathcal{O}(\rho^{\alpha_0-1}).$$

By assumption,  $r_0 > 0$  sufficiently small such that

$$U_{2r_0}(\xi_0) \cap U_{2r_0}(\xi_1) = \emptyset$$

and for l = 0, 1, if  $\xi_l \in \Sigma$ ,  $U_{2r_0}(\xi_l) \subset \Sigma$ .

$$\langle P\Psi^i_{\tau^0,\xi_0}, P\Psi^j_{\tau^1,\xi_1}\rangle = \int_{\Sigma\backslash U_{2r_0}(\xi_0)} + \int_{\Sigma\cap U_{2r_0}(\xi_0)} \rho^2 \chi_{\xi_0} e^{-\varphi_\xi} e^{U_{\tau^0,\xi_0}} \Psi^i_{\tau^0,\xi_0} P\Psi^j_{\tau^1,\xi_1} dv_g.$$

As  $\rho \to 0$ , we have

$$\int_{\Sigma \setminus U_{2r_0}(\xi_0)} \rho^2 \chi_{\xi_0}(x) e^{-\varphi_{\xi}(x)} e^{U_{\tau^0,\xi_0}} \Psi_{\tau^0,\xi_0}^i P \Psi_{\tau^1,\xi_1}^j dv_g = \mathcal{O}(\rho^2 \|P\Psi_{\tau^1,\xi_1}^j\|) = \mathcal{O}(\rho).$$

By Lemma B.3, for  $j \neq 0$ 

$$\int_{U_{2r_{0}}(\xi_{0})\cap\Sigma} \rho^{2} \chi_{\xi_{0}} e^{-\varphi_{\xi_{0}}} e^{U_{\tau^{0},\xi_{0}}} \Psi_{\tau^{0},\xi_{0}}^{i} P \Psi_{\tau^{1},\xi_{1}}^{j} dv_{g}$$

$$= \int_{U_{2r_{0}}(\xi_{0})} \rho^{2} \chi_{\xi_{0}} e^{-\varphi_{\xi_{0}}} e^{U_{\tau^{0},\xi_{0}}} \Psi_{\tau^{0},\xi_{0}}^{i} \left( \chi_{\xi_{1}} \frac{4y_{\xi_{1}}(x)_{j}}{\tau^{2}\rho^{2} + |y_{\xi_{1}}(x)|^{2}} + \varrho(\xi_{1}) H^{j}(x,\xi_{1}) + \mathcal{O}(\rho^{\alpha_{0}}) \right)$$

$$= \varrho(\xi_{i}) H^{j}(\xi_{0},\xi_{1}) \int_{U_{2r_{0}}(\xi_{0})} \rho^{2} \chi_{\xi_{0}} e^{-\varphi_{\xi_{0}}} e^{U_{\tau^{0},\xi_{0}}} \Psi_{\tau^{0},\xi_{0}}^{i} dv_{g}$$

$$+ \mathcal{O}\left( \int_{U_{2r_{0}}(\xi_{0})} \rho^{2} \chi_{\xi_{0}} e^{-\varphi_{\xi_{0}}} e^{U_{\tau^{0},\xi_{0}}} \Psi_{\tau^{0},\xi_{0}}^{i} (|y_{\xi_{0}}| + \rho^{\alpha_{0}}) dv_{g} \right)$$

$$= \mathcal{O}(\rho^{\alpha_{0}-1});$$

for j = 0,

$$\int_{U_{2r_0}(\xi_0)\cap\Sigma} \rho^2 \chi_{\xi_0} e^{-\varphi_{\xi_0}} e^{U_{\tau^0,\xi_0}} \Psi_{\tau^0,\xi_0}^i P \Psi_{\tau^1,\xi_1}^j dv_g 
= \int_{U_{2r_0}(\xi_0)} \rho^2 \chi_{\xi_0} e^{-\varphi_{\xi_0}} e^{U_{\tau^0,\xi_0}} \Psi_{\tau^0,\xi_0}^i \left( -\chi_{\xi_1} \frac{4\tau\rho^2}{\tau^2\rho^2 + |y_{\xi_1}|^2} + \mathcal{O}(\rho^{\alpha_0+1}) \right) 
= \mathcal{O}(\rho^{\alpha_0}).$$

Therefore for any  $\xi_1 \neq \xi_0$ ,

$$\langle P\Psi^i_{\tau^0,\xi_0}, P\Psi^j_{\tau^1,\xi_1}\rangle = \begin{cases} \mathcal{O}(\rho^{\alpha_0}) & \text{ when } i \text{ or } j=0\\ \mathcal{O}(\rho^{\alpha_0-1}) & \text{ otherwise} \end{cases}.$$

Remark B.2. Analogue to the proof in Lemma C.1, for any  $\alpha_0 \in (0,1)$ , we have as  $\rho \to 0$  for j, i = 1, 2 for  $\xi \in \mathring{\Sigma}$  and i, j = 0, 1 for  $\xi \in \partial \Sigma$ ,

$$\langle P\Psi_{\tau,\xi}^{i}, \partial_{\xi_{j}}PU_{\tau,\xi} \rangle = \frac{8\varrho(\xi)D_{i}}{\pi\tau^{2}\rho^{2}}\delta_{ij} + \mathcal{O}(\rho^{\alpha_{0}-1}),$$

and

$$\langle P\Psi^{i}_{\tau^{0},\xi_{0}}, \partial_{\xi_{j}}PU_{\tau^{1},\xi_{1}}\rangle = \begin{cases} \mathcal{O}(\rho^{\alpha_{0}}) & \text{when } i \text{ or } j = 0\\ \mathcal{O}(\rho^{\alpha_{0}-1}) & \text{otherwise} \end{cases},$$

where three different points  $\xi, \xi_0, \xi_1 \in \Sigma$  and uniformly in  $\tau, \tau^0, \tau^1$  are bounded away from zero and the  $\delta_{ij}$  is the Kronecker symbol, and  $D_0 = \int_{\mathbb{R}^2} \frac{1-|y|^2}{(1+|y|^2)^4} dy$ ,  $D_1 = D_2 = \int_{\mathbb{R}^2} \frac{|y|^2}{(1+|y|^2)^4} dy$ .

 $\mathbf{C}$ 

In the next two parts, we consider  $\xi = (\xi_1, \dots, \xi_m)$  in a compact subset of  $\mathcal{D}$ .

This part gives some technical lemmas to prove Proposition 3.1 which reduces the problem into a finite-dimensional problem.

**Lemma C.1.** Let  $\xi = (\xi_1, \dots, \xi_m)$  in a compact subset of  $\mathring{\Sigma}^k \times (\partial \Sigma)^{m-k} \setminus \Delta$ . For any  $p \geq 1$ , there is a positive constant c := c(p) such that for any  $\rho > 0$ ,

$$\left| \rho^2 V e^{\sum_{i=1}^m PU_i} - \rho^2 \sum_{i=1}^m \chi_i e^{U_i} \right|_{L^p(\Sigma)} \le c \rho^{\frac{2-p}{p}},$$

where  $p \in (1,2)$ .

*Proof.* Let  $\mathcal{D} \subset \mathscr{D}$  be a compact subset. Then there exists  $\delta > 0$  such that  $\mathcal{D} \subset M_{\delta}$ . There is a uniform  $r_0 > 0$  for any  $\xi \in M_{\delta}$ . By calculation,

$$\int_{\Sigma} |\rho^{2} V e^{\sum_{i=1}^{m} PU_{i}} - \rho^{2} \sum_{i=1}^{m} \chi_{i} e^{U_{i}}|^{p} dv_{g}$$

$$= \sum_{i=1}^{m} \int_{\Sigma \cap U_{2r_{0}}(\xi)} |\rho^{2} V e^{\sum_{i=1}^{m} PU_{i}} - \rho^{2} \sum_{h=1}^{m} \chi_{h} e^{U_{h}}|^{p} dv_{g}$$

$$+ \int_{\Sigma \setminus \bigcup_{i=1}^{m} U_{2r_{0}}(\xi)} |\rho^{2} V e^{\sum_{i=1}^{m} PU_{i}} - \rho^{2} \sum_{h=1}^{m} \chi_{h} e^{U_{h}}|^{p} dv_{g},$$

and as  $\rho \to 0$ ,  $\int_{\Sigma \setminus \bigcup_{i=1}^m U_{2r_0}(\xi)} |\rho^2 V e^{\sum_{i=1}^m P U_i} - \rho^2 \sum_{h=1}^m \chi_h e^{U_h}|^p dv_g(x) = \mathcal{O}(\rho^{2p})$ . By Lemma B.1, for any  $x \in U_{2r_0}(\xi_h)$ 

$$\sum_{i=1}^{m} PU_{i} - \chi_{h}U_{h}$$

$$= \left(\sum_{i \neq h} \varrho(\xi_{i})G^{g}(\xi_{h}, \xi_{i}) + \varrho(\xi_{h})H^{g}(\xi_{h}, \xi_{h}) - \log(8\tau_{h}^{2})\right) + \mathcal{O}(|y_{\xi_{h}}(x)| + \rho^{1+\alpha_{0}})$$

$$= -\log V(\xi_{h}) + \mathcal{O}(\rho^{1+\alpha_{0}} + |y_{\xi_{h}}(x)|).$$

Hence,

$$\int_{U_{2r_{0}}(\xi_{h})\cap\Sigma} |\rho^{2}V e^{\sum_{i=1}^{m} PU_{i}} - \rho^{2}\chi_{h}e^{U_{h}}|^{p}dv_{g}(x) 
= \int_{U_{r_{0}}(\xi_{h})\cap\Sigma} \left| \rho^{2}e^{U_{h}} (e^{\sum_{i=1}^{m} PU_{i}-\chi_{h}U_{h}+\log V(x)} - 1) \right|^{p} dv_{g}(x) + \mathcal{O}(\rho^{2p}) 
= \mathcal{O}\left(\int_{U_{r_{0}}(\xi_{h})\cap\Sigma} \rho^{2p}e^{pU_{h}} (|y_{\xi_{h}}(x)| + \rho^{1+\alpha_{0}})^{p}dv_{g}(x)\right) + \mathcal{O}(\rho^{2p}) 
= \mathcal{O}\left(\int_{B_{r_{0}}^{\xi_{h}}} \left(\frac{8\tau_{h}^{2}\rho^{2}(|y| + \rho^{1+\alpha_{0}})}{(\tau_{h}^{2}\rho^{2} + |y|^{2})^{2}}\right)^{p} dy + \rho^{2p}\right) 
= \mathcal{O}(\rho^{2-p}),$$

where  $p \in (1,2)$ .

**Lemma C.2.** For any  $p \ge 1$  and r > 1, there are positive constants  $c_1, c_2$  such that for any  $\rho > 0$ , the following estimates hold for any  $\phi_1, \phi_2 \in \overline{\mathbb{H}}^1$ .

$$(29) |\rho^2 V e^{\sum_{i=1}^m PU_i} (e^{\phi_1} - 1 - \phi_1)|_{L^p(\Sigma)} \le c_1 e^{c_2 ||\phi_1||^2} \rho^{\frac{(2-2pr)}{pr}} ||\phi_1||^2$$

and

(30) 
$$|\rho^{2}Ve^{\sum_{i=1}^{m}PU_{i}}(e^{\phi_{1}}-e^{\phi_{2}}-(\phi_{1}-\phi_{2}))|_{L^{p}(\Sigma)}$$

$$\leq c_{1}e^{c_{2}(\|\phi_{1}\|^{2}+\|\phi_{2}\|^{2})}\rho^{\frac{(2-2pr)}{pr}}(\|\phi_{1}\|+\|\phi_{2}\|)\|\phi_{1}-\phi_{2}\|.$$

*Proof.* By the mean value theorem, for some  $s \in (0,1)$ 

$$|(e^{\phi_1} - e^{\phi_2} - (\phi_1 - \phi_2))| \le |e^{s\phi_1 + (1-s)\phi_2} - 1| |\phi_1 - \phi_2| \le e^{|\phi_1| + |\phi_2|} |\phi_1 - \phi_2| (|\phi_1| + |\phi_2|).$$

Applying the Hölder inequality, Sobolev inequality and Moser-Trudinger inequality, we deduce that

$$\left( \int_{\Sigma} V^{p} e^{p\sum_{i=1}^{m} PU_{i}} |e^{\phi_{1}} - e^{\phi_{2}} - (\phi_{1} - \phi_{2})|^{p} dv_{g} \right)^{1/p}$$

$$\leq C \sum_{h=1}^{2} \left( \int_{\Sigma} V^{p} e^{p\sum_{i=1}^{m} PU_{i}} (e^{|\phi_{1}| + |\phi_{2}|} |\phi_{1} - \phi_{2}| |\phi_{h}|)^{p} dv_{g} \right)^{1/p}$$

$$\leq C \sum_{h=1}^{2} \left( \int_{\Sigma} V^{pr} e^{pr\sum_{i=1}^{m} PU_{i}} dv_{g} \right)^{\frac{1}{pr}} \left( \int_{\Sigma} e^{ps(|\phi_{1}| + |\phi_{2}|)} dv_{g} \right)^{\frac{1}{ps}}$$

$$\left( \int_{\Sigma} |\phi_{1} - \phi_{2}|^{pt} |\phi_{h}|^{pt} dv_{g} \right)^{\frac{1}{pt}}$$

$$\leq C \sum_{h=1}^{2} \left( \int_{\Sigma} V^{pr} e^{pr\sum_{i=1}^{m} PU_{i}} dv_{g}(x) \right)^{\frac{1}{pr}} e^{\frac{ps}{8\pi} (\|\phi_{1}\|^{2} + \|\phi_{2}\|^{2})} \|\phi_{1} - \phi_{2}\| \|\phi_{h}\|,$$

where 
$$r, s, t \in (1, +\infty)$$
,  $\frac{1}{r} + \frac{1}{s} + \frac{1}{t} = 1$ . By Lemma B.1,  

$$\int_{\bigcup_{i=1}^{m} U_{2r_0}(\xi_i)} V^{pr} e^{pr \sum_{i=1}^{m} PU_i} dv_g(x)$$

$$= \sum_{i=1}^{m} \int_{U_{2r_0}(\xi_i)} \exp \left\{ pr \chi_i U_i + pr \left( \sum_{h \neq i} G^g(\xi_i, \xi_h) + \varrho(\xi_i) H^g(\xi_i, \xi_i) + \log V(\xi_i) - \log(8\tau_i^2) \right) + \mathcal{O}(\rho^{1+\alpha_0} + |y_{\xi_i}(x)|) \right\} dv_g(x)$$

$$\leq C \left( \sum_{i=1}^{m} \int_{U_{2r_0}(\xi_i)} e^{pr \chi_i U_i} (1 + \mathcal{O}(\rho^{1+\alpha_0} + |y_{\xi_i}(x)|)) dv_g(x) \right)$$

$$\leq C \left( \sum_{i=1}^{m} \int_{B_{2r_0}^{\xi_i}} e^{\hat{\varphi}_{\xi_i}(y)} \left( \frac{8\tau_i^2}{(\tau_i^2 \rho^2 + |y|^2)^2} \right)^{pr} (1 + \mathcal{O}(\rho^{1+\alpha_0} + |y|)) dy \right)$$

$$\leq C \rho^{2-4pr}$$

By the definition of  $PU_i$ ,  $PU_i = \mathcal{O}(1)$  constant in  $\Sigma \setminus U_{2r_0}(\xi_i)$ . It follows that

$$\sum_{\Sigma \setminus \bigcup_{i=1}^m U_{2r_0}(\xi_i)} e^{pr \sum_{i=1}^m PU_i} = \mathcal{O}(1).$$

Therefore the estimate (30) holds and if we take  $\phi_2 \equiv 0$ , we obtain the estimate (29).

D

**Proof of Lemma 3.1.** Assume the conclusion in Lemma 3.1 does not hold. Then there exists  $\xi \in M_{\delta} \subset \mathscr{D}$  for some small  $\delta > 0$ , a sequence  $\rho_n \to 0$  and  $\phi_n \in K_{\xi}^{\perp}$  with  $\|\phi_n\| = 1$  and  $\|L_{\xi}^{\rho_n}(\phi)\| = o(\frac{1}{|\log \rho_n|})$ . To simplify the notations, we use  $\rho$  instead of  $\rho_n$  and  $\phi$  instead of  $\phi_n$ .

(31) 
$$\phi - i^*(\rho^2 V e^{\sum_{i=1}^m PU_i} \phi) = \psi + w,$$

where  $\psi \in K_{\xi}^{\perp}$  and  $w \in K_{\xi}$ . Then  $\|\psi\| = o(\frac{1}{|\log \rho|}) \to 0(\rho \to 0)$ . It is equivalent that  $\phi$  solves the following problem in the weak sense,

$$\begin{cases} (-\Delta_g + \beta)\phi = \rho^2 V e^{\sum_{i=1}^m PU_i} \phi - \overline{\rho^2 V e^{\sum_{i=1}^m PU_i} \phi} + (-\Delta_g + \beta)(\psi + w), & \text{in } \mathring{\Sigma}, \\ \partial_{\nu_g} \phi = 0, & \text{on } \partial \Sigma. \end{cases}$$

Step 1.  $||w|| = o(1)(\rho \to 0)$ .

Given that  $w \in K_{\xi}$ , we have  $w = \sum_{i=1}^{m} \sum_{j=1}^{J_i} c_{ij} P \Psi_i^j$ . Consider the inner product of equation (31) with  $P \Psi_{i'}^{j'}$ , leading to the following equation:

$$\begin{split} &\langle \phi, P\Psi_{i'}^{j'} \rangle - \int_{\Sigma} P\Psi_{i'}^{j'} \left( \rho^2 V e^{\sum_{i=1}^m PU_i} \phi - \frac{1}{|\Sigma|_g} \int_{\Sigma} \rho^2 V e^{\sum_{i=1}^m PU_i} \phi dv_g \right) dv_g \\ &= &\langle \psi, P\Psi_{i'}^{j'} \rangle + \langle w, P\Psi_{i'}^{j'} \rangle. \end{split}$$

Since  $P\Psi_{i'}^{j'} \in \overline{H}^1$  and  $\phi \in K_{\xi}^{\perp}$ , we have  $\int_{\Sigma} P\Psi_{i'}^{j'} dv_g = 0$  and  $\langle \psi, P\Psi_{i'}^{j'} \rangle = \langle \phi, P\Psi_{i'}^{j'} \rangle = 0$ . It follows

(32) 
$$-\rho^{2} \int_{\Sigma} V e^{\sum_{i=1}^{m} PU_{i}} \phi P \Psi_{i'}^{j'} dv_{g} = \sum_{i=1}^{m} \sum_{j=1}^{J_{i}} c_{ij} \langle P \Psi_{i}^{j}, P \Psi_{i'}^{j'} \rangle.$$

Applying Lemma B.4, the right-hand side of the equation (32) equals

$$\frac{8\varrho(\xi_{i'})D_1}{\pi\tau_{i'}^2\rho^2}c_{i'j'} + \mathcal{O}(\rho^{\alpha_0-1}\sum_{i=1}^m\sum_{j=1}^{J_i}|c_{ij}|).$$

The left-hand side of equation (32) can be expanded as follows:

$$\int_{\Sigma} \rho^{2} \left( \sum_{i=1}^{m} \chi_{i} e^{U_{i}} - V e^{\sum_{i=1}^{m} PU_{i}} \right) P \Psi_{i'}^{j'} \phi dv_{g} - \sum_{i=1}^{m} \int_{\Sigma} \rho^{2} \chi_{i} e^{U_{i}} (P \Psi_{i'}^{j'} - \chi_{i'} \Psi_{i'}^{j'}) \phi dv_{g} - \rho^{2} \int_{\Sigma} \chi_{i'}^{2} (-e^{-\varphi_{i'}} + 1) e^{U_{i'}} \Psi_{i'}^{j'} \phi dv_{g} - \rho^{2} \int_{\Sigma} \chi_{i'}^{2} e^{-\varphi_{i'}} e^{U_{i'}} \Psi_{i'}^{j'} \phi dv_{g}.$$

Since  $\|\phi\| = 1$  and  $\phi \in K_{\xi}^{\perp}$ ,  $\int_{\Sigma} \rho^2 \chi_{i'}^2 e^{-\varphi_{i'}} e^{U_{i'}} \Psi_{i'}^{j'} \phi = \mathcal{O}(\rho^2) + \langle P \Psi_{i'}^{j'}, \phi \rangle = \mathcal{O}(\rho^2)$ . By calculation, we have

$$\left| \rho^2 \int_{\Sigma} (e^{-\varphi_{i'}} - 1) \chi_{i'} e^{U_{i'}} \Psi_{i'}^{j'} dv_g \right| \leq \mathcal{O}\left( \int_{|y| \le 2r_0} \frac{\tau_i^2 \rho^2 |y|^3 dy}{(\tau_i^2 \rho^2 + |y|^2)^3} \right) = \mathcal{O}(\rho).$$

Applying Lemma B.4 and Lemma C.1,

$$\left| \int_{\Sigma} \rho^{2} \left( \sum_{i=1}^{m} \chi_{i} e^{U_{i}} - V e^{\sum_{i=1}^{m} P U_{i}} \right) P \Psi_{i'}^{j'} \phi dv_{g} \right|$$

$$\leq C \left| \rho^{2} \left( \sum_{i=1}^{m} \chi_{i} e^{U_{i}} - V e^{\sum_{i=1}^{m} P U_{i}} \right) \right|_{L^{p}(\Sigma)} |\phi|_{L^{q}(\Sigma)} ||P \Psi_{i'}^{j'}||$$

$$\leq \mathcal{O}(\rho^{\frac{2-p}{p}-1}) = \mathcal{O}(\rho^{\frac{2(1-p)}{p}}),$$

where  $\frac{1}{p} + \frac{1}{q} < 1$  and C > 0 is a constant. Further, Lemma B.3 implies  $P\Psi_{i'}^{j'} - \chi_{i'}\Psi_{i'}^{j'} = \mathcal{O}(1)(\rho \to 0)$ . And applying Lemma B.2, for any  $i = 1, \dots, m$ 

$$\left| \rho^2 \int_{\Sigma} \chi_i e^{U_i} \phi(P\Psi_{i'}^{j'} - \chi_{i'}\Psi_{i'}^{j'}) dv_g \right| \leq \mathcal{O}(|\rho^2 \chi_i e^{U_i}|_{L^p(\Sigma)} |\phi|_{L^q(\Sigma)})$$

$$\leq \mathcal{O}(\rho^{\frac{2(1-p)}{p}}) (\rho \to 0).$$

Combining these estimates, we conclude

$$\frac{8\varrho(\xi_{i'})D_{i'}}{\pi\tau_{i'}^2\rho^2}c_{i'j'} + \mathcal{O}\left(\rho^{\alpha_0-1}\sum_{i=1}^m\sum_{j=1}^{J_i}|c_{ij}|\right) = \mathcal{O}(\rho^{\frac{2(1-p)}{p}})(\rho\to 0).$$

Then  $|c_{i'j'}| = \mathcal{O}(\rho^{\frac{2}{p}})(\rho \to 0)$ , where  $p \in (1,2)$ . So

(33) 
$$\sum_{i=1}^{m} \sum_{j=1}^{J_i} |c_{ij}| = \mathcal{O}(\rho^{\frac{2}{p}})$$

by the arbitrariness of i' and j'. Lemma B.4 and (33) yield that

$$||w||^2 = \left|\left|\sum_{i=1}^m \sum_{j=1}^{J_i} c_{ij} P \Psi_i^j\right|\right|^2 = \mathcal{O}\left(\sum_{i=1}^m \sum_{j=1}^{J_i} |c_{ij}|^2 \frac{1}{\rho^2} + \mathcal{O}(\rho^{\alpha_0 - 1})\right) \le \mathcal{O}(\rho^{\frac{4}{p} - 2}).$$

Hence, it follows that as  $\rho \to 0$ ,  $||w|| = \mathcal{O}(\rho^{\frac{2-p}{p}}) \to 0$  for any  $p \in (1,2)$ .

Step 2. 
$$\langle \phi, P\Psi_i^0 \rangle \to 0 (\rho \to 0)$$
.

Following the construction in [16] and [15], we define

$$\omega_i(y) = \frac{4}{3\tau_i} \log(\tau_i^2 \rho^2 + |y|^2) \frac{\tau_i^2 \rho^2 - |y|^2}{\tau_i^2 \rho^2 + |y|^2} + \frac{8}{3\tau_i} \frac{\tau_i^2 \rho^2}{\tau_i^2 \rho^2 + |y|^2},$$

and

$$t_i(y) = -2\frac{\tau_i^2 \rho^2}{\tau_i^2 \rho^2 + |y|^2}.$$

And we observe that  $\omega_i(y)$ ,  $t_i(y)$  satisfies the following equations respectively,

$$-\Delta\omega_{i} - \rho^{2}e^{u_{\tau_{i},0}}\omega_{i} = \rho^{2}e^{u_{\tau_{i},0}}\psi_{\tau_{i},0}^{0} \text{ in } \mathbb{R}^{2}$$

and

$$-\Delta t_i - \rho^2 e^{u_{\tau_i,0}} t_i = \rho^2 e^{u_{\tau_i,0}} \text{ in } \mathbb{R}^2.$$

$$\int_{\mathbb{R}^2} |\nabla \omega_i|^2 = M_i^2 (1 + o(1)) (\log \rho)^2, \quad \int_{\mathbb{R}^2} |\nabla t_i|^2 = O(1), \text{ as } \rho \to 0$$

with  $M_i = \frac{32}{3\tau_i} \left( \int_{\mathbb{R}^2} \frac{|y|^2}{(1+|y|^2)^4} \right)^{1/2}$ . Let

$$u_i(x) = \chi_i(x) \left( \omega_i(y_{\xi_i}(x)) + \frac{2\varrho(\xi_i)}{3\tau_i} H^g(\xi_i, \xi_i) t_i(y_{\xi_i}(x)) \right), \text{ for all } x \in U_{2r_0}(\xi_i).$$

The projection  $Pu_i \in \overline{\mathbf{H}}^1$  from  $u_i$  is given by

(34) 
$$\begin{cases} (-\Delta_g + \beta)Pu_i = -\chi_i \Delta_g u_i(x) + \overline{\chi_i \Delta_g u_i(x)} & x \in \mathring{\Sigma} \\ \partial_{\nu_g} Pu_i = 0 & x \in \partial \Sigma \\ \int_{\Sigma} Pu_i = 0 \end{cases}$$

Let us consider  $\eta_i = u_i - Pu_i + \frac{2\varrho(\xi_i)}{3\tau_i}H^g(x,\xi_i)$ . The integral of  $\eta_i$  over  $\Sigma$  is given by  $\int_{\Sigma} \eta_i dv_g = \mathcal{O}(\rho^2 \log^2 \rho)$ . If  $\xi_i \in \mathring{\Sigma}$ , we have  $\partial_{\nu_g} \eta_i \equiv 0$  in  $\partial \Sigma$ . For  $\xi_i \in \partial \Sigma$ ,  $|\partial_{\nu_g} \eta_i(x)|_{L^p(\partial \Sigma)} = \mathcal{O}(\rho^{\frac{1}{p}} |\log \rho|)$ . In view of  $\int_{\mathbb{R}^2} \frac{1-|y|^2}{(1+|y|^2)^3} \log(1+|y|^2) dy = -\frac{\pi}{2}$ , and  $\int_{\mathbb{R}^2} \frac{2}{(1+|y|^2)^3} dy = \int_{\mathbb{R}^2} \frac{1}{(1+|y|^2)^2} dy = \pi$ ,

$$|(-\Delta_g + \beta)\eta_i|_{L^p(\Sigma)} = \mathcal{O}(\rho^{\frac{1}{p}}|\log \rho|).$$

By the  $L^p$ -theory,

$$\|\eta_i - \overline{\eta_i}\|_{W^{2,p}(\Sigma)} \le C\left(\left|\partial_{\nu_g}\eta_i\right|_{L^p(\partial\Sigma)} + \left|\left(-\Delta_g + \beta\right)\eta_i\right|_{L^p(\Sigma)}\right) \le C\rho^{\frac{1}{p}}\left|\log\rho\right|,$$

for any p > 1. Applying Sobolev's embedding theory,  $|\eta_i - \overline{\eta_i}|_{C^{\gamma}(\Sigma)} \leq C \rho^{\frac{1}{p}} |\log \rho|$ , for any  $\gamma \in (0, 2(1 - \frac{1}{p}))$ . Choosing  $p \in (1, 2]$ , we deduce that

$$(35) |\eta_i| \le \mathcal{O}(\rho^{\frac{1}{p}}|\log \rho|).$$

Moreover, for any  $x \in \Sigma \setminus \{\xi_i\}$ , the following inequality holds:

(36) 
$$\left| Pu_i(x) - \frac{2\varrho(\xi_i)}{3\tau_i} G^g(x, \xi_i) \right| \le \mathcal{O}(\rho^{\frac{1}{p}} |\log \rho|).$$

Additionally,  $||Pu_i||^2$  is computed directly as

$$||Pu_i||^2 = \langle Pu_i, Pu_i \rangle = -\int_{\Sigma} \chi_i \left( u_i + \frac{2\varrho(\xi_i)}{3\tau_i} H_{\xi_i}^g + \mathcal{O}(\rho^{\frac{1}{p}} |\log \rho|) \right) \Delta_g u_i$$
  
=  $\mathcal{O}(|\log \rho|^2)(\rho \to 0).$ 

Thus

(37) 
$$||Pu_i|| = \mathcal{O}(|\log \rho|)(\rho \to 0).$$

Applying  $Pu_i$  as a test function for (31),

$$\langle Pu_i, \phi \rangle - \int_{\Sigma} \rho^2 V e^{\sum_{h=1}^m PU_h} \phi Pu_i dv_g = \langle Pu_i, w + \psi \rangle.$$

Considering

$$|\langle Pu_i, w + \psi \rangle| \le ||Pu_i||(||w|| + ||h||) \le ||Pu_i||o\left(\frac{1}{|\log \rho|}\right) = o(1),$$

we deduce that

(38) 
$$\langle Pu_i, \phi \rangle - \int_{\Sigma} \rho^2 V e^{\sum_{h=1}^m PU_h} \phi Pu_i dv_g(x) = o(1).$$

By (34) and  $\|\phi\| = 1$  with the Hölder inequality,

$$(39) \qquad \langle Pu_{i}, \phi \rangle = \int_{\Sigma} (-\chi_{i} \Delta_{g} u_{i} + \overline{\chi_{i}} \Delta_{g} u_{i}) \phi dv_{g}$$

$$= \int_{\Sigma} \rho^{2} e^{U_{i}} u_{i} dv_{g} + \int_{\Sigma} \rho^{2} e^{-\varphi_{i}} e^{U_{i}} \Psi_{i}^{0} dv_{g}$$

$$+ \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} H^{g}(\xi_{i}, \xi_{i}) \rho^{2} \chi_{i} e^{-\varphi_{i}} e^{U_{i}} \phi dv_{g} + \mathcal{O}(\rho^{2})$$

$$= \int_{\Sigma} \rho^{2} e^{U_{i}} u_{i} dv_{g} + \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} H^{g}(\xi_{i}, \xi_{i}) \rho^{2} \chi_{i} (e^{-\varphi_{i}} - 1) e^{U_{i}} \phi dv_{g}$$

$$+ \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} (H^{g}(\xi_{i}, \xi_{i}) - H^{g}(x, \xi_{i})) \rho^{2} \chi_{i} e^{U_{i}} \phi dv_{g}$$

$$+ \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} H^{g}(x, \xi_{i}) \rho^{2} \chi_{i} e^{U_{i}} \phi dv_{g} + \langle P\psi_{i}^{0}, \phi \rangle + \mathcal{O}(\rho^{2})$$

$$= \int_{\Sigma} \rho^{2} e^{U_{i}} u_{i} dv_{g} + \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} H^{g}(x, \xi_{i}) \rho^{2} \chi_{i} e^{U_{i}} \phi dv_{g} + \langle P\Psi_{i}^{0}, \phi \rangle + \mathcal{O}(\rho^{\frac{2-q}{q}}),$$

for any  $q \in (1, 2)$ . On the other hand, (B.2) and (35) with the Hölder inequality yield that

$$(40) \qquad \int_{\Sigma} \rho^{2} V e^{\sum_{h=1}^{m} P U_{h}} \phi P u_{i} dv_{g} = \int_{\Sigma} \rho^{2} \chi_{i} V e^{U_{i}} (1 + \mathcal{O}(|y_{\xi_{i}}(x)| + \rho^{\alpha_{0}+1}))$$

$$\cdot \left( u_{i} + \frac{2\varrho(\xi_{i})}{3\tau_{i}} H_{\xi_{i}}^{g} + \mathcal{O}(\rho^{\frac{1}{p}} |\log \rho|) \right) \phi dv_{g}$$

$$= \int_{\Sigma} \rho^{2} e^{U_{i}} u_{i} dv_{g} + \int_{\Sigma} \frac{2\varrho(\xi_{i})}{3\tau_{i}} H^{g}(x, \xi_{i}) \rho^{2} \chi_{i} e^{U_{i}} \phi dv_{g}$$

$$+ \mathcal{O}(\rho^{\frac{1}{p}+2\left(\frac{1}{s}-1\right)}),$$

for any  $s \in (1,2)$ . We choose s, p sufficiently close to 1 such that  $\frac{1}{p} + 2(\frac{1}{s} - 1) > 0$ . Then (39) and (40) imply that  $\langle P\Psi_i^0, \phi \rangle = o(1)$ , as  $\rho \to 0$ .

Step 3. Construct a contradiction.

Define the following space for  $\xi = (\xi_1, \dots, \xi_m) \in M_\delta$ . We denote that  $\mathbb{R}_i = \mathbb{R}^2$  if  $1 \leq i \leq k$ ;  $\mathbb{R}_i = \mathbb{R}_+^2 := \{y \in \mathbb{R}^2 : y_2 \geq 0\}$  if  $k+1 \leq i \leq m$ . Let  $\pi_N$  be the stereographic projection through the north pole for the standard unit sphere in  $\mathbb{R}^3$ . We denote that  $S_i = \pi_N(\mathbb{R}_i)$  for  $i = 1, \dots, m$ . We define

$$L_i := \left\{ \Psi : \left| \frac{\Psi}{1 + |y|^2} \right|_{L^2(\mathbb{R}_i)} < +\infty \right\},$$

and

$$H_i := \left\{ \Psi : |\nabla \Psi|_{L^2(\mathbb{R}_i)} + \left| \frac{\Psi}{1 + |y|^2} \right|_{L^2(\mathbb{R}_i)} < \infty \right\}$$

The associated norms are defined as the following,

$$\|\Psi\|_{L_i} := \left|\frac{\Psi}{1+|y|^2}\right|_{L^2(\mathbb{R}_i)} \text{ and } \|\Psi\|_{H_i} := |\nabla\Psi|_{L^2(\mathbb{R}_i)} + \left|\frac{\Psi}{1+|y|^2}\right|_{L^2(\mathbb{R}_i)}.$$

The maps

(41) 
$$L_i \to L^2(S_i) : \Psi \mapsto \Psi \circ \pi_N$$

and  $H_i \to H^1(S_i) : \Psi \mapsto \Psi \circ \pi_N$  are isometric. Let  $\Omega_i^{\rho} := \frac{1}{\tau_i \rho} B_{2r_0}^{\xi_i}, \phi_i^{\rho}(x) = \phi(y_{\xi_i}^{-1}(\tau_i \rho y))$  and  $\chi_i^{\rho}(y) = \chi(\tau_i \rho |y|)$ . Consider

$$\tilde{\phi}_i^{\rho} = \begin{cases} \phi_i^{\rho} \chi_i^{\rho} & y \in \Omega_i^{\rho} \\ 0 & y \in \mathbb{R}_i \setminus \Omega_i^{\rho} \end{cases}.$$

By Lemma C.1, we have

$$\sum_{h=1}^{m} \rho^{2} \int_{\Sigma} \chi_{h} e^{U_{h}} \phi^{2} dv_{g} = \rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \phi^{2} dv_{g} 
+ \mathcal{O} \left( \int_{\Sigma} \rho^{2} | \sum_{h=1}^{m} \chi_{h} e^{U_{h}} - V e^{\sum_{h=1}^{m} PU_{h}} | \phi^{2} dv_{g} \right) 
\leq \rho^{2} \int_{\Sigma} e^{\sum_{h=1}^{m} PU_{h}} \phi^{2} 
+ \mathcal{O} \left( \left| \rho^{2} \left( \sum_{h=1}^{m} \chi_{h} e^{U_{h}} - V e^{\sum_{h=1}^{m} PU_{h}} \right) \right|_{L^{p}(\Sigma)} |\phi^{2}|_{L^{q}(\Sigma)} \right) 
= \rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \phi^{2} dv_{g} + \mathcal{O}(\rho^{\frac{2-p}{p}}) 
= \rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \phi^{2} dv_{g} + o(1)(\rho \to 0),$$

where  $p \in (1,2)$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . On the other hand, we take the inner product of (31) with  $\phi$ , since  $\|\phi\| = 1$  and  $\|\psi\| = o(\frac{1}{\log \rho})$ ,

$$\rho^2 \int_{\Sigma} V e^{\sum_{h=1}^m PU_h} \phi^2 dv_g = \langle \phi, \phi \rangle - \langle w + \psi, \phi \rangle = 1 + o(1).$$

$$\sum_{i=1}^{m} \rho^{2} \int_{\Sigma} \chi_{i} e^{U_{i}} \phi^{2} dv_{g} = \sum_{i=1}^{m} \int_{B_{2r_{0}}^{\xi_{i}}} \frac{8\tau_{i}^{2} \rho^{2} \chi^{2}(|y|/r_{0})}{(\tau_{i}^{2} \rho^{2} + |y|^{2})^{2}} (\phi \circ y_{\xi_{i}}^{-1}(\tau_{i} \rho y))^{2} dy + \mathcal{O}(\rho^{2})$$

$$= 8 \sum_{i=1}^{m} \int_{\mathbb{R}_{i}} \frac{|\tilde{\phi}_{i}^{\rho}(y)|^{2}}{(1 + |y|^{2})^{2}} dy + \mathcal{O}(\rho^{2}).$$

$$\begin{split} \int_{\Omega_i^{\rho}} |\nabla \widetilde{\phi}_i^{\rho}|^2 dy &= \int_{\frac{1}{\tau_i \rho} B_{2r_0}^{\xi_i}} |\chi_i^{\rho} \nabla \phi_i^{\rho} + \phi_i^{\rho} \nabla \chi_i^{\rho}|^2 dy \\ &= \mathcal{O}\left(\int_{B_{2r_0}^{\xi_i}} (|\nabla \phi_i(y_{\xi_i}^{-1}(y))|^2 + |\phi(y_{\xi_i}^{-1}(y))|^2) dy\right) \\ &= \mathcal{O}\left(\int_{\Sigma} |\nabla \phi|_g^2 dv_g + \int_{\Sigma} e^{-\varphi_i} |\phi(x)|^2 dv_g\right) \\ &= \mathcal{O}\left(||\phi||\right) = \mathcal{O}(1). \end{split}$$

Hence  $\tilde{\phi}_i^{\rho}$  is bounded in  $H_i$ . We observe that  $H_i$  compactly embeds into  $L_i$ . Up to a subsequence, as  $\rho \to 0$ ,  $\tilde{\phi}_i^{\rho} \to \tilde{\phi}_i^0$  weakly in  $H_i$  and strongly in  $L_i$ .

(42) 
$$\sum_{i=1}^{m} \|\tilde{\phi}_{i}^{0}\|_{L_{i}}^{2} = \frac{1}{8}.$$

For any  $h \in C_c^{\infty}(\mathbb{R}^2)$ , assume that supp  $h \subset B_{R_0}(0)$ . If  $\tau_i \rho < \frac{r_0}{R_0}$ , then supp  $\nabla \chi \left(\frac{|y|}{r_0}\right) \cap \sup h\left(\frac{1}{\tau_i \rho}y\right) = \emptyset$ . For any  $\Phi \in \overline{\mathbb{H}}^1$ ,

$$(43) 0 = \int_{B_{2r_0}^{\xi_i}} \Phi \circ y_{\xi_i}^{-1}(y) \nabla \chi \left(\frac{|y|}{r_0}\right) \cdot \nabla h \left(\frac{1}{\tau_i \rho} y\right) dy$$
$$= \int_{B_{2r_0}^{\xi_i}} h \left(\frac{1}{\tau_i \rho} y\right) \nabla (\Phi \circ y_{\xi_i}^{-1}(y)) \cdot \nabla \chi \left(\frac{|y|}{r_0}\right) dy.$$

In (43), we take  $\Phi = \phi$ , w and  $\psi$ , respectively.

Claim. for any  $||h|| := (\int_{\mathbb{R}^2} |\nabla h|^2 + |h|^2)^{\frac{1}{2}} \le 1$  and  $h \in C_c^{\infty}(\mathbb{R}^2)$ ,

(44) 
$$\int_{\Sigma} \chi_i h^2 \left( \frac{1}{\tau_i \rho} y_{\xi_i}(x) \right) dv_g(x) = O(\rho^2) (\rho \to 0)$$

and

(45) 
$$\int_{\Sigma} \chi_i \left| \nabla h \left( \frac{1}{\tau_i \rho} y_{\xi_i}(x) \right) \right|_q^2 dv_g(x) = \mathcal{O}(1)(\rho \to 0).$$

Indeed.

$$\int_{\Sigma} \chi_{i} h^{2} \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(y) \right) dv_{g}(x) = \int_{B_{2\tau_{0}}^{\xi_{i}}} e^{\hat{\varphi}_{i}(y)} \chi \left( \frac{|y|}{r_{0}} \right) h^{2} \left( \frac{1}{\tau_{i}\rho} y \right) dy \\
\leq \mathcal{O} \left( \rho^{2} \int_{\Omega_{i}^{\rho}} |h(y)|^{2} dy \right) \\
\leq \mathcal{O}(\rho^{2}) ||h|| = \mathcal{O}(\rho^{2}). \\
\int_{\Sigma} \chi_{i} \left| \nabla h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(y) \right) \right|_{g}^{2} dv_{g}(x) = \int_{B_{2\tau_{0}}^{\xi_{i}}} \chi \left( \frac{|y|}{r_{0}} \right) \left| \nabla h \left( \frac{1}{\tau_{i}\rho} y \right) \right| dy \\
\leq \mathcal{O} \left( \int_{\Omega_{i}^{\rho}} |\nabla h(y)| dy \right) \leq \mathcal{O}(1).$$

The claim above concluded.

Combining the result in Step 1 and  $\|\psi\| = o(\frac{1}{|\log \rho|})$ 

(46) 
$$||w|| + ||\psi|| = o(1)(\rho \to 0).$$

Assume that  $0 < \rho < \frac{r_0}{\tau_i R_0}$ , as  $\rho \to 0$ 

$$\begin{split} &\int_{\mathbb{R}_{i}} \nabla \tilde{\phi}_{i}^{\rho} \nabla h dy = \int_{B_{2r_{0}}^{\xi_{i}}} \nabla \left( \chi \left( \frac{|y|}{r_{0}} \right) \phi \circ y_{\xi_{i}}^{-1}(y) \right) \cdot \nabla h \left( \frac{1}{\tau_{i}\rho} y \right) dy \\ \overset{(43)}{=} &\int_{B_{2r_{0}}^{\xi_{i}}} \nabla \phi \circ y_{\xi_{i}}^{-1}(y) \cdot \nabla \left( \chi \left( \frac{|y|}{r_{0}} \right) h \left( \frac{1}{\tau_{i}\rho} y \right) \right) dy \\ &= &\int_{\Sigma} \left\langle \nabla \phi, \nabla \left( \chi_{i}(x) h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) \right) \right\rangle_{g} dv_{g} \\ \overset{(31)}{=} &-\beta \int_{\Sigma} \chi_{i} h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) \phi dv_{g} + \int_{\Sigma} \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} PU_{h}} \phi h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) dv_{g}(x) \\ &- \frac{1}{|\Sigma|_{g}} \int_{\Sigma} \rho^{2} V e^{\sum_{h=1}^{m} PU_{h}} \phi dv_{g}(x) \int_{\Sigma} \chi_{i} h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) dv_{g}(x) \\ &+ \int_{\Sigma} \left\langle \nabla (w + \psi), \nabla \left( \chi_{i}(x) h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) \right) \right\rangle_{g} dv_{g} \\ &= &- \tau_{i}^{2} \rho^{2} \beta \int_{\mathbb{R}_{i}} \tilde{\phi}_{i}^{\rho}(y) h(y) dy + \int_{\Sigma} \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} PU_{h}} \phi h \left( \frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x) \right) dv_{g}(x) \\ &- \tau_{i}^{2} \rho^{2} \int_{\Sigma} \rho^{2} V e^{\sum_{h=1}^{m} PU_{h}} \phi dv_{g}(x) \int_{\mathbb{R}_{i}} \chi \left( \frac{\tau_{i}\rho|y|}{r_{0}} \right) e^{\varphi_{i}(\tau_{i}\rho y)} h(y) dy + o(1), \end{split}$$

for any  $h \in C_c^{\infty}(\mathbb{R}^2)$  with  $||h|| \leq 1$ . By the Hölder inequality and Lemma C.1,

$$\left| \int_{\mathbb{R}_i} \tilde{\phi}_i^{\rho}(y) h(y) dy \right| \leq \|\tilde{\phi}_i^{\rho}(y)\|_{L_i} \left( \int_{\mathbb{R}_i} (1 + |y|^2)^2 |h(y)| dy \right)^{\frac{1}{2}} \leq C \|\tilde{\phi}_i^{\rho}(y)\|_{L_i} \|h\|,$$

where C > 0 is a constant depending on  $R_0$ , and

$$\left| \rho^{2} \int_{\Sigma} \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} P U_{h}} \phi \right| \leq \rho^{2} \left| \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} P U_{h}} \right|_{L^{p}(\Sigma)} \|\phi\| 
\leq \rho^{2} \left( \left| \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} P U_{h}} - \rho^{2} \sum_{i=1}^{m} \chi_{i} e^{U_{i}} \right|_{L^{p}(\Sigma)} + \left| \rho^{2} \sum_{i=1}^{m} \chi_{i} e^{U_{i}} \right|_{L^{p}(\Sigma)} \right) \|\phi\| 
= \mathcal{O} \left( \rho^{\frac{2}{p}} \|\phi\| \right).$$

Applying Lemma B.1 and (13),

$$\int_{\Sigma} \rho^{2} \chi_{i} V e^{\sum_{h=1}^{m} P U_{h}} \phi h\left(\frac{1}{\tau_{i}\rho} y_{\xi_{i}}(x)\right) dv_{g}(x) 
= \int_{U_{2r_{0}}(\xi_{i})} \frac{8\tau_{i}^{2}\rho^{2} \chi_{i}}{\left(\tau_{i}^{2}\rho^{2} + |y_{\xi_{i}}|^{2}\right)^{2}} \exp\left\{-\log(8\tau_{i}^{2}) + \varrho\left(\xi_{i}\right) H\left(\xi_{i}, \xi_{i}\right) \right. 
\left. + \sum_{h \neq i} \varrho\left(\xi_{h}\right) G\left(\xi_{i}, \xi_{h}\right) + \log V\left(\xi_{i}\right) + \mathcal{O}\left(|y_{\xi_{i}}| + \rho^{1+\alpha_{0}}\right)\right\} \phi h\left(\frac{1}{\tau_{i}\rho} y_{\xi_{i}}\right) dv_{g} 
= \int_{B_{r_{0}}^{\xi_{i}}} e^{\hat{\varphi}_{i}(y)} \frac{8\tau_{i}^{2}\rho^{2}}{\left(\tau_{i}^{2}\rho^{2} + |y|^{2}\right)^{2}} \phi \circ y_{\xi_{i}}^{-1}(y) \chi\left(\frac{|y|}{r_{0}}\right) h\left(\frac{1}{\tau_{i}\rho} y\right) dy + o(1) 
= \int_{\mathbb{R}_{i}} \frac{8}{(1+|y|^{2})^{2}} \tilde{\phi}_{i}^{\rho} h(y) dy + o(1).$$

Then  $\tilde{\phi}_i^0$  is a distributional solution for the equation

(48) 
$$-\Delta U = \frac{8}{(1+|y|^2)^2} U \text{ in } \mathbb{R}_i \text{ with } \int_{\mathbb{R}^2} |\nabla U|^2 dy < \infty,$$

with boundary condition  $\partial_{\nu_0}U=0$  on  $\partial\mathbb{R}_i$ , where  $\nu_0$  is the unit outward normal of  $\partial\mathbb{R}_i$ . By the regularity theory,  $\tilde{\phi}_i^0$  is a smooth solution. It is well-known that any solutions to problem (48) are in the following form,  $\tilde{\phi}_i^0(y)=\frac{a_0^i(1-|y|^2)}{1+|y|^2}+\sum_{j=1}^{J_i}\frac{a_j^iy_j}{1+|y|^2}$ , where  $a_j^i\in\mathbb{R}$  for  $i=1,\cdots,m,j=0,\cdots,J_i$  (see Lemma D.1. of [16]).

Applying the result from Step 2.,

$$\frac{16}{\tau_{i}} \int_{\mathbb{R}_{i}} \frac{|y|^{2} - 1}{(|y|^{2} + 1)^{3}} \tilde{\phi}_{i}^{0}(y) dy = \lim_{\rho \to 0} \frac{16}{\tau_{i}} \int_{\Omega_{i}^{\rho}} \frac{|y|^{2} - 1}{(|y|^{2} + 1)^{3}} \phi_{i}^{\rho} \chi_{i}^{\rho} dy$$

$$= \lim_{\rho \to 0} 16 \tau_{i} \rho^{2} \int_{B_{2\tau_{0}}^{\xi_{i}}} \frac{|y|^{2} - \tau_{i}^{2} \rho^{2}}{(|y|^{2} + \tau_{i}^{2} \rho^{2})^{3}} \phi \circ y_{\xi_{i}}^{-1}(y) \chi\left(\frac{|y|}{r_{0}}\right) dy$$

$$= \lim_{\rho \to 0} \int_{B_{2\tau_{0}}^{\xi_{i}}} \rho^{2} e^{u_{\tau_{i},0}} \psi_{\tau_{i},0}^{0} \phi \circ y_{\xi_{i}}^{-1}(y) \chi(|y|) dy$$

$$= \lim_{\rho \to 0} \int_{\Sigma} \rho^{2} \chi_{i} e^{-\varphi_{i}} e^{U_{i}} \Psi_{i}^{0} \phi dv_{g}$$

$$= \lim_{\rho \to 0} \langle P\Psi_{i}^{0}, \phi \rangle = 0.$$

For any  $i = 1, \dots, m$  and  $j = 1, \dots, J_i$ ,

$$\begin{split} &\frac{32}{\tau_{i}\rho}\int_{\mathbb{R}_{i}}\frac{y_{j}}{(|y|^{2}+1)^{3}}\tilde{\phi}_{i}^{0}dy = \lim_{\rho\to 0}\frac{32}{\tau_{i}\rho}\int_{\Omega_{i}^{\rho}}\frac{y_{j}}{(|y|^{2}+1)^{3}}\phi_{i}^{\rho}\chi_{i}^{\rho}dy \\ &= \lim_{\rho\to 0}32\tau_{i}^{2}\rho^{2}\int_{B_{2r_{0}}^{\xi_{i}}}\frac{y_{j}}{(|y|^{2}+\tau_{i}^{2}\rho^{2})^{3}}\phi\circ y_{\xi_{i}}^{-1}(y)\chi\left(\frac{|y|}{r_{0}}\right)dy \\ &= \lim_{\rho\to 0}\int_{B_{2r_{0}}^{\xi_{i}}}\rho^{2}\chi\left(\frac{|y|}{r_{0}}\right)e^{u_{\tau_{i},0}}\psi_{\tau_{i},0}^{j}\phi\circ y_{\xi_{i}}^{-1}(y)dy \\ &= \lim_{\rho\to 0}\int_{U_{2r_{0}}(\xi_{i})}\rho^{2}\chi_{i}e^{U_{i}}e^{-\varphi_{i}}\Psi_{i}^{j}\phi(x)dv_{g} \\ &= \lim_{\rho\to 0}\langle P\Psi_{i}^{j},\phi\rangle = 0. \end{split}$$

Thus for any  $i=1,\dots,m, j=1,\dots,J_i$   $\int_{\mathbb{R}_i} \frac{|y|^2-1}{(|y|^2+1)^3} \tilde{\phi}_i^0 dy = \int_{\mathbb{R}_i} \frac{y_j}{(|y|^2+1)^3} \tilde{\phi}_i^0 dy = 0$ . It indicates that  $\tilde{\phi}_i^0 \equiv 0$ , which contradicts to (42).

 $\mathbf{E}$ 

The last part gives some technical lemmas to obtain the  $C^1$ -expansion of the reduced functional  $\tilde{E}_{\rho}$ .

**Lemma E.1.** As  $\rho \to 0$ , the following asymptotic expansions hold

$$\langle PU_i, PU_i \rangle = \varrho(\xi_i)(6\log 2 - 4\log \rho - 2\log(8\tau_i^2) + \varrho(\xi_i)H^g(\xi_i, \xi_i) - 2) + \mathcal{O}(\rho|\log \rho|),$$

and for any  $i \neq j \langle PU_i, \nabla PU_j \rangle = \varrho(\xi_i)\varrho(\xi_j)G^g(\xi_i, \xi_j) + \mathcal{O}(\rho)$ .

*Proof.* Applying Lemma B.1 and (13), as  $\rho \to 0$ 

$$\begin{split} \langle PU_i, PU_i \rangle &= \int_{\Sigma} |\nabla PU_i|_g^2 + \beta |PU_i|^2 dv_g = \rho^2 \int_{\Sigma} \chi_i e^{-\varphi_i} e^{U_i} PU_i dv_g \\ &= \int_{U_{r_0}(\xi_i)} \frac{8\tau_i^2 \rho^2}{(\tau_i^2 \rho^2 + |y_{\xi_i}(x)|^2)^2} e^{-\varphi_i} \left( \log \frac{1}{(\tau_i^2 \rho^2 + |y_{\xi_i}(x)|^2)^2} + \varrho(\xi_i) H^g(\xi_i, \xi_i) \right. \\ &\quad + \mathcal{O}(|y_{\xi_i}(x)| + \rho^{1+\alpha_0})) \, dv_g(x) + \mathcal{O}(\rho^2) \\ &= \int_{B_{r_0}^{\xi_i}} \frac{8\tau_i^2 \rho^2}{(\tau_i^2 \rho^2 + |y|^2)^2} \left( \log \frac{\tau_i^4 \rho^4}{(\tau_i^2 \rho^2 + |y|^2)^2} - 2\log(\tau_i^2 \rho^2) + \varrho(\xi_i) H^g(\xi_i, \xi_i) \right. \\ &\quad + \mathcal{O}(|y| + \rho^{1+\alpha_0})) \, dy + \mathcal{O}(\rho^2) \\ &= \varrho(\xi_i) (6 \log 2 - 4 \log \rho - 2 \log(8\tau_i^2) + \varrho(\xi_i) H^g(\xi_i, \xi_i) - 2) + \mathcal{O}(\rho |\log \rho|), \end{split}$$

where we applied the fact that for any r > 0, as  $\rho \to 0$ ,  $\int_{|y| < r} \frac{\rho^2}{(\rho^2 + |y|^2)^2} dy = \pi - \frac{\pi \rho^2}{r^2} + \frac{\pi \rho^4}{(r^2 + \rho^2)r^2}$  and  $\int_{|y| < r} \frac{\rho^2 \log(\frac{\rho^2 + |y|^2}{\rho^2})}{(\rho^2 + |y|^2)^2} dy = \pi + \frac{\pi \rho^2 \log(\rho^2)}{r^2} + \mathcal{O}(\rho^2)$ . For any  $i \neq j$ , by Lemma B.1,

$$\langle PU_{i}, PU_{j} \rangle = \rho^{2} \int_{\Sigma} \chi_{i} e^{-\varphi_{i}} e^{U_{i}} PU_{j} dv_{g}$$

$$= \int_{U_{2r_{0}}(\xi_{i})} \frac{8\tau_{i}^{2} \rho^{2}}{(\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}(x)|^{2})^{2}} e^{-\varphi_{i}(x)} (\varrho(\xi_{j}) G^{g}(\xi_{i}, \xi_{j}) + \mathcal{O}(|y_{\xi_{i}}(x)| + \rho^{1+\alpha_{0}}))$$

$$+ \mathcal{O}(\rho^{2} ||PU_{j}||)$$

$$= 8\varrho(\xi_{j}) G^{g}(\xi_{i}, \xi_{j}) \int_{B_{2r_{0}}^{\xi_{i}}} \frac{\tau_{i}^{2} \rho^{2}}{(\tau_{i}^{2} \rho^{2} + |y|^{2})^{2}} dy + \mathcal{O}(\rho)$$

$$= \varrho(\xi_{i}) \varrho(\xi_{j}) G^{g}(\xi_{i}, \xi_{j}) + \mathcal{O}(\rho)(\rho \to 0).$$

Lemma E.2.

(49) 
$$\rho^2 \int_{\Sigma} V e^{\sum_{i=1}^m PU_i} = \sum_{i=1}^m \varrho(\xi_i) + o(1)(\rho \to 0).$$

*Proof.* Applying Lemma B.1 and (13),

$$\begin{split} & \rho^2 \int_{\Sigma} V e^{\sum_{i=1}^m P U_i} dv_g \\ &= \sum_{i=1}^m \rho^2 \int_{U_{2r_0}(\xi_i)} e^{\chi_i U_i + \varrho(\xi_i) H^g(x,\xi_i) - \log 8\tau_i^2 + \sum_{j \neq i} \varrho(\xi_j) G^g(x,\xi_j) + \mathcal{O}(\rho^{1+\alpha_0})} dv_g(x) \\ &+ \mathcal{O}(\rho^2) \\ &= \sum_{i=1}^m \int_{U_{r_0}(\xi_i)} \frac{8\tau_i^2 \rho^2 e^{\varrho(\xi_i) H^g(\xi_i,\xi_i) - \log(8\tau_i^2) + \log V(\xi_i) + \sum_{j \neq i} \varrho(\xi_j) G^g(\xi_i,\xi_j)}{(\tau_i^2 \rho^2 + |y_\xi(x)|^2)^2} \\ & (1 + \mathcal{O}(|y_\xi(x)| + \rho^{1+\alpha_0})) dv_g(x) dv_g(x) + \mathcal{O}(\rho^2) \\ &= \sum_{i=1}^m \int_{B_{r_0}^{\xi_i}} \frac{8\tau_i^2 \rho^2 e^{\hat{\varphi}_i(y)}}{(\tau_i^2 \rho^2 + |y|^2)^2} (1 + \mathcal{O}(|y| + \rho^{1+\alpha_0})) + \mathcal{O}(\rho^2) \\ &= \sum_{i=1}^m \int_{\frac{1}{\tau_i \rho} B_{r_0}^{\xi_i}} (1 + \mathcal{O}(\rho|y|)) (1 + \mathcal{O}(\rho|y| + \rho^{1+\alpha_0})) \frac{8}{(1 + |y|^2)^2} dy \\ &+ \mathcal{O}(\rho^2) \\ &= \sum_{i=1}^m \varrho(\xi_i) + \mathcal{O}(\rho) (\rho \to 0). \end{split}$$

**Lemma E.3.** Let  $i, h = 1, 2, \dots, m$ . And if  $\xi_i \in \mathring{\Sigma}$ , j = 1, 2 and if  $\xi_i \in \partial \Sigma$ , j = 1. Then as  $\rho \to 0$ ,

$$\rho^{2} \int_{\Sigma} \chi_{h} e^{U_{h}} \partial_{(\xi_{i})_{j}} PU_{i} dv_{g}$$

$$= \frac{1}{2} \delta_{ih} \varrho(\xi_{i})^{2} \partial_{(\xi_{i})_{j}} H^{g}(\xi_{i}, \xi_{i}) + (1 - \delta_{ih}) \varrho(\xi_{i}) \varrho(\xi_{h}) \partial_{(\xi_{i})_{j}} G^{g}(\xi_{h}, \xi_{i}) + \mathcal{O}(\rho).$$

Proof.  $\rho^2 \int_{\Sigma} \chi_h e^{U_h} \partial_{(\xi_i)_j} PU_i = \rho^2 \left( \int_{\Sigma \cap U_{2r_0}(\xi_h)} + \int_{\Sigma \setminus U_{2r_0}(\xi_h)} \right) \chi_h e^{U_h} \partial_{(\xi_i)_j} PU_i$ . We observe that  $\int_{\Sigma \setminus U_{2r_0}(\xi_h)} \rho^2 \chi_h e^{U_h} \partial_{(\xi_i)_j} PU_i = 0$ . For  $h \neq i$ ,  $U_{2r_0}(\xi_h) \cap U_{2r_0}(\xi_i) = \emptyset$  by the choice of  $r_0$ . We notice that

$$\begin{aligned} \partial_{(\xi_{i})_{j}}|y_{\xi}(x)|^{2}|_{x=y_{\xi_{i}}^{-1}(y)} &= -2\langle (y_{\xi_{i}})_{*}\partial_{(\xi_{i})_{j}}y_{\xi}^{-1}(y), y \rangle \\ &= -2\langle (y_{\xi_{i}})_{*}\partial_{(\xi_{i})_{j}}(\xi_{i} + y + \mathcal{O}(|y|^{2})), y \rangle \\ &= -2y_{j} + \mathcal{O}(|y|^{3})(|y| \to 0). \end{aligned}$$

In this case, by Remark B.1

$$\int_{U_{2r_0}(\xi_h)} \rho^2 \chi_h e^{U_h} \partial_{(\xi_i)_j} P U_i dv_g = \int_{U_{2r_0}(\xi_h)} \rho^2 \chi_h e^{U_h} (\partial_{(\xi_i)_j} (\chi_i U_i) + \varrho(\xi_i) \partial_{(\xi_i)_j} H_{\xi_i}^g + \mathcal{O}(\rho^{\alpha_0}))$$

$$= \varrho(\xi_h) \varrho(\xi_i) \partial_{(\xi_i)_j} H^g(\xi_h, \xi_i) + o(1).$$

Claim. as  $\rho \to 0$ ,

$$\int_{U_{2\tau_0}(\xi_h)} \rho^2 \chi_i e^{U_i} \frac{2\partial_{(\xi_i)_j} |y_{\xi_i}(x)|^2}{\tau_i^2 \rho^2 + |y_{\xi_i}(x)|^2} dv_g(x) = \mathcal{O}(\rho^2) + \int_{U_{\tau_0}(\xi_h)} \rho^2 e^{U_i} \frac{4(-y_{\xi_i}(x)_j + \mathcal{O}(|y_{\xi_i}(x)|^3))}{\tau_i^2 \rho^2 + |y_{\xi_i}(x)|^2} dv_g(x) \\
= o(1).$$

Indeed, if  $\xi_i \in \mathring{\Sigma}$ , we have  $|e^{\hat{\varphi}_i(y)} - 1| = |\hat{\varphi}_{\xi_i}(y)| + \mathcal{O}(|\hat{\varphi}_{\xi_i}(y)|^2) = \mathcal{O}(|y|^2)$  as  $y \to 0$ .

$$\int_{U_{\tau_0}(\xi_i)\cap\Sigma} \rho^2 \chi_i e^{U_i} \frac{2\partial_{(\xi_i)_j} |y_{\xi_i}(x)|^2}{\tau_i^2 \rho^2 + |y_{\xi_i}(x)|^2} dv_g(x) = \int_{B_{\tau_0}^{\xi_i}} \rho^2 e^{\hat{\varphi}_{\xi_i}(y)} \frac{32\tau_i^2 \rho^2 (-y_j + \mathcal{O}(|y|^2))}{(\tau_i^2 \rho^2 + |y|^2)^3} dy + \mathcal{O}(\rho^2)$$

$$= \int_{B_{\tau_0}^{\xi_i}} \rho^2 (1 + \mathcal{O}(|y|^2)) \frac{-32\tau_i^2 y_j + \mathcal{O}(|y|^3)}{(\tau_i^2 \rho^2 + |y|^2)^3} dy$$

$$= \mathcal{O}(\rho).$$

So, the claim above is concluded. For  $i, h = 1, 2, \dots, m$  and  $j = 1, \dots, J_i$ . By Remark B.1,

$$\int_{\Sigma} \rho^{2} \chi_{i} e^{U_{i}} \partial_{(\xi_{i})_{j}} P U_{i} dv_{g}$$

$$= \int_{\Sigma} \frac{8\tau_{i}^{2} \rho^{2} \chi_{i}}{(\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}|^{2})^{2}} \left( \chi_{i} \frac{2\partial_{(\xi_{i})_{j}} |y_{\xi}|^{2}}{\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}|^{2}} + \varrho(\xi_{i}) \partial_{(\xi_{i})_{j}} H_{\xi_{i}}^{g} + \mathcal{O}(\rho^{\alpha_{0}}) \right) dv_{g}$$

$$= \int_{U_{r_{0}}(\xi_{i})} \rho^{2} \chi_{i}(x) e^{U_{i}(x)} \frac{2\partial_{(\xi_{i})_{j}} |y_{\xi}(x)|^{2}}{\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}(x)|^{2}} dv_{g}(x)$$

$$+ \frac{1}{2} \varrho(\xi_{i}) \partial_{(\xi_{i})_{j}} H^{g}(\xi_{i}, \xi_{i}) \int_{U_{r_{0}}(\xi_{i})} \frac{8\tau_{i}^{2} \rho^{2}}{(\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}(x)|^{2})^{2}} dv_{g}(x) + \mathcal{O}(\rho^{\alpha_{0}})$$

$$= \frac{1}{2} \varrho(\xi_{i})^{2} \partial_{(\xi_{i})_{j}} H^{g}(\xi_{i}, \xi_{i}) + o(1).$$

For  $i \neq h$ , via Lemma B.2,

$$\int_{U_{2r_{0}}(\xi_{h})\cap\Sigma} \rho^{2} \chi_{h} e^{U_{h}} \partial_{(\xi_{i})_{j}} P U_{i} dv_{g}$$

$$= \int_{U_{2r_{0}}(\xi_{h})\cap\Sigma} \frac{8\tau_{h}^{2} \rho^{2} \chi_{h}}{(\tau_{h}^{2} \rho^{2} + |y_{\xi_{h}}|^{2})^{2}} \left( \chi_{i} \frac{2\partial_{(\xi_{i})_{j}} |y_{\xi_{i}}|^{2}}{\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}|^{2}} + \varrho(\xi_{i}) \partial_{(\xi_{i})_{j}} H_{\xi_{i}}^{g} + \mathcal{O}(\rho^{\alpha_{0}}) \right) dv_{g}$$

$$= \int_{U_{2r_{0}}(\xi_{h})\cap\Sigma} \chi_{h}(x) \frac{8\tau_{h}^{2} \rho^{2}}{(\tau_{h}^{2} \rho^{2} + |y_{\xi_{h}}|^{2})^{2}} \left( \varrho(\xi_{i}) \partial_{(\xi_{i})_{j}} G^{g}(\cdot, \xi_{i}) + \mathcal{O}(\rho^{\alpha_{0}}) \right) dv_{g}$$

$$= \varrho(\xi_{i}) \varrho(\xi_{h}) \partial_{(\xi_{i})_{j}} G^{g}(\xi_{h}, \xi_{i}) + \mathcal{O}(\rho^{\alpha_{0}}).$$

Combining all the estimates above, Lemma E.3 is concluded.

**Lemma E.4.** Let i = 1, 2, ..., m and  $j = 1, ..., J_i$ . As  $\rho \to 0$ ,

$$\rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \partial_{(\xi_{i})_{j}} PU_{i} dv_{g}$$

$$= \partial_{(\xi_{i})_{j}} \left( \sum_{h=1}^{m} \varrho(\xi_{h})^{2} H^{g}(\xi_{h}, \xi_{h}) + \sum_{l \neq h} G^{g}(\xi_{h}, \xi_{l}) + \sum_{h=1}^{m} \varrho(\xi_{h}) \log V(\xi_{h}) \right) + o(1).$$

*Proof.* First, we divide the integral into three parts to calculate, i.e.,

$$\begin{split} & \rho^2 \int_{\Sigma} V e^{\sum_{h=1}^m P U_h} \partial_{(\xi_i)_j} P U_i dv_g \\ = & \rho^2 \left( \int_{\Sigma \setminus \cup_{h=1}^m U_{2r_0}(\xi_h)} + \int_{U_{2r_0}(\xi_i) \cap \Sigma} + \int_{\cup_{l \neq i} U_{2r_0}(\xi_l)} \right) V e^{\sum_{h=1}^m P U_h} \partial_{(\xi_i)_j} P U_i dv_g. \end{split}$$

By Remark B.1, as  $\rho \to 0$   $\int_{\Sigma \setminus \bigcup_{h=1}^{m} U_{2r_0}(\xi_h)} \rho^2 V e^{\sum_{h=1}^{m} P U_h} \partial_{(\xi_i)_j} P U_i dv_g$  $= \mathcal{O}\left(\rho^2 \int_{\Sigma \setminus \bigcup_{h=1}^{m} U_{2r_0}(\xi_h)} \left| \partial_{\xi_j} (\chi_{\xi} U_{\tau,\xi}) + \varrho(\xi) \partial_{\xi_j} H_{\xi}^g + \mathcal{O}(\rho^{\alpha_0}) \right| dv_g\right) = \mathcal{O}(\rho^2)$ . We observe that for any  $i = 1, 2, \cdots, m$  and  $j = 1, \cdots, J_i$ ,  $\partial_{(\xi_i)_j} H^g(\xi_i, \xi_i) = 2\partial_{x_j} H^g(x, \xi_i)|_{x=\xi_i}$ ,  $e^{\hat{\varphi}_i(y)} = 1 + \mathcal{O}(|y|^2)(|y| \to 0)$ , and  $\partial_{(\xi_i)_j} |y_{\xi}(x)|^2|_{x=y_{\xi_i}^{-1}(y)} = -2y_j + \mathcal{O}(|y|^3)(|y| \to 0)$ . Applying Lemma B.1 and Remark B.1 with (13),

$$\begin{split} &\int_{U_{2r_0}(\xi_i)} \rho^2 V e^{\sum_{h=1}^m PU_h} \partial_{(\xi_i)_j} PU_i dv_g \\ &= \int_{U_{2r_0}(\xi_i)} \left( \frac{\rho^2 V e^{\varrho(\xi_i) H_{\xi_i}^g + \sum_{l \neq i} \varrho(\xi_l) G^g(\cdot, \xi_l) + \mathcal{O}(\rho^{1+\alpha_0})}}{(\tau_i^2 \rho^2 + |y_{\xi_i}|^2)^2} \right) \left( \frac{2\chi_i \partial_{(\xi_i)_j} |y_{\xi_i}|^2}{(\tau_i^2 \rho^2 + |y_{\xi_i}|^2)} \right. \\ &\quad + \varrho(\xi_l) \partial_{(\xi_l)_j} H_{\xi_l}^g + \mathcal{O}(\rho^{\alpha_0}) \right) dv_g \\ &= \int_{B_{\tau_0}^{\xi_i}} \frac{8\tau_i^2 \rho^2 e^{\beta \xi_i (y)}}{(\tau_i^2 \rho^2 + |y|^2)^2} \exp\left\{ \varrho(\xi_i) H^g(y_{\xi_i}^{-1}(y), \xi_i) + \sum_{h \neq i} \varrho(\xi_h) G^g(y_{\xi_i}^{-1}(y), \xi_h) \right. \\ &\quad + \log V(y_{\xi_i}^{-1}(y)) - \log(8\tau_i^2) + \mathcal{O}(|y^2 + \rho^{1+\alpha_0}) \right\} \left( \frac{2\partial_{(\xi_l)_j} |y_{\xi_l}(x)|^2}{(\tau_i^2 \rho^2 + |y_{\xi_l}(x)|^2)} \Big|_{x=y_{\xi_l}^{-1}(y)} \right. \\ &\quad + \frac{1}{2} \varrho(\xi_i) \partial_{(\xi_l)_j} H^g(\xi_i, \xi_i) + \mathcal{O}(|y| + \rho^{\alpha_0}) \right) dy + \mathcal{O}(\rho^2) \\ &= \int_{\frac{1}{\tau_i \rho} B_{\tau_0}^{\xi_i}} \frac{8}{(1 + |y|^2)^2} (1 + \mathcal{O}(\rho^2 |y|^2)) \left( 1 + \frac{1}{2} \tau_i \rho \sum_{s=1}^2 \varrho(\xi_i) \partial_{(\xi_l)_s} H^g(\xi_i, \xi_i) y_s \right. \\ &\quad + \tau_i \rho \sum_{h \neq i} \varrho(\xi_h) \sum_{s=1}^2 \partial_{(\xi_i)_s} G^g(\xi_i, \xi_h) y_s + \tau_i \rho \sum_{s=1}^2 \partial_{(\xi_i)_s} \log V(\xi_i) y_s \\ &\quad + \mathcal{O}(\tau_i^2 \rho^2 |y|^2 + \rho^{1+\alpha_0}) \right) \cdot \left( \frac{-4y_j}{1 + |y|^2} + \frac{\varrho(\xi_i)}{2} \partial_{(\xi_l)_j} H^g(\xi_i, \xi_i) \right. \\ &\quad + \mathcal{O}(\rho^{\alpha_0} + \rho |y|) \right) dy + \mathcal{O}(\rho^2) \\ &= \frac{1}{2} \varrho(\xi_i)^2 \partial_{(\xi_l)_j} H^g(\xi_i, \xi_i) + \frac{1}{2} \varrho(\xi_i)^2 \partial_{(\xi_l)_j} H^g(\xi_i, \xi_i) \\ &\quad + \sum_{h \neq i} \varrho(\xi_l) \varrho(\xi_h) \partial_{(\xi_l)_j} G^g(\xi_i, \xi_h) + \varrho(\xi_l) \partial_{(\xi_l)_j} \log V(\xi_l) + o(1) \\ &= \varrho(\xi_l)^2 \partial_{(\xi_l)_j} H^g(\xi_i, \xi_i) + \sum_{h \neq i} \varrho(\xi_l) \varrho(\xi_h) \partial_{(\xi_l)_j} G^g(\xi_i, \xi_h) \\ &\quad + \varrho(\xi_l) \partial_{(\xi_l)_j} \log V(\xi_l) + o(1), \end{aligned}$$

where we applied  $\int_{\mathbb{R}^2} \frac{1}{(1+|y|^2)^2} dy = \pi = 2 \int_{\mathbb{R}^2} \frac{|y|^2}{(1+|y|^2)^3} dy$ .

For any  $h \neq i$ , analogue to the proof for h = i, we can obtain

$$\int_{U_{2r_0}(\xi_h)} \rho^2 V e^{\sum_{h=1}^m PU_h} \partial_{(\xi_i)_j} PU_i dv_g = \varrho(\xi_i) \varrho(\xi_h) \partial_{(\xi_i)_j} G^g(\xi_h, \xi_i) + o(1).$$

Combining the estimates above,

$$\rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \partial_{(\xi_{i})_{j}} PU_{i} dv_{g}$$

$$= \partial_{(\xi_{i})_{j}} \left( \sum_{h=1}^{m} \varrho(\xi_{h})^{2} H^{g}(\xi_{h}, \xi_{h}) + \sum_{l \neq h} \varrho(\xi_{h}) \varrho(\xi_{l}) G^{g}(\xi_{h}, \xi_{l}) + \sum_{h=1}^{m} \varrho(\xi_{h}) \log V(\xi_{h}) \right)$$

$$+o(1).$$

**Lemma E.5.** Let  $i, h = 1, \dots, m$ .. Then as  $\rho \to 0$ ,

$$\left| \rho^2 \chi_h e^{U_h} \left( \partial_{(\xi_i)_j} P U_i - \chi_i \partial_{(\xi_i)_j} U_i \right) \right|_{L^p(\Sigma)} \le O\left( \rho^{\frac{2(1-p)}{p}} \right).$$

*Proof.* By Remark B.1,  $\partial_{(\xi_i)_j} PU_i - \chi_i \partial_{(\xi_i)_j} U_i = \mathcal{O}(1)$ . Then applying Lemma B.2,

$$\left| \rho^2 \chi_h e^{U_h} \left( \partial_{(\xi_i)_j} P U_i - \chi_i \partial_{(\xi_i)_j} U_i \right) \right|_{L^p(\Sigma)} \le O \left( \left| \rho^2 \chi_h e^{U_h} \right|_{L^p(\Sigma)} \right) = O \left( \rho^{\frac{2(1-p)}{p}} \right).$$

**Lemma E.6.** Let  $\xi = (\xi_1, \dots, \xi_m) \in \mathcal{D}$ . Let  $\phi \in K_{\xi}^{\perp}$  and  $\|\phi\| \leq \mathcal{O}(\rho^{\frac{2-p}{p}} |\log \rho|)$ , where  $p \in (1, \frac{6}{5})$ . Then for  $i = 1, 2, \dots, m$ ,

(50) 
$$\left\langle \sum_{h=1}^{m} PU_h + \phi - i^*(\rho^2 V e^{\sum_{h=1}^{m} PU_h + \phi}), \partial_{(\xi_i)_j} PU_i \right\rangle = -\frac{1}{2} \frac{\partial \mathcal{F}}{\partial(\xi_i)_j}(\xi) + o(1)(\rho \to 0).$$

which is uniformly for  $\xi$  in any compact subset of  $\mathscr{D}$ .

*Proof.* For  $y = y_{\xi_i}(x) \ \partial_{(\xi_i)_j} |y_{\xi_i}(x)|^2 = -2y_j + \mathcal{O}(|y|^3)$ . Since  $\|\phi\| = o(1)$  and  $\langle P\Psi_j^i, \phi \rangle = 0$ , we have

$$\langle \phi, \partial_{(\xi_{i})_{j}} P U_{i} \rangle = \int_{\Sigma} \rho^{2} e^{-\varphi_{i}} e^{U_{i}} \phi \partial_{(\xi_{i})_{j}} \chi_{i} dv_{g} + \int_{\Sigma} \rho^{2} e^{-\varphi_{i}} \chi_{i} e^{U_{i}} \phi \partial_{(\xi_{i})_{j}} U_{i} dv_{g}$$

$$+ \int_{\Sigma} \rho^{2} \chi_{i} e^{U_{i}} \phi \partial_{(\xi_{i})_{j}} e^{-\varphi_{i}} dv_{g}$$

$$= \int_{\Sigma} \rho^{2} \chi_{i} e^{-\varphi_{i}} e^{U_{i}} \phi \Psi_{i}^{j} dv_{g} + \mathcal{O}\left(\int_{B_{2r_{0}}^{\xi_{i}}} \frac{\tau_{i}^{2} \rho^{2} \chi\left(\frac{|y|}{r_{0}}\right) \left(\tau_{i}^{2} \rho^{2} |y|^{2} + |y|^{4} + |y|^{3}\right)}{\left(\tau_{i}^{2} \rho^{2} + |y|^{2}\right)^{3}} |\phi| dv_{g}\right)$$

$$= \langle \phi, P \Psi_{i}^{j} \rangle + o(1) = o(1).$$

Considering that  $\int_{\Sigma} \partial_{(\xi_i)_j} PU_i dv_g = 0$  and  $\chi_i \chi_h \equiv 0$  for any  $i \neq h$ , we have

$$\left\langle \sum_{h=1}^{m} PU_{h} + \phi - i^{*}(\rho^{2}Ve^{\sum_{h=1}^{m} PU_{h} + \phi}), \partial_{(\xi_{i})_{j}}PU_{i} \right\rangle$$

$$= \sum_{h=1}^{m} \langle PU_{h}, \partial_{(\xi_{i})_{j}}PU_{i} \rangle + \langle \phi, \partial_{(\xi_{i})_{j}}PU_{i} \rangle - \rho^{2} \int_{\Sigma} Ve^{\sum_{h=1}^{m} PU_{h} + \phi} \partial_{(\xi_{i})_{j}}PU_{i}dv_{g}$$

$$\stackrel{(51)}{=} \sum_{h=1}^{m} \int_{\Sigma} \rho^{2} \chi_{h}e^{-\varphi_{h}}e^{U_{h}} \partial_{(\xi_{i})_{j}}PU_{i}dv_{g} - \rho^{2} \int_{\Sigma} Ve^{\sum_{h=1}^{m} PU_{h}} (e^{\phi} - \phi - 1) \partial_{(\xi_{i})_{j}}PU_{i}dv_{g}$$

$$-\rho^{2} \int_{\Sigma} Ve^{\sum_{h=1}^{m} PU_{h}} - \rho^{2} \int_{\Sigma} \chi_{i}^{2}e^{U_{i}} \phi \partial_{(\xi_{i})_{j}}U_{i}dv_{g}$$

$$-\sum_{h\neq i} \rho^{2} \int_{\Sigma} \chi_{h}e^{U_{h}} \phi \chi_{i} \partial_{(\xi_{i})_{j}}U_{i}dv_{g} - \rho^{2} \int_{\Sigma} Ve^{\sum_{h=1}^{m} PU_{h}} \partial_{(\xi_{i})_{j}}PU_{i}dv_{g} + o(1).$$

Let  $q \ge 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$  and for any r > 1. By Lemma B.4 and Lemma C.2,

$$\left| \rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} (e^{\phi} - \phi - 1) \partial_{(\xi_{i})_{j}} PU_{i} dv_{g} \right|$$

$$\leq \left| \rho^{2} h e^{\sum_{h=1}^{m} PU_{h}} (e^{\phi} - \phi - 1) \right|_{L^{p}(\Sigma)} \left| \partial_{(\xi_{i})_{j}} PU_{i} \right|_{L^{q}(\Sigma)}$$

$$\leq c \|\phi\|^{2} \rho^{\frac{2-2pr}{pr}} \left| \partial_{(\xi_{i})_{j}} PU_{i} \right|_{L^{q}(\Sigma)}$$

$$\leq c \|\phi\|^{2} \rho^{\frac{2-3pr}{pr}}.$$

By Lemma E.5,

$$\left| \rho^2 \int_{\Sigma} \sum_{h=1}^m \chi_h e^{U_h} \phi(\chi_i \partial_{(\xi_i)_j} U_i - \partial_{(\xi_i)_j} P U_i) dv_g \right|$$

$$\leq c \sum_{h=1}^m \|\phi\| |\rho^2 \chi_h e^{U_h} (\chi_i \partial_{(\xi_i)_j} U_i - \partial_{(\xi_i)_j} P U_i)|_{L^p(\Sigma)}$$

$$\leq c \|\phi\| \rho^{\frac{2(1-p)}{p}}.$$

By Lemma C.1,

$$\left| \rho^{2} \int_{\Sigma} \left( \sum_{h=1}^{m} \chi_{h} e^{U_{h}} - V e^{\sum_{h=1}^{m} P U_{h}} \right) \rho^{2} \phi \partial_{(\xi_{i})_{j}} P U_{i} \right|$$

$$\leq c \|\phi\| \left| \sum_{h=1}^{m} \chi_{h} e^{U_{h}} - \rho^{2} V e^{\sum_{h=1}^{m} P U_{h}} \right|_{L^{p}} \|\partial_{(\xi_{i})_{j}} P U_{i}\|$$

$$\leq c \|\phi\| \rho^{\frac{2-p}{p}} \frac{1}{\rho} = c \|\phi\| \rho^{\frac{2(1-p)}{p}}.$$

In view of  $\partial_{(\xi_i)_i} |y_{\xi_i}(x)|^2 = -2y_{\xi_i}(x)_i + \mathcal{O}(|y_{\xi_i}(x)|^3)$  as  $x \to \xi_i$ , as  $\rho \to 0$ 

$$\rho^{2} \int_{\Sigma} \chi_{i} e^{U_{i}} \phi \chi_{i} \partial_{(\xi_{i})_{j}} U_{i} dv_{g}$$

$$= \rho^{2} \int_{\Sigma} e^{U_{i}} \phi \chi_{i} e^{-\varphi_{i}} (1 + \mathcal{O}(|y_{\xi_{i}}|^{2})) \left( P \Psi_{i}^{j} + \mathcal{O}\left(\frac{|y_{\xi_{i}}|^{3}}{\tau_{i}^{2} \rho^{2} + |y_{\xi_{i}}|^{2}} \right) \right) dv_{g} + \mathcal{O}(\rho^{2})$$

$$= \langle \phi, P \Psi_{j}^{i} \rangle + \mathcal{O}(\rho) = o(1).$$

On the other hand, applying Lemma E.3 and Lemma E.4,

$$\sum_{h=1}^{m} \rho^{2} \int_{\Sigma} \chi_{h} e^{U_{h}} \partial_{(\xi_{i})_{j}} PU_{i} - \rho^{2} \int_{\Sigma} V e^{\sum_{h=1}^{m} PU_{h}} \partial_{(\xi_{i})_{j}} PU_{i} = -\frac{1}{2} \partial_{(\xi_{i})_{j}} \mathcal{F}_{k,m}(\xi) + o(1).$$

For any  $p \in (1, \frac{6}{5})$ , take r > 1 sufficiently close to 1 such that  $2(\frac{2-p}{p}) + \frac{2-3pr}{pr} > 0$ . Thus  $\rho \to 0$ 

$$\left\langle \sum_{h=1}^{m} PU_{h} + \phi - i^{*}(\rho^{2}Ve^{\sum_{h=1}^{m} PU_{h} + \phi}), \partial_{(\xi_{i})_{j}} PU_{i} \right\rangle = -\frac{1}{2}\partial_{(\xi_{i})_{j}} \mathcal{F}_{k,m}(\xi) + o(1).$$

## References

- [1] Oscar Agudelo and Angela Pistoia. Boundary concentration phenomena for the higher-dimensional Keller-Segel system. *Calc. Var.*, 55:132, 2016.
- [2] Daniele Bartolucci, Changfeng Gui, Yeyao Hu, Aleks Jevnikar, and Wen Yang. Mean field equations on tori: existence and uniqueness of evenly symmetric blow-up solutions. *Discrete Contin. Dyn. Syst.*, 40(6):3093–3116, 2020.
- [3] Luca Battaglia. A general existence result for stationary solutions to the Keller-Segel system. *Discrete Contin. Dyn. Syst.*, 39(2):905–926, 2019.
- [4] Piotr Biler. Local and global solvability of some parabolic system modeling chemotaxis. *Adv. Math. Sci. Appl.*, 8:715–743, 1998.
- [5] Luis A. Caffarelli and Yisong Yang. Vortex condensation in the Chern-Simons Higgs model: an existence theorem. *Commun. Math. Phys.*, 168(2):321–336, 1995.
- [6] Emanuele Caglioti, Pierre-Louis Lions, Carlo Marchioro, and Mario Pulvirenti. A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. *Commun. Math. Phys.*, 143:501–525, 1992.

- [7] Emanuele Caglioti, Pierre-Louis Lions, Carlo Marchioro, and Mario Pulvirenti. A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description ii. *Commun. Math. Phys.*, 174:229–260, 1995.
- [8] Sun-Yung Alice Chang, Matthew Gursky, and Paul Yang. The scalar curvature equation on 2- and 3-spheres. Calc. Var. Partial Differ. Equ., 1:205–229, 1993.
- [9] Sun-Yung Alice Chang and Paul Yang. Prescribing Gaussian curvature on  $S^2$ . Acta Math., 159:215–259, 1987.
- [10] Sun-Yung Alice Chang and Paul C. Yang. Conformal deformation of metrics on  $S^2$ . J. Differ. Geom., 27(2):259–296, 1988.
- [11] Shiing-shen Chern. An elementary proof of the existence of isothermal parameters on a surface. *Proc. Amer. Math. Soc.*, 6:771–782, 1955.
- [12] Chang Kung Ching and Liujia Quan. On Nirenberg's Problem. *Int. J. Math.*, 4(1):35–58, 1993.
- [13] Manuel del Pino, Michal Kowalczyk, and Monica Musso. Singular limits in Liouville-type equations. Calc. Var. Partial Differ. Equ., 24(1):47–81, 2005.
- [14] Manuel del Pino, Angela Pistoia, and Giusi Vaira. Large mass boundary condensation patterns in the stationary Keller-Segel system. *J. Differential Equations*, 261:3414–3462, 2016.
- [15] Pierpaolo Esposito and Pablo Figueroa. Singular mean field equations on compact Riemann surfaces. *Nonlinear Anal.*, 111:33–65, 2014.
- [16] Pierpaolo Esposito, Massimo Grossi, and Angela Pistoia. On the existence of blowing-up solutions for a mean field equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 22(2):227–257, 2005.
- [17] Pablo Figueroa. Bubbling solutions for mean field equations with variable intensities on compact Riemann surfaces. *JAMA*, 152(2):507–555, 2024.
- [18] Herbert Gajewski, Klaus Zacharias, and Konrad Gröger. Global behaviour of a reactiondiffusion system modelling chemotaxis. Math. Nachr., 195(1):77–114, 1998.
- [19] David Gilbarg and Neil S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Berlin, Heidelberg, 2 edition, 2001.

- [20] Zhengni Hu, Thomas Bartsch, and Mohameden Ahmedou. Blow-up solutions for mean field equations with neumann boundary conditions on Riemann surfaces, 2024. Preprint on arXiv.
- [21] Willi Jäger and Stephan Luckhaus. On explosions of solutions to a system of partial differential equations modelling chemotaxis. *Trans. Am. Math. Soc.*, 329(2):819–824, 1992.
- [22] Jerry Kazdan and Frank Warner. Curvature functions for compact 2-manifolds. *Ann. Math.*, 99:14–47, 1974.
- [23] Evelyn F. Keller and Lee A. Segel. Initiation of slime mold aggregation viewed as an instability. *J. Theor. Biol.*, 26(3):399–415, 1970.
- [24] Michael K.-H. Kiessling. Statistical mechanics of classical particles with logarithmic interactions. *Commun. Pure Appl. Math.*, 46:27–56, 1993.
- [25] Yanyan Li. On a singularly perturbed elliptic equation. Adv. Differ. Equ., 2:955–980, 1997.
- [26] Giacomo Nardi. Schauder estimate for solutions of poisson's equation with Neumann boundary condition. L'Enseign. Math., 60(3):421–435, 2015.
- [27] Margarida Nolasco and Gabriella Tarantello. Double vortex condensates in the Chern-Simons-Higgs theory. Calc. Var. Partial Differ. Equ., 9:31–94, 1999.
- [28] Manuel del Pino and Juncheng Wei. Collapsing steady states of the Keller-Segel system. Nonlinearity, 19(3):661–684, 2006.
- [29] Angela Pistoia and Giusi Vaira. Steady states with unbounded mass of the Keller-Segel system. *Proc. Roy. Soc. Edinburgh Sect. A*, 145(1):203–222, 2015.
- [30] Renate Schaaf. Stationary solutions of chemotaxis systems. Trans. Amer. Math. Soc., 292:531–556, 1985.
- [31] Takasi Senba and Takashi Suzuki. Some structures of the solution set for a stationary system of chemotaxis. Adv. Math. Sci. Appl., 10:191–224, 2000.
- [32] Gabriella Tarantello. Multiple condensate solutions for the Chern-Simons Higgs theory. J. Math. Phys., 37(8):3769–3796, 1996.
- [33] Gabriella Tarantello. Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. *Discrete Contin. Dyn. Syst.*, 28:31–973, 2010.

- [34] Guofang Wang and Juncheng Wei. Steady state solutions of a reaction-diffusion system modeling chemotaxis. *Math. Nachr.*, 233(1):221–236, 2002.
- [35] Yisong Yang. Solitons in Field Theory and Nonlinear Analysis. Springer, Berlin, 2001.
- [36] Yunyan Yang. Extremal functions for Moser-Trudinger inequalities on 2-dimensional compact Riemannian manifolds with boundary. *Int. J. Math.*, 17(3):313–330, 2006.
- [37] Yunyan Yang and Jie Zhou. Blow-up analysis involving isothermal coordinates on the boundary of compact Riemann surface. *J. Math. Anal. Appl.*, 504(2):125440, 2021.

Thomas Bartsch, Mohameden Ahmedou, Zhengni Hu Mathematisches Institut Universität Giessen Arndtstr. 2 35392 Giessen, Germany Thomas.Bartsch@math.uni-giessen.de Mohameden.Ahmedou@math.uni-giessen.de Zhengni.Hu@math.uni-giessen.de