# Anti-seizure medication load is not correlated with early termination of seizure spread

Nathan Evans<sup>1\*</sup>, Sarah J. Gascoigne<sup>1</sup>, Guillermo M. Besné<sup>1</sup>, Chris Thornton<sup>1</sup>,

Gabrielle M. Schroeder<sup>1</sup>, Fahmida A Chowdhury<sup>3</sup>, Beate Diehl<sup>3</sup>, John S Duncan<sup>3</sup>,

Andrew W McEvoy<sup>3</sup>, Anna Miserocchi<sup>3</sup>, Jane de Tisi<sup>3</sup>,

Peter N. Taylor<sup>1,2,3</sup>, Yujiang Wang<sup>1,2,3</sup>

September 4, 2024

1. CNNP Lab (www.cnnp-lab.com), Interdisciplinary Computing and Complex BioSystems Group, School of Computing, Newcastle University, Newcastle upon Tyne, United Kingdom

2. Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom

3. UCL Queen Square Institute of Neurology, Queen Square, London, United Kingdom

\* n.evans5@newcastle.ac.uk

### 1 Abstract

Anti-seizure medications (ASMs) are the mainstay of treatment for epilepsy, yet their effect on seizure spread is not fully understood. Higher ASM doses have been associated with shorter and less severe seizures. Our objective was to test if this effect was due to limiting seizure spread through early termination of otherwise unchanged seizures.

We retrospectively examined intracranial EEG (iEEG) recordings in 15 subjects that underwent ASM tapering during pre-surgical monitoring. We estimated ASM plasma concentrations based on pharmaco-kinetic modelling. In each subject, we identified seizures that followed the same onset and initial spread patterns, but some seizures terminated early (truncated seizures), and other seizures continued to spread (continuing seizures). We compared ASM concentrations at the times of truncated seizures and continuing seizures.

We found no substantial difference between ASM concentrations when truncated vs. continuing seizures occurred (Mean difference = 4%, sd = 29%, p=0.6).

Our results indicate that ASM did not appear to halt established seizures in this cohort. Further research is needed to understand how ASM may modulate seizure duration and severity.

## 2 Introduction

Anti-seizure medications (ASMs) are the most common treatment for epilepsy; however, the effect of ASMs on seizure onset and spread patterns is not fully understood. Analysis of ictal intracranial EEG (iEEG) has shown an individual-level tendency for seizures to follow specific spread patterns (Karoly et al., 2018; Schroeder et al., 2020). However not every seizure will spread in the same way; instead, there are variations in the pattern. Further, some subjects have truncated seizures that share the same onset and initial spread as other continuing seizures, but terminate early (Schroeder et al., 2022; Wagner et al., 2015). It was our objective to test if ASMs limit seizure spread through early termination of otherwise unchanged seizures.

Tapering ASMs during iEEG implantation is a common approach to provoke seizures during presurgical monitoring to help identify seizure onset regions (Consales et al., 2021). While medication tapering is performed for clinical and monitoring purposes, it can also provide unique insights into seizure dynamics (Mukamel and Fried, 2012). We hypothesised that truncated seizures are more likely to occur in the presence of higher doses of ASMs, and that continuing seizures are associated with lower doses. To test this hypothesis, we retrospectively analysed data collected from individuals who were undergoing the medication tapering procedure during iEEG monitoring.

### 3 Methods

#### 3.1 Study design

We retrospectively analysed iEEG data from subjects undergoing presurgical evaluation for medically refractory focal epilepsy at the National Hospital for Neurology and Neuroscience. Data were collected for clinical purposes and independent of this study. Our retrospective and anonymised data were obtained from the National Epilepsy & Neurology Database (NEN), composed of individuals with focal epilepsy from the National Hospital for Neurology and Neurosurgery, and analysed with the approval of Newcastle University Ethics Committee (27742/2022).

Subjects were included if they had undergone iEEG, had subsequent epilepsy surgery, associated pre-op and post-op T1w imaging information, and iEEG implantation information between 2008

and 2019. Subjects were additionally required to have had ASM tapering with available data on the time and dose of each medication throughout their time in the EMU. We also required that subjects had at least two visually confirmed seizures that had activity captured by our automatic detection algorithms, discussed in Section 3.4. Finally, we matched pairs of truncated and continuing seizures within subjects, and only those subjects with at least one pair of seizures were retained for our analysis. Fifteen subjects met all inclusion criteria.

#### 3.2 Subject metadata

	All	Tapered	Tapered with matched pairs
count	69	42	15
Age	32(15.4)	31(14.0)	32(13.8)
Sex (M/F)	33/36	20/22	7/8
Disease duration	20(15.0)	17.5(14.0)	20(13.5)
TLE/eTLE	31/38	27/15	9/6

Table 1 shows the demographic information for the subjects included in this study.

**Table 1: Metadata of subjects.** Clinical metadata and demographics split by whether ASM tapering was performed during epilepsy monitoring. Data is presented in the following format:

- Count: number of subjects.
- Age (yrs): subject age in years. median(interquartile range).
- Sex: subject sex. male/female
- Disease duration (yrs): Time between epilepsy diagnosis and recording in years. median(interquartile range)
- Diagnosis: Temporal Lobe Epilepsy/extratemporal Lobe Epilepsy.

## 3.3 Anti-seizure medication tapering and blood plasma concentration estimation

For each subject that underwent ASM tapering, we modelled their ASM plasma concentrations to estimate the relative ASM levels at the time of each seizure event.

Each medication dose taken by each subject was recorded during the initial monitoring. A continuous estimation of the plasma concentration for each medication was generated from the intake information using known pharmaco-kinetics. We modelled first order absorption and elimination using equation 1. The parameters used were specific to each medication, and the justification for each parameter used is explained in more detail in Besne et al. (2024).

We applied a 24 hour rolling average to compute plasma concentrations to account for inaccuracies in recorded intake time. For each ASM, the level was normalised as a ratio of the maximum level, ranging from 1 (maximum ASM concentration during the recording) to 0 for all medication types. A single combined ASM load was calculated by adding the ASM loads together for each

$$C = \frac{F * D * Ka}{Vd * (Ka - Ke)} * (e^{-\mathrm{Ke}^{*}\mathrm{t}} - e^{-\mathrm{Ka}^{*}\mathrm{t}})$$
(1)

**Equation 1:** Pharmaco-kinetic equation for plasma concentration of a single dose oral intake with firstorder absorption and elimination. C, plasma concentration. F, bio-availability. D, dose at intake. Ka, absorption constant. Ke, elimination constant. t, time since intake. Vd, volume distribution.

time point, then dividing by the initial pre-tapering load, so each subject starts with a combined ASM load of 1.

#### 3.4 EEG preprocessing and ictal activity detection

Intracranial EEG (iEEG) was recorded as part of the presurgical workup. Seizure timings were extracted from clinical reports and visually confirmed. For each seizure, iEEG segments containing ictal and 120 seconds of pre-ictal activity were extracted. Prior to analysis, all data were resampled to 512 Hz. An iterative noise detection algorithm was used to identify pre-ictal noise, which was confirmed by visual inspection. On a within-subject basis, channels identified as noisy were removed from all seizures. When seizures occurred in close succession (seizure cluster), we retained only the lead seizure. Recordings were re-referenced to a common average reference, notch filtered at 50 Hz (and harmonics up to 200 Hz) with a 2 Hz window to remove line noise. Recordings were then band-pass filtered between 0.5 and 200 Hz. All filters used a fourth-order, zero-phase shift Butterworth filter.

Our method for detecting ictal activity was an extension of the previously-described seizure "imprint" algorithm (Gascoigne et al., 2024). Briefly, we first computed eight EEG signal features: energy, line length, and band power in 6 distinct bands ( $\delta, \theta, \alpha, \beta$ , low- $\gamma$ , and high- $\gamma$ ) in one second windows with a 7/8th second overlap. The distribution in all features across all pre-ictal windows was then used to calculate the level of abnormality for each window in the ictal segments. By thresholding the abnormality, this approach produces a binary "imprint" of the seizure showing where seizure activity (abnormal activity compared to pre-ictal) has been detected at each time point in each channel (Fig. 1 A). The details of the algorithm are documented in code form for one example seizure on Github.

#### 3.5 Matching early terminating seizures with continuing seizures

We aimed to match truncated seizures with continuing seizures that share the same onset and initial propagation patterns (Fig. 1A for an example pair). As seizures do not always propagate at the same rate (Michael Wenzel et al., 2017; Schroeder et al., 2020, 2022), we used dynamic time warping (DTW) to compares sequences, in this case imprints, without penalising differences in the speed of progression. Two versions of this algorithm were used, open ended DTW (OE-DTW) and classical DTW. Classical DTW compares the entirety of both imprints, and OE-DTW allows for one imprint to be cut short if that provides a better match (Fig. 1B). We defined a truncated-continuing pair as one in which the OE-DTW algorithm improved on the classical DTW algorithm by at least 20%.

#### 3.6 Statistical analyses

To test if ASM tapering during an iEEG implantation made pairs of truncated/continuing seizures more likely to occur, we compared the percentage of subjects that had truncated pairs detected in the tapered and non-tapered groups. A  $\chi^2$  test was performed to determine if there was a significant difference between these two groups.

To determine if there were any differences between ASM concentrations at the time of truncated/continuing seizures across subjects, we estimated the ASM concentration of the time of each seizure (Fig. 2A). Then, for every truncated/continuing seizure pair, we calculated the difference between the ASM concentrations at the time that those seizures occurred (Fig. 2B&C). For each subject, we took the median of these ASM differences for each pair detected, creating a distribution across the whole cohort (Fig. 2D). We used a paired sample t-test to determine if there was a significant difference in ASM dose between the truncated and continuing seizures.

#### 3.7 Code and data availability

All analysis was performed in MATLAB version R2023B. Anonymised imprint and ASM intake schedule, along with analysis code will be available on Github: https://github.com/NathanMEvans/Temporal-Modulation-truncation upon acceptance.



Figure 1: Truncated-continuing pair matching. A) Examples of continuing (top) and truncated (bottom) seizures from the same subject. Each EEG signal (left) is pre-processed, features extracted and then thresholded to generate a binary imprint (right). This imprint is then trimmed to the times of minimum and maximum spread (shown with black dotted lines). B) Simplified illustration of how the open ended dynamic time warping (OE-DTW) algorithm compares the trimmed imprints from section A. For each time point in each imprint, the minimum cumulative distance from the start of both seizures is found. The OE-DTW algorithm (red) finds a minimum distance using every time point in imprint A, and any amount of imprint B. The proportion of imprint B used for this comparison is marked with a dotted red line. The DTW algorithm (blue) must include the whole of both imprints. C) This process is repeated, comparing every seizure recorded from this subject to every other. This produces a matrix of distances for both the DTW and OE-DTW algorithms. D) For each seizure pair the % improvement is above the threshold of 20%, the seizure pair is labelled a truncated-continuing pair, shown with a black outline. For this example subject, the 13<sup>th</sup> seizure was matched as a truncated seizure to the 1<sup>st</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 7<sup>th</sup>, 10<sup>th</sup> and 12<sup>th</sup> seizures, with the OE-DTW algorithm providing no improvement for the vast majority of other seizure pairs.

## 4 Results

## 4.1 Truncation detection no more likely to occur in subjects undergoing medication tapering

Of the 42 subjects with ASM tapering, 15 (36%) had truncated pairs detected, whilst 7 (26%) of 27 subjects that did not undergo tapering had truncated pairs detected. A  $\chi^2$  test of independence showed that there was no significant association between tapering and truncation pair occurrence  $(\chi^2_{1,N=69} = 0.725, p = 0.40)$ , suggesting that factors other than ASM load affect the extent of seizure propagation.

## 4.2 Higher medication load is not associated with occurrence of truncated seizures

We calculated the difference between the ASM load at the time of truncated and continuing seizures (Fig. 2). For each subject we computed a median difference to create a distribution across subjects (mean = 0.04, SD = 0.29). Using a paired-sample t-test, we found no evidence that the mean of the distribution differed from zero (t(14) = 0.54, p = 0.60), therefore we have no evidence of a difference in ASM level when truncated *vs.* continuing seizures occurred.



**Figure 2: Drug load difference across subjects.** A) Modelled Combined ASM load plotted against time for an example subject (green). Seizures (red) shown at the time and ASM load that they occurred at. B) Difference between ASM load for each seizure when compared to each other seizure. Each black square shows where seizure X(row) was found to be a truncated version of seizure Y(row). C) Difference between truncated and continuing ASM load for each matched pair. D) Distribution of median differences of ASM load across all subjects. The mean of the distribution is marked with a pink line.

## 5 Discussion

We investigated the associations between ASM plasma concentration and seizure spread dynamics. Specifically, we compared ASM loads to the occurrence of truncated or continuing seizures, and we found no evidence of any consistent association. Furthermore, the proportion of subjects with drug tapering who had truncated pairs detected was not significantly different from those without tapering. These findings did not support our initial hypothesis, and suggest that the association seen between medication load and seizure severity is not driven by early seizure termination, but by another mechanism.

Considering seizure spread patterns, shorter seizure durations could be a result of a seizure progressing faster while following the same spread patterns, or following a separate spread pattern that takes less time to progress from onset to termination. Different patterns could also be less severe in the extent of their spread. Previous studies (Marciani and Gotman, 1986; Spencer et al., 1981)found that medication withdrawal does not affect the onset localisation for surgery in most cases, but in individuals in whom medication did affect onsets (Engel and Crandall, 1983; Napolitano and Orriols, 2013). A better understanding of how and when this occurs could be helpful when estimating the true epileptogenic zone.

Other unestablished factors could promote the occurrence of truncated seizures. Previous work (Donos et al., 2018) has shown that clinical seizure type can be partially predicted from the initial dynamics of the seizure onset. It may be possible to use a similar approach to predict when a seizure will be truncated.

Increasing ASM doses is the current first-line approach for controlling seizures, but many people already take the highest tolerable dose. Therefore, other approaches, such as neuromodulation with vagus nerve stimulation may be helpful in some individuals (Hampel et al., 2015). The mechanisms of action is not fully understood, but are likely to involve blocking propagation in epileptic networks (Ghosh et al., 2023). It is possible that neuromodulation promotes early termination of seizures in a similar manner to the seemingly naturally occurring truncated seizures observed in this study.

iEEG implantations give a unique insight into brain states and seizure dynamics that could be leveraged to understand many treatments. However, iEEG data obtained from those undergoing presurgical evaluation may not be representative of the wider epilepsy population. Less invasive modalities, such as scalp EEG and magnetoencephalography (MEG), may provide more generalisable insights into the relationship between ASMs and seizure severity. An additional challenge when investigating the impact of ASMs on seizure dynamics is disentangling the impact of changing ASM doses from other time varying factors that are known to impact seizure dynamics such as fatigue, anxiety and disturbance of the sleep-wake cycle (Philippa J Karoly et al., 2021; Schroeder et al., 2020). These external factors make identifying the impact of tapering ASM challenging, especially when the duration of iEEG studies is relatively short.

In conclusion, more research into the effect of ASMs on seizure spread patterns is needed. Investigating truncated seizures recorded from pre-surgical iEEG may improve our understanding of the mechanisms modulating seizure termination, and aid in the conception of new therapeutic approaches.

### 6 Acknowledgements

We thank members of the Computational Neurology, Neuroscience & Psychiatry Lab (www.cnnplab.com) for discussions on the analysis and manuscript; P.N.T. and Y.W. are both supported by UKRI Future Leaders Fellowships (MR/T04294X/1, MR/V026569/1). JSD, JdT are supported by the NIHR UCLH/UCL Biomedical Research Centre.

## References

- The national epilepsy and neurology database. https://www.hra.nhs.uk/ planning-and-improving-research/application-summaries/research-summaries/ national-epilepsy-neurology-database/. Accessed: 2024-02-07.
- Guillermo M. Besne, Nathan Evans, Mariella Panagiotopoulou, Billy Smith, Fahmida A. Chowdhury, Beate Diehl, John S. Duncan, Andrew W. McEvoy, Anna Miserocchi, Jane de Tisi, Mathew Walker, Peter N. Taylor, Chris Thornton, and Yujiang Wang. Anti-seizure medication tapering is associated with delta band power reduction in a dose, region and time-dependent manner, May 2024. URL http://arxiv.org/abs/2405.01385. arXiv:2405.01385 [q-bio].
- Alessandro Consales, Sara Casciato, Sofia Asioli, Carmen Barba, Massimo Caulo, Gabriella Colicchio, Massimo Cossu, Luca de Palma, Alessandra Morano, Giampaolo Vatti, Flavio Villani, Nelia Zamponi, Laura Tassi, Giancarlo Di Gennaro, and Carlo Efisio Marras. The surgical treatment of epilepsy. *Neurological Sciences*, 42(6):2249–2260, June 2021. ISSN 1590-3478. doi: 10.1007/s10072-021-05198-y. URL https://doi.org/10.1007/s10072-021-05198-y.
- Cristian Donos, Mihai Dragos Maliia, Matthias Dümpelmann, and Andreas Schulze-Bonhage. Seizure onset predicts its type. *Epilepsia*, 59(3):650–660, 2018. ISSN 1528-1167. doi: 10.1111/ epi.13997. URL https://onlinelibrary.wiley.com/doi/abs/10.1111/epi.13997. \_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/epi.13997.
- J. Engel and P. H. Crandall. Falsely localizing ictal onsets with depth EEG telemetry during anticonvulsant withdrawal. *Epilepsia*, 24(3):344–355, June 1983. ISSN 0013-9580. doi: 10.1111/ j.1528-1157.1983.tb04898.x.
- Sarah J. Gascoigne, Nathan Evans, Gerard Hall, Csaba Kozma, Mariella Panagiotopoulou, Gabrielle M. Schroeder, Callum Simpson, Christopher Thornton, Frances Turner, Heather Woodhouse, Jess Blickwedel, Fahmida A. Chowdhury, Beate Diehl, John S. Duncan, Ryan Faulder, Rhys H. Thomas, Kevin Wilson, Peter N. Taylor, and Yujiang Wang. Incomplete resection of the intracranial electroencephalographic seizure onset zone is not associ-

ated with postsurgical outcomes. *Epilepsia*, n/a(n/a), July 2024. ISSN 1528-1167. doi: 10.1111/epi.18061. URL https://onlinelibrary.wiley.com/doi/abs/10.1111/epi.18061. \_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/epi.18061.

- Shampa Ghosh, Jitendra Kumar Sinha, Soumya Ghosh, Hitaishi Sharma, Rakesh Bhaskar, and Kannan Badri Narayanan. A Comprehensive Review of Emerging Trends and Innovative Therapies in Epilepsy Management. *Brain Sciences*, 13(9):1305, September 2023. ISSN 2076-3425. doi: 10.3390/brainsci13091305. URL https://www.ncbi.nlm.nih.gov/pmc/articles/ PMC10527076/.
- Kevin G. Hampel, Hartmut Vatter, Christian E. Elger, and Rainer Surges. Cardiac-based vagus nerve stimulation reduced seizure duration in a patient with refractory epilepsy. *Seizure*, 26: 81–85, March 2015. ISSN 1532-2688. doi: 10.1016/j.seizure.2015.02.004.
- Philippa J. Karoly, Levin Kuhlmann, Daniel Soudry, David B. Grayden, Mark J. Cook, and Dean R. Freestone. Seizure pathways: A model-based investigation. *PLOS Computational Biology*, 14(10):e1006403, October 2018. ISSN 1553-7358. doi: 10.1371/journal.pcbi. 1006403. URL https://journals.plos.org/ploscompbiol/article?id=10.1371/journal. pcbi.1006403. Publisher: Public Library of Science.
- M. G. Marciani and J. Gotman. Effects of drug withdrawal on location of seizure onset. *Epilepsia*, 27(4):423–431, 1986. ISSN 0013-9580. doi: 10.1111/j.1528-1157.1986.tb03562.x.
- Michael Wenzel, Michael Wenzel, Jordan P. Hamm, Jordan P. Hamm, Darcy S. Peterka, Darcy S. Peterka, Rafael Yuste, and Rafael Yuste. Reliable and Elastic Propagation of Cortical Seizures In Vivo. *Cell Reports*, 19(13):2681–2693, June 2017. doi: 10.1016/j.celrep.2017.05.090. MAG ID: 2729704653.
- Roy Mukamel and Itzhak Fried. Human intracranial recordings and cognitive neuroscience. Annual Review of Psychology, 63:511–537, 2012. ISSN 1545-2085. doi: 10.1146/ annurev-psych-120709-145401.

- Cayetano E. Napolitano and Miguel A. Orriols. Changing patterns of propagation in a superrefractory status of the temporal lobe. Over 900 seizures recorded over nearly one year. *Epilepsy* & Behavior Case Reports, 1:126–131, 2013. ISSN 2213-3232. doi: 10.1016/j.ebcr.2013.07.001.
- Philippa J Karoly, Philippa J. Karoly, Vikram R. Rao, Vikram R. Rao, Nicholas M. Gregg, Nicholas M. Gregg, Greg Worrell, Gregory A. Worrell, Christophe Bernard, Christophe Bernard, Mark Cook, Mark J. Cook, Maxime O. Baud, and Maxime O. Baud. Cycles in epilepsy. *Nature Reviews Neurology*, 17(5):267–284, 2021. doi: 10.1038/s41582-021-00464-1. MAG ID: 3138016489.
- Gabrielle M. Schroeder, Beate Diehl, Fahmida A. Chowdhury, John S. Duncan, Jane de Tisi, Andrew J. Trevelyan, Rob Forsyth, Andrew Jackson, Peter N. Taylor, and Yujiang Wang. Seizure pathways change on circadian and slower timescales in individual patients with focal epilepsy. *Proceedings of the National Academy of Sciences*, 117(20):11048–11058, May 2020. doi: 10. 1073/pnas.1922084117. URL https://www.pnas.org/doi/full/10.1073/pnas.1922084117. Publisher: Proceedings of the National Academy of Sciences.
- Gabrielle M Schroeder, Fahmida A Chowdhury, Mark J Cook, Beate Diehl, John S. Duncan, Philippa J Karoly, Peter N Taylor, and Yujiang Wang. Multiple mechanisms shape the relationship between pathway and duration of focal seizures. *Brain communications*, July 2022. doi: 10.1093/braincomms/fcac173. MAG ID: 4283828369.
- Susan S. Spencer, Dennis D. Spencer, Pétér D. Williamson, and Richard H. Mattson. Ictal Effects of Anticonvulsant Medication Withdrawal in Epileptic Patients. *Epilepsia*, 22(3):297–307, June 1981. ISSN 1528-1167. doi: 10.1111/j.1528-1157.1981.tb04113.x. URL https://onlinelibrary.wiley.com/doi/10.1111/j.1528-1157.1981.tb04113.x. Publisher: John Wiley & Sons, Ltd.
- Fabien B. Wagner, Emad N. Eskandar, G. Rees Cosgrove, Joseph R. Madsen, Andrew S. Blum, N. Stevenson Potter, Leigh R. Hochberg, Sydney S. Cash, and Wilson Truccolo. Microscale Spatiotemporal Dynamics during Neocortical Propagation of Human Focal Seizures. *NeuroImage*,

122:114-130, November 2015. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2015.08.019. URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4618174/.