Keldysh's theorem revisited

Johannes M. Schumacher[∗]

December 23, 2024

Abstract

In a variety of applications, the problem comes up of describing the principal part of the inverse of a holomorphic operator at an eigenvalue in terms of left and right root functions associated to the eigenvalue. Such a description was given by Keldysh in 1951. His theorem, the proof of which was published only in 1971, is a fundamental result in the local spectral theory of operator-valued functions. Here we present a streamlined derivation in the matrix case, and we extend Keldysh's theorem by means of a new principal part formula. Special attention is given to the semisimple case (first-order poles).

Keywords: matrix-valued function, root functions, principal part, semisimplicity condition, residue formula.

MSC: 47A56, 15A54, 15A24.

1 Introduction

In 1951, M.V. Keldysh gave a representation theorem for the principal part, at a given point in the complex plane, of operator-valued functions of the form $(I + A(\lambda))^{-1}$, where $A(\lambda)$ is a compact linear operator that depends holomorphically on the complex parameter λ . Keldysh's work was primarily motivated by applications to the dynamic analysis of flexible structures, in particular aircraft. Similar applications have motivated another early stream of research, exemplified by [Frazer et al. \(1938\)](#page-18-0) and [Lancaster \(1966\)](#page-18-1), which concentrated on matrices (rather than operators) that depend on a complex parameter.^{[1](#page-0-0)} In more recent years, while the local spectral theory of matrix-valued functions has remained highly relevant in the study of vibrating structures, applications in other areas have emerged as well. Perhaps most notably, a key result in [Engle and Granger](#page-17-0) [\(1987\)](#page-17-0), one of the publications for which the authors received the Nobel Prize in Economics in 2003, is a representation theorem for the principal part of the inverse of a holomorphic matrix at a particular point of interest in the complex plane. In the further development of the literature on cointegration in econometric theory, the local spectral theory of matrix-valued functions has continued to play a central role; see for example [la Cour \(1998\)](#page-18-2), [Johansen \(2008\)](#page-18-3), and [Faliva and Zoia](#page-17-1)

[∗]Amsterdam School of Economics, University of Amsterdam, Amsterdam, the Netherlands. E-mail: <j.m.schumacher@uva.nl>. ORCID 0000-0001-5753-3412.

¹Holomorphic operator-valued functions are classically used to describe relations between forces and displacements in a mechanical structure as a function of a complex (Fourier transform) parameter. The matrix case appears when attention is focused on forces and displacements at finitely many points; the vibration of the structure itself may then still be described by a partial differential equation.

[\(2009\)](#page-17-1). For the use of Keldysh's theorem in the numerical analysis of nonlinear eigenvalue problems, one may consult for instance Mennicken and Möller (2003) and Güttel and Tisseur (2017). [Kozlov and Maz'ya \(2013,](#page-18-6) p. 6), in their monograph on differential equations with operator coefficients, refer to the theorem as a fundamental result. Generally speaking, Keldysh's theorem can play a key role in situations where transform analysis is applied in a multivariate setting; for recent examples in the context of Markov-modulated stochastic processes, see [Beare et al. \(2022](#page-17-2)) and [Beare and Toda \(2024](#page-17-3)).

Keldysh's 1951 paper is a short note without proofs, containing the representation theorem as well as a completeness theorem for extended systems of eigenvectors. Proofs of both theorems were provided by Keldysh in a manuscript that was written in 1950 and circulated in 1951, but that was published only in 1971.^{[2](#page-1-0)} Later authors have given alternative proofs of (versions of) Keldysh's representation theorem; see for instance [Gohberg and Sigal \(1971\)](#page-18-7), Mennicken and Möller (1984). Some partial results have been rediscovered independently, as discussed later in this paper. Text-book treatments can be found in Mennicken and Möller (2003) and [Kozlov and Maz'ya \(2013\)](#page-18-6), both in an operator setting.

For many applications, the matrix case is sufficient. In this case, the theory has its own characteristics; techniques from commutative algebra can be used with ease, and it is natural to treat left and right eigenvectors on an equal footing. We present below a derivation of Keldysh's theorem in the matrix case using an approach that might be termed geometric, since the treatment is centered around a certain sequence of subspaces associated to a singularity of a matrix-valued function. The classical theory may in this way become more accessible to researchers interested in matrix-valued functions. A remark that can be made about the principal part expression given by Keldysh is that it does not treat left and right root spaces symmetrically. In this paper, a new principal part formula is obtained (Thm. [6.6\)](#page-12-0) that does give both an equal status.

2 Problem setting

Let Ω be a fixed open subset of the complex plane. The ring of holomorphic functions and the field of meromorphic functions defined on Ω are denoted by H and by M, respectively. Throughout the paper, attention is focused on a fixed point $\lambda_0 \in \Omega$. The ring of locally holomorphic functions (i.e., functions that are holomorphic in a neighborhood of λ_0) is denoted by \mathcal{H}_0 . Due to the fact that zeros of holomorphic functions are isolated, a function $y \in \mathcal{H}_0$ is a unit in \mathcal{H}_0 (i.e., has an inverse in \mathcal{H}_0) if and only if $y(\lambda_0) \neq 0$.

A matrix $M \in \mathcal{M}^{n \times n}$ is said to be *invertible* when it has an inverse in $\mathcal{M}^{n \times n}$, or, equivalently, when det M is not identically zero. A matrix $M \in \mathcal{H}_0^{n \times n}$ that has an inverse in $\mathcal{H}_0^{n \times n}$ is said to be unimodular (with respect to the ring \mathcal{H}_0). It follows from Cramer's rule that $M \in \mathcal{H}_0^{n \times n}$ is unimodular if and only if det M is a unit in \mathcal{H}_0 , i.e., if and only if the constant matrix $M(\lambda_0) \in$ $\mathbb{C}^{n \times n}$ is nonsingular. We will say that a matrix $M \in \mathcal{H}_0^{n \times m}$ is *left (right)* unimodular when it has a left (right) inverse in $\mathcal{H}_0^{m \times n}$; this happens if and only if the constant matrix $M(\lambda_0)$ has full column (row) rank.

 2 See the editor's note in [Keldysh \(1971](#page-18-9)).

For notational convenience, define $\chi_0 \in \mathcal{H}$ by

$$
\chi_0(\lambda) = \lambda - \lambda_0 \qquad (\lambda \in \Omega). \tag{1}
$$

The Laurent expansion around λ_0 of a matrix $M \in \mathcal{M}^{n \times m}$ can then be written as

$$
M = \sum_{j=-s}^{\infty} M_j \chi_0^j \tag{2}
$$

with $M_j \in \mathbb{C}^{n \times n}$ $(j = -s, -s + 1, \dots)$. The meromorphic matrix M is said to have a pole at λ_0 if there exists $j < 0$ such that $M_j \neq 0$, and the *order* of the pole at λ_0 is the smallest value of j for which this holds. Matrices in $\mathcal{H}_0^{n \times n}$ are said to have pole order 0 at λ_0 . The coefficient of χ_0^{-1} in the Laurent series expansion is called the *residue* of M at λ_0 . The part of the sum in [\(2\)](#page-2-0) that corresponds to indices $j < 0$ is called the *principal part* of M at λ_0 . The same terminology may be used when $m = 1$ or $n = 1$ and M is viewed as a column or row vector, rather than as a matrix.

The notation " $\dot{=}$ " is used in this paper to express *equality of principal parts at* λ_0 . In other words, given $M_1, M_2 \in \mathcal{M}^{n \times m}$, we define

$$
M_1 \doteq M_2 \quad \Leftrightarrow \quad \text{there exists } H \in \mathcal{H}_0^{n \times m} \text{ such that } M_2 = M_1 + H. \tag{3}
$$

This could also be written in a more standard way as $M_1 = M_2$ mod $\mathcal{H}_0^{n \times m}$, but it is convenient to use a shorter form. The notation in [\(3\)](#page-2-1) is applied in the same way to vectors in \mathcal{M}^n . We will also use the quotient space $\mathcal{M}^n/\mathcal{H}^n_0$ consisting of equivalence classes of vector functions, where two meromorphic vectors are taken to be equivalent when their difference is holomorphic in a neighborhood of λ_0 . In other words, the equivalence relation is the one given by equality of principal parts at λ_0 . The equivalence class of $y \in \mathcal{M}^n$ will be denoted by [y].

To avoid repetitions, throughout the paper the symbol T will denote a matrix in $\mathcal{H}^{n \times n}$ that is invertible as a meromorphic matrix and that has a singularity at λ_0 , i.e., we have det $T(\lambda_0) = 0$ while $\det T$ is not identically 0. It follows from Cramer's rule that T^{-1} is meromorphic. The order of the pole of T^{-1} at λ_0 is denoted by s, i.e.,

$$
s = \min\{j \ge 0 \mid \chi_0^j T^{-1} \in \mathcal{H}_0^{n \times n}\}.
$$
\n(4)

We also write

$$
r = \dim \ker T(\lambda_0). \tag{5}
$$

The number r is called the *geometric multiplicity* of the root of the holomorphic matrix T at the point λ_0 . The algebraic multiplicity of the root of T at λ_0 is the multiplicity of λ_0 as a root of the scalar function $\det T$.

3 Partial multiplicities and root functions

In the ring H_0 , the product of two elements can only be 0 if at least one of them is 0, and an element is divisible by another if and only if the multiplicity of the zero at λ_0 of the first is equal to or higher than the multiplicity of the zero at λ_0 of the second. These properties imply in particular that \mathcal{H}_0 is a principal ideal domain. One can therefore use the canonical form for matrices over principal ideal domains that was given by H.J.S. Smith in 1861 (see for instance [MacDuffee \(1946](#page-18-10), Thm. 26.2)).

Theorem 3.1 (Smith form w.r.t. \mathcal{H}_0). There exist unimodular matrices $U_L, U_R \in \mathcal{H}_0^{n \times n}$ and uniquely determined nonnegative integers m_1, \ldots, m_n such that

$$
U_L T U_R = \text{diag}(\chi_0^{m_1}, \dots, \chi_0^{m_n}) \qquad (m_1 \geq \dots \geq m_n). \qquad (6)
$$

The numbers m_i are called the *partial multiplicities* of the root of T at λ_0 . It follows from [\(6\)](#page-3-0) that the algebraic multiplicity is the sum of the partial multiplicities. Since the pole order of T at λ_0 is the same as that of any matrix obtained from T by left and right unimodular transformations, we have $s = \max_{1 \leq i \leq n} m_i = m_1$. Note also that $m_r > 0$, and $m_i = 0$ for $i = r + 1, \ldots, n$. It will be convenient to define a diagonal matrix containing only the diagonal elements in the Smith form that correspond to positive partial multiplicities:

$$
\Delta = \text{diag}(\chi_0^{m_1}, \dots, \chi_0^{m_r}) \in \mathcal{H}_0^{r \times r}.
$$
\n⁽⁷⁾

There is a constructive algorithm by which any given holomorphic matrix can be reduced to Smith form. The algorithm is laborious, however, and gives little insight in the way in which the invariants m_i relate to the coefficients in the power series expansion of T around λ_0 . Fortunately, an alternative way of determining the partial multiplicities is available. Define a sequence of subspaces of \mathbb{C}^n as follows:^{[3](#page-3-1)}

$$
L_j = \{ y(\lambda_0) \mid y \in \mathcal{H}_0^n, \ Ty \in \chi_0^j \mathcal{H}_0^n \} \qquad (j = 0, 1, 2, ...). \tag{8}
$$

Note that $L_j \supset L_{j+1}$ for all j, since $\chi_0^{j+1} \mathcal{H}_0^n \subset \chi_0^j \mathcal{H}_0^n$ for all j. Also, $L_1 = \ker T(\lambda_0)$, and $L_j = \{0\}$ for $j > s$. Define a corresponding nonincreasing sequence of nonnegative integers by

$$
\ell_j = \dim L_j \qquad (j = 0, 1, 2, \dots). \tag{9}
$$

Lemma 3.2. The integers ℓ_j defined in [\(9\)](#page-3-2) are invariants under left and right multiplication of T by unimodular matrices.

Proof. Fix $j \geq 0$. Invariance under left multiplication by unimodular matrices is immediate. To prove the other half of the claim, let a unimodular matrix $U \in \mathcal{H}_0^{n \times n}$ be given, and take $y^0 \in L_j(TU)$. There exists $\tilde{y} \in \mathcal{H}_0^n$ such that $\tilde{y}(\lambda_0) = y^0$ and $TU\tilde{y} \in \chi_0^j\mathcal{H}_0^n$. Defining $y \in \mathcal{H}_0^n$ by $y = U\tilde{y}$, we have $Ty \in \chi_0^j \mathcal{H}_0^n$, so that $y(\lambda_0) \in L_j(T)$. Since $y^0 = \tilde{y}(\lambda_0) = U^{-1}(\lambda_0)y(\lambda_0)$, this shows that $L_j(TU) \subset U^{-1}(\lambda_0)L_j(T)$. Similarly we have $L_j(T) \subset U(\lambda_0)L_j(TU)$, and it follows that dim $L_i(T) = \dim L_i(TU)$. 口

By reduction to Smith form, one finds the following.

³We will also write $L_i(T)$ in cases where T needs to be specified. Mennicken and Möller (2003, p. 14) give a different but equivalent definition.

Proposition 3.3. For all $j \geq 0$, we have

$$
\ell_j = \#\{1 \le i \le n \mid m_i \ge j\} \tag{10}
$$

where the integers m_i are given by [\(6\)](#page-3-0).

Because the integers m_i are in nonincreasing order, we can also write $\ell_j = \max\{i \geq 1 \mid m_i \geq j\}$ instead of [\(10\)](#page-4-0). Therefore, $m_i \geq j$ if and only if $\ell_j \geq i$. This shows that $(m_i)_{i=1,2,...}$ relates to $(\ell_j)_{j=1,2,...}$ in the same way as $(\ell_j)_{j=1,2,...}$ relates to $(m_i)_{i=1,2,...}$. Consequently, we have

$$
m_i = \max\{j \ge 1 \mid \ell_j \ge i\} = \#\{j \ge 1 \mid \ell_j \ge i\}
$$

 $(i = 1, ..., n).$ (11)

From [\(10\)](#page-4-0) it also follows that $\#\{i \mid m_i = j\} = \ell_j - \ell_{j+1}$, so that

$$
\sum_{i=1}^{r} m_i = \sum_{j=1}^{s} j(\ell_j - \ell_{j+1}) = \sum_{j=1}^{s} \ell_j.
$$
 (12)

The sequences (ℓ_1, \ldots, ℓ_s) and (m_1, \ldots, m_r) are *conjugate partitions* of their common sum in the sense of enumerative combinatorics; see for instance [Andrews \(1998,](#page-17-4) Def. 1.8).

From [\(11\)](#page-4-1), one sees that the partial multiplicities m_i can be determined by computing the dimensions of the subspaces L_j $(j = 1, \ldots, s)$.^{[4](#page-4-2)} The following proposition shows that finding these dimensions is a matter of solving finite systems of linear equations stated in terms of the first s coefficients in the power series expansion of T around λ_0 .

Proposition 3.4. For $j \geq 1$, a vector $y^0 \in \mathbb{C}^n$ belongs to the subspace L_j if and only if there exist vectors $y^1, \ldots, y^{j-1} \in \mathbb{C}^n$ such that

$$
\sum_{p=0}^{k} \frac{1}{p!} T^{(p)}(\lambda_0) y^{k-p} = 0 \quad \text{for all } 0 \le k \le j-1.
$$
 (13)

Proof. For a vector $y \in \mathcal{H}_0^n$ with power series expansion $y = \sum_{j=0}^{\infty} y^j \chi_0^j$ around λ_0 , the left hand side of [\(13\)](#page-4-3) gives the k-th coefficient of the power series expansion of Ty around λ_0 . Therefore, if vectors y^1, \ldots, y^{j-1} are given such that [\(13\)](#page-4-3) is satisfied, then the function defined by $y =$ $\sum_{p=0}^{j-1} \chi_0^p y^p$ satisfies $y(\lambda_0) = y^0$ and $Ty \in \chi_0^j \mathcal{H}_0^n$. Conversely, take $y^0 \in L_j$, and let $y \in \mathcal{H}_0^n$ be such that $Ty \in \chi_0^j H_0^n$. Since $(Ty)^{(k)}(\lambda_0) = 0$ for $0 \le k \le j-1$, the vectors defined for $p = 1, \ldots, j-1$ by $y^p = y^{(p)}(\lambda_0)/p!$ satisfy [\(13\)](#page-4-3). □

The following definition is standard in the literature; see for instance Mennicken and Möller $(2003, \text{ Def. } 1.6).$ $(2003, \text{ Def. } 1.6).$ ^{[5](#page-4-4)}

Definition 3.5. A function $y \in \mathcal{H}^n$ is said to be a *root function* for T at λ_0 if $y(\lambda_0) \neq 0$ and $T(\lambda_0)y(\lambda_0)=0$. The order of the zero of Ty at λ_0 is called the multiplicity of y (with respect to T at λ_0), and is denoted by $\nu(y)$.

⁴Consequently, one might *define* the partial multiplicities m_i by [\(11\)](#page-4-1), rather than via the Smith form. This is the route taken in Mennicken and Möller (2003). In the work of [Keldysh \(1951](#page-18-11), [1971](#page-18-9)), partial multiplicities appear as the maximal multiplicities that appear in systems of root functions whose evaluations at λ_0 form a basis for the nullspace of $T(\lambda_0)$; compare Prop. [4.6](#page-7-0) below.

⁵One might use \mathcal{H}_0 rather than \mathcal{H} in the definition of root functions without causing harm. See also Lemma [3.6.](#page-5-0)

[Gohberg and Sigal \(1971\)](#page-18-7) attribute the introduction of root functions to Kreĭn and Trofimov (1969) and [Palant \(1970\)](#page-18-13) independently.^{[6](#page-5-1)} [Keldysh \(1951,](#page-18-11) [1971\)](#page-18-9) instead uses "systems of eigenvectors and associated vectors". Such a system appears as the sequence of coefficients of a polynomial root function.^{[7](#page-5-2)} The coefficient corresponding to the constant term of the polynomial is called an eigenvector, since it must belong to ker $T(\lambda_0)$, and the other coefficients are called associated vectors. In part of the literature (for instance [Bart et al. \(1979\)](#page-17-5)), a system of eigenvectors and associated vectors is called a Jordan chain.

In Def. [3.5,](#page-4-5) root functions are required to be holomorphic throughout the domain Ω , whereas, in the definition of the subspaces L_j in [\(8\)](#page-3-3), locally holomorphic functions were used. The following lemma (which formalizes a remark in Mennicken and Möller $(2003, p. 14)$) shows that this difference is inconsequential.

Lemma 3.6. A vector $y^0 \in \mathbb{C}^n$ with $y^0 \neq 0$ belongs to the subspace L_j if and only if there exists a root function $y \in \mathcal{H}^n$ of multiplicity j such that $y(\lambda_0) = y^0$.

Proof. If $y^0 \in L_j$, then, by definition, there exists $\tilde{y} \in \mathcal{H}_0^n$ such that $\tilde{y}(\lambda_0) = y^0$ and $T\tilde{y} \in$ $\chi_0^j \mathcal{H}_0^n$. For a function $y \in \mathcal{H}^n$ to be such that $y(\lambda_0) = y_0$ and $Ty \in \chi_0^j \mathcal{H}_0^n$, it is sufficient that $y^{(k)}(\lambda_0) = \tilde{y}^{(k)}(\lambda_0)$ for $k = 0, 1, \ldots, j$. These requirements can indeed be satisfied; in particular, one can choose y to be a polynomial. This proves the "only if" part of the claim; the "if" part is trivial. \Box

4 Canonical matrices

For any matrix $M \in \mathcal{H}_0^{n \times m}$, one can unambiguously define an induced mapping \overline{M} from the quotient space $\mathcal{M}^m/\mathcal{H}_0^m$ to the quotient space $\mathcal{M}^n/\mathcal{H}_0^n$ by

$$
\overline{M}: [y] \mapsto [My] \qquad (y \in \mathcal{M}^m). \tag{14}
$$

This is a linear mapping. The relevance of the induced mapping to the concept of root functions is indicated by the fact that a vector $y \in \mathcal{H}_0^n$ is a root function for T at λ_0 with multiplicity at least j if and only if $[\chi_0^{-j}y] \in \ker \overline{T}$. To connect the dimension of $\ker \overline{T}$ to the multiplicities defined in [\(6\)](#page-3-0), we need the following simple lemma.

Lemma 4.1. If $M \in \mathcal{H}_0^{n \times m}$ is left unimodular, then the induced mapping \overline{M} : $\mathcal{M}^m/\mathcal{H}_0^m \to$ $\mathcal{M}^n/\mathcal{H}_0^n$ is injective.

Proof. To prove the claim, we have to show that, if $y \in \mathcal{M}^m$ is such that $My \in \mathcal{H}_0^n$, then $y \in \mathcal{H}_0^m$. But this is immediate, since M has a left inverse in $\mathcal{H}_0^{m \times n}$. П

Proposition 4.2. We have

$$
\dim \ker \overline{T} = \sum_{i=1}^{r} m_i.
$$
\n(15)

 6 The latter paper is as given in the list of references of [Gohberg and Sigal \(1971](#page-18-7)), but they actually refer to it as being authored by Macaev and Palant. Perhaps they had [Macaev and Palant \(1966\)](#page-18-14) in mind. The source cited by Mennicken and Möller (2003) for the notion of root functions is [Trofimov \(1968](#page-18-15)).

⁷Requiring root functions to be polynomial is possible (see the proof of Lemma [3.6\)](#page-5-0); this restriction is often viewed as inconvenient, however.

Proof. If $U \in \mathcal{H}_0^{n \times n}$ is unimodular, then, for any $[y] \in \text{ker } \overline{T}$, we have $[U^{-1}y] \in \text{ker } \overline{T}U$ and $\overline{U}[U^{-1}y] = [y]$. This shows that the induced mapping \overline{U} : $\mathcal{M}^n/\mathcal{H}_0^n \to \mathcal{M}^n/\mathcal{H}_0^n$ maps the space ker \overline{TU} onto the space ker \overline{T} . Since \overline{U} is injective by the above lemma, it follows that dim ker \overline{TU} = dim ker \overline{T} . Again by the lemma above, we also have ker $\overline{UT} = \ker \overline{T}$ when U is unimodular. We can therefore use the Smith form to conclude that [\(15\)](#page-5-3) holds. \square

Proposition 4.3. If $\{y_1, \ldots, y_q\}$ is a collection of root functions for T at λ_0 such that the constant vectors $\{y_1(\lambda_0), \ldots, y_q(\lambda_0)\}\$ are linearly independent, then the sum of the multiplicities of these root functions does not exceed the algebraic multiplicity of the root of T at λ_0 .

Proof. Let $Y \in \mathcal{H}_0^{n \times q}$ be defined as the matrix with columns y_i $(i = 1, ..., q)$, write $\nu_i = \nu(y_i)$, and define $D = \text{diag}(\chi_0^{\nu_1}, \ldots, \chi_0^{\nu_q})$. Note that Y is left unimodular. By the definition of root functions, we have $T Y D^{-1} \in \mathcal{H}_0^{n \times q}$, which implies that the space $(Y D^{-1} \mathcal{H}_0^q)/\mathcal{H}_0^n \subset \mathcal{M}^n/\mathcal{H}_0^n$ is a subspace of ker \overline{T} . Since the induced mapping \overline{Y} is injective on $\mathcal{M}^q/\mathcal{H}^q_0$ by Lemma [4.1,](#page-5-4) the dimension of $(YD^{-1}\mathcal{H}_0^q)/\mathcal{H}_0^n$ is the same as the dimension of $(D^{-1}\mathcal{H}_0^q)/\mathcal{H}_0^q$. The matrix D induces a bijective mapping from $(D^{-1}\mathcal{H}_0^q)/\mathcal{H}_0^q$ to $\mathcal{H}_0^n/(D\mathcal{H}_0^n)$. The dimension of the latter space is equal to $\sum_{i=1}^q \nu_i$, since it has a basis consisting of equivalence classes modulo $D\mathcal{H}_0^n$ of the form $[\chi_0^j e_i]$, where e_i denotes the *i*-th unit vector in \mathcal{H}_0^q $(i = 1, ..., q)$ and $0 \leq j \leq \nu_i - 1$. It follows that

$$
\sum_{i=1}^{q} \nu_i = \dim (D^{-1} \mathcal{H}_0^q) / \mathcal{H}_0^q = \dim (Y D^{-1} \mathcal{H}_0^q) / \mathcal{H}_0^q \le \dim \ker \overline{T} = \sum_{i=1}^{r} m_i.
$$

The proposition motivates the following definition.

Definition 4.4. A *canonical system of root functions* for T at λ_0 is a collection $\{y_1, \ldots, y_r\}$ of root functions for T at λ_0 such that the following conditions are satisfied:

- (i) the vectors $y_1(\lambda_0), \ldots, y_r(\lambda_0) \in \mathbb{C}^n$ are linearly independent;
- (ii) $\sum_{i=1}^r \nu(y_i) = \sum_{i=1}^r m_i;$
- (iii) the multiplicities $\nu(y_i)$ are in nonincreasing order, i.e., $\nu(y_i) \geq \nu(y_k)$ whenever $i \leq k$.

Condition (iii) can of course always be satisfied by reordering the root functions if necessary; the requirement just serves to simplify the notation.

We shall say that a basis $\{y_1^0, \ldots, y_r^0\}$ of ker $T(\lambda_0)$ is *adapted* to the nonincreasing sequence $\{L_j\}_{1\leq j\leq s}$ when, for each $j=1,\ldots,s$, the subspace L_j is spanned by the vectors $y_1^0,\ldots,y_{\ell_j}^0$ where $\ell_j = \dim L_j$. Clearly, one can construct such a basis by starting with a basis for the smallest subspace L_s , then extending it to a basis for the next larger subspace L_{s-1} , and so on.

Proposition 4.5. A canonical system of root functions for T at λ_0 exists.

Proof. Take a basis $\{y_1^0, \ldots, y_r^0\}$ for ker $T(\lambda_0)$ that is adapted to the sequence L_1, \ldots, L_s . By Lemma [3.6,](#page-5-0) there are root functions y_1, \ldots, y_r such that $y_i(\lambda_0) = y_i^0$ for $i = 1, \ldots, r$ and $\nu(y_i) \geq j$ when $y_i^0 \in L_j$. Because of the adaptedness of the basis, the root functions that are constructed in this way contain ℓ_{m_1} elements with multiplicity at least $m_1 = s, \ell_{m_2} - \ell_{m_1}$ elements with multiplicity at least m_2 , and so on. Therefore, we have (using (12))

$$
\sum_{i=1}^{r} \nu(y_i) \ge \sum_{j=1}^{s} j(\ell_j - \ell_{j+1}) = \sum_{j=1}^{s} \ell_j = \sum_{i=1}^{r} m_i.
$$
 (16)

By Prop. [4.3,](#page-6-0) equality holds, and the system $\{y_1, \ldots, y_r\}$ is canonical.

Proposition 4.6. If $\{y_1, \ldots, y_r\}$ is a canonical system, then $\nu(y_i) = m_i$ for $i = 1, \ldots, r$.

Proof. Write $\nu_i := \nu(y_i)$. For $j = 1, ..., s$, define $\hat{\ell}_j = \#\{1 \leq i \leq r \mid \nu_i \geq j\}$. The sequences (ν_1,\ldots,ν_r) and $(\hat{\ell}_1,\ldots,\hat{\ell}_s)$ are conjugate partitions, just as (m_1,\ldots,m_r) and (ℓ_1,\ldots,ℓ_s) are. Moreover, since $y_i(\lambda_0) \in L_j$ when $\nu_i \geq j$, we have $\hat{\ell}_j \leq \ell_j$ for all $j = 1, \ldots, s$. It follows that

$$
\nu_i = \#\{1 \le j \le s \mid \hat{\ell}_j \ge i\} \le \#\{1 \le j \le s \mid \ell_j \ge i\} = m_i \qquad (i = 1, \dots, r).
$$

Consequently, the canonicity condition $\sum_{i=1}^{r} \nu_i = \sum_{i=1}^{r} m_i$ can hold only when $\nu_i = m_i$ for all $i=1,\ldots,r.$ \Box

It is convenient to employ a matrix format for canonical systems of root functions.

Definition 4.7. A matrix $Y \in \mathcal{H}^{n \times r}$ is said to be a *right canonical matrix* (for T at λ_0) when its columns form a canonical system of root functions for T at λ_0 .

Since the root functions in a canonical system are ordered by nonincreasing multiplicity, we have the following.

Lemma 4.8. If Y is a right canonical matrix for T at λ_0 , then, for each $j = 1, \ldots, s$, the first ℓ_j columns of $Y(\lambda_0)$ form a basis for the subspace L_j .

Recall the definition of the matrix Δ in [\(7\)](#page-3-4) and the notation $\dot{=}$ introduced in [\(3\)](#page-2-1).

Proposition 4.9. A matrix $Y \in \mathcal{H}^{n \times r}$ is a right canonical matrix for T at λ_0 if and only if the following two conditions are satisfied:

- (i) $Y(\lambda_0) \in \mathbb{C}^{n \times r}$ has full column rank;
- (*ii*) $TY\Delta^{-1} \doteq 0$.

Proof. Suppose that conditions (i) and (ii) hold. Condition (ii) is equivalent to the statement that there exists a matrix $H \in \mathcal{H}_0^{n \times r}$ such that $TY = H\Delta$; i.e., $Ty_i = h_i \chi_0^{m_i}$ for $i = 1, \ldots, r$, where $y_i \in \mathcal{H}^n$ and $h_i \in \mathcal{H}_0^n$ denote the *i*-th columns of Y and H respectively. This implies that y_i is a root function for T at λ_0 , with multiplicity $\nu(y_i) \geq m_i$. By condition (i) and Prop. [4.3,](#page-6-0) the sum of the multiplicities of the root functions y_i is equal to $\sum_{i=1}^r m_i$, and hence $\{y_1, \ldots, y_r\}$ is a canonical system. Conversely, if $Y \in \mathcal{H}^{n \times r}$ is right canonical, then, for each $i = 1, \ldots, r$, the *i*-th column of Y satisfies $Ty_i \in \chi_0^{m_i} \mathcal{H}_0^n$ by Prop. [4.6.](#page-7-0) This implies that (ii) holds, and (i) holds by definition. \Box

The following proposition rephrases and extends Lemma A.9.3 of [Kozlov and Maz'ya \(2013\)](#page-18-6) in transposed form. The proof uses a modification of the argument in the proof of Thm. 1.2 in [Gohberg et al. \(1993\)](#page-18-16).

$$
\Box
$$

Proposition 4.10. Let Y be a right canonical matrix for T at λ_0 . Then the constant matrix $(TY\Delta^{-1})(\lambda_0) \in \mathbb{C}^{n \times r}$ has full column rank, and the following direct sum decomposition holds:

$$
\mathbb{C}^n = \operatorname{im} T(\lambda_0) \oplus \operatorname{im} (TY\Delta^{-1})(\lambda_0). \tag{17}
$$

Proof. For a proof by contradiction, suppose there exist $y^0 \in \mathbb{C}^n$ and $x \in \mathbb{C}^r$, with $x \neq 0$, such that $(TY\Delta^{-1})(\lambda_0)x = T(\lambda_0)y^0$. Define $k = \min\{j \geq 1 \mid \chi_0^j \Delta^{-1}x \in \mathcal{H}_0^r\}$. From $\chi_0^k \Delta^{-1}x \in \mathcal{H}_0^n$ it follows that $x_i = 0$ for all indices i such that $m_i \geq k+1$, i.e., $1 \leq i \leq \ell_{k+1}$. Write \hat{x} := $(\chi_0^k \Delta^{-1} x)(\lambda_0)$. Note that $\hat{x} \neq 0$, because otherwise we would have $\chi_0^{k-1} \Delta^{-1} x \in \mathcal{H}_0^r$ and k would not be minimal. Define $y = \chi_0^k (Y \Delta^{-1} x - y^0) \in \mathcal{H}_0^n$. The assumption $(TY \Delta^{-1})(\lambda_0) x = T(\lambda_0) y^0$ implies that $TY\Delta^{-1}x - Ty^0 \in \chi_0\mathcal{H}_0^n$, and hence $Ty = \chi_0^k(TY\Delta^{-1}x - Ty^0) \in \chi_0^{k+1}\mathcal{H}_0^n$. It follows that $y(\lambda_0) \in L_{k+1}$. On the other hand, $y(\lambda_0) = Y(\lambda_0)\hat{x}$, and the vector \hat{x} inherits from x the property that all of its entries with indices $i \leq \ell_{k+1}$ are 0. In view of Lemma [4.8,](#page-7-1) we have a contradiction. It follows that the matrix $(TY\Delta^{-1})(\lambda_0)$ has full column rank (for this, take $y^0 = 0$) and that $\text{im } T(\lambda_0) \cap \text{im } (TY\Delta^{-1})(\lambda_0) = \{0\}.$ The proof is completed by noting that the dimension of im $T(\lambda_0)$ is $n-r$, while dim im $(TY\Delta^{-1})(\lambda_0)=r$ by the full column rank property. \Box

5 Keldysh's theorem

A holomorphic row vector v of length n is said to be a *left root function* for T at λ when $v(\lambda_0) \neq 0$ and $v(\lambda_0)T(\lambda_0) = 0$. The notion of a left canonical matrix for T at λ_0 is defined analogously to Def. [4.7,](#page-7-2) and one has the following characterization.

Proposition 5.1. A matrix $V \in \mathcal{H}^{r \times n}$ is a left canonical matrix for T at λ_0 if and only if $V(\lambda_0) \in \mathbb{C}^{r \times n}$ has full row rank, and $\Delta^{-1} V T \doteq 0$.

Prop. [4.10](#page-8-0) translates to left canonical matrices as follows.

Proposition 5.2. Let V be a left canonical matrix for T at λ_0 . Then the constant matrix $(\Delta^{-1}VT)(\lambda_0)$ has full row rank, and the following direct sum decomposition holds:

$$
\mathbb{C}^n = \ker T(\lambda_0) \oplus \ker(\Delta^{-1}VT)(\lambda_0). \tag{18}
$$

For later use, we note the following corollary.

Corollary 5.3. Let V and Y be left and right canonical matrices, respectively, for T at λ_0 . Then the matrices $\Delta^{-1} V T Y \in \mathcal{H}_0^{r \times r}$ and $V T Y \Delta^{-1} \in \mathcal{H}_0^{r \times r}$ are unimodular.

Proof. From the definition of right canonical matrices, it follows that $Y(\lambda_0) \in \mathbb{C}^{n \times r}$ is a basis matrix for ker $T(\lambda_0)$. Since the subspace ker $\Delta^{-1}VT(\lambda_0)$ is complementary to ker $T(\lambda_0)$ by Prop. [5.2,](#page-8-1) the matrix $(\Delta^{-1}VTY)(\lambda_0) = (\Delta^{-1}VT)(\lambda_0)Y(\lambda_0) \in \mathbb{C}^{r \times r}$ is nonsingular. Consequently, $\Delta^{-1} V T Y$ is unimodular. The proof for $V T Y \Delta^{-1}$ is analogous. \Box

The Smith form may be used to show that the principal part of T^{-1} at λ_0 can be described in terms of suitably chosen left and right canonical matrices for T at λ_0 .

Theorem 5.4. There exist left and right canonical matrices V and Y for T at λ_0 such that $T^{-1} \doteq Y \Delta^{-1} V.$

Proof. There are unimodular matrices U_L , $U_R \in \mathcal{H}_0^{n \times n}$ such that, with conformable partitioning,

$$
T = \begin{bmatrix} U_L^1 & U_L^2 \end{bmatrix} \begin{bmatrix} \Delta & 0 \\ 0 & I_{n-r} \end{bmatrix} \begin{bmatrix} U_R^1 \\ U_R^2 \end{bmatrix} . \tag{19}
$$

Write, again with conformable partitioning,

$$
U_L^{-1} = \begin{bmatrix} V \\ \tilde{V} \end{bmatrix}, \qquad U_R^{-1} = \begin{bmatrix} Y & \tilde{Y} \end{bmatrix}.
$$
 (20)

We have

$$
T^{-1} = \begin{bmatrix} Y & \tilde{Y} \end{bmatrix} \begin{bmatrix} \Delta^{-1} & 0 \\ 0 & I_{n-r} \end{bmatrix} \begin{bmatrix} V \\ \tilde{V} \end{bmatrix} = Y\Delta^{-1}V + \tilde{Y}\tilde{V} \doteq Y\Delta^{-1}V. \tag{21}
$$

From the fact that $[Y(\lambda_0) \ \ \tilde{Y}(\lambda_0)]$ is nonsingular it follows that $Y(\lambda_0)$ has full column rank r. Multiplying from the left by T and from the right by U_L^1 , one finds from [\(21\)](#page-9-0) that $TY\Delta^{-1} \doteq 0$, so that Y is right canonical by Prop. [5.1.](#page-8-2) Left canonicity of V is proved analogously. □

Remark 5.5. For future reference, note that from [\(19\)](#page-9-1) and [\(20\)](#page-9-2) it also follows that

$$
\begin{bmatrix} \Delta & 0 \\ 0 & I_{n-r} \end{bmatrix} = \begin{bmatrix} V \\ \tilde{V} \end{bmatrix} T \begin{bmatrix} Y & \tilde{Y} \end{bmatrix} = \begin{bmatrix} VTY & VT\tilde{Y} \\ \tilde{V}TY & \tilde{V}T\tilde{Y} \end{bmatrix}.
$$
 (22)

In particular, we have $VTY = \Delta$.

Thm. [5.4](#page-9-3) is the matrix version of Thm. 7.1 in [Gohberg and Sigal \(1971\)](#page-18-7), and the proof as given above follows their argument. The result given in [Keldysh \(1951](#page-18-11), [1971](#page-18-9)) has a stronger matrix version. We write it as follows.

Theorem 5.6 (Keldysh 1951). For any right canonical matrix Y for T at λ_0 , there exists a left canonical matrix V for T at λ_0 , determined uniquely up to addition of a matrix of the form ΔH with $H \in \mathcal{H}^{r \times n}$, such that

$$
T^{-1} \doteq Y\Delta^{-1}V.\tag{23}
$$

For any such matrix V , one has

$$
\Delta^{-1} V T Y \Delta^{-1} \doteq \Delta^{-1}.
$$
\n(24)

Condition (24) is called the *biorthogonality condition*. It is stated here in essentially the form given by [Kozlov and Maz'ya \(2013,](#page-18-6) Thm. A.10.1). These authors also prove (Thm. A.10.2) that the condition [\(24\)](#page-9-4) is equivalent to the quadruply indexed formulation in terms of systems of eigenvectors and associated vectors as given by [Keldysh \(1951](#page-18-11), [1971](#page-18-9)) and by Mennicken and Möller [\(2003\)](#page-18-4).

To prove the theorem, one can make use of the following lemma. The proof as given below is a rephrasing of part of the argument in the proof of Thm. 1.5.4 in Mennicken and Möller (2003).

Lemma 5.7. Let $Y \in \mathcal{H}^{n \times r}$ be a right canonical matrix for T at λ_0 , and suppose $V \in \mathcal{H}^{r \times n}$ is such that $T^{-1} - Y\Delta^{-1}V$ has pole order $j \geq 1$ at λ_0 . Then there exists a matrix $\tilde{V} \in \mathcal{H}^{r \times n}$ such that $T^{-1} - Y\Delta^{-1}\tilde{V}$ has pole order at most $j-1$ at λ_0 .

Proof. Write $A = T^{-1} - Y\Delta^{-1}V$. By assumption, there are matrices $A_1, \ldots, A_j \in \mathbb{C}^{n \times n}$ such that $A = A_1 \chi_0^{-1} + \cdots + A_j \chi_0^{-j}$. For any $x \in \mathbb{C}^n$, we have $A_j x = (\chi_0^j Ax)(\lambda_0) \in L_j$, since $T\chi_0^jAx = \chi_0^j(I - TY\Delta^{-1}V)x \in \chi_0^j\mathcal{H}_0^n$ by Prop. [4.9.](#page-7-3) This implies, by Lemma [4.8,](#page-7-1) that each of the columns of A_j can be written as a linear combination of the first ℓ_j columns of $Y(\lambda_0)$. Let the submatrix of Y formed by these columns be denoted by $Y_{1:\ell_j}$, and let $M \in \mathbb{C}^{\ell_j \times n}$ be such that $A_j = Y_{1:\ell_j}(\lambda_0)M$. Then the pole order of $A - \chi_0^{-j}Y_{1:\ell_j}M$ at λ_0 is at most $j-1$. Now define

$$
\tilde{V} = V + \chi_0^{-j} \Delta \begin{bmatrix} M \\ 0_{(r-\ell_j)\times n} \end{bmatrix}
$$

.

Note that $\tilde{V} \in \mathcal{H}^{r \times n}$, since $m_i \geq j$ for $i \leq \ell_j$. Moreover,

$$
T^{-1} - Y\Delta^{-1}\tilde{V} = A - Y\Delta^{-1}\chi_0^{-j}\Delta\begin{bmatrix} M \\ 0 \end{bmatrix} = A - \chi_0^{-j}Y_{1:\ell_j}M
$$

so that \tilde{V} has the required properties.

Keldysh's theorem, in the matrix case, can now be proved as follows.

Proof. Existence of $V \in \mathcal{H}^{r \times n}$ such that [\(23\)](#page-9-5) holds follows by repeated application of Lemma [5.7,](#page-10-0) starting with $V = 0$. For the rest of the proof, note that Y is left unimodular by Prop. [4.9;](#page-7-3) let $Z \in \mathcal{H}_0^{r \times n}$ be a left inverse. The biorthogonality condition [\(24\)](#page-9-4) is obtained by multiplying both sides of [\(23\)](#page-9-5) from the left by Z and from the right by $TY\Delta^{-1} \in \mathcal{H}_0^{n \times r}$. Next, if V satisfies (23), then clearly the same holds for any matrix \hat{V} of the form $\hat{V} = V + \Delta H$ where $H \in \mathcal{H}^{r \times n}$. Conversely, suppose that $V, \hat{V} \in \mathcal{H}^{r \times n}$ both satisfy [\(23\)](#page-9-5). Define $H = \Delta^{-1}(\hat{V} - V)$; then $\hat{V} = V + \Delta H$. We have $YH = 0$; multiplication from the left by Z shows that H belongs to $\mathcal{H}_0^{r \times n}$. Since H is of the form $H = \Delta^{-1}\tilde{V}$ with $\tilde{V} \in \mathcal{H}^{r \times n}$, and Δ only has zeros at λ_0 , it follows that in fact $H \in \mathcal{H}^{r \times n}$.

By multiplying both sides of (23) from the left by Z and from the right by T, one finds that $\Delta^{-1}VT = 0$. Furthermore, the biorthogonality relation [\(24\)](#page-9-4) means that there exists a matrix $H \in \mathcal{H}_0^{r \times r}$ such that $\Delta^{-1} = \Delta^{-1} V T Y \Delta^{-1} + H$, which implies that $V T Y \Delta^{-1} = I_r - \Delta H$. Since $\Delta(\lambda_0) = 0$ and $TY\Delta^{-1} \in \mathcal{H}_0^{n \times r}$, it follows that $V(\lambda_0)(TY\Delta^{-1})(\lambda_0) = I_r$, which implies that $V(\lambda_0)$ has full row rank. Canonicity of V now follows from Prop. [5.1.](#page-8-2) \Box

6 Extensions

[Kozlov and Maz'ya \(2013,](#page-18-6) Thm. A.10.1) present Keldysh's theorem with an additional statement concerning the uniqueness of solutions of the biorthogonality condition [\(24\)](#page-9-4), read as an equation for the left canonical matrix V when the right canonical matrix Y is given. They restrict the choice of left and right canonical matrices V and Y to the ones for which $\Delta^{-1}V$ and $Y\Delta^{-1}$ are polynomial matrices in χ_0^{-1} . This restriction is avoided in the version below. Also, the proof provides more detailed than is given in [Kozlov and Maz'ya \(2013](#page-18-6)). For notational convenience,

we look at the biorthogonality condition as an equation for the right canonical matrix when the left canonical matrix is given. The following simple observation will be used.

Lemma 6.1. Let $M \in \mathcal{H}_0^{p \times n}$ be left unimodular, and let $j \geq 1$. If $y \in \mathcal{H}_0^n$ satisfies $My \in \chi_0^j \mathcal{H}_0^p$, then $y \in \chi_0^j \mathcal{H}_0^n$.

Proof. Let $\tilde{M} \in \mathcal{H}_0^{n \times p}$ be a left inverse of M. For $y \in \mathcal{H}_0^n$ such that $My \in \chi_0^j \mathcal{H}_0^p$, we have $y = \tilde{M}(My) \in \chi_0^j \mathcal{H}_0^n.$ □

Theorem 6.2. For any left canonical matrix V for T at λ_0 , there exists a right canonical matrix Y for T at λ_0 such that the biorthogonality condition [\(24\)](#page-9-4) holds. This matrix is determined uniquely up to addition of a matrix of the form $H\Delta$ with $H \in \mathcal{H}^{n \times r}$. Formula [\(23\)](#page-9-5) for the principal part of T^{-1} at λ_0 holds with any such matrix.

Proof. Existence has already been shown in Thm. [5.6.](#page-9-6) Relation [\(24\)](#page-9-4) is equivalent to existence of a matrix $\tilde{H} \in \mathcal{H}_0^{r \times r}$ such that $\Delta^{-1} VTY = I + \tilde{H}\Delta$. To prove uniqueness of right canonical matrices Y that satisfy this equation, up to an additive term of the form $H\Delta$ with $H \in \mathcal{H}^{n \times r}$, we need to show that, if (i) $\Delta^{-1} V T Y = \tilde{H} \Delta$ where $\tilde{H} \in \mathcal{H}_0^{r \times r}$ and (ii) Y is the difference of two right canonical matrices, then $Y = H\Delta$ for some $H \in \mathcal{H}^{n \times r}$. Condition (i) can be restated in columnwise fashion as

$$
\Delta^{-1} V T y_i \in \chi_0^{m_i} \mathcal{H}_0^r \qquad (i = 1, \dots, r)
$$

where y_i denotes the *i*-th column of Y. Condition (ii) implies that $Ty_i \in \chi_0^{m_i} \mathcal{H}_0^n$ for all $i = 1, \ldots, r$. By Prop. [5.2](#page-8-1) and Lemma [6.1,](#page-11-0) combination of the two conditions leads to the conclusion that $y_i \in \chi_0^{m_i} \mathcal{H}_0^n$ for all $i = 1, \ldots, r$. In other words, $Y\Delta^{-1}$ does not have a singularity at λ_0 . Since Δ only has zeros at λ_0 and $Y \in \mathcal{H}^{n \times r}$, it follows that in fact $Y\Delta^{-1}$ has no singularities at all in the domain Ω ; i.e., $Y = H\Delta$ for some $H \in \mathcal{H}^{n \times r}$.

Finally, let Y be a right canonical matrix for which the biorthogonality condition [\(24\)](#page-9-4) holds. From the version of Thm. [5.6](#page-9-6) in which the left (rather than the right) canonical matrix is taken to be given, it follows that there exists a right canonical matrix \tilde{Y} that satisfies the relation T^{-1} = $\tilde{Y}\Delta^{-1}V$ as well as the biorthogonality condition. By what has been proved in the previous paragraph, there exists $H \in \mathcal{H}^{n \times n}$ such that $Y = \tilde{Y} + H\Delta$. From the fact that $T^{-1} \doteq \tilde{Y}\Delta^{-1}V$, it then follows that we also have $T^{-1} \doteq Y\Delta^{-1}V$. \Box

[Gohberg and Sigal \(1971\)](#page-18-7) give a principal part formula in which both the right and the left canonical matrix are supposed to be chosen in a particular way. In the version of [Keldysh \(1951](#page-18-11), [1971\)](#page-18-9), the right canonical matrix can be arbitrarily chosen, but the left canonical matrix needs to be selected in a special way to match the choice of the right canonical matrix. One may wonder whether it is possible to give a principal part formula in which both the left and the right canonical matrix can be arbitrarily chosen. This question is answered in the positive below. First, we show an alternative form of the biorthogonality condition.

Proposition 6.3. Let $V \in \mathcal{H}^{r \times n}$ and $Y \in \mathcal{H}^{n \times r}$ be left and right canonical matrices, respectively, for T at λ_0 . Then the matrix VTY is invertible, and the biorthogonality relation [\(24\)](#page-9-4) holds if and only if

$$
(VTY)^{-1} \doteq \Delta^{-1}.
$$
\n⁽²⁵⁾

Proof. It has been shown in Cor. [5.3](#page-8-3) that $\Delta^{-1} VTY$ is unimodular; hence, VTY is invertible. Suppose now that [\(24\)](#page-9-4) holds, so that there exists a matrix $H \in \mathcal{H}_0^{r \times r}$ such that $\Delta^{-1} V T Y \Delta^{-1} =$ $\Delta^{-1} + H$. From this one finds, upon multiplying from the left by $(\Delta^{-1} V T Y)^{-1}$, that

$$
\Delta^{-1} = (\Delta^{-1} V T Y)^{-1} \Delta^{-1} + (\Delta^{-1} V T Y)^{-1} H = (V T Y)^{-1} + (\Delta^{-1} V T Y)^{-1} H
$$

which implies that $\Delta^{-1} \doteq (VTY)^{-1}$.

Conversely, suppose there exists a matrix $H \in \mathcal{H}_0^{r \times r}$ such that $(VTY)^{-1} = \Delta^{-1} + H$. Multiplying from the right by Δ , one finds $I + H\Delta = (VTY)^{-1}\Delta = (\Delta^{-1}VTY)^{-1}$, so that $\Delta^{-1} V T Y = (I + H \Delta)^{-1}$. We can then write

$$
\Delta^{-1}VTY\Delta^{-1} = (I + H\Delta)^{-1}\Delta^{-1} = (I + H\Delta)^{-1}((I + H\Delta)\Delta^{-1} - H) = \Delta^{-1} - (I + H\Delta)^{-1}H
$$

which shows that $\Delta^{-1} V T Y \Delta^{-1} \doteq \Delta^{-1}$.

The following two lemmas will be used in the proof of the main novel result of this paper.

Lemma 6.4. If $M \in \mathcal{H}_0^{r \times n}$ is right unimodular, then there exists a unimodular matrix $U_R =$ $[U_R^1 \ U_R^2]$ in $\mathcal{H}_0^{n \times (r + (n-r))}$ such that $M[U_R^1 \ U_R^2] = [I_r \ 0].$

Proof. Since M is right unimodular, the matrix $M(\lambda_0) \in \mathbb{C}^{r \times n}$ has full row rank. Consequently, we can choose a matrix $\tilde{M}_0 \in \mathbb{C}^{(n-r)\times n}$ such that the matrix $\hat{M}_0 \in \mathbb{C}^{n\times n}$ obtained by stacking $M(\lambda_0)$ and \tilde{M}_0 is nonsingular. The matrix $\hat{M} \in \mathcal{H}_0^{n \times n}$ obtained by stacking M and \tilde{M}_0 is then unimodular. Define $U_R = \hat{M}^{-1} \in \mathcal{H}_0^{n \times n}$, and partition this matrix as $U_R = [U_R^1 \quad U_R^2]$ with $U_R^1 \in \mathcal{H}_0^{n \times r}$. We have

$$
\begin{bmatrix} M \\ \tilde{M}_0 \end{bmatrix} \begin{bmatrix} U_R^1 & U_R^2 \end{bmatrix} = \begin{bmatrix} I_r & 0 \\ 0 & I_{n-r} \end{bmatrix}
$$

so that in particular $M[U_R^1 \ U_R^2] = [I_r \ 0].$

Lemma 6.5. Let $M \in \mathbb{C}^{n \times n}$, and write dim ker $M = r$. Let $N_L \in \mathbb{C}^{(n-r)\times n}$ and $N_R \in \mathbb{C}^{n \times (n-r)}$ be matrices such that N_R has full column rank, ker $N_L \cap \text{im } M = \{0\}$, and $\text{im } N_R \cap \text{ker } M = \{0\}$. Then the matrix $N_L M N_R \in \mathbb{C}^{(n-r)\times(n-r)}$ is nonsingular.

Proof. Suppose $x \in \mathbb{C}^n$ is such that $N_L M N_R x = 0$. Then $M N_R x \in \text{im } M \cap \text{ker } N_L = \{0\}$, so that $MN_Rx = 0$. Consequently, $N_Rx \in \text{ker } M \cap \text{im } N_R = \{0\}$, so that $N_Rx = 0$ and hence $x = 0$. \Box

Theorem 6.6. Let $V \in \mathcal{H}^{r \times n}$ and $Y \in \mathcal{H}^{n \times r}$ be left and right canonical matrices, respectively, for T at λ_0 . Then $VTY \in \mathcal{H}^{r \times r}$ is invertible, and the principal part of T^{-1} at λ_0 is given by

$$
T^{-1} \doteq Y(VTY)^{-1}V.\tag{26}
$$

Proof. The claim concerning invertibility of VTY was proven already in Prop. [6.3.](#page-11-1) Since $\Delta^{-1}VT \in$ $\mathcal{H}_0^{r\times n}$ is right unimodular, it follows from Lemma [6.4](#page-12-1) that there exists a left unimodular matrix $\tilde{Y} \in \mathcal{H}_0^{n \times (n-r)}$ such that $\Delta^{-1} V T \tilde{Y} = 0$. In particular, $(\Delta^{-1} V T)(\lambda_0) \tilde{Y}(\lambda_0) = 0$. Because $\tilde{Y}(\lambda_0)$ has full column rank, it follows that im $\tilde{Y}(\lambda_0) = \ker (\Delta^{-1} V T)(\lambda_0)$. Then Prop. [5.2](#page-8-1) shows that $\lim \tilde{Y}(\lambda_0) \cap \ker T(\lambda_0) = \{0\}.$ Since $Y(\lambda_0)$ is a basis matrix for ker $T(\lambda_0)$, this means that the composite matrix $[Y(\lambda_0) \ \ \tilde{Y}(\lambda_0)] \in \mathbb{C}^{n \times n}$ is nonsingular, and hence the matrix $[Y \ \ \tilde{Y}] \in \mathcal{H}_0^{n \times n}$ is

 \Box

 \Box

unimodular. From the fact that $\Delta^{-1}VT\tilde{Y}=0$, it also follows that $VT\tilde{Y}=0$. Similarly, one can find a right unimodular matrix $\tilde{V} \in \mathcal{H}_0^{(n-r)\times n}$ such that $\text{im } T(\lambda_0) \cap \text{ker } \tilde{V}(\lambda_0) = \{0\}$, the composite matrix in $\mathcal{H}_0^{n \times n}$ formed from V and \tilde{V} is unimodular, and $\tilde{V}TY = 0$. It follows from Lemma [6.5](#page-12-2) that the matrix $\tilde{V}(\lambda_0)T(\lambda_0)\tilde{Y}(\lambda_0) \in \mathbb{C}^{(n-r)\times(n-r)}$ is nonsingular, so that $\tilde{V}T\tilde{Y} \in \mathcal{H}_0^{(n-r)\times(n-r)}$ is unimodular. From

$$
\begin{bmatrix} V \\ \tilde{V} \end{bmatrix} T \begin{bmatrix} Y & \tilde{Y} \end{bmatrix} = \begin{bmatrix} VTY & 0 \\ 0 & \tilde{V}T\tilde{Y} \end{bmatrix}
$$

we have

$$
T^{-1} = \begin{bmatrix} Y & \tilde{Y} \end{bmatrix} \begin{bmatrix} (VTY)^{-1} & 0 \\ 0 & (\tilde{V}T\tilde{Y})^{-1} \end{bmatrix} \begin{bmatrix} V \\ \tilde{V} \end{bmatrix} = Y(VTY)^{-1}V + \tilde{Y}(\tilde{V}T\tilde{Y})^{-1}\tilde{V}.
$$

This proves the claim, since $\tilde{Y}(\tilde{V}T\tilde{Y})^{-1}\tilde{V}$ is holomorphic in a neighborhood of λ_0 .

 \Box

The above result shows that both the condition $VTY = \Delta$ (as in Thm. [5.4,](#page-9-3) see Remark [5.5\)](#page-9-7) and the condition $(VTY)^{-1} \doteq \Delta^{-1}$ (as in Thm. [5.6](#page-9-6) and Thm. [6.2,](#page-11-2) see Prop. [6.3\)](#page-11-1) are sufficient for Keldysh's principal part formula [\(23\)](#page-9-5) to hold.

While formula [\(26\)](#page-12-3) is simple and nicely symmetric, in a computation one may prefer to use one of the alternative forms $Y(\Delta^{-1}VTY)^{-1}\Delta^{-1}V$ or $Y\Delta^{-1}(VTY\Delta^{-1})^{-1}V$. The reason for this is that the matrices $\Delta^{-1}VTY$ and $VTY\Delta^{-1}$ are unimodular as shown in Cor. [5.3,](#page-8-3) so that the inverses (or just the first terms in their power series expansions around λ_0 , as needed) can be computed relatively easily.

7 The semisimple case

The pole of T at λ_0 is said to be *semisimple* when it is of order 1; i.e., $\chi_0 T^{-1} \in \mathcal{H}_0^{n \times n}$. In several applications, it is of interest to characterize semisimplicity in terms of root spaces associated to T at λ_0 , and to have an expression for the residue of T^{-1} in the semisimple case. Conditions for semisimplicity are not discussed explicitly by [Keldysh \(1951](#page-18-11), [1971\)](#page-18-9), but he does prove [\(Keldysh](#page-18-9), [1971,](#page-18-9) p. 21) that the pole order is equal to the maximal multiplicity of root functions. Hence, a necessary and sufficient condition for semisimplicity is that all root functions have multiplicity 1. In terms of the partial multiplicities m_i defined in [\(6\)](#page-3-0), semisimplicity holds if and only if $m_i = 1$ for all $i = 1, \ldots, r$. A characterization in terms of the first two coefficients of the power series expansion of T around λ_0 can be derived as follows.

Proposition 7.1. The pole of T^{-1} at λ_0 is semisimple if and only if

$$
\ker T(\lambda_0) \cap (T'(\lambda_0))^{-1}(\operatorname{im} T(\lambda_0)) = \{0\}.
$$
 (27)

Proof. As noted below Thm. [3.1,](#page-3-5) the pole order of T^{-1} at λ_0 satisfies $s = \max_{i=1,\dots,r} m_i = m_1$. By Prop. [3.3,](#page-4-7) we have $m_1 = 1$ if and only if $\ell_2 = 0$, where $\ell_2 = \dim L_2$, and L_2 is defined by [\(8\)](#page-3-3). According to Prop. [3.4,](#page-4-8) the subspace L_2 consists of all vectors y^0 that satisfy (i) $T(\lambda_0)y^0 = 0$, and (ii) there exists $y^1 \in \mathbb{C}^n$ such that $T(\lambda_0)y^1 + T'(\lambda_0)y^0 = 0$. In other words, L_2 is given by the expression at the left hand side of [\(27\)](#page-13-0). □ Alternative characterizations follow from this via the following lemma.

Lemma 7.2. Let matrices $M, N \in \mathbb{C}^{n \times n}$ be given. Write $r = \dim \ker M$, and let $Z_L \in \mathbb{C}^{r \times n}$ and $Z_R \in \mathbb{C}^{n \times r}$ be basis matrices for the left and right null spaces of M respectively. The following statements are equivalent.

- (*i*) ker $M \cap N^{-1}(\text{im } M) = \{0\}.$
- (ii) ker $M \cap \text{ker } N = \{0\}$ and $\text{im } M \cap N(\text{ker } M) = \{0\}.$
- (iii) $\mathbb{C}^n = \text{im } M \oplus N(\text{ker } M)$.
- (iv) The matrix $Z_L N Z_R \in \mathbb{C}^{r \times r}$ is nonsingular.

Proof. Assume that (i) holds. Then ker $N \cap \text{ker } M = \{0\}$, since ker $N \subset N^{-1}(\text{im } M)$. Take $x \in$ im $M \cap N(\ker M)$, and let $y \in \ker M$ be such that $x = Ny$. We then have $y \in \ker M \cap N^{-1}(\text{im }M)$, so that $y = 0$ and hence $x = 0$. This proves that (i) implies (ii). Now, assume that (ii) holds. Take $x \in \text{ker } M \cap N^{-1}(\text{im } M)$. Then $Nx \in \text{im } M \cap N(\text{ker } M)$ so that $Nx = 0$. It follows that $x \in \text{ker } N \cap \text{ker } M$; hence, $x = 0$. This shows that (ii) implies (i). Equivalence of (ii) and (iii) follows by noting that

$$
\dim(\operatorname{im} M + N(\ker M)) = \dim \operatorname{im} M + \dim N(\ker M)) - \dim(\operatorname{im} M \cap N(\ker M))
$$

=
$$
\dim \operatorname{im} M + \dim \ker M - \dim(\ker N \cap \ker M) - \dim(\operatorname{im} M \cap N(\ker M))
$$

=
$$
n - (\dim(\ker N \cap \ker M) + \dim(\operatorname{im} M \cap N(\ker M))).
$$

Finally, observe that

$$
\dim \operatorname{im} Z_L N Z_R = \dim N(\operatorname{im} Z_R) - \dim(\ker Z_L \cap N(\operatorname{im} Z_R))
$$

=
$$
\dim N(\ker M) - \dim(\operatorname{im} M \cap N(\ker M))
$$

=
$$
r - (\dim(\ker N \cap \ker M) + \dim(\operatorname{im} M \cap N(\ker M))).
$$

Since the matrix Z_LNZ_R has size $r \times r$, it follows that (ii) and (iv) are equivalent.

□

Proposition 7.3. Let Z_L and Z_R be basis matrices for the left and right null spaces of $T(\lambda_0)$ respectively. Each of the following conditions is necessary and sufficient for semisimplicity of the pole of T^{-1} at λ_0 .

- (i) $\mathbb{C}^n = \operatorname{im} T(\lambda_0) \oplus T'(\lambda_0)(\ker T(\lambda_0)).$
- (*ii*) ker $T(\lambda_0) \cap \ker T'(\lambda_0) = \{0\}$ and $\operatorname{im} T(\lambda_0) \cap T'(\lambda_0)(\ker T(\lambda_0)) = \{0\}.$
- (iii) The matrix $Z_L T'(\lambda_0) Z_R \in \mathbb{C}^{r \times r}$ is nonsingular.

The necessity of condition (iii) can also be derived from Cor. [5.3.](#page-8-3) Indeed, in the semisimple case, the corollary implies that the matrix $\chi_0^{-1}Z_RTZ_L \in \mathcal{H}_0^{r\times r}$ is unimodular. Since $(\chi_0^{-1}Z_RTZ_L)(\lambda_0)$ = $Z_R T'(\lambda_0) Z_L$, it follows that $Z_R T'(\lambda_0) Z_L$ is nonsingular.

If semisimplicity holds, the principal part of T^{-1} at λ_0 consists only of the residue, denoted by $Res(T^{-1}; \lambda_0)$. A formula for the residue can be obtained from Keldysh's theorem by specializing the notions of left and right canonical matrices to the semisimple case. For a first-order pole, canonical matrices can be constructed using constant vectors only. A matrix in $\mathbb{C}^{r\times n}$ ($\mathbb{C}^{n\times r}$) is left (right) canonical if and only if it is a basis matrix for the left (right) null space of $T(\lambda_0)$. From Thm. [5.6,](#page-9-6) one then has the following corollary.

Corollary 7.4. If the pole of T^{-1} at λ_0 is semisimple, then, given any basis matrix $Z_R \in \mathbb{C}^{n \times r}$ for the right null space of $T(\lambda_0)$, there exists a uniquely determined basis matrix $Z_L \in \mathbb{C}^{r \times n}$ for the left null space of $T(\lambda_0)$ such that the residue of T^{-1} at λ_0 is given by $Z_R Z_L$. The matrix Z_L satisfies

$$
Z_L T'(\lambda_0) Z_R = I. \tag{28}
$$

Proof. Let Z_R be a basis matrix for ker $T(\lambda_0)$. Thm. [5.6](#page-9-6) implies that there exists a uniquely determined constant matrix Z_L such that $T^{-1} \doteq Z_R \Delta^{-1} Z_L = \chi_0^{-1} Z_R Z_L$. This shows that the residue of T^{-1} at λ_0 is given by $Z_R Z_L$. Since $Z_L T(\lambda_0) Z_R = 0$, it follows from the biorthogonality condition [\(24\)](#page-9-4) that

$$
\chi_0^{-1}I \doteq \chi_0^{-2}Z_LTZ_R \doteq \chi_0^{-1}Z_LT'(\lambda_0)Z_R.
$$

This implies [\(28\)](#page-15-0).

One can now reason as follows. Let a basis matrix $Z_R \in \mathbb{C}^{n \times r}$ for ker $T(\lambda_0)$ be given, and let $Z_L \in \mathbb{C}^{r \times n}$ be an *arbitrary* left basis matrix for the left null space of $T(\lambda_0)$. Let \tilde{Z}_L denote the left basis matrix obtained from Cor. [7.4.](#page-15-1) Since Z_L and \tilde{Z}_L are both left basis matrices, there exists a nonsingular matrix $M \in \mathbb{C}^{r \times r}$ such that $\tilde{Z}_L = M Z_L$. Equation [\(28\)](#page-15-0) then implies that $MZ_LT'(\lambda_0)Z_R = I$, and hence $M = (Z_LT'(\lambda_0)Z_R)^{-1}$. It follows that $\tilde{Z}_L = (Z_LT'(\lambda_0)Z_R)^{-1}Z_L$. This leads to the next corollary.

Corollary 7.5. Suppose that the pole of T^{-1} at λ_0 is semisimple. Let $Z_L \in \mathbb{C}^{r \times n}$ and $Z_R \in \mathbb{C}^{n \times r}$ be basis matrices for the left and the right null spaces of $T(\lambda_0)$ respectively. Then the matrix $Z_L T'(\lambda_0) Z_R \in \mathbb{C}^{r \times r}$ is nonsingular, and the residue of T^{-1} at λ_0 is given by

$$
Res(T^{-1}; \lambda_0) = Z_R (Z_L T'(\lambda_0) Z_R)^{-1} Z_L.
$$
\n(29)

Semisimplicity conditions and corresponding residue formulas have a somewhat complicated history. As noted before, conditions for semisimplicity are not given explicitly in [Keldysh \(1951](#page-18-11), [1971\)](#page-18-9). For polynomial matrices T with nonsingular leading coefficient matrix, [Lancaster \(1966](#page-18-1), Thm. 4.3) proves that a necessary condition for the pole of T^{-1} at λ_0 to be semisimple is that there exists basis matrices Z_L and Z_R for the left and right null spaces of $Z(\lambda_0)$ such that $Z_LT'(\lambda_0)Z_R =$ I, and that the residue is then given by $Z_R Z_L$. The proof uses a linearization technique.^{[8](#page-15-2)} Working in an infinite-dimensional setting and also using a linearization technique, [Howland \(1971\)](#page-18-17) shows that (i) and (ii) in Prop. [7.3](#page-14-0) are both necessary and sufficient for semisimplicity. This extends earlier work of [Steinberg \(1968](#page-18-18)). Neither Howland nor Steinberg gives a formula for the residue, and both do not refer to Keldysh's work. In the matrix setting, [Schumacher \(1985,](#page-18-19) [1986](#page-18-20)) gives

 8 Linearization is meant here in the sense that a linear pencil of matrices is used to represent the given polynomial matrix T . The assumption that the highest-order coefficient matrix of T is nonsingular is expedient in this context since it allows a straightforward generalization of the companion form, a standard tool in the theory of higher-order scalar differential equations.

condition (iii) as well as the residue formula [\(29\)](#page-15-3), with a derivation on the basis of the Smith form. The 1951 paper by Keldysh is referred to, unfortunately with the incorrect claim that the normalization factor $(Z_L T'(\lambda_0) Z_R)^{-1}$ is missing in that paper.^{[9](#page-16-0)} Condition (iii) also appears in Mennicken and Möller $(2003, Prop. 1.7.3)$,^{[10](#page-16-1)} with a derivation on the basis of Keldysh's theorem.

The pole of T at λ_0 is said to be *simple* when in addition to $s = 1$ we also have $r = 1$, i.e., the algebraic multiplicity of the pole is equal to 1. In this case, Keldysh's theorem Thm. [5.6](#page-9-6) implies that, for any right null vector y^0 of $T(\lambda_0)$, there exists a left null vector v^0 of $T(\lambda_0)$ such that $T^{-1} = \chi_0^{-1} y^0 v^0$. Since left null vectors are determined up to a nonzero scalar factor, it is immediate from the biorthogonality condition $\chi_0^{-2}v^0Ty^0 = \chi_0^{-1}$ that the vector v^0 required in Keldysh's theorem should be chosen such that $v^{0}T'(\lambda_{0})y^{0} = 1$. [Beyn \(2012\)](#page-17-6) gives this result and refers to it as Keldysh's theorem for simple eigenvalues.

The residue formula [\(29\)](#page-15-3) for semisimple poles is akin to the principal part formula [\(26\)](#page-12-3). One can derive the former from the latter in the following way. Let it be given that the pole of T at λ_0 is semisimple, and let Z_L and Z_R be basis matrices for the left and right null spaces of $T(\lambda_0)$ respectively. We have

$$
(Z_R T Z_L)^{-1} = (\chi_0^{-1} Z_R T Z_L)^{-1} \chi_0^{-1} \doteq (Z_R T'(\lambda_0) Z_L)^{-1} \chi_0^{-1}.
$$

Applying [\(26\)](#page-12-3), one arrives at the residue formula [\(29\)](#page-15-3).

As a final comment, we note that, while the semisimplicity condition in Prop. [7.3\(](#page-14-0)iii) and the residue formula [\(29\)](#page-15-3) can be obtained from Keldysh's theorem as shown above, a short derivation from the Smith form is possible. The proof below summarizes the argument in [Schumacher \(1985](#page-18-19), [1986\)](#page-18-20).

Proposition 7.6. Let Z_L and Z_R be basis matrices for the left and right null spaces of $T(\lambda_0)$ respectively. The pole of T at λ_0 is semisimple if and only if the matrix $Z_L T'(\lambda_0) Z_R \in \mathbb{C}^{r \times r}$ is nonsingular, and in that case the residue of T^{-1} at λ_0 is given by [\(29\)](#page-15-3).

Proof. Let $\tilde{T} = U_L T U_R$ where U_L , $U_R \in \mathcal{H}_0^{n \times n}$ are unimodular. Since $\tilde{T}(\lambda_0) = U_L(\lambda_0) T(\lambda_0) U_R(\lambda_0)$ where $U_L(\lambda_0)$ and $U_R(\lambda_0)$ are nonsingular, we have dim ker $\tilde{T}(\lambda_0) = \dim \ker T(\lambda_0) = r$. The pole of \tilde{T}^{-1} at λ_0 is semisimple if and only if the same holds for T^{-1} , and in that case we have

$$
Res(\tilde{T}^{-1}; \lambda_0) = (U_R(\lambda_0))^{-1} Res(T^{-1}; \lambda_0) (U_L(\lambda_0))^{-1}.
$$
\n(30)

Let $Z_L(\tilde{Z}_L) \in \mathbb{C}^{r \times n}$ and $Z_R(\tilde{Z}_R) \in \mathbb{C}^{n \times r}$ be basis matrices for the left and right null spaces of $T(\lambda_0)$ $(\tilde{T}(\lambda_0))$. Since $U_R(\lambda_0)\tilde{Z}_R$ and Z_R are both basis matrices for ker $T(\lambda_0)$, there exists a nonsingular matrix $M_R \in \mathbb{C}^{r \times r}$ such that $U_R(\lambda_0) \tilde{Z}_R = Z_R M_R$. Likewise, there exists a nonsingular

⁹This calls for an explanation. When writing the cited publications, I was unaware of [Keldysh \(1971](#page-18-9)) and had no direct access to [Keldysh \(1951](#page-18-11)), but I did know about the existence of the latter paper from [Gohberg and Sigal](#page-18-7) [\(1971](#page-18-7)). However, Gohberg and Sigal cite Keldysh's theorem in a version in which the biorthogonality condition is omitted. From the fact that the normalization factor does not appear in their paper, I wrongly concluded that it had also not been given by Keldysh. While the residue formula [\(29\)](#page-15-3) is not stated explicitly in [Keldysh \(1951](#page-18-11)), using the biorthogonality condition one can, as shown above, derive it from Keldysh's theorem, including the normalization factor.

¹⁰The proposition in fact states as a condition that there should exist basis matrices Z_L and Z_R such that $Z_L T'(\lambda_0) Z_R$ is nonsingular. Since left (right) basis matrices are defined uniquely up to left (right) multiplication by a nonsingular matrix, it is clear that the condition holds for a given pair of basis matrices (Z_L, Z_R) if and only if it holds for all such pairs.

matrix $M_L \in \mathbb{C}^{r \times r}$ such that $\tilde{Z}_L U_L(\lambda_0) = M_L Z_L$. Hence,

$$
\tilde{Z}_L \tilde{T}'(\lambda_0) \tilde{Z}_R = \tilde{Z}_L \big(U'_L(\lambda_0) T(\lambda_0) U_R(\lambda_0) + U_L(\lambda_0) T'(\lambda_0) U_R(\lambda_0) + U_L(\lambda_0) T(\lambda_0) U'_R(\lambda_0) \big) \tilde{Z}_R \n= M_L Z_L T'(\lambda_0) Z_R M_R.
$$

It follows that the matrix $\tilde{Z}_L \tilde{T}'(\lambda_0) \tilde{Z}_R \in \mathbb{C}^{r \times r}$ is nonsingular if and only if the same is true for $Z_L T'(\lambda_0) Z_R$. If this holds, then it follows by comparing [\(30\)](#page-16-2) with

$$
\tilde{Z}_R(\tilde{Z}_L\tilde{T}'(\lambda_0)\tilde{Z}_R)^{-1}\tilde{Z}_L = (U_R(\lambda_0))^{-1}Z_R(Z_LT'(\lambda_0)Z_R)^{-1}Z_L(U_L(\lambda_0))^{-1}
$$
(31)

that the residue formula [\(29\)](#page-15-3) is valid for T if and only if it is valid for \tilde{T} . To prove the claims of the proposition, it therefore suffices to verify them for a matrix in Smith form; basis matrices for left and right null spaces may be arbitrarily chosen. Therefore, suppose the matrix T is given by

$$
T(\lambda) = \begin{bmatrix} \Delta(\lambda) & 0 \\ 0 & I_{n-r} \end{bmatrix} \qquad (\lambda \in \Omega)
$$

with $\Delta = \text{diag}(\chi_0^{m_1}, \ldots, \chi_0^{m_r})$ as in [\(7\)](#page-3-4). Basis matrices for the left and right null spaces of T at λ_0 are given by

$$
Z_L = \begin{bmatrix} I_r & 0 \end{bmatrix}, \qquad Z_R = \begin{bmatrix} I_r \\ 0 \end{bmatrix}.
$$

We have $Z_L T'(\lambda_0) Z_R = \Delta'(\lambda_0)$. This matrix is nonsingular if and only if $m_i = 1$ for all $i = 1, ..., r$, i.e., if and only if the pole of T^{-1} at λ_0 is semisimple. If this holds, then $\Delta = \chi_0 I_r$ so that $Z_L T'(\lambda_0) Z_R = Z_L Z_R = I_r$, and

$$
T^{-1} = \begin{bmatrix} \chi_0^{-1} I_r & 0 \\ 0 & I_{n-r} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & I_{n-r} \end{bmatrix} + \chi_0^{-1} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} = \chi_0^{-1} Z_R (Z_L T'(\lambda_0) Z_R)^{-1} Z_L.
$$

References

- George E. Andrews. The Theory of Partitions. Cambridge University Press, Cambridge, UK, 1998.
- H. Bart, I. Gohberg, and M.A. Kaashoek. *Minimal Factorization of Matrix and Operator Func*tions. Birkhäuser, Basel, 1979.
- Brendan K. Beare and Alexis Akira Toda. The effect of reducible Markov modulation on tail probabilities in models of random growth. arXiv:2405.13556, 2024.
- Brendan K. Beare, Won-Ki Seo, and Alexis Akira Toda. Tail behavior of stopped Lévy processes with Markov modulation. Econometric Theory, 38:986–1013, 2022.
- Wolf-Jürgen Beyn. An integral method for solving nonlinear eigenvalue problems. Linear Algebra and Its Applications, 436:3839–3863, 2012.
- Robert F. Engle and Clive W.J. Granger. Co-integration and error correction: representation, estimation, and testing. Econometrica, 55:251–276, 1987.
- Mario Faliva and Maria Grazia Zoia. Dynamic Model Analysis: Advanced Matrix Methods and Unit-root Econometrics Representation Theorems (2nd ed.). Springer, Berlin, 2009.
- R.A. Frazer, W.J. Duncan, and A.R. Collar. Elementary Matrices and Some Applications to Dynamics and Differential Equations. Cambridge University Press, Cambridge, UK, 1938.
- I. Gohberg, M.A. Kaashoek, and F. van Schagen. On the local theory of regular analytic matrix functions. Linear Algebra and Its Applications, 182:9–25, 1993.
- I.C. Gohberg and E.I. Sigal. An operator generalization of the logarithmic residue theorem and the theorem of Rouché. Mathematics of the $USSR - Sbornik$, 13(4):603-625, 1971.
- Stefan Güttel and Françoise Tisseur. The nonlinear eigenvalue problem. Acta Numerica, 26:1–94, 2017.
- James S. Howland. Simple poles of operator-valued functions. Journal of Mathematical Analysis and Applications, 36:12–21, 1971.
- Søren Johansen. Representation of cointegrated autoregressive processes with application to fractional processes. Econometric Reviews, 28:121–145, 2008.
- M.V. Keldysh. On the characteristic values and characteristic functions of certain classes of nonselfadjoint equations. Doklady Akademii Nauk SSSR, 77:11–14, 1951. (Russian.) An English translation is available as an appendix in: A.S. Markus, Introduction to the Spectral Theory of Polynomial Operator Pencils. Translated by H.H. McFaden. Translations of Mathematical Monographs, vol. 71, American Mathematical Society, Providence, RI, 1988.
- M.V. Keldysh. On the completeness of the eigenfunctions of some classes of non-selfadjoint linear operators. Russian Mathematical Surveys, 26:15–44, 1971.
- Vladimir Kozlov and Vladimir Maz'ya. Differential Equations with Operator Coefficients. Springer, Berlin, 2013.
- S.G. Kreĭn and V.P. Trofimov. Holomorphic operator-valued functions of several complex variables. Functional Analysis and Its Applications, 3:330–331, 1969.
- Lisbeth la Cour. A parametric characterization of integrated vector autoregressive (VAR) processes. Econometric Theory, 14:187–199, 1998.
- Peter Lancaster. Lambda-matrices and Vibrating Systems. Pergamon Press, Oxford, UK, 1966.
- V.I. Macaev and Yu.A. Palant. Distribution of the spectrum of a polynomial operator pencil. Doklady Akademii Nauk Armjanskii SSR, 42:257–261, 1966. (Russian).
- C.C. MacDuffee. The Theory of Matrices. Chelsea, New York, 1946. Corrected reprint of first edition, Springer, 1933.
- Reinhard Mennicken and Manfred Möller. Root functions, eigenvectors, associated vectors and the inverse of a holomorphic operator function. Archiv der Mathematik, 42:455–463, 1984.
- Reinhard Mennicken and Manfred Möller. Non-self-adjoint Boundary Eigenvalue Problems. Elsevier, Amsterdam, 2003.
- Yu.A. Palant. On a method of testing for the multiple completeness of a system of eigenvectors and associated vectors of a polynomial operator bundle. Vestnik Khar'kovskogo Gosudarstvennogo Universiteta, (4) 34:1-13, 1970. (Russian.)
- J.M. Schumacher. Dynamic analysis of flexible spacecraft: some mathematical aspects. ESTEC Working Paper 1403, European Space Research and Technology Centre, Noordwijk, the Netherlands, February 1985.
- J.M. Schumacher. Residue formulas for meromorphic matrices. In C.I. Byrnes and A. Lindquist, editors, Computational and Combinatorial Methods in Systems Theory, pages 97–112. North-Holland, Amsterdam, 1986.
- Stanly Steinberg. Meromorphic families of compact operators. Archive for Rational Mechanics and Analysis, 31:372–379, 1968.
- V.P. Trofimov. The root subspaces of operators that depend analytically on a parameter. Matematicheskie Issledovaniya, 3(9):117–125, 1968. (Russian).