

LIE IDEALS IN PROPERLY INFINITE  $C^*$ -ALGEBRAS

HANNES THIEL

ABSTRACT. We show that every Lie ideal in a unital, properly infinite  $C^*$ -algebra is commutator equivalent to a unique two-sided ideal. It follows that the Lie ideal structure of such a  $C^*$ -algebra is concisely encoded by its lattice of two-sided ideals. This answers a question of Robert in this setting.

We obtain similar structure results for Lie ideals in unital, real rank zero  $C^*$ -algebras without characters. As an application, we show that every Lie ideal in a von Neumann algebra is related to a unique two-sided ideal, which solves a problem of Brešar, Kissin, and Shulman.

## 1. INTRODUCTION

A *Lie ideal* in a  $C^*$ -algebra  $A$  is a linear subspace  $L \subseteq A$  such that for every  $a \in A$  and  $x \in L$  the commutator  $[a, x] := ax - xa$  belongs to  $L$ . The study of Lie ideals in  $C^*$ -algebras has a long history, and we refer to Section 2 in [Mar10] for an overview of available results and directions of active research.

Immediate examples of Lie ideals in a  $C^*$ -algebra  $A$  are all subspaces contained in the center  $Z(A)$  and all subspaces containing the commutator subspace  $[A, A]$ . If  $A$  is unital and simple, then this does in fact describes all Lie ideals – a consequence of Herstein’s description of Lie ideals in simple rings [Her55]. Since the center of a unital, simple  $C^*$ -algebra  $A$  contains only scalar multiples of the identity, a subspace  $L$  is a Lie ideal in  $A$  if and only if  $L = \{0\}$ ,  $L = \mathbb{C}1$ , or  $[A, A] \subseteq L$ .

Pop [Pop02] showed that a unital  $C^*$ -algebra satisfies  $A = [A, A]$  if and only if  $A$  has no tracial states. Consequently, a simple, unital  $C^*$ -algebra  $A$  without tracial states has exactly three Lie ideals:  $\{0\}$ ,  $\mathbb{C}1$  and  $A$ . Cuntz and Pedersen [CP79] gave a description of the closure  $\overline{[A, A]}$  as the intersection of the kernels of all tracial states on  $A$ . In particular, a simple, unital  $C^*$ -algebra  $A$  with a unique tracial state has exactly four closed Lie ideals:  $\{0\}$ ,  $\mathbb{C}1$ ,  $\overline{[A, A]}$  and  $A$ . This was first observed by Marcoux and Murphy [MM98], generalizing the case of UHF-algebras, which was first proved by Marcoux [Mar95].

Ng and Robert [NR16] showed that  $[A, A]$  is closed in every pure, exact  $C^*$ -algebra. It follows that a simple, unital, exact, pure  $C^*$ -algebra  $A$  with a unique tracial state has exactly four Lie ideals:  $\{0\}$ ,  $\mathbb{C}1$ ,  $[A, A]$  and  $A$ . This applies for example to UHF-algebras, to irrational rotation algebras, and to the Jiang-Su algebra (where it was first shown by Ng [Ng12]). It also applies to many, and possibly all, simple, reduced group  $C^*$ -algebras. Indeed, if  $G$  is a countable, exact, acylindrically hyperbolic group with trivial finite radical and with the rapid decay property, then  $C_{\text{red}}^*(G)$  has strict comparison [AGEP24], and therefore is pure [APTV24], and has a unique tracial state [BKKO17].

---

*Date:* 23 December 2024.

*2020 Mathematics Subject Classification.* Primary 46L05, 46L10. Secondary 16W10, 17B60, 47B47.

*Key words and phrases.* Lie ideals,  $C^*$ -algebras, properly infinite, von Neumann algebras, commutators, square-zero elements.

The first named author was partially supported by the Knut and Alice Wallenberg Foundation (KAW 2021.0140).

Turning towards Lie ideals in non-simple  $C^*$ -algebras, it is clear that the structure of the lattice of two-sided ideals will play a role. A systematic study of the interplay between Lie and two-sided ideals was initiated by Brešar, Kissin, and Shulman [BKS08]. We first introduce the necessary notation. Following the usual convention, given subspaces  $M, N \subseteq A$ , we use  $[M, N]$  to denote the subspace of  $A$  generated by the set of commutators  $[x, y]$  for  $x \in M$  and  $y \in N$ . Further, given a subspace  $K \subseteq A$ , we set

$$T(K) := \{a \in A : [A, a] \subseteq K\}.$$

This ‘derived subspace’ was first considered by Herstein [Her55], and it is denoted by  $N(K)$  in [BKS08].

Given a two-sided ideal  $I \subseteq A$ , Brešar, Kissin, and Shulman observed that a subspace  $L \subseteq A$  is a Lie ideal whenever

$$[A, I] \subseteq L \subseteq T([A, I]).$$

We say that a Lie ideal  $L$  is *embraced by*  $I$  if it satisfies these inclusions. The Lie ideals that are embraced by some two-sided ideal are thought of as ‘unsurprising’. A major question is if there exist any Lie ideals that do not arise this way:

**Problem A** (Brešar, Kissin, and Shulman [BKS08, p. 74]). Given a  $C^*$ -algebra  $A$ , is every Lie ideal of  $A$  embraced by some two-sided ideal of  $A$ ?

To study this problem, two further notions were introduced in [BKS08]: Given a Lie ideal  $L \subseteq A$  and a two-sided ideal  $I \subseteq A$ , we say that  $L$  is *commutator equivalent to*  $I$  if

$$[A, I] = [A, L].$$

Further, we say that  $L$  is *related to*  $I$  if

$$[A, I] \subseteq L \subseteq T(I).$$

It is easy to see that these constitute a sufficient and a necessary conditions for embracement. Indeed, if  $L$  is commutator equivalent to  $I$ , then  $L$  is embraced by  $I$ . Further, if  $L$  is embraced by  $I$ , then  $L$  is related to  $I$ .

It was shown in [BKS08, Theorem 5.27] that every Lie ideal  $L$  in a  $C^*$ -algebra  $A$  is topologically commutator equivalent to a two-sided ideal  $I$  in the sense that  $[A, L]$  and  $[A, I]$  have the same closure. This leads to a description of all *closed* Lie ideals in terms of *closed* two-sided ideals. However, this does not show that all closed Lie ideals in a  $C^*$ -algebra are commutator equivalent to (let alone embraced by or related to) a two-sided ideal. For general non-closed Lie ideals even less is known, and Problem A as well as the following stronger problem remain open:

**Problem B** (Robert [Rob16, Question 1.12]). Given a  $C^*$ -algebra  $A$ , is every Lie ideal of  $A$  commutator equivalent to some two-sided ideal of  $A$ ?

The results mentioned at the beginning of the introduction show that every Lie ideal in a unital, simple  $C^*$ -algebra  $A$  is commutator equivalent to a two-sided ideal, which gives positive solutions to Problems A and B in this setting. Indeed, the only two-sided ideals are  $\{0\}$  and  $A$ , and a Lie ideal is commutator equivalent to  $\{0\}$  if and only if it is contained in  $Z(A)$ , while a Lie ideal it is commutator equivalent to  $A$  if and only if it contains  $[A, A]$ . (This uses that  $[A, A] = [A, [A, A]]$ , which follows for example from [GT24, Corollary 3.4].)

In [FMS82], Fong, Miers and Sourour showed that every Lie ideal in the von Neumann algebra of bounded, linear operators on a separable Hilbert space is embraced by a two-sided ideal. This was substantially extended by Brešar, Kissin, and Shulman, who showed in [BKS08, Theorem 5.19] that every Lie ideal in a von

Neumann algebra is commutator equivalent to a two-sided ideal, thereby providing positive solutions to Problems A and B for von Neumann algebras.

Our main result is a positive solution to Problem B for unital, properly infinite C\*-algebras.

**Theorem C (5.3).** Let  $L$  be a Lie ideal in a unital, properly infinite C\*-algebra  $A$ . Then  $L$  is commutator equivalent to the two-sided ideal  $I := A[A, L]A$ , and hence also embraced by  $I$ , and related to  $I$ . Further,  $I$  is the only two-sided ideal of  $A$  to which  $L$  is related.

The proof of Theorem C is inspired by a result of Marcoux [Mar95], who showed that if  $B$  is a unital C\*-algebra and  $n \geq 2$ , then every Lie ideal in the matrix algebra  $A = M_n(B)$  is related to a two-sided ideal in  $A$ . Our main technical advancement is a generalization of this result to unital C\*-algebras  $A$  that admit a unital \*-homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ .

The next result summarizes Theorems 2.7, 2.8, 3.8, and 3.10."

**Theorem D.** Let  $A$  be a unital C\*-algebra that admits a unital \*-homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ . The following statements hold:

- (1) Given a Lie ideal  $L \subseteq A$ , for the two-sided ideal  $I = A[A, L]A$ , we have

$$[A, I] = [A, [A, L]] \subseteq [A, L] \subseteq I, \quad \text{and} \quad [A, I] \subseteq L \subseteq T(I).$$

Further,  $I$  is the only two-sided ideal of  $A$  to which  $L$  is related.

- (2) A Lie ideal  $L \subseteq A$  is embraced by some two-sided ideal (which then necessarily is  $A[A, L]A$ ) if and only if  $L$  is commutator equivalent to some two-sided ideal, if and only if  $[A, L] = [A, [A, L]]$ .
- (3) Given a two-sided ideal  $I \subseteq A$ , we have

$$I = A[A, I]A, \quad \text{and} \quad [A, I] = [A, [A, I]].$$

- (4) There is a natural bijection between the set  $\mathcal{I}$  of two-sided ideals in  $A$  and the set

$$\mathcal{L} = \{L \subseteq A \text{ Lie ideal} : L = [A, L]\}.$$

given by the maps  $I \mapsto [A, I]$  and  $L \mapsto A[A, L]A$ .

We note that Theorem D applies to all unital, properly infinite C\*-algebras, which then yields Theorem C. It also applies to unital, real rank zero C\*-algebras without characters, and in particular to all von Neumann algebras with zero commutative summand, which we use to recover the solution to Problem B for von Neumann algebras from [BKS08]. Our approach also shows that a Lie ideal in a von Neumann algebra without commutative summand is related to a unique two-sided ideal, which solves [BKS08, Problem 5.21].

**Theorem E (6.1).** Let  $L$  be a Lie ideal in a von Neumann algebra  $M$ . Then  $L$  is commutator equivalent to the two-sided ideal  $I := M[M, L]M$ , and hence also embraced by  $I$ , and related to  $I$ .

If  $M$  has zero commutative summand, then  $I$  is the only two-sided ideal of  $M$  to which  $L$  is related.

**Conventions.** Given a C\*-algebra, we use  $A_+$  to denote its collection of positive elements, and we use  $\tilde{A}$  to denote the minimal unitization. The absolute value of an element  $x \in A$  is defined as  $|x| := (x^*x)^{\frac{1}{2}}$ .

**Acknowledgements.** The author thanks Leonel Robert for valuable comments on the Dixmier property for C\*-algebras, and Matej Brešar for useful feedback on an earlier version of this paper.

## 2. LIE IDEALS IN $C^*$ -ALGEBRAS WITH WEAKLY DIVISIBLE UNIT

In this section, we prove that every Lie ideal  $L$  in a  $C^*$ -algebra  $A$  that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$  is related to a unique two-sided ideal  $I$  in  $A$ ; see Theorem 2.7. We further show that  $L$  is commutator equivalent to  $I$  if and only if  $[A, L] = [A, [A, L]]$ ; see Theorem 2.8.

Following [PR04, Definition 5.1], a non-zero projection  $p$  in a  $C^*$ -algebra is said to be *weakly divisible of degree  $n$*  if there exists a unital  $*$ -homomorphism

$$M_{n_1}(\mathbb{C}) \oplus M_{n_2}(\mathbb{C}) \oplus \dots \oplus M_{n_r}(\mathbb{C}) \rightarrow pAp$$

for some natural numbers  $r \geq 1$  and  $n_1, \dots, n_r \geq n$ . The  $*$ -homomorphism may additionally be assumed to be injective.

As noted in [PR04, p.164], every matrix algebra  $M_n(\mathbb{C})$  for  $n \geq 2$  admits a unital (not necessarily injective)  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ , and it follows that a projection  $p$  weakly divisible of degree 2 if and only if there exists a unital (not necessarily injective)  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow pAp$ . We deduce that the unit in a  $C^*$ -algebra  $A$  is weakly divisible of degree 2 if and only if there exists an injective  $*$ -homomorphism

$$M_2(\mathbb{C}) \rightarrow A, \quad M_3(\mathbb{C}) \rightarrow A, \quad \text{or} \quad M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A.$$

In the first two cases, we get  $A \cong M_2(B)$  and  $A \cong M_3(C)$  for suitable unital  $C^*$ -algebras  $B$  and  $C$ , and for such matrix algebras it was shown by Marcoux [Mar95, Theorem 2.6] that every Lie ideal is related to a two-sided ideal (which is also unique, as shown in [BKS08, Corollary 4.18]).

Our main contribution to Theorem 2.7 is therefore in the case that  $A$  admits a unital (injective)  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$  but no unital  $*$ -homomorphism  $M_n(\mathbb{C}) \rightarrow A$  for  $n = 2$  or  $n = 3$  (or actually any  $n \geq 2$ ). The Cuntz algebra  $\mathcal{O}_\infty$  is an important such example.

Our approach is inspired by the methods developed by Marcoux [Mar95] to describe Lie ideals in matrix algebras, as well as the techniques of Brešar, Kissin, and Shulman [BKS08, Section 3] to study Lie ideals in algebras that decompose with respect to an idempotent. However, the result does not simply follow by combining these methods, for example by applying [BKS08, Theorem 3.4] for a suitable idempotent in the  $C^*$ -algebra. Specifically, given a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , we do not decompose  $A$  with respect to the projection  $p$  given by the image of the unit of  $M_2(\mathbb{C})$ , in which case  $pAp + (1-p)A(1-p)$  is a direct sum of matrix algebras (whence its Lie ideals are related to two-sided ideals), but it is unclear if (3.17) in [BKS08, Theorem 3.4] holds (or if  $p$  is locally cyclic in the sense of [BKS08, Definition 3.5]). Instead, we use a ‘mixed’ decomposition with respect to a projection  $g$  that is given by the image of the sum of a rank-one projection in  $M_2(\mathbb{C})$  with a rank-one projection in  $M_3(\mathbb{C})$ , as explained in Paragraph 2.1. In this case, one can show that  $g$  is locally cyclic, which is the idea underlying the proof that  $A_{12}LA_{12} \subseteq [A, [A, L]]$  in Lemma 2.2. On the other hand, the algebra  $gAg + (1-g)A(1-g)$  is not a sum of matrix algebras and it is therefore not immediate that its Lie ideals are related to two-sided ideals.

**2.1.** Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism

$$\varphi: M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A.$$

Let  $(e_{ij})_{i,j=1,2}$  denote the image of matrix units in  $M_2(\mathbb{C})$  under  $\varphi$ , and similarly let  $(f_{ij})_{i,j=1,2,3}$  denote the image of matrix units in  $M_3(\mathbb{C})$  under  $\varphi$ . We do not assume that  $\varphi$  is injective, and therefore the  $e_{ij}$  or  $f_{ij}$  may be zero.

Set

$$g := e_{11} + f_{11}, \quad \text{and} \quad h := e_{22} + f_{22} + f_{33}.$$

Then  $1 = g + h$ , and we view elements in  $A$  as operator matrices with respect to the decomposition induces by  $g$  and  $h$ . More formally, we set

$$A_{11} := gAg, \quad A_{12} := gAh, \quad A_{21} := hAg, \quad \text{and} \quad A_{22} := hAh,$$

and every element  $a$  in  $A$  can be written uniquely as a sum  $a = a_{11} + a_{12} + a_{21} + a_{22}$  with  $a_{ij} \in A_{ij}$  for  $i, j = 1, 2$ .

**Lemma 2.2.** *Retain the notation from Paragraph 2.1, and let  $L \subseteq A$  be a Lie ideal. Then the sets*

$$A_{11}LA_{22}, \quad A_{22}LA_{11}, \quad A_{12}LA_{12}, \quad \text{and} \quad A_{21}LA_{21}$$

are all contained in  $[A, [A, L]]$ , and thus also in  $[A, L]$  and in  $L$ .

*Proof.* We verify that  $A_{11}LA_{22} \subseteq [A, [A, L]]$ . We first show that  $gLh \subseteq L$ . To see this, let  $x \in L$ . Then

$$gx - xg = [g, x] \in [A, L], \quad \text{and} \quad gx - 2gxx + xg = [g, [g, x]] \in [A, [A, L]] \subseteq [A, L].$$

Adding these expressions, we get

$$2gxh = 2gx - 2gxx \in [A, L],$$

and using that  $hg = 0$  we get

$$gxh = [\frac{1}{2}g, gxh] \in [A, L],$$

We have shown that  $gLh \subseteq L$ .

Now, given  $x \in L$  and  $a, b \in A$ , we have

$$gagxhbh = [[gag, gxh], hbh] \in [[A, L], A] = [A, [A, L]].$$

We have shown that  $A_{11}LA_{22} \subseteq [A, [A, L]]$ .

Analogously, one verifies that  $A_{22}LA_{11} \subseteq [A, [A, L]]$ .

We verify that  $A_{12}LA_{12} \subseteq [A, [A, L]]$ . Given  $v \in A_{12}$  and  $y \in L$ , we have

$$(1) \quad v xv = [-\frac{1}{2}v, [v, y]] \in [A, [A, L]].$$

Set

$$v_1 := e_{12} + f_{12}, \quad w_1 := e_{21} + f_{21}, \quad v_2 := e_{12} + f_{13}, \quad \text{and} \quad w_2 := e_{21} + f_{31}.$$

We have

$$v_1 w_1 = v_2 w_2 = e_{11} + f_{11} = g, \quad w_1 v_1 = e_{22} + f_{22}, \quad \text{and} \quad w_2 v_2 = e_{22} + f_{33}.$$

Now, given  $x \in L$  and  $a, b \in A$ , set

$$b' = b(\frac{1}{2}e_{22} + f_{22} + f_{33}).$$

Then  $b'$  belongs to  $gAh$  and we have

$$b = b'(2e_{22} + f_{22} + f_{33}).$$

We have already proved that  $gAgLhAh \subseteq L$ , which gives

$$w_1 a x b' w_1 \in L, \quad \text{and} \quad w_2 a x b' w_2 \in L.$$

Applying (1) for  $y = w_1 a x b' w_1$  and  $v = v_1$ , we get

$$a x b' (e_{22} + f_{22}) = v_1 (w_1 a x b' w_1) v_1 \in [A, [A, L]].$$

Analogously, we obtain that

$$a x b' (e_{22} + f_{33}) = v_2 (w_2 a x b' w_2) v_2 \in [A, [A, L]],$$

and consequently

$$a x b = a x b' (2e_{22} + f_{22} + f_{33}) \in [A, [A, L]].$$

We have shown that  $A_{12}LA_{12} \subseteq [A, [A, L]]$ .

Analogously, one verifies that  $A_{21}LA_{21} \subseteq [A, [A, L]]$ . □

**Lemma 2.3.** *Retain the notation from Paragraph 2.1, and let  $L \subseteq A$  be a Lie ideal. Then*

$$AgLhA = AhLgA.$$

*Proof.* Using at the first step that  $A = AgA = AhA$  since  $g$  and  $h$  are full, and applying at the third step that  $hAgLhAg \subseteq L$  by Lemma 2.2, we get

$$AgLhA = AhAgLhAgA = Ah(hAgLhAg)gA \subseteq AhLgA.$$

Analogously, we have

$$AhLgA = AgAhLgAhA = Ag(gAhLgAh)hA \subseteq AgLhA,$$

as desired.  $\square$

**Lemma 2.4.** *Retain the notation from Paragraph 2.1, and let  $L \subseteq A$  be a Lie ideal. Then*

$$[A, L] \subseteq AgLhA.$$

*Proof.* To simplify notation, we set  $I = AgLhA$ . We have

$$[A, gLh] \subseteq AgLh + gLhA \subseteq I.$$

Further, using Lemma 2.3 at the third step, we have

$$[A, hLg] \subseteq AhLg + hLgA \subseteq AhLgA = AgLhA = I.$$

Thus, given  $x \in L$ , we have  $[A, gxh + hxg] \subseteq I$ . Note that  $gxh, hxg \in L$  by Lemma 2.2. It follows that

$$gxg + hxx = x - gxh + hxg \in L.$$

We need to show that  $[A, gxg + hxx] \subseteq I$ . Using additivity of the Lie bracket and that  $I$  is an additive subgroup, it suffices to verify that  $[A_{ij}, gxg + hxx] \subseteq I$  for  $i, j = 1, 2$ .

*We show that  $[A_{12}, gxg + hxx] \subseteq I$ .* Given  $a \in A$ , we have

$$[gah, gxg + hxx] = gahxx - xgag.$$

Using that  $gxg + hxx \in L$ , we have  $[gah, gxg + hxx] \in L$ , and thus

$$[gah, gxg + hxx] \in L \cap gAh = gLh \subseteq AgLhA = I.$$

We have shown that  $[A_{12}, gxg + hxx] \subseteq I$ .

Similarly, and using also Lemma 2.3 at the last step, we get

$$[hag, gxg + hxx] = hagxg - hxhag \in L \cap hAg = hLg \subseteq AhLgA = I$$

for every  $a \in A$ . This shows that  $[A_{21}, gxg + hxx] \subseteq I$ .

*We show that  $[A_{11}, gxg + hxx] \subseteq I$ .* Given  $a \in A$ , set  $y = gagxg - gxgag$ . Then

$$y = gagxg - gxgag = [gag, gxg + hxx] \in [A, L] \subseteq L.$$

Then

$$y(e_{12} + f_{12}) = [y, e_{12} + f_{12}] \in [L, A] \subseteq L$$

and thus

$$y(e_{12} + f_{12}) \in L \cap gAh = gLh \subseteq I.$$

We deduce that

$$y = y(e_{12} + f_{12})(e_{21} + f_{21}) \in IA = I.$$

*We show that  $[A_{22}, gxg + hxx] \subseteq I$ .* Given  $b \in A$ , set  $z = hbxh - hxxbh$ . Then

$$z = hbxh - hxxbh = [hbh, gxg + hxx] \in L \cap A_{22}.$$

Then

$$(e_{12} + f_{12})z = [e_{12} + f_{12}, z] \in L \cap gAh \subseteq I, \quad \text{and} \quad f_{13}z = [f_{13}, z] \in I.$$

We deduce that

$$z = (e_{21} + f_{21})(e_{12} + f_{12})z + f_{31}f_{13}z \in AI + AI = I,$$

as desired.  $\square$

**Proposition 2.5.** *Retain the notation from Paragraph 2.1, and let  $L \subseteq A$  be a Lie ideal. Then*

$$A[A, L] = A[A, L]A = AgLhA = AhLgA.$$

*Proof.* In general, if  $M \subseteq A$  is a Lie ideal, then  $AMA = AM$ ; see, for example, [GLT24, Lemma 3.2]. Since  $[A, L]$  is a Lie ideal, we get  $A[A, L] = A[A, L]A$ .

By Lemma 2.4, we have  $[A, L] \subseteq AgLhA$ , and thus  $A[A, L]A \subseteq AgLhA$ . By Lemma 2.3, we have  $AgLhA = AhLgA$ . Finally, by Lemma 2.2, we have  $gLh \subseteq [A, L]$ , and thus  $AgLhA \subseteq A[A, L]A$ .  $\square$

**Lemma 2.6.** *Retain the notation from Paragraph 2.1, and let  $L \subseteq A$  be a Lie ideal, and set  $I = A[A, L]A$ . Then*

$$[A, I] \subseteq [A, [A, L]].$$

*Proof.* For each  $i, j = 1, 2$ , set  $I_{ij} := I \cap A_{ij}$ . Since  $I$  is a two-sided ideal, we have  $I = I_{11} + I_{12} + I_{21} + I_{22}$ . By Proposition 2.5, we have  $I = AgLhA$ , and thus

$$I = AgLhA = \underbrace{gAgLhAg}_{I_{11}} + \underbrace{gAgLhAh}_{I_{12}} + \underbrace{hAgLhAg}_{I_{21}} + \underbrace{hAgLhAh}_{I_{22}}.$$

By additivity of the Lie bracket and since  $[A, L]$  is an additive subgroup, it suffices to show that  $[A, I_{ij}] \subseteq [A, [A, L]]$  for each  $i, j = 1, 2$ .

By Lemma 2.2, we have

$$I_{12} = gAgLhAh \subseteq [A, L], \quad \text{and} \quad I_{21} = hAgLhAg \subseteq [A, L],$$

and thus  $[A, I_{12}] \subseteq [A, [A, L]]$  and  $[A, I_{21}] \subseteq [A, [A, L]]$ .

We show that  $[A, I_{11}] \subseteq [A, [A, L]]$ . It suffices to verify that  $[A_{ij}, I_{11}] \subseteq [A, [A, L]]$  for each  $i, j = 1, 2$ . Using that  $hg = 0$ , and using Lemma 2.2 at the last step, we get

$$[A_{12}, I_{11}] = [gAh, gAgLhAg] = -(gAgLhAg)(gAh) = gAgLhAgAh \subseteq [A, [A, L]].$$

Similarly, we have

$$[A_{21}, I_{11}] = [hAg, gAgLhAg] = (hAg)(gAgLhAg) = hAgAgLhAg \subseteq [A, [A, L]].$$

We also have  $[A_{22}, I_{11}] = \{0\} \subseteq [A, [A, L]]$ .

Further, using Lemma 2.2 at the last step, we have

$$I_{11} = (gAgLh)Ag \subseteq [gAgLh, Ag] + Ag(gAgLh) \subseteq [A, L] + AgLh.$$

Using again Lemma 2.2 at the last step, we get

$$\begin{aligned} [A_{11}, I_{11}] &\subseteq [A_{11}, [A, L]] + [A_{11}, AgLh] \subseteq [A, [A, L]] + [gAg, AgLh] \\ &\subseteq [A, [A, L]] + gAgAgLh \subseteq [A, [A, L]]. \end{aligned}$$

We have verified that  $[A, I_{11}] \subseteq [A, [A, L]]$ .

Analogously, one shows that  $[A, I_{22}] \subseteq [A, [A, L]]$ , which completes the proof.  $\square$

The next result verifies statement (1) in Theorem D.

**Theorem 2.7.** *Let  $A$  be a unital C\*-algebra that admits a unital \*-homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $L \subseteq A$  be a (not necessarily closed) Lie ideal. Then, for the two-sided ideal  $I := A[A, L]A$ , we have*

$$(2) \quad [A, I] = [A, [A, L]] \subseteq [A, L] \subseteq I, \quad \text{and} \quad [A, I] \subseteq L \subseteq T(I).$$

*Further,  $I$  is the only two-sided ideal of  $A$  to which  $L$  is related.*

*Proof.* Using Lemma 2.4 at the first step, and Proposition 2.5 at the second step, we have

$$[A, L] \subseteq AgLhA = I.$$

This implies that  $[A, [A, L]] \subseteq [A, I]$ , and the converse inclusion  $[A, I] \subseteq [A, [A, L]]$  is shown in Lemma 2.6. Using that  $L$  is a Lie ideal, we have  $[A, L] \subseteq L$ , and therefore  $[A, [A, L]] \subseteq [A, L]$ . This shows the left chain of inclusions in (2).

The inclusion  $[A, I] \subseteq L$  follows using that  $[A, I] \subseteq [A, L]$  and  $[A, L] \subseteq L$ . Finally, since  $[A, L] \subseteq I$ , we have  $L \subseteq T(I)$ .

To show uniqueness of  $I$ , let  $J$  be a two-sided ideal that is related to  $L$ , that is, such that

$$[A, J] \subseteq L \subseteq T(J).$$

Then  $[A, L] \subseteq J$ , and since  $I$  is the two-sided ideal generated by  $[A, L]$  we get  $I \subseteq J$ .

Applying Lemma 2.2 at the first step for  $J$ , considered as a Lie ideal, we get

$$gJh \subseteq [A, J] \subseteq L,$$

and thus

$$gJh \subseteq L \cap gAh = gLh.$$

Since  $g$  and  $h$  are full, we have  $A = AgA$  and  $A = AhA$ , and consequently

$$J = AJA = AgAJAhA = AgJhA \subseteq AgLhA = I.$$

In conclusion, we get  $I = J$ , as desired.  $\square$

The next result verifies statement (2) in Theorem D.

**Theorem 2.8.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $L \subseteq A$  be a (not necessarily closed) Lie ideal. Then the following statements are equivalent:*

- (1) *The Lie ideal  $L$  is commutator equivalent to the two-sided ideal  $A[A, L]A$ .*
- (2) *The Lie ideal  $L$  is embraced by some two-sided ideal.*
- (3) *We have  $[A, L] = [A, [A, L]]$ .*

*Proof.* It is clear that (1) implies (2). Assuming (2), let us show that (3) holds. Set  $I := A[A, L]A$ , and let  $J \subseteq A$  be some two-sided ideal that embraces  $L$ . Then  $L$  is also related to  $J$ . It follows that  $I = J$ , since  $I$  is the unique two-sided ideal to which  $L$  is related, by Theorem 2.7. It follows that  $L \subseteq T([A, I])$ . Using this at the first step, and using Theorem 2.7 at the second step, we get

$$[A, L] \subseteq [A, I] = [A, [A, L]],$$

which verifies (3).

Assuming (3), let us show that (1) holds. Using that assumption at the first step, and Theorem 2.7 at the second step, we get

$$[A, L] = [A, [A, L]] = [A, I],$$

which verifies (1).  $\square$

The next result is essentially contained in [PR04, Proposition 5.7]. We only show how to remove the separability assumption. A *character* on a  $C^*$ -algebra  $A$  is one-dimensional, irreducible representation, that is, a surjective  $*$ -homomorphism  $A \rightarrow \mathbb{C}$ .

**Proposition 2.9** (Perera, Rørdam). *Let  $A$  be a unital  $C^*$ -algebra of real rank zero. Then  $A$  has no characters if and only if  $A$  admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ .*



*Proof.* The backwards implication is clear. To show the forward implication, assume that  $A$  has no characters. By [KR15, Lemma 3.5], there exists a *separable* sub- $C^*$ -algebra  $A_0 \subseteq A$  containing the unit of  $A$ . Using that real rank zero is separably inheritable (see [Bla06, Section II.8.5]), we obtain a *separable* sub- $C^*$ -algebra  $B \subseteq A$  of real rank zero with  $A_0 \subseteq B$ . It follows that  $B$  has no characters, which allows us to apply [PR04, Proposition 5.7] for  $B$ . We obtain a unital  $*$ -homomorphism from  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$  to  $B$ , and thus also to  $A$ .  $\square$

**Example 2.10.** Let  $A$  be a unital  $C^*$ -algebra of real rank zero that has no characters. Then Theorem 2.7 applies to  $A$ . In particular, every Lie ideal in  $A$  is related to a (unique) two-sided ideal.

We expect that the following question has a positive answer.

**Question 2.11.** Let  $A$  be a unital  $C^*$ -algebra that has no characters. Is every Lie ideal in  $A$  related to a (unique) two-sided ideal?

### 3. TWO-SIDED IDEALS IN $C^*$ -ALGEBRAS WITH WEAKLY DIVISIBLE UNIT

In this section, we prove that every two-sided ideal  $I$  in a  $C^*$ -algebra  $A$  that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$  satisfies  $I = A[A, I]A$ , and we describe  $[A, I]$  as the subspace spanned by certain square-zero elements in  $I$ ; see Theorem 3.8. If  $I$  is a Dixmier ideal (in which case there is a well-defined ideal  $I^{\frac{1}{2}}$  satisfying  $(I^{\frac{1}{2}})^2 = I$ ), then  $I = [A, I] + [A, I^{\frac{1}{2}}]^2$ ; see Theorem 3.14. In this case every element in  $I$  is a sum of square-zero elements in  $I$  and of products of pairs of square-zero elements in  $I^{\frac{1}{2}}$ .

The next definition generalizes the notion of orthogonally factorizable square-zero elements in a ring ([GT23, Definition 5.1]) to a relative notion with respect to a two-sided ideal.

**Definition 3.1.** Let  $I$  be a two-sided ideal in a  $C^*$ -algebra  $A$ . We let  $N_2(I)$  denote the square-zero elements in  $I$ , that is,

$$N_2(I) := \{x \in I : x^2 = 0\}.$$

We further set

$$FN_2(A, I) := \{x \in I : \text{there exist } a, b \in A, y \in I \text{ with } x = ayb \text{ and } ba = 0\}.$$

**Lemma 3.2.** Let  $I \subseteq A$  be a two-sided ideal in a  $C^*$ -algebra, and let  $x \in FN_2(A, I)$ . Then there exist  $a, b \in A_+$  and  $y \in FN_2(A, I) \subseteq I$  such that

$$x = ayb, \quad \text{and} \quad ab = ba = 0.$$

*Proof.* By definition, there are  $c, d \in A$  and  $z \in I$  such that

$$x = czd, \quad \text{and} \quad dc = 0.$$

Let  $c = v|c|$  and  $d = w|d|$  the polar decompositions in  $A^{**}$  and set

$$a := |c^*|^{\frac{1}{2}}, \quad b := |d|^{\frac{1}{2}}, \quad \text{and} \quad y := |c^*|^{\frac{1}{2}}vxw|d|^{\frac{1}{2}}.$$

By properties of the polar decomposition (see, for example, [GKT23, Proposition 2.1]), we have  $|c^*|^{\frac{1}{2}}v, w|d|^{\frac{1}{2}} \in A$ , and thus  $y \in A$ . Further, we have  $x = ayb$ .

Since  $dc = 0$ , we have  $(d^*d)^n(cc^*)^m = 0$  for every  $m, n \geq 1$ . It follows that  $p(d^*d)q(cc^*) = 0$  for every polynomials  $p$  and  $q$  with vanishing constant terms. Using functional calculus, we get  $f(d^*d)g(cc^*) = 0$  for every continuous functions  $f, g: \mathbb{R} \rightarrow \mathbb{R}$  with  $f(0) = g(0) = 0$ . In particular, we get  $|d|^{\frac{1}{2}}|c^*|^{\frac{1}{2}} = 0$ . Thus, we have  $ba = 0$ , and then also  $ab = 0$ .

To show that  $y$  belongs to  $\text{FN}_2(A, I)$ , consider the decomposition

$$y = |c^*|^{\frac{1}{4}} (|c^*|^{\frac{1}{4}} v x w |d|^{\frac{1}{4}}) |d|^{\frac{1}{4}}.$$

Then  $|c^*|^{\frac{1}{4}} v x w |d|^{\frac{1}{4}}$  belongs to  $I$ , and we have also seen above that  $|c^*|^{\frac{1}{4}}$  and  $|d|^{\frac{1}{4}}$  have zero product, whence  $y \in \text{FN}_2(A, I)$ .  $\square$

The next result is analogous to [GT23, Lemma 5.2].

**Lemma 3.3.** *Let  $I \subseteq A$  be a two-sided ideal in a  $C^*$ -algebra. Then*

$$\text{FN}_2(A, I) \subseteq [A, \text{FN}_2(A, I)] \subseteq [A, [A, I]] \subseteq [A, I], \quad \text{and} \quad \text{FN}_2(A, I) \subseteq \text{N}_2(I).$$

*Proof.* Let  $x \in \text{FN}_2(A, I)$ . By Lemma 3.2, we can pick  $a, b \in A_+$  and  $y \in I$  such that  $x = ayb$  and  $ba = 0$ . As in the proof of Lemma 3.2, we see that  $ba^{\frac{1}{2}} = 0$ . It follows that  $a^{\frac{1}{2}}yb \in \text{FN}_2(A, I)$  and then

$$x = ayb = [a^{\frac{1}{2}}, a^{\frac{1}{2}}yb] \in [A, \text{FN}_2(A, I)].$$

Since  $\text{FN}_2(A, I)$  is a subset of  $I$ , we have  $[A, \text{FN}_2(A, I)] \subseteq [A, I]$ . Combining these results, and using at the last step that  $[A, I]$  is a Lie ideal, we get

$$\text{FN}_2(A, I) \subseteq [A, \text{FN}_2(A, I)] \subseteq [A, [A, \text{FN}_2(A, I)]] \subseteq [A, [A, I]] \subseteq [A, I].$$

Given  $x = ayb \in \text{FN}_2(A, I)$ , with  $a, b \in A$  and  $y \in I$  such that  $ba = 0$ , we have

$$x^2 = (ayb)(ayb) = 0,$$

and thus  $x \in \text{N}_2(I)$ .  $\square$

**Remark 3.4.** We note that Definition 3.1 and [GT23, Definition 5.1] agree when considering the ideal  $I = A$  in a  $C^*$ -algebra  $A$ . Indeed, an element  $x$  in  $A$  belongs to  $\text{FN}_2(A)$  in the sense of [GT23, Definition 5.1] if there exist  $s, t \in A$  such that  $x = st$  and  $ts = 0$ , while  $x$  belongs to  $\text{FN}_2(A, A)$  as defined in Definition 3.1 if there exist  $a, y, b \in A$  such that  $x = ayb$  and  $ba = 0$ .

Given a factorization  $x = ayb$  with  $ba = 0$ , we can use  $s = a$  and  $t = yb$ . Conversely, given a factorization  $x = st$  with  $ts = 0$ , it follows that  $|t|^{\frac{1}{2}}s = 0$ . We consider the polar decomposition  $t = v|t|$  in  $A^{**}$ . Then  $v|t|^{\frac{1}{2}}$  belongs to  $A$ , and we can use  $a = s$ ,  $y = v|t|^{\frac{1}{2}}$  and  $b = |t|^{\frac{1}{2}}$ .

Given a subset  $V$  in a complex vector space, we use  $\text{span}_{\mathbb{C}} V$  to denote the linear subspace generated by  $V$ . Following [GT23, Definition 2.6], we say that a  $\mathbb{C}$ -algebra  $A$  is *zero-product balanced* if for all  $a, b, c \in A$  we have

$$ab \otimes c - a \otimes bc \in \text{span}_{\mathbb{C}} \{u \otimes v \in A \otimes_{\mathbb{C}} A : uv = 0\} \subseteq A \otimes_{\mathbb{C}} A.$$

We see that the class of  $C^*$ -algebras considered in this section are zero-product balanced:

**Lemma 3.5.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ . Then  $A$  is zero-product balanced as a  $\mathbb{C}$ -algebra.*

*Proof.* As in Paragraph 2.1, let  $(e_{ij})_{i,j=1,2}$  and  $(f_{ij})_{i,j=1,2,3}$  be the images in  $A$  of matrix units in  $M_2(\mathbb{C})$  and  $M_3(\mathbb{C})$ . Then  $e_{11} + f_{11}$  and  $e_{22} + f_{22}$  are two full orthogonal projections in  $A$ . This allows us to apply [Rob16, Corollary 3.8], which shows that  $A$  is generated by its projections as a  $\mathbb{C}$ -algebra, and thus is zero-product balanced by [GT23, Proposition 3.6].  $\square$

A two-sided ideal  $I \subseteq A$  in a  $C^*$ -algebra is said to be *idempotent* if  $I = I^2$ , that is, if every element in  $I$  is of the form  $x_1y_1 + \dots + x_ny_n$  for some  $n \in \mathbb{N}$  and suitable  $x_j, y_j \in I$ .

**Proposition 3.6.** *Let  $A$  be a  $C^*$ -algebra that is zero-product balanced as a  $\mathbb{C}$ -algebra, and let  $I \subseteq A$  be a two-sided ideal such that  $I = AIA$  (for example,  $A$  is unital or  $I$  is idempotent). Then*

$$[A, I] = [A, [A, I]] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I).$$

*Proof.* The inclusions ‘ $\supseteq$ ’ hold by Lemma 3.3. The proof of the converse inclusion is based on a combination of the methods used in the proofs of Proposition 2.14 and Theorem 5.3 in [GT23], which in turn are inspired by that of [Bre21, Theorem 9.1]. To verify that  $[A, I] \subseteq \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$ , let  $x \in A$  and  $y \in I$ . Set  $F := \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$ , and define

$$\varphi_y: A \times A \rightarrow I/F, \quad \varphi_y(a, b) := bya + F. \quad (a, b \in A)$$

Note that  $\varphi_y$  is a well-defined  $\mathbb{C}$ -bilinear map. Further,  $\varphi_y$  preserves zero-products since if  $a, b \in A$  satisfy  $ab = 0$ , then  $bya \in \text{FN}_2(A, I) \subseteq F$ . Using [GT23, Proposition 2.8] at the second step, we obtain that

$$cyab + F = \varphi_y(ab, c) = \varphi_y(a, bc) = bcya + F$$

and thus

$$bcya - cyab \in F = \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$$

for all  $a, b, c \in A$ . The assumption that  $I = AIA$  allows us to pick  $n$  and  $c_j, a_j \in A$  and  $y_j \in I$  for  $j = 1, \dots, n$  such that  $y = \sum_j c_j y_j a_j$ . Then

$$[x, y] = \sum_{j=1}^n [x, c_j y_j a_j] = \sum_{j=1}^n (x c_j y_j a_j - c_j y_j a_j x) \in \text{span}_{\mathbb{C}} \text{FN}_2(A, I),$$

which implies that  $[A, I] \subseteq \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$ .  $\square$

**Question 3.7.** Does there exist a two-sided ideal  $I$  in a  $C^*$ -algebra  $A$  such that  $[A, [A, I]] \neq [A, I]$ ?

The next result verifies statement (3) in Theorem D.

**Theorem 3.8.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $I \subseteq A$  be a (not necessarily closed) two-sided ideal. Then*

$$I = A[A, I]A = A[A, I], \quad \text{and} \quad [A, I] = [A, [A, I]] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I).$$

*Proof.* Considering  $I$  as a Lie ideal in  $A$ , it follows from Theorem 2.7 that  $A[A, I]A$  is the unique two-sided ideal in  $A$  to which  $I$  is related. Since  $I$  is clearly related to itself, we get  $I = A[A, I]A$ . The inclusion  $A[A, I] \subseteq A[A, I]A$  holds since  $A$  is unital. The converse inclusion holds since  $[A, I]$  is a Lie ideal in  $A$ ; see, for example, [GLT24, Lemma 3.2].

Now the equality  $[A, I] = [A, [A, I]]$  also follows from Theorem 2.7. Further, since  $A$  is zero-product balanced by Lemma 3.5, we have  $[A, I] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$  by Proposition 3.6.  $\square$

**Remark 3.9.** Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ . Applying Theorem 3.8 for the ideal  $A$ , we get

$$A = A[A, A]A,$$

which means that  $A$  is generated by its commutators as an ideal.

In fact,  $A$  is even generated by its commutators as a ring. Indeed, by [GT25, Theorem 5.15], every element in  $A$  is a sum of a commutator and the product of two commutators, and also the sum of three elements that are products of commutators.

The next result verifies statement (4) in Theorem D.

**Theorem 3.10.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ . There is a natural bijection between the set  $\mathcal{I}$  of two-sided ideals in  $A$  and the set*

$$\mathcal{L} := \{L \subseteq A \text{ Lie ideal} : L = [A, L]\}$$

given by the maps  $I \mapsto [A, I]$  and  $L \mapsto A[A, L]A$ .

*Proof.* We denote the maps by  $\alpha: \mathcal{I} \rightarrow \mathcal{L}$  and  $\beta: \mathcal{L} \rightarrow \mathcal{I}$ . Given  $I \in \mathcal{I}$ , using Theorem 3.8 at the second step, we have

$$\alpha(I) = [A, I] = [A, [A, I]] = [A, \alpha(I)]$$

which shows that  $\alpha$  is well-defined.

Next, we show that  $\alpha$  and  $\beta$  are inverses of each other. First, given  $I \in \mathcal{I}$ , using Theorem 3.8 at the last step, we get

$$\beta(\alpha(I)) = \beta([A, I]) = A[A, I]A = I.$$

Second, given  $L \in \mathcal{L}$ , it follows from  $L = [A, L]$  that  $[A, L] = [A, [A, L]]$ , and thus  $L = [A, [A, L]]$ . Using this at the last step, and using Theorem 2.7 at the third step, we get

$$\alpha(\beta(L)) = \alpha(A[A, L]A) = [A, A[A, L]A] = [A, [A, L]] = L,$$

as desired, □

Robert showed in [Rob16, Lemma 2.1] that every nilpotent element in a  $C^*$ -algebra is a sum of commutators. With view towards Theorem 3.8, we ask:

**Question 3.11.** Let  $I$  be a (not necessarily closed) two-sided ideal in a  $C^*$ -algebra  $A$ . Does every nilpotent element in  $I$  belong to  $[A, I]$ ?

We give positive answers to Question 3.11 for arbitrary two-sided ideals in von Neumann algebras (Proposition 6.2), and for semiprime two-sided ideals in  $C^*$ -algebras (Proposition 4.5).

Following [GKT23, Definition 3.2], a *Dixmier ideal* in a  $C^*$ -algebra  $A$  is a two-sided ideal  $I \subseteq A$  such that  $I$  is positively spanned (that is,  $I = \text{span}_{\mathbb{C}}(I \cap A_+)$ ) and hereditary (that is, if  $a, b \in A_+$  satisfy  $a \leq b$  and  $b \in I$ , then  $a \in I$ ) and strongly invariant (that is, if  $x \in A$  satisfies  $x^*x \in I$ , then  $xx^* \in I$ ). Two-sided ideals in von Neumann algebras are automatically Dixmier ideals, which was essentially shown by Dixmier (see [GKT23, Proposition 3.4]).

Given a Dixmier ideal  $I$  and  $s \in (0, \infty)$ , there is a unique Dixmier ideal whose set of positive elements is  $\{a^s : a \in I \cap A_+\}$ , and we use  $I^s$  to denote this ideal; see Proposition 3.7 and Definition 3.8 in [GKT23]. We note that for  $n \in \mathbb{N}$  the ring-theoretic definition of  $I^n$  as the two-sided ideal generated by  $x_1 \cdots x_n$  for  $x_1, \dots, x_n \in I$  agrees with the ‘new’ definition of  $I^n$  as the linear span of elements  $a^n$  for  $a \in I \cap A_+$ . This leads to a well-behaved theory of roots and powers for Dixmier ideals ([GKT23, Theorem 3.9]), which we recall for the convenience of the reader.

**Theorem 3.12** (Gardella, Kitamura, Thiel). *Let  $I \subseteq A$  be a Dixmier ideal in a  $C^*$ -algebra. Then*

$$(I^s)^t = I^{st} = (I^t)^s, \quad \text{and} \quad I^s I^t = I^{s+t} = I^t I^s$$

for all  $s, t \in (0, \infty)$ .

**Lemma 3.13.** *Let  $I \subseteq A$  be a Dixmier ideal in a  $C^*$ -algebra, let  $x \in I$  with polar decomposition  $x = v|x|$  in  $A^{**}$ , and let  $s \in (0, \infty)$ . Then*

$$v|x|^s, |x|^s, |x^*|^s, |x|^s v^* \in I^s.$$

*Proof.* Since  $I$  is a Dixmier ideal, we have  $x^* \in I$ , and thus  $x^*x, xx^* \in I^2$ . Then by Theorem 3.12, we get

$$|x| = (x^*x)^{\frac{1}{2}} \in (I^2)^{\frac{1}{2}} = I, \quad \text{and} \quad |x^*| = (xx^*)^{\frac{1}{2}} \in (I^2)^{\frac{1}{2}} = I.$$

It follows that  $|x|^s, |x^*|^s \in I^s$ .

By [GKT23, Proposition 3.10], given a Dixmier ideal  $J \subseteq A$ , an element  $y \in A$  belongs to  $J^{\frac{1}{2}}$  if and only if  $y^*y \in J$ . Applying this for the Dixmier ideal  $I^{2s}$ , and using that

$$(v|x|^s)^*(v|x|^s) = |x|^{2s} \in I^{2s},$$

we deduce that  $v|x|^s \in (I^{2s})^{\frac{1}{2}} = I^s$ . Since  $I^s$  is a Dixmier ideal, we further have  $|x|^s v^* = (v|x|^s)^* \in I^s$ .  $\square$

The next result showcases a situation when every element in a Dixmier ideal  $I$  is a sum of square-zero element in  $I$  and products of pairs of square-zero elements in  $I^{\frac{1}{2}}$ . In Theorem 6.4, we strengthen this result for ideals in von Neumann algebras.

**Theorem 3.14.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $I \subseteq A$  be a Dixmier ideal. Then*

$$I = [A, I] + [A, I^{\frac{1}{2}}]^2.$$

*Proof.* We first show that

$$A \cdot \text{FN}_2(A, I) \subseteq [A, I] + [A, I^{\frac{1}{2}}]^2.$$

Let  $c \in A$  and  $x \in \text{FN}_2(A, I)$ . By Lemma 3.2, we can pick  $a, b \in A_+$  and  $y \in I$  such that  $x = ayb$  and  $ab = ba = 0$ . Set  $z := yb \in I$ , and let  $z = v|z|$  be the polar decomposition in  $A^{**}$ . Then

$$cx = [ca^{\frac{1}{2}}, a^{\frac{1}{2}}z] + a^{\frac{1}{2}}zca^{\frac{1}{2}} = [ca^{\frac{1}{2}}, a^{\frac{1}{2}}z] + (a^{\frac{1}{2}}v|z|^{\frac{1}{2}})(|z|^{\frac{1}{2}}ca^{\frac{1}{2}}).$$

By Lemma 3.13,  $v|z|^{\frac{1}{2}}$  and  $|z|^{\frac{1}{2}}$  belong to  $I^{\frac{1}{2}}$ . Further, since  $za = 0$ , we get  $|z|^{\frac{1}{2}}a^{\frac{1}{2}} = 0$ , and hence

$$a^{\frac{1}{2}}v|z|^{\frac{1}{2}} = [a^{\frac{1}{2}}, v|z|^{\frac{1}{2}}] \in [A, I^{\frac{1}{2}}], \quad \text{and} \quad |z|^{\frac{1}{2}}ca^{\frac{1}{2}} = [-ca^{\frac{1}{2}}, |z|^{\frac{1}{2}}] \in [A, I^{\frac{1}{2}}].$$

It follows that  $cx$  belongs to  $[A, I] + [A, I^{\frac{1}{2}}]^2$ .

Using Theorem 3.8 at the first and second step, and using the above at the last step, we get

$$I = A[A, I] = \text{span}_{\mathbb{C}} \left( A \cdot \text{FN}_2(A, I) \right) \subseteq [A, I] + [A, I^{\frac{1}{2}}]^2.$$

On the other hand, we clearly have  $[A, I] \subseteq I$ . Using Theorem 3.12, we also have  $[A, I^{\frac{1}{2}}]^2 \subseteq (I^{\frac{1}{2}})^2 = I$ .  $\square$

#### 4. SQUARE-ZERO ELEMENTS IN SEMIPRIME IDEALS IN C\*-ALGEBRAS

In this section, we study the subspace  $N$  generated by the square-zero elements in a semiprime ideal in a  $C^*$ -algebra  $A$ . We show that  $ANA$ , the two-sided ideal generated by  $N$ , satisfies  $ANA = [A, N] + N^2$ ; see Theorem 4.3. This generalizes results from [GT24, Section 4].

We then characterize semiprimeness for two-sided ideals in a  $C^*$ -algebra whose unit is weakly divisible of degree 2; see Theorem 4.6. In particular, such an ideal is semiprime if and only if it is generated by its commutators as an ideal (or as a ring).

We begin with some preparatory results about commutators involving Dixmier ideals, which might be of use in future work.

**Lemma 4.1.** *Let  $I \subseteq A$  be a Dixmier ideal in a  $C^*$ -algebra. Then*

$$[A, I] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq I.$$

*Proof.* By Theorem 3.12, we have  $I = I^{\frac{1}{2}}I^{\frac{1}{2}}$ . Using that  $[a, xy] = [ax, y] + [ya, x]$  for  $a \in A$  and  $x, y \in I$ , it follows that

$$[A, I] = [A, I^{\frac{1}{2}}I^{\frac{1}{2}}] \subseteq [AI^{\frac{1}{2}}, I^{\frac{1}{2}}] + [I^{\frac{1}{2}}A, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Further, we have  $[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq I^{\frac{1}{2}}I^{\frac{1}{2}} = I$ .  $\square$

Recall that  $N_2(I)$  denotes the set of square-zero elements in an ideal  $I$ .

**Lemma 4.2.** *Let  $I \subseteq A$  be a Dixmier ideal in a  $C^*$ -algebra. Then*

$$N_2(I) \subseteq \text{span}_{\mathbb{C}} [N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})].$$

*Proof.* Let  $x \in N_2(I)$ , and let  $x = v|x|$  be the polar decomposition in  $A^{**}$ . We have  $x = |x^*|v = |x^*|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$ . Since  $x^2 = 0$ , the elements  $|x|$  and  $|x^*|$  are orthogonal, and hence so are  $|x|^{\frac{1}{2}}$  and  $|x^*|^{\frac{1}{2}}$ . Therefore

$$x = [\frac{1}{2}v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}}].$$

By Lemma 3.13, the element  $\frac{1}{2}v|x|^{\frac{1}{2}}$  belongs to  $I^{\frac{1}{2}}$ , and thus  $\frac{1}{2}v|x|^{\frac{1}{2}} \in N_2(I^{\frac{1}{2}})$ . Set

$$y := |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}} + v|x|^{\frac{1}{2}} - v^*|x^*|^{\frac{1}{2}}.$$

It is elementary (but tedious) to check that  $y^2 = 0$ . Note that  $y$  belongs to  $I^{\frac{1}{2}}$  by Lemma 3.13, and thus  $y \in N_2(I^{\frac{1}{2}})$ . Using that  $v|x|^{\frac{1}{2}}$  and  $v^*|x^*|^{\frac{1}{2}}$  also belong to  $N_2(I^{\frac{1}{2}})$ , it follows that

$$|x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}} = y - v|x|^{\frac{1}{2}} + v^*|x^*|^{\frac{1}{2}} \in \text{span}_{\mathbb{C}} N_2(I^{\frac{1}{2}}),$$

and thus

$$x = [\frac{1}{2}v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}}] \in \text{span}_{\mathbb{C}} [N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})],$$

as desired.  $\square$

A two-sided ideal  $I$  in a ring  $R$  is said to be *semiprime* if for all two-sided ideals  $J \subseteq R$  we have  $J \subseteq I$  whenever  $J^2 \subseteq I$ . We refer to [Lam01, Section 10] for details. In [GKT23, Section 5], together with Gardella and Kitamura, we showed that a two-sided ideal  $I$  in a  $C^*$ -algebra is semiprime if and only if it is idempotent (that is,  $I = I^2$ ), if and only if it is a Dixmier ideal and satisfies  $I = I^{\frac{1}{2}}$ .

The next result is a generalization of [GT24, Theorem 4.2] from the study of square-zero elements in  $C^*$ -algebras to square-zero elements in semiprime ideals in  $C^*$ -algebras. Recall that  $\tilde{A}$  denotes the minimal unitization of a  $C^*$ -algebra  $A$ .

**Theorem 4.3.** *Let  $I \subseteq A$  be a semiprime ideal in a  $C^*$ -algebra, let  $N$  denote the subspace generated by  $N_2(I)$ , and let  $J := \tilde{A}N\tilde{A}$  denote the two-sided ideal of  $A$  generated by  $N$ . Then  $J$  is semiprime, and we have*

$$N = [N, N] \subseteq N^2, \quad J = [J, J]^2 = AN = [A, N]^2 = [A, N] + N^2,$$

and

$$[J, J] = [A, J] = [A, [A, N]] = [[J, J], [J, J]] = [[A, N], [A, N]].$$

*Proof.* We will use that  $I$  is a Dixmier ideal satisfying  $I = I^{\frac{1}{2}}$ . Applying Lemma 4.2 at the second step, we have

$$N = \text{span}_{\mathbb{C}} N_2(I) \subseteq \text{span}_{\mathbb{C}} [N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})] = \text{span}_{\mathbb{C}} [N_2(I), N_2(I)] = [N, N].$$

The inclusion  $[N, N] \subseteq N^2$  follows since every commutator in  $[N, N]$  belongs to  $N^2$ . To verify that  $[N, N] \subseteq N$ , let  $x, y \in N_2(I)$ . Then

$$[x, y] = (1+x)y(1-x) - y + xyx.$$

Each of the three summands is a square-zero element in  $I$ , which shows  $[x, y] \in N$ .

We have

$$J := \tilde{A}N\tilde{A} \subseteq \tilde{A}N^2\tilde{A} \subseteq (\tilde{A}N)(N\tilde{A}) \subseteq J^2,$$

which shows that  $J$  is idempotent and therefore semiprime by [GKT23, Theorem A].

The inclusion  $AN \subseteq \tilde{A}N\tilde{A} = J$  is clear. Conversely, given  $a \in \tilde{A}$ ,  $x \in N_2(I)$  and a unitary  $u \in \tilde{A}$ , using that  $u^*xu \in N_2(I)$  and that  $N \subseteq N^2$ , we have

$$axu = au(u^*xu) \in \tilde{A}N_2(I) \subseteq \tilde{A}N \subseteq \tilde{A}N^2 \subseteq AN.$$

Since every element in  $\tilde{A}$  is a linear combination of four unitaries, we get  $J = AN$ .

The inclusion  $[A, N] + N^2 \subseteq \tilde{A}N\tilde{A} = J$  is clear. For the converse inclusion, we use a similar argument as in the proof of Theorem 3.14 to show that  $AN \subseteq [A, N] + N^2$ . Given  $a \in A$  and  $x \in N_2(I)$ , let  $x = v|x|$  be the polar decomposition in  $A^{**}$ . Since  $I$  is semiprime, it is Dixmier and  $I = I^{\frac{1}{2}} = I^{\frac{1}{4}}$ . By Lemma 3.13, we have  $v|x|^{\frac{1}{2}} \in I^{\frac{1}{2}}$  and thus  $v|x|^{\frac{1}{2}} \in N_2(I)$ . Similarly, we have  $v|x|^{\frac{1}{4}}, |x|^{\frac{1}{4}}a|x|^{\frac{1}{2}} \in N_2(I)$ . Using that  $x = |x^*|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$ , we get

$$ax = [a|x^*|^{\frac{1}{2}}, v|x|^{\frac{1}{2}}] + (v|x|^{\frac{1}{4}})(|x|^{\frac{1}{4}}a|x|^{\frac{1}{2}}) \in [A, N_2(I)] + N_2(I)^2.$$

Using that  $J = AN$ , we get  $J = [A, N] + N^2$ .

The inclusion  $[A, N]^2 \subseteq \tilde{A}N\tilde{A} = J$  is clear. Conversely, using that  $N = [N, N]$ , we have

$$[A, N] = [A, [N, N]] \subseteq [N, [A, N]] \subseteq [[A, N], [A, N]] \subseteq [A, N]^2.$$

We also have  $N^2 = [N, N]^2 \subseteq [A, N]^2$  and thus

$$J = [A, N] + N^2 \subseteq [A, N]^2.$$

The inclusion  $[J, J] \subseteq [A, J]$  is clear. Using that  $J = J^2$  and  $[a, xy] = [ax, y] + [ya, x]$  for  $a \in A$  and  $xy \in J$ , we have

$$[A, J] = [A, J^2] \subseteq [AJ, J] + [JA, J] \subseteq [J, J].$$

This shows that  $[J, J] = [A, J]$ . We get

$$[[A, N], [A, N]] \subseteq [A, [A, N]] \subseteq [A, J] = [J, J].$$

We have

$$[[A, N], N^2] \subseteq [A, N^2] \subseteq [AN, N] + [NA, N] \subseteq [A, N].$$

and similarly  $[N^2, N^2] \subseteq [A, N^2] \subseteq [A, N]$ . We also have

$$[A, N] = [A, [N, N]] \subseteq [N, [A, N]] = [[N, N], [A, N]] \subseteq [[A, N], [A, N]].$$

Using that  $J = [A, N] + N^2$ , it follows that

$$\begin{aligned} [J, J] &\subseteq [[A, N], [A, N]] + [N^2, [A, N]] + [[A, N], N^2] + [N^2, N^2] \\ &\subseteq [[A, N], [A, N]]. \end{aligned}$$

Using that  $[A, N] \subseteq [A, J] = [J, J]$ , we also have

$$[J, J] = [[A, N], [A, N]] \subseteq [[J, J], [J, J]].$$

Since the converse inclusion is clear.

Finally, since  $[A, N] \subseteq [J, J]$  and  $J = [A, N]^2$ , we have  $J = [J, J]^2$ .  $\square$

**Lemma 4.4.** *Let  $I \subseteq A$  be a Dixmier ideal in a C\*-algebra, and let  $x \in I$  be nilpotent. Then  $x \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$ .*

*Proof.* As in the proof of [Rob16, Lemma 2.1], we proceed by induction on the degree of nilpotency. To start, let  $x \in I$  be a square-zero element. By Lemma 3.13, the elements  $v|x|^{\frac{1}{2}}$  and  $|x|^{\frac{1}{2}}$  belong to  $I^{\frac{1}{2}}$ , and thus

$$x = [v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}}] \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Next, assume that for some  $k \geq 2$  we have shown that every element  $x \in I$  with  $x^k = 0$  belongs to  $[I^{\frac{1}{2}}, I^{\frac{1}{2}}]$ . Let  $x \in I$  satisfy  $x^{k+1} = 0$ , and let  $x = v|x|$  be the polar decomposition of  $x$  in  $A^{**}$ . Set  $y := |x|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$ , which belongs to  $A$ . As in the proof of [Rob16, Lemma 2.1], one shows that  $y^k = 0$ . By Lemma 3.13, we have  $v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} \in I^{\frac{1}{2}}$ . By assumption of the induction, we have  $y \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$ , and thus

$$x = [v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}}] + y \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$$

as desired.  $\square$

**Proposition 4.5.** *Let  $I \subseteq A$  be a semiprime ideal in a  $C^*$ -algebra, and let  $x \in I$  be nilpotent. Then  $x \in [A, I]$ .*

*Proof.* Since  $I$  is semiprime, it is a Dixmier ideal and satisfies  $I = I^{\frac{1}{2}}$ . Applying Lemma 4.4, we have

$$x \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq [A, I^{\frac{1}{2}}] = [A, I],$$

as desired.  $\square$

**Theorem 4.6.** *Let  $A$  be a unital  $C^*$ -algebra that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $I \subseteq A$  be a two-sided ideal. The following statements are equivalent:*

- (1) *The ideal  $I \subseteq A$  is semiprime.*
- (2) *We have  $I = A[I, I]A$ , that is,  $I$  is generated by its commutators as an ideal in  $A$ .*
- (3) *We have  $I = \tilde{I}[I, I]\tilde{I}$ , that is,  $I$  is generated by its commutators as an ideal.*
- (4) *We have  $I = [I, I]^2$ .*
- (5) *We have  $[I, I] = [[I, I], [I, I]]$ .*

*Proof.* To show that (2) implies (1), assume that  $I$  is generated by  $[I, I]$  as an ideal in  $A$ . We clearly have  $[I, I] \subseteq I^2$ , and therefore

$$I = A[I, I]A \subseteq AI^2A = I^2.$$

It follows that  $I$  is idempotent, and thus semiprime by [GKT23, Theorem A].

It is clear that (4) implies (3), which in turn implies (2).

Assume that  $I$  is semiprime, and let  $N$  denote the subspace generated by the square-zero elements in  $I$ . Applying Theorem 3.8, we have

$$I = A[A, I]A = \text{span}_{\mathbb{C}} A \text{FN}_2(A, I)A \subseteq ANA.$$

Now it follows from Theorem 4.3 that

$$I = ANA = [ANA, ANA]^2 = [I, I]^2,$$

which verifies (4), and that

$$[I, I] = [ANA, ANA] = [[ANA, ANA], [ANA, ANA]] = [[I, I], [I, I]], \quad \text{and}$$

which verifies (5).

Finally, assuming (5) let us verify (1). Using that  $[a, xy] = [ax, y] + [ya, x]$  for  $a \in A$  and  $x, y \in I$ , we have

$$[A, I^2] \subseteq [AI, I] + [IA, I] \subseteq [I, I].$$

Applying Theorem 3.8 for the ideal  $I^2$ , we get

$$I^2 = A[A, I^2]A \subseteq A[I, I]A \subseteq A[[I, I], [I, I]]A \subseteq I^4.$$



It follows that  $I^2 = I^4$ , and thus semiprime by [GKT23, Theorem A].  $\square$

## 5. LIE IDEALS IN PROPERLY INFINITE C\*-ALGEBRAS

Let  $A$  be a  $C^*$ -algebra with a unit that is weakly divisible of degree 2, that is, that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ , and let  $L \subseteq A$  be a (not necessarily closed) Lie ideal. We have seen in Theorem 2.7 that  $L$  is related to the two-sided ideal  $I := A[A, L]A$ . Further, by Theorem 2.8,  $L$  is commutator equivalent to  $I$  if and only if  $[A, L] = [A, [A, L]]$ .

This raises the question when Lie ideals  $L$  in  $A$  satisfy  $[A, L] = [A, [A, L]]$ . A sufficient condition is that  $A = Z(A) + [A, A]$ , where  $Z(A)$  denotes the center of  $A$ ; see Lemma 5.1. We observe in Proposition 5.2 that this condition is satisfied in a number of situations, in particular for all properly infinite  $C^*$ -algebras.

We then prove the main result of the paper: Every Lie ideal in a unital, properly infinite  $C^*$ -algebra is commutator equivalent to a (unique) two-sided ideal; see Theorem 5.3.

**Lemma 5.1.** *Let  $A$  be a unital  $C^*$ -algebra such that  $A = Z(A) + [A, A]$ , and let  $L \subseteq A$  be a Lie ideal. Then*

$$[A, L] = [A, [A, L]].$$

*Proof.* The inclusion ' $\supseteq$ ' is clear. Conversely, using the assumption at the first step, and using the Jacobi identity at the last step, we get

$$[A, L] = [Z(A) + [A, A], L] = [[A, A], L] \subseteq [A, [A, L]],$$

as desired.  $\square$

Following Winter [Win12], a  $C^*$ -algebra  $A$  is said to be *pure* if its Cuntz semigroup  $\text{Cu}(A)$  is almost unperforated and almost divisible. Pure  $C^*$ -algebras form a robust class [APTV24] that includes every  $C^*$ -algebra that tensorially absorbs the Jiang-Su algebra  $\mathcal{Z}$ , that is,  $A \cong A \otimes \mathcal{Z}$ . By [APT18, Theorem 7.3.11], a  $C^*$ -algebra  $A$  is pure if and only if  $\text{Cu}(A) \cong \text{Cu}(A) \otimes \text{Cu}(\mathcal{Z})$ .

By a *quasitrace* of a  $C^*$ -algebra  $A$  we mean a lower-semicontinuous,  $[0, \infty]$ -valued 2-quasitrace. By Haagerup's theorem, every bounded quasitrace on a unital, exact  $C^*$ -algebra is a trace, and this was later extended by Kirchberg, who showed that arbitrary quasitraces on exact  $C^*$ -algebras are traces.

A unital  $C^*$ -algebra  $A$  is *weakly central* if the center separates maximal ideals: Maximal ideals  $M_1, M_2 \subseteq A$  satisfy  $M_1 = M_2$  whenever  $M_1 \cap Z(A) = M_2 \cap Z(A)$ . By Vesterstrøm's theorem,  $A$  is weakly central if and only if  $A$  has the *centre-quotient property*: For every closed, two-sided ideal  $I \subseteq A$ , the quotient map  $A \rightarrow A/I$  maps  $Z(A)$  onto  $Z(A/I)$ .

A unital  $C^*$ -algebra  $A$  has the *Dixmier property* if for every element  $a \in A$  the closed convex hull of the unitary orbit  $\{uau^* : u \in A \text{ unitary}\}$  has nonempty intersection with the center  $Z(A)$ . It was shown in [ART17, Theorem 1.1] that this property holds if and only if  $A$  is weakly central, every simple quotient of  $A$  has at most one tracial state, and every extreme tracial state of  $A$  factors through a simple quotient. Dixmier showed that every von Neumann algebra has the property named after him.

**Proposition 5.2.** *Let  $A$  be a unital  $C^*$ -algebra. Then  $A = Z(A) + [A, A]$  in each of the following cases:*

- (1) *If  $A$  is pure, every quasitrace on  $A$  is a trace, and  $A$  has the Dixmier property.*
- (2) *If  $A$  has no tracial states (in particular, if  $A$  is properly infinite).*
- (3) *If  $A$  is a von Neumann algebra.*

*Proof.* (1) If  $A$  has the Dixmier property, then every element in  $A$  is a Dixmier element in the sense of [AGR23], and thus it follows from [AGR23, Lemma 4.10] that  $A = Z(A) + \overline{[A, A]}$ .

If  $A$  is pure and every quasitrace on  $A$  is a trace, then  $[A, A]$  is closed by [NR16, Theorem 1.1]. Combining both results, we get  $A = Z(A) + [A, A]$ .

(2) By [Pop02, Theorem 1], we have  $A = [A, A]$  if (and only if)  $A$  has no tracial states.

(3) It was shown in the proof of [BKS08, Theorem 5.19] that every von Neumann algebra  $A$  satisfies  $A = Z(A) + [A, A]$ .  $\square$

**Theorem 5.3.** *Let  $A$  be a unital, properly infinite  $C^*$ -algebra, and let  $L \subseteq A$  be a Lie ideal. Then for the two-sided ideal  $I := A[A, L]A$ , we have*

$$[A, L] = [A, [A, L]] = [A, I].$$

*In particular,  $L$  is commutator equivalent to  $I$ , and hence also embraced by  $I$ , and related to  $I$ .*

*Further,  $I$  is the only two-sided ideal of  $A$  to which  $L$  is related.*

*Proof.* Since  $A$  is unital and properly infinite, there exists a unital  $*$ -homomorphism from the Cuntz algebra  $\mathcal{O}_\infty$  to  $A$ ; see [Bla06, Proposition III.1.3.3]. Further, it is well-known that  $\mathcal{O}_\infty$  admits a unital  $*$ -homomorphism from  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$ , using for example Proposition 2.9 and that  $\mathcal{O}_\infty$  is a unital  $C^*$ -algebra of real rank zero that admits no characters. It follows that  $A$  admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ .

We may therefore apply Theorem 2.7 and obtain that  $L$  is related to  $I$ , and that  $I$  is the only two-sided ideal to which  $L$  is related. Further, since  $A$  is unital and properly infinite, we have  $A = [A, A]$  by Pop's theorem [Pop02], and thus  $[A, L] = [A, [A, L]]$  by Lemma 5.1. It therefore follows from Theorem 2.8 that  $L$  is commutator equivalent to  $I$  (and thus also embraced by  $I$ ).  $\square$

**Corollary 5.4.** *Let  $A$  be a unital, properly infinite  $C^*$ -algebra. Then an additive subgroup  $L \subseteq A$  is a Lie ideal if and only if  $[A, I] \subseteq L \subseteq T([A, I])$  for some (unique) two-sided ideal  $I \subseteq A$ . We get a decomposition of the family  $\mathcal{L}$  of all Lie ideals in  $A$  as:*

$$\mathcal{L} = \bigsqcup_{I \in \mathcal{I}} \{L \subseteq A \text{ additive subgroup} : [A, I] \subseteq L \subseteq T([A, I])\}$$

*where  $\mathcal{I}$  denotes the lattice of two-sided ideals in  $A$ .*

**Remark 5.5.** The proof of Theorem 5.3 can easily be adapted to show the following: In a  $C^*$ -algebra  $A$  that admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$  and such that  $A = Z(A) + [A, A]$ , every Lie ideal  $L$  is commutator equivalent to the two-sided ideal  $A[A, L]A$ .

We note that a unital  $C^*$ -algebra admits no characters whenever it is pure or has no tracial states. Therefore, with view towards Question 2.11 and Proposition 5.2, we expect that every Lie ideal  $L$  is commutator equivalent to  $A[A, L]A$  whenever  $A$  is a unital, pure,  $C^*$ -algebra with the Dixmier property and such that every quasitrace on  $A$  is a trace, or whenever  $A$  is unital and has no tracial states.

**Question 5.6.** Given Lie ideals  $L, M \subseteq A$  in a unital, properly infinite  $C^*$ -algebra, what is the relationship between the two-sided ideals generated by  $[L, M]$ , by  $[A, [L, M]]$ , and by  $[A, L]$  and  $[A, M]$ ?

## 6. LIE IDEALS AND TWO-SIDED IDEALS IN VON NEUMANN ALGEBRAS

In this section, we show that every (not necessarily closed) Lie ideal  $L$  in a von Neumann algebra  $M$  is commutator equivalent to the two-sided ideal  $I :=$

$M[M, L]M$ , that is,  $[M, L] = [M, I]$ ; see Theorem 6.1. This recovers [BKS08, Theorem 5.19]. Moreover, if  $M$  has zero commutative summand, then we show that  $I$  is the only two-sided ideal to which  $L$  is related, which solves [BKS08, Problem 5.21].

Given a two-sided ideal  $I$  in a von Neumann algebra with zero commutative summand, every element in  $I$  is a sum of products of pairs of square-zero elements in  $I^{\frac{1}{2}}$ ; see Theorem 6.4.

**Theorem 6.1.** *Let  $M$  be a von Neumann algebra, and let  $L \subseteq M$  be a Lie ideal. Then, for the two-sided ideal  $I := M[M, L]M$ , we have*

$$[M, L] = [M, [M, L]] = [M, I].$$

*In particular,  $L$  is commutator equivalent to  $I$ , and hence also embraced by  $I$ , and related to  $I$ .*

*If  $M$  has zero commutative summand, then  $I$  is the only two-sided ideal of  $M$  to which  $L$  is related.*

*Proof.* Let  $M = M_0 \oplus M_1$  be the (unique) decomposition of  $M$  into a commutative summand  $M_0$  and a summand  $M_1$  that admits no characters. By Proposition 2.9, there is a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M_1$ . Set

$$L_1 := M_1 \cap L, \quad \text{and} \quad I_1 := M_1 \cap I.$$

Then  $L_1$  is a Lie ideal in  $M_1$ , and we have

$$[M, L] = [M_1, L_1], \quad \text{and} \quad I = I_1 = M_1[M_1, L_1]M_1.$$

We may apply Theorem 2.7 for the Lie ideal  $L_1$  in  $M_1$  and obtain that  $L_1$  is related to  $I_1$ , and that  $I_1$  is the only two-sided ideal of  $M_1$  to which  $L_1$  is related. It follows that  $L$  is related to  $I$ , and if  $M_0$  is zero, then  $I$  is also unique with this property. (If  $M_0$  is nonzero, then  $M_0 \oplus I$  is another two-sided ideal of  $M$  to which  $L$  is related.)

As shown in the proof of [BKS08, Theorem 5.19], we have  $M = Z(M) + [M, M]$ . It follows that  $[M, L] = [M, [M, L]]$  by Lemma 5.1, and then

$$[M, L] = [M, [M, L]] = [M, I].$$

by Theorem 2.8. □

Recall that  $N_2(I)$  denotes the square-zero elements, and we use  $\text{Nil}(I)$  to denote the set of nilpotent elements in an ideal  $I$ . Recall the definition of  $\text{FN}_2(M, I)$  from Definition 3.1.

**Proposition 6.2.** *Let  $I \subseteq M$  be a two-sided ideal in a von Neumann algebra. Then*

$$[M, I] = \text{span}_{\mathbb{C}} N_2(I) = \text{span}_{\mathbb{C}} \text{Nil}(I) = \text{span}_{\mathbb{C}} \text{FN}_2(M, I).$$

*Proof.* Let  $M = M_0 \oplus M_1$  denote the (unique) decomposition of  $M$  into a commutative summand  $M_0$  and a summand  $M_1$  that admits no characters. Then

$$\begin{aligned} [M, I] &= [M_1, M_1 \cap I], \quad N_2(I) = N_2(M_1 \cap I), \\ \text{Nil}(I) &= \text{Nil}(M_1 \cap I), \quad \text{and} \quad \text{FN}_2(M, I) = \text{FN}_2(M_1, M_1 \cap I), \end{aligned}$$

which allows us to reduce the problem to studying the ideal  $M_1 \cap I$  in  $M_1$ .

Without loss of generality, we may therefore assume that  $M$  has zero commutative summand. Then  $M$  admits a unital  $*$ -homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M$  by Proposition 2.9. Using Theorem 3.8 at the first step, we get

$$[M, I] = \text{span}_{\mathbb{C}} \text{FN}_2(M, I) \subseteq \text{span}_{\mathbb{C}} N_2(I) \subseteq \text{span}_{\mathbb{C}} \text{Nil}(I).$$

It remains to show that  $\text{Nil}(I) \subseteq [M, I]$ . As in the proof of Lemma 4.4 (and [Rob16, Lemma 2.1]), we proceed by induction on the degree of nilpotency. To

start, let  $x \in I$  be a square-zero element. Let  $x = v|x|$  be the polar decomposition in  $M$ . Then  $|x| = v^*x \in I$ . Since  $x^2 = 0$ , we have  $|x|v = 0$ , and thus

$$x = [v, |x|] \in [M, I].$$

Next, assume that for some  $k \geq 2$  we have shown that every element  $x \in I$  with  $x^k = 0$  belongs to  $[M, I]$ . Let  $x \in I$  satisfy  $x^{k+1} = 0$ . Again, let  $x = v|x|$  be the polar decomposition in  $M$ . We will show that the element  $y := |x|v$  satisfies  $y^k = 0$ . Since  $x^{k+1} = 0$  and  $|x| = v^*v|x|$ , we have

$$y^k|x| = |x|v \cdots |x|v|x| = v^*v|x|v \cdots |x|v|x| = v^*x^{k+1} = 0.$$

For every polynomial  $p$  with vanishing constant term it follows that  $y^k p(|x|) = 0$ . Using functional calculus, we get  $y^k|x|^{\frac{1}{n}}$  for every  $n \geq 1$ . The sequence  $(|x|^{\frac{1}{n}})_n$  converges in the weak\*-topology of  $M$  to  $v^*v$ , which is the support projection of  $|x|$ . We deduce that  $y^k v^*v = 0$ . Using that  $v = vv^*v$ , we get

$$y^k = y^{k-1}|x|v = y^{k-1}|x|vv^*v = y^k v^*v = 0.$$

By assumption of the induction, we have  $y \in [M, I]$ , and thus

$$x = [v, |x|] + |x|v \in [M, I],$$

as desired.  $\square$

**Remark 6.3.** Let  $H$  be a separable, infinite-dimensional Hilbert space, and let  $I$  be a two-sided ideal in the von Neumann algebra  $\mathcal{B}(H)$ . Given a nilpotent element  $x \in I$ , it follows from Proposition 6.2 that  $x \in [\mathcal{B}(H), I]$ , that is,  $x$  is a finite sum of commutators of an element in  $\mathcal{B}(H)$  and an element in  $I$ . Dykema and Krishnaswamy-Usha showed in [DKU18, Proposition 3.1] that one summand suffices, that is, we have  $x = [y, z]$  for some  $y \in \mathcal{B}(H)$  and  $z \in I$ . Does their result hold for nilpotent elements in two-sided ideals in arbitrary von Neumann algebras?

**Theorem 6.4.** *Let  $M$  be a von Neumann algebra with zero commutative summand, and let  $I \subseteq M$  be a two-sided ideal. Then*

$$I = [M, I^{\frac{1}{2}}]^2 = [I^{\frac{1}{4}}, I^{\frac{1}{4}}]^2,$$

and

$$[M, I] \subseteq [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]] \subseteq [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

*Proof.* We will use that  $I$  is a Dixmier ideal, since two-sided ideals in von Neumann algebras are automatically Dixmier ideals ([GKT23, Proposition 3.4]). Using Proposition 6.2 at the first and last step, and using Lemma 4.2 at the second step, we have

$$[M, I] = \text{span}_{\mathbb{C}} N_2(I) \subseteq \text{span}_{\mathbb{C}} [N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})] = [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]].$$

Using Lemma 4.1, we then get

$$[M, I] \subseteq [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]] \subseteq [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$$

By Proposition 2.9, there is a unital \*-homomorphism  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M$ . We can therefore apply Theorem 3.14 at the first step, and together with the above we obtain that

$$I = [M, I] + [M, I^{\frac{1}{2}}]^2 \subseteq [M, I^{\frac{1}{2}}]^2.$$

Applying Lemma 4.1 for the Dixmier ideal  $I^{\frac{1}{2}}$ , and using Theorem 3.12 at the last step, we get

$$I \subseteq [M, I^{\frac{1}{2}}]^2 \subseteq [I^{\frac{1}{4}}, I^{\frac{1}{4}}]^2 \subseteq (I^{\frac{1}{2}})^2 = I,$$

as desired.  $\square$

**Remark 6.5.** It is likely that the inclusions involving  $[M, I]$  in Theorem 6.4 are equalities. The key question is whether  $[I^{\frac{1}{2}}, I^{\frac{1}{2}}]$  is a subset of  $[M, I]$ . Using that  $I$  is a Dixmier ideal and therefore admits a well-behaved theory of roots and powers (Theorem 3.12), we have

$$[I^{\frac{1}{2}}, I^{\frac{1}{2}}] = [I^{\frac{1}{4}} I^{\frac{1}{4}}, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{4}}, I^{\frac{1}{4}} I^{\frac{1}{2}}] + [I^{\frac{1}{2}}, I^{\frac{1}{2}} I^{\frac{1}{4}}] = [I^{\frac{1}{4}}, I^{\frac{3}{4}}],$$

and by an analogous argument  $[I^{\frac{1}{4}}, I^{\frac{3}{4}}] \subseteq [I^{\frac{1}{8}}, I^{\frac{7}{8}}]$ . Inductively, it follows that  $[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{2^n}}, I^{\frac{2^n-1}{2^n}}]$  for every  $n \geq 1$ , and thus

$$[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq \bigcap_{n \geq 1} [I^{\frac{1}{2^n}}, I^{\frac{2^n-1}{2^n}}] \subseteq \bigcap_{\varepsilon > 0} [M, I^{1-\varepsilon}].$$

It is, however, not clear if  $\bigcap_{\varepsilon > 0} [M, I^{1-\varepsilon}] = [M, I]$ .

For two-sided ideals  $I, J \subseteq \mathcal{B}(H)$ , it was shown in [DFWW04, Theorem 5.10] that  $[\mathcal{B}(H), IJ] = [I, J]$ . In particular, we have  $[\mathcal{B}(H), I] = [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$ , and we obtain the following consequence of Theorem 6.4:

**Corollary 6.6.** *Let  $H$  be a separable, infinite-dimensional Hilbert space, and let  $I$  be a two-sided ideal in  $\mathcal{B}(H)$ . Then*

$$[\mathcal{B}(H), I] = [[\mathcal{B}(H), I^{\frac{1}{2}}], [\mathcal{B}(H), I^{\frac{1}{2}}]] = [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] = [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

**Question 6.7.** Let  $I, J \subseteq M$  be two-sided ideals in a von Neumann algebra. Do we have  $[M, IJ] = [I, J]$ ?

## REFERENCES

- [AGEP24] T. AMRUTAM, D. GAO, S. K. ELAYAVALLI, and G. PATCHELL, Strict comparison in reduced group  $C^*$ -algebras, preprint (arXiv:2412.06031), 2024.
- [APT18] R. ANTOINE, F. PERERA, and H. THIEL, Tensor products and regularity properties of Cuntz semigroups, *Mem. Amer. Math. Soc.* **251** (2018), viii+191.
- [APTV24] R. ANTOINE, F. PERERA, H. THIEL, and E. VILALTA, Pure  $C^*$ -algebras, preprint (arXiv:2406.11052 [math.OA]), 2024.
- [ART17] R. ARCHBOLD, L. ROBERT, and A. TIKUISIS, The Dixmier property and tracial states for  $C^*$ -algebras, *J. Funct. Anal.* **273** (2017), 2655–2718.
- [AGR23] R. J. ARCHBOLD, I. GOGIĆ, and L. ROBERT, Local variants of the Dixmier property and weak centrality for  $C^*$ -algebras, *Int. Math. Res. Not. IMRN* (2023), 1483–1513.
- [Bla06] B. BLACKADAR, *Operator algebras, Encyclopaedia of Mathematical Sciences* **122**, Springer-Verlag, Berlin, 2006, Theory of  $C^*$ -algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.
- [BKKO17] E. BREUILLARD, M. KALANTAR, M. KENNEDY, and N. OZAWA,  $C^*$ -simplicity and the unique trace property for discrete groups, *Publ. Math. Inst. Hautes Études Sci.* **126** (2017), 35–71.
- [Bre21] M. BREŠAR, *Zero product determined algebras, Frontiers in Mathematics*, Birkhäuser/Springer, Cham, 2021.
- [BKS08] M. BREŠAR, E. KISSIN, and V. S. SHULMAN, Lie ideals: from pure algebra to  $C^*$ -algebras, *J. Reine Angew. Math.* **623** (2008), 73–121.
- [CP79] J. CUNTZ and G. K. PEDERSEN, Equivalence and traces on  $C^*$ -algebras, *J. Funct. Anal.* **33** (1979), 135–164.
- [DFWW04] K. DYKEMA, T. FIGIEL, G. WEISS, and M. WODZICKI, Commutator structure of operator ideals, *Adv. Math.* **185** (2004), 1–79.
- [DKU18] K. DYKEMA and A. KRISHNASWAMY-USHA, Nilpotent elements of operator ideals as single commutators, *Proc. Amer. Math. Soc.* **146** (2018), 3031–3037.
- [FMS82] C. K. FONG, C. R. MIERS, and A. R. SOUROY, Lie and Jordan ideals of operators on Hilbert space, *Proc. Amer. Math. Soc.* **84** (1982), 516–520.
- [GKT23] E. GARDELLA, K. KITAMURA, and H. THIEL, Semiprime ideals in  $C^*$ -algebras, *J. Eur. Math. Soc.* (to appear), preprint (arXiv:2311.17480), 2023.
- [GLT24] E. GARDELLA, T.-K. LEE, and H. THIEL, Fully noncentral Lie ideals and invariant additive subgroups in rings, preprint (arXiv:2409.03362), 2024.
- [GT23] E. GARDELLA and H. THIEL, Zero-product balanced algebras, *Linear Algebra Appl.* **670** (2023), 121–153.

- [GT24] E. GARDELLA and H. THIEL, Prime ideals in  $C^*$ -algebras and applications to Lie theory, *Proc. Amer. Math. Soc.* **152** (2024), 3647–3656.
- [GT25] E. GARDELLA and H. THIEL, Rings and  $C^*$ -algebras generated by commutators, *J. Algebra* **662** (2025), 214–241.
- [Her55] I. N. HERSTEIN, On the Lie and Jordan rings of a simple associative ring, *Amer. J. Math.* **77** (1955), 279–285.
- [KR15] E. KIRCHBERG and M. RØRDAM, When central sequence  $C^*$ -algebras have characters, *Internat. J. Math.* **26** (2015), 1550049, 32.
- [Lam01] T. Y. LAM, *A first course in noncommutative rings*, second ed., *Graduate Texts in Mathematics* **131**, Springer-Verlag, New York, 2001.
- [Mar10] L. W. MARCOUX, Projections, commutators and Lie ideals in  $C^*$ -algebras, *Math. Proc. R. Ir. Acad.* **110A** (2010), 31–55.
- [MM98] L. W. MARCOUX and G. J. MURPHY, Unitarily-invariant linear spaces in  $C^*$ -algebras, *Proc. Amer. Math. Soc.* **126** (1998), 3597–3605.
- [Mar95] L. MARCOUX, On the closed Lie ideals of certain  $C^*$ -algebras, *Integral Equations Operator Theory* **22** (1995), 463–475.
- [Ng12] P. W. NG, Commutators in the Jiang-Su algebra, *Internat. J. Math.* **23** (2012), 1250113, 29.
- [NR16] P. W. NG and L. ROBERT, Sums of commutators in pure  $C^*$ -algebras, *Münster J. Math.* **9** (2016), 121–154.
- [PR04] F. PERERA and M. RØRDAM, AF-embeddings into  $C^*$ -algebras of real rank zero, *J. Funct. Anal.* **217** (2004), 142–170.
- [Pop02] C. POP, Finite sums of commutators, *Proc. Amer. Math. Soc.* **130** (2002), 3039–3041.
- [Rob16] L. ROBERT, On the Lie ideals of  $C^*$ -algebras, *J. Operator Theory* **75** (2016), 387–408.
- [Win12] W. WINTER, Nuclear dimension and  $\mathcal{Z}$ -stability of pure  $C^*$ -algebras, *Invent. Math.* **187** (2012), 259–342.

HANNES THIEL, DEPARTMENT OF MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY AND UNIVERSITY OF GOTHENBURG, GOTHENBURG SE-412 96, SWEDEN.

*Email address:* [hannes.thiel@chalmers.se](mailto:hannes.thiel@chalmers.se)

*URL:* [www.hannesthiel.org](http://www.hannesthiel.org)