

The contribution of Narrow-Line Seyfert 1 galaxies to the soft X-ray background

G. Hasinger^a I. Lehmann^{a,b} M. Schmidt^c J.E. Gunn^d
D.P. Schneider^b R. Giacconi^e J. Trümper^f G. Zamorani^g

^a*Astrophysikalisches Institut Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany*

^b*Department of Astronomy & Astrophysics, 525 Davey Lab, The Pennsylvania State University, University Park, Pennsylvania, PA 16802, USA*

^c*California Institute of Technology, Pasadena, CA 91125, USA*

^d*Princeton University Observatory, Peyton Hall, Princeton, NJ 08540, USA*

^e*Associated Universities, Inc. 1400 16th Street, NW, Suite 730 Washington, DC 20036, USA*

^f*Max-Planck-Institute für extraterrestrische Physik, Karl-Schwarzschild-Str. 1, 85748 Garching bei München, Germany*

^g*Osservatorio Astronomico, Via Ranzani 1, I-40127, Bologna, Italy*

Abstract

The ROSAT Ultradeep HRI survey in the Lockman Hole contains a complete sample of 91 X-ray sources with fluxes in the 0.5-2 keV band larger than $1.2 \cdot 10^{-15}$ erg cm^{-2} s^{-1} , where over 75 % of the sources are quasars or Seyfert galaxies. During the course of our optical identification work, we have obtained optical spectra of 67 narrow emission line galaxies (NELG), which are physically not associated with the X-ray sources. We have derived the equivalent width (EW) and the full width at half maximum (FWHM) for the most prominent emission lines of 41 quasars and Seyfert galaxies taken from the ROSAT Deep Survey (RDS), which has a flux limit of $5.5 \cdot 10^{-15}$ erg cm^{-2} s^{-1} in the 0.5-2.0 keV band. Furthermore we have obtained the EW and FWHM values of the field NELGs. Here we present the spectroscopic discrimination between RDS Seyfert galaxies and field galaxies (NELG). The analysis of the emission lines has revealed that a single object out of 69 spectroscopically identified AGN fits the optical criteria of Narrow-Line Seyfert 1 galaxies (NLS1). This may indicate that NLS1 contribute only marginally to the soft X-ray background, but we can not exclude a possible larger contribution.

Key words: galaxies: active; quasars: general; quasars: absorption lines; X-rays: galaxies

1 The ROSAT Deep Surveys in the Lockman Hole

The most sensitive ROSAT surveys consist of a 207 ksec ROSAT PSPC exposure, a 205 ksec HRI raster scan and a total 1112 ksec HRI exposure of a 0.3 deg^2 area in the Lockman Hole region. The HRI images are the basis for the Ultradeep HRI Survey (Hasinger et al. 1999). At a flux limit of $10^{-15} \text{ erg s}^{-1}$ in the 0.5-2.0 keV energy band, the HRI survey has resolved about 70-80% of the soft X-ray background into discrete sources. The ROSAT Deep Survey (RDS), based on the PSPC image, includes a statistically complete sample of 50 X-ray sources with fluxes in the 0.5-2.0 keV band greater than $5.5 \cdot 10^{-15} \text{ erg s}^{-1}$ (Hasinger et al. 1998). The spectroscopic identification of the RDS using the Keck and Palomar telescopes have shown that about 75% of the sources are quasars and Seyfert galaxies (Schmidt et al. 1998, Lehmann et al. 2000).

Both surveys contain 91 X-ray sources, where the faintest sources reach a flux of $1.2 \cdot 10^{-15} \text{ erg s}^{-1}$ in the 0.5-2.0 keV band. Recent optical/infrared work has led to an identification of 88 of the 91 X-ray sources, confirming a high fraction of AGNs (72 objects). This is the largest fraction of AGNs found in any previous X-ray survey (Boyle et al. 1995, Georgantopolous et al. 1996, Bower et al. 1996 and McHardy et al. 1998). Among our AGNs is the most distant X-ray selected quasar at a redshift of 4.45 (Schneider et al. 1998). Groups and clusters of galaxies ($\sim 10\%$) form the second most abundant class of objects. One X-ray source has been classified as a narrow emission line galaxy (NELG), whereas some deep ROSAT PSPC surveys found a significantly larger fraction of NELGs (Boyle et al. 1995, McHardy et al. 1998). We see no evidence that NELGs or other classes of objects dominate the soft X-ray counts at faint fluxes.

2 Optical spectral properties of RDS quasars and Seyfert galaxies

In the following we discuss the emission line properties of the 41 spectroscopically confirmed RDS quasars and Seyfert galaxies. 33 X-ray sources have been identified with quasars and Seyfert 1 galaxies in the redshift range between 0.08 and 2.83. Their optical spectra show broad emission lines (FWHM $> 1500 \text{ km s}^{-1}$) of $\text{Ly}\alpha$ $\lambda 1216$, C IV $\lambda 1548$, C III] $\lambda 1908$ and Mg II $\lambda 2798$ at medium and large redshifts, or of Balmer lines ($\text{H}\alpha$ $\lambda 6563$, $\text{H}\beta$ $\lambda 4861$) at lower redshifts. Most of them can be classified as type I AGN, whereas the large Balmer decrement of two Seyfert galaxies having broad $\text{H}\alpha$ or $\text{H}\beta$ emission lines indicates type II AGN (see object 59A in Fig. 1).

The optical spectra of six RDS AGNs show only narrow emission lines with FWHM $< 1500 \text{ km s}^{-1}$ indicating a type II AGN. When their redshifts are




Fig. 2. Keck LRIS spectra of the narrow emission line AGN 26A ($z = 0.616$) from the RDS sample and the narrow emission line field galaxy (NELG) 14C at $z = 0.119$. The spectra of 26A and the NELG show a prominent galaxy continuum above the Ca H+K $\lambda\lambda 3934/3968$ break. The high-ionization [Ne V] $\lambda 3426$ emission line in the spectrum of 26A confirms the AGN nature of this object.

the RIXOS-sample (Puchnarewicz et al. 1997) and the CRSS-sample (Boyle et al. 1997), and the mean values derived from several emission line and UV/optical selected AGN samples (Schmidt et al. 1986).

lines (1)	RDS AGN EW/FWHM				Field NELG EW/FWHM			
	mean (2)	σ (3)	error (4)	n (5)	mean (6)	σ (7)	error (8)	n (9)
[Ne V] $\lambda 3426$	4/680	4/250	1/90	9/8	-	-	-	-
[O II] $\lambda 3727$	12/440	11/200	3/60	17/11	30/290	28/120	4/30	61/18
[Ne III] $\lambda 3868$	5/820	4/400	2/180	8/5	4/770	2/100	1/60	8/3
[O III] $\lambda 4959$	6/380	5/270	2/120	8/5	8/600	8/320	2/130	21/6
[O III] $\lambda 5007$	20/310	28/80	9/30	9/5	26/290	30/100	6/40	29/7

Tab. 1. Rest frame EW [\AA] and FWHM [km s^{-1}] values of the forbidden narrow emission lines of the RDS AGN in comparison with those of the field NELGs. High ionization [Ne V] $\lambda 3426$ emission lines have been detected only in AGN.

The mean RDS EW values of the forbidden narrow emission lines, found for Seyfert galaxies at redshifts below 1, are slightly smaller than those from the AGN comparison samples (Lehmann et al. 2000). The RDS EW of [Ne III] $\lambda 3868$ and the [O III] $\lambda\lambda 4959/5007$ lines seem to be in better agreement with those from field NELGs (Tab. 1) than with those from other AGN samples. The optical spectra of most RDS AGNs, which contain forbidden narrow emission lines, show several typical galaxy absorption lines, eg., Ca H+K $\lambda\lambda 3934/3968$, CH G $\lambda 4301$ (see 26A in Fig. 2). A strong continuum contribution produced by the host galaxy could account for these less stronger forbidden lines in our RDS AGNs.

4 Contribution of NLS1 to the soft X-ray background

Although the majority of our faint X-ray sources have been spectroscopically identified with quasars and Seyfert galaxies, only one out of 69 AGNs matches the optical criteria of narrow line Seyfert 1 galaxies taken from Osterbrock & Pogge (1985) and more precisely defined by Goodrich (1989). These somewhat subjective criteria are as follows. 1) NLS1 have slightly broader Balmer lines ($< 2000 \text{ km s}^{-1}$) in comparison to the forbidden lines such as [O II] $\lambda 3727$ or [O III] $\lambda\lambda 4959/5007$. 2) The flux ratio [O III] $\lambda 5007$ to $H\beta$ $\lambda 4861$ is < 3 , a level which allows to discriminate between NLS1 and Seyfert 2 galaxies (Shuder & Osterbrock 1981). 3) Emission lines of Fe II or of high ionization [Fe VII] $\lambda 6087$ and [Fe X] $\lambda 6375$ are often present.

The spectrum of the NLS1 37A (Fig. 3) shows prominent Fe II bumps at 4500–4600 Å and at 5250–5350 Å (in the rest frame). The FWHM of the broad component of the $H\beta$ $\lambda 4861$ emission line is slightly above the limit for NLS1 ($2070 \pm 20 \text{ km s}^{-1}$). The flux ratio [O III] $\lambda 5007/H\beta$ $\lambda 4861$ is clearly below 3. The NLS1 37A at $z = 0.462$ belongs to the highest-redshift objects of this class. The bright soft X-ray selected sample of Grupe et al. (1999) contains for example only 4 objects (12%) above $z = 0.3$.




Fig. 3. Keck LRIS spectrum of the NLS1 galaxy 37A ($z = 0.467$). The symbol "x" marks a region of an atmospheric absorption band ($\sim 7600 \text{ \AA}$).

The fraction of NLS1 in X-ray selected AGN samples (see for an overview Grupe in these proceedings) has to be considered with caution. NLS1 at redshifts above 0.65 could be missed due to the limited wavelength coverage of the optical spectra ($\lambda < 8200 \text{ \AA}$). The $H\beta/[O III]$ region is covered in 13 of 69 spectroscopically identified AGNs from the ROSAT Deep Surveys. Most of our AGNs show only broad UV emission lines, eg., [C III] $\lambda 1908$ or Mg II $\lambda 2798$, at high redshifts. Recently, Rodriguez-Pascual et al. (1997) have detected broad components of strong UV emission lines (eg., $L\alpha$ $\lambda 1216$ and C IV $\lambda 1548$) in several NLS1. If this is a common property we can not exclude a significantly larger fraction of NLS1 in high-redshift AGN samples.

But even the fraction of NLS1 in some low-redshift AGN samples, eg. 7% of

the ROSAT Bright Survey (Schwope et al. 2000) or 22% of the RIXOS AGN sample (Puchnarewicz et al. 1997), could result from a different classification scheme of NLS1 objects (upper limit for $\text{FWHM}_{H\beta} < 1500$ or 2000 km s^{-1}). Without well-controlled samples at low and high redshifts it is not possible to obtain reliable statements about the cosmological evolution of NLS1.

The fraction of NLS1 found in the ROSAT Deep Surveys ($\sim 1\%$) may indicate a marginal contribution of NLS1 to the soft X-ray background. Nevertheless we have to consider this as a lower limit so far.

References

- [1] R.G. Bower, G. Hasinger, F.J. Castander, et al., *MNRAS* **281** (1996) 59.
- [2] B.J. Boyle, R.G. McMahon, B.J. Wilkes, et al., *MNRAS* **272** (1995) 462.
- [3] B.J. Boyle, B.J. Wilkes, M. Elvis, et al., *MNRAS* **285** (1997) 511.
- [4] I. Georgantopoulos I., Steward G.C., Shanks T., et al., *MNRAS* **280** (1996) 276.
- [5] R.W. Goodrich, *ApJ* **342** 224.
- [6] D. Grupe, K. Beuermann, K. Mannheim, H.-C. Thomas, *A&A* **350** (1999) 805.
- [7] G. Hasinger G., R. Burg, R. Giacconi, et al., *A&A* **329** (1998) 482.
- [8] G. Hasinger, I. Lehmann, R. Giacconi, et al., *MPE-Report* **272** (1999) 199.
- [9] I. Lehmann, G. Hasinger, M. Schmidt, et al., *A&A* **354** (2000) 34.
- [10] I. McHardy, L.R. Jones, M.R. Merrifield, et al., *MNRAS* **295** (1998) 641.
- [11] D.E. Osterbrock, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, *University Science Books* (1981)
- [12] D.E. Osterbrock, R.W. Pogge, *ApJ* **297** (1985) 166
- [13] E.M. Puchnarewicz, K.O. Mason, F.J. Carrera, et al., *MNRAS* **291** (1997) 177.
- [14] P.M. Rodriguez-Pacual, J.M. Mas-Hesse, M. Santos-Lleo, *A&A* **327** (1997) 72.
- [15] M. Schmidt M., D.P. Schneider, J.E. Gunn, *ApJ* **306** (1986) 411.
- [16] M. Schmidt, G. Hasinger, J.E. Gunn, et al., *A&A* **329** (1998) 495.
- [19] D.P. Schneider, M. Schmidt, G. Hasinger, et al. em *AJ* **115**, 1230
- [18] J.M. Shuder, D.E. Osterbrock, *ApJ* **250**, 55
- [19] A. Schwope, G. Hasinger, I. Lehmann, et al., *AN* **321** (2000) 1