Dipole anisotropies of IRAS galaxies and the contribution of a large-scale local void

Kenji Tomita

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

tomita@yukawa.kyoto-u.ac.jp

ABSTRACT

Recent observations of dipole anisotropies show that the velocity of the Local Group $(v_{\rm G})$ induced by the clustering of IRAS galaxies has an amplitude and direction similar to those of the velocity of Cosmic Microwave Background dipole anisotropy $(v_{\rm CMB})$, but the difference $|v_{\rm G} - v_{\rm CMB}|$ is still ~ 170 km/s, which is about 28% of $|v_{\rm CMB}|$. Here we consider the possibility that the origin of this difference comes from a hypothetical large-scale local void, with which we can account for the accelerating behavior of type Ia supernovae due to the spatial inhomogeneity of the Hubble constant without dark energies and derive the constraint to the model parameters of the local void. It is found as a result that the distance between the Local Group and the center of the void must be $(10 - 20)h^{-1}$ Mpc, whose accurate value depends on the background model parameters.

Subject headings: cosmic microwave background - cosmology: large scale structure of the universe - observations

1. Introduction

From the observed dipole component of temperature anisotropies in the cosmic microwave background radiation (CMB), our peculiar motion relative to the CMB frame has been determined because the dipole anisotropy is connected with the motion by a Doppler effect. The velocity of the Local Group relative to the CMB frame is found to be

$$\boldsymbol{v}_{\text{CMB}} = (-25.2, -545.4, 276.5) = 612.0 \ (-0.041, -0.89, 0.45) \ \text{km s}^{-1},$$
(1)

where the x-axis and z-axis are towards the Galactic center and the Galactic pole, respectively, and the total velocity $|\boldsymbol{v}_{\text{CMB}}|$ is 612 ± 22 in the direction of $(l, b) = (268 \pm 3, 27 \pm 3)$ (Rowan-Robinson et al. 2000; Lawrence 1999).

The peculiar motion is, on the other hand, caused gravitationally by the inhomogeneous matter distribution around us. Unitil now the latter velocity of the Local Group $v_{\rm G}$ has been derived on the basis of linear gravitational instability using matter distributions that have been taken from

optical survey of galaxies, the IRAS galactic redshift survey and cluster surveys, for a long time (Yahil et al 1986; Meiksin & Davis 1986; Lahav 1987; Rowan-Robinson et al. 1990; Strauss et al. 1992; Schmoldt et al. 1999; Branchini et al. 1999; Rowan-Robinson et al. 2000). The problem of the convergence of the velocity has also been discussed in many studies (Juszkiewicz, Vittorio and Wyse 1990; Scaramella, Vettolani & Zamorani 1994; Plionis & Kolokotronis 1998).

The most recent determination of $v_{\rm G}$ was given by Rowan-Robinson et al. (2000) using the Point Source Catalogue (PSCz) IRAS galactic redshift survey, which was analyzed out to a distance $300h^{-1}$ Mpc $(H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1})$:

$$v_{\rm G} \equiv v(r_{\rm LG}) = 697.7 \ n_{\rm G} \ {\rm km \ s}^{-1},$$
 (2)

where $\mathbf{n}_{\rm G} \equiv (-0.257, -0.811, 0.482)$ represents a directional unit vector. The above two velocities $\mathbf{v}_{\rm CMB}$ and $\mathbf{v}_{\rm G}$ have similar amplitudes and directions, but, when inhomogeneities out to $300h^{-1}$ Mpc are taken into account, the difference $|\mathbf{v}_{\rm G} - \mathbf{v}_{\rm CMB}|$ reaches ~ 170 km/s, which is about 28% of $|\mathbf{v}_{\rm CMB}|$. The origin of this difference may come from a hypothetical object behind the Galactic center, inhomogeneous biasing or nonlinear correction to perturbation equations, as was indicated by Rowan-Robinson et al. (2000).

In this paper we investigate the contribution of a hypothetical local void in which the inner Hubble constant $(H_0^{\rm I})$ is larger than the outer Hubble constant $(H_0^{\rm II})$ and the radius of the boundary is ~ $200(h^{\rm I})^{-1}$ Mpc, where $H_0^{\rm I} = 100h^{\rm I}$ km s⁻¹ Mpc⁻¹ (see Fig. 1). It is assumed that we are near the center of the inner region (e.g. Tomita (2000a,b)).

Here let us briefly describe the motivations for introducing the local void and the significance of its model parameters. Firstly, the observed values of the Hubble constant appear to be inhomogeneous: the local value $(H_0^{\rm I})$ in the neighborhood is about 72 km s⁻¹ Mpc⁻¹ (Sakai et al. 2000; Freedman et al. 2001), but the global values (H_0^{II}) in remote regions are smaller than H_0^{I} , by factors which reach even 20%. The global values have been derived using gravitational lensing (Keeton & Kochanek 1997; Courbin et al. 1997; Fassnacht et al. 1999; Williams 2000; Tada & Futamase 2001; Kochanek 2002a,b,c) and the Sunyaev-Zeldovich effect (Reese et al. 2000), although they depend on cosmological models and include some uncertainties. This inhomogeneity of the Hubble constant strongly suggests the existence of a local void (on scales with the redshift z < 0.1), because, if $H_0^{\rm I} > H_0^{\rm II}$ locally, the central region must necessarily be underdense with respect to the total matter density. A local void on scales ~ 70 Mpc has once be studied by Zehavi et al. (1998) in terms of the Hubble bubble. At present, the existence of an underdense region on such scales is not clear from optical observations, athough it has so often been suggested on scales ~ 300 Mpc (Marinoni et al 1999; Marzke et al. 1998; Folkes et al. 1999; Zucca et al. 1997), and so we should expect the situation that the distribution of luminous matter (such as galaxies and clusters) is nearly homogeneous, while that of dark matter is locally inhomogeneous. This situation.

which seems to be contradictory may be realized through the difference of the feedback system in hierarchical galaxy formation between the central (local) region and the outer (global) region (Tomita 2002), since in the low-density region, more galaxies are produced owing to less supernova explosions and photoionization, than in the high-density region (Kauffmann et al. 1999; Benson et al. 2001; Sommerville et al. 2001).

Second, the accelerating behavior of high-redshift supernovae (SNIa) was found by two groups, the High-z SN Search Team (Schmidt et al. 1998; Riess et al. 1998, 1999) and the Supernova Cosmology Project (Perlmutter et al. 1999). Its origin has usually been explained by considering the role of nonzero cosmological constant Λ or dark energies in homogeneous cosmological models. Because the existence of nonzero Λ or dark energies is not evident theoretically, however, the search for other explanations may be significant. In the model with $H_0^{\rm I} > H_0^{\rm II}$, on the other hand, the underdense region (the local void) plays the role of vacuum, and supernovae there show accelerating behaviors, even though we have no Λ and dark energies. In our recent papers (Tomita 2001a,b), it was shown that, for larger $(H_0^{\rm I} - H_0^{\rm II})/H_0^{\rm II}$, the magnitude (m) and redshift (z) relation of highredshift supernovae can be well reproduced in the models with smaller Λ , and, if $(H_0^{\rm I} - H_0^{\rm II})/H_0^{\rm II} =$ 0.2, the relation can be quantitatively explained in the model which reduces to the Einstein-de Sitter model. The best radius of the boundary for fitting the m-z relation is found to be ~ 200 Mpc. Moreover the latest data of a supernova with $z \sim 1.7$ (Riess et al. 2001) are naturally consistent with this last model with $\Lambda = 0$ (Tomita 2001b). For cosmological parameters in the inner region, we take the low-density value ($\Omega_0^{\rm I} \sim 0.3$), corresponding to the statistical analyses for local observations of galaxies and clusters. In the outer region, spatially flat models are chosen in accord with the recent results of CMB observations (Lange et al. 2001; Stomper et al. 2001; Pryke et al. 2001), and our models with small or vanishing Λ are found to be compatible with their data in the weak prior.

Third, bulk flow on the scale more than 100 Mpc was found by by Hudson et al. (1999) and Willick (1999). It is impossible to account for its appearance as one of the linear perturbations in homogeneous cosmological models. In a model with a local void on scales ~ 200 Mpc, however, we can interpret the bulk flow as the nonlinear deviation of the local flow from the global expansion of the universe. This point is also one of the motivations for our inhomogeneous cosmological model.

If a smooth power spectrum is assumed for the density perturbations, it is to be noticed here that the underdense nonlinear structure itself as a local void on scales ~ 200 Mpc is a unique object which evolved gravitationally from primordial adiabatic perturbations but whose amplitude deviates remarkably from that of average linear perturbations, so that it can be found at most only a few times within our horizon, and we cannot see anything similar in the neighborhood outside our local void. If similar structures were found more frequently, we would have a power spectrum with a sharp spike corresponding to their size.

Owing to the gap of Hubble constants, an inner comoving observer has the additional velocity $v_o [= 100(h^{\rm I} - h^{\rm II})R_o \text{ km/s}]$ relative to the CMB (outer) frame, if the distance between the observer

and the center of the inner region is R_o Mpc. In a previous paper (Tomita 2000b) we derived the dipole moment D of CMB anisotropy caused by the additional velocity, and the Doppler velocity v_d corresponding to D, which depends on the model parameters in the inner and outer regions. If a certain condition for the parameters is satisfied, this velocity v_d gives the above-mentioned difference, so that we may have

$$\boldsymbol{v}_d = 166.6 \ \boldsymbol{n}_d \ \mathrm{km \ s}^{-1},$$
 (3)

where $\mathbf{n}_d \equiv (0.925, 0.124, 0.360)$ is also a directional unit vector. In §2 the expressions for the peculiar velocity in the linear approximation are derived in our model with a local void, and their difference from the homogeneous case is shown. In §3 the derivation of D and v_d is briefly accounted for and their numerical values are shown, and the above condition is discussed. Some mistakes found in a table in the previous paper are corrected there. In §4, it is shown that the bulk flows can be caused as the sum of the large-scale peculiar velocity and the additional velocity v_o . Section 5 is dedicated to concluding remarks.

2. Peculiar velocity induced by inhomogeneities

Our simple model with a local void consists of the inner homogeneous region $(V_{\rm I})$ and the outer homogeneous region $(V_{\rm II})$, whose boundary is spherical. The line elements are expressed as

$$ds^{2} = g^{j}_{\mu\nu} (dx^{j})^{\mu} (dx^{j})^{\nu} = -c^{2} (dt^{j})^{2} + [a^{j}(t^{j})]^{2} \Big\{ d(\chi^{j})^{2} + [f^{j}(\chi^{j})]^{2} d\Omega^{2} \Big\},$$
(4)

where $j \ (= I \text{ or II})$ represents the regions, $f^j(\chi^j) = \sin \chi^j, \chi^j$ and $\sinh \chi^j$ for $k^j = 1, 0, -1$, respectively, and $d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2$. The boundary shell is a time-like hypersurface Σ given as $\chi^{\rm I} = \chi^{\rm I}_1$ and $\chi^{\rm II} = \chi^{\rm II}_1$.

The background models in $V^{\rm I}$ and $V^{\rm II}$ are specified by model parameters H_0^j , Ω_0^j and λ_0^j for $j = {\rm I}$ and II, where $\Omega_0^j \equiv [8\pi G/3(H_0^j)^2](\rho_0)^j$ and $\lambda_0^j \equiv \Lambda c^2/(H_0^j)^2$, and expressed using $y^j (\equiv a^j/(a_0)^j)$ and $\tau^j (\equiv H_0^j t^j)$ as

$$dy^{j}/d\tau^{j} = (y^{j})^{-1/2} P_{j}(y^{j}),$$
(5)

where

$$P_j(y^j) \equiv [\Omega_0^j + \lambda_0^j (y^j)^3 + (1 - \Omega_0^j - \lambda_0^j) y^j]^{1/2}$$
(6)

and a_0^j is given by

$$a_0^j H_0^j = 1/\sqrt{1 - \Omega_0^j - \lambda_0^j}.$$
(7)

Their Hubble constants and density parameters in the two regions are assumed to satisfy the relations $H_0^{\rm I} > H_0^{\rm II}$ and $\Omega_0^{\rm I} < \Omega_0^{\rm II}$.

In this section we consider local behaviors of matter motion in the Newtonian treatment. In terms of the common time coordinate t and Cartesian coordinates X, the equations of mass continuity, motion and gravitation are

$$\frac{\partial \rho}{\partial t} + \nabla_{\boldsymbol{X}}(\rho \boldsymbol{V}) = 0, \tag{8}$$

$$\frac{\partial \boldsymbol{V}}{\partial t} + (\boldsymbol{V} \cdot \nabla_{\boldsymbol{X}})\boldsymbol{V} + \nabla_{\boldsymbol{X}} \Phi = 0, \qquad (9)$$

$$\nabla_{\boldsymbol{X}}^2 \Phi = 4\pi G\rho,\tag{10}$$

where ρ , V and Φ are the mass density field, the velocity field and the gravitational potential, and X represents the proper distance. The background motion is expressed using uniform densities $(\rho = \bar{\rho})$ and velocities $(V \propto X)$ in both regions. In order to describe local inhomogeneous motions, let us next use for X the comoving coordinates r in the inner region, where $X = a_{\rm I}r$ and consider linear density and velocity perturbations $\delta_{\rm I} \equiv (\rho - \bar{\rho}_{\rm I})/\bar{\rho}_{\rm I}$ and v in the inner region, where $\bar{\rho}_{\rm I}$ is the background (unperturbed) density in the inner region. Then the first two equations reduce to

$$\frac{\partial \delta_{\mathrm{I}}}{\partial t} + \frac{1}{a_{\mathrm{I}}} \nabla \boldsymbol{r} \cdot \boldsymbol{v} = 0, \tag{11}$$

$$\frac{\partial \boldsymbol{v}}{\partial t} + \frac{\dot{a}_{\mathrm{I}}}{a_{\mathrm{I}}} \boldsymbol{v} + \frac{1}{a_{\mathrm{I}}} \nabla_{\boldsymbol{r}} \phi = 0.$$
(12)

The gravitational equation is

$$\frac{1}{a_{\mathrm{I}}^2} \nabla \boldsymbol{r}^2 \phi = 4\pi G(\rho - \bar{\rho}),\tag{13}$$

which holds for ρ in both regions and the background density $\bar{\rho}$ is represented by $\bar{\rho}_{\rm I}, \bar{\rho}_b$ and $\bar{\rho}_{\rm II}$ in the inner region, the shell, and the outer region, respectively. The solution for equation (13) is expressed as

$$\phi = -Ga_1^2 \int \frac{\bar{\rho}_1 \delta_1}{|\boldsymbol{r} - \boldsymbol{r}'|} d^3 \boldsymbol{r}'$$
(14)

or

$$\phi = -Ga_{\mathrm{I}}^{2} \Big[\int_{\mathrm{I}} \frac{\bar{\rho}_{\mathrm{I}} \delta_{\mathrm{I}}}{|\boldsymbol{r} - \boldsymbol{r}'|} d^{3}\boldsymbol{r}' + \frac{1}{a_{\mathrm{I}}} \int_{\mathrm{shell}} \frac{(\delta\rho)_{b} r_{b}^{2}}{|\boldsymbol{r} - \boldsymbol{r}'_{\mathrm{b}}|} \sin \theta' d\theta' d\varphi' + \int_{\mathrm{II}} \frac{(\delta\rho)_{\mathrm{II}}}{|\boldsymbol{r} - \boldsymbol{r}'|} d^{3}\boldsymbol{r}' \Big],$$
(15)

where \mathbf{r}_b and $(\delta \rho)_b (\equiv (\rho - \bar{\rho})_b)$ are the coordinate and density perturbation in the boundary shell, respectively, and $(\delta \rho)_{\text{II}} \equiv \rho - \bar{\rho}_{\text{II}}$.

Solving Eqs. (11) and (12), we obtain the velocity field in the inner region:

$$\boldsymbol{v}(\boldsymbol{r}) = \frac{a_{\mathrm{I}}}{4\pi} \int \frac{\dot{\delta}_{\mathrm{I}}(\boldsymbol{r}-\boldsymbol{r}')d^{3}\boldsymbol{r}'}{|\boldsymbol{r}-\boldsymbol{r}'|^{3}} + \frac{G}{a_{\mathrm{I}}} \int_{\mathrm{shell}}^{t} \int_{0}^{t} a_{\mathrm{I}}(\delta\rho)_{b} dt \frac{r_{b}^{2}(\boldsymbol{r}-\boldsymbol{r}_{b}')}{|\boldsymbol{r}-\boldsymbol{r}_{b}'|^{3}} \sin\theta' d\theta' d\varphi' + \frac{G}{a_{\mathrm{I}}} \int_{\mathrm{II}} \int_{0}^{t} \bar{\rho}_{\mathrm{I}} a_{\mathrm{I}}^{2} \delta_{\mathrm{I}} \Big[\frac{(\delta\rho)_{\mathrm{II}}}{(\delta\rho)_{\mathrm{I}}} - 1 \Big] dt \frac{(\boldsymbol{r}-\boldsymbol{r}')}{|\boldsymbol{r}-\boldsymbol{r}'|^{3}} d^{3}\boldsymbol{r}',$$
(16)

where the first integral in the right-hand side of equation (16) is performed in both regions and we used the equation for density perturbation $\delta_{\rm I}$:

$$\frac{\partial^2 \delta_{\rm I}}{\partial t^2} + 2 \frac{\dot{a}_{\rm I}}{a_{\rm I}} \frac{\partial \delta_{\rm I}}{\partial t} = 4\pi G \bar{\rho}_{\rm I} \delta_{\rm I}. \tag{17}$$

If we adopt the solution $D_{\rm I}$ in the growing mode in equation (17), and use $f^{\rm I} \equiv d \ln D_{\rm I}/d \ln a_{\rm I}$, we obtain

$$\boldsymbol{v}(\boldsymbol{r}) = \frac{a_{\mathrm{I}}}{4\pi} H_0^{\mathrm{I}} f^{\mathrm{I}} \int \frac{\delta_{\mathrm{I}}(\boldsymbol{r} - \boldsymbol{r}') d^3 \boldsymbol{r}'}{|\boldsymbol{r} - \boldsymbol{r}'|^3} + Res, \qquad (18)$$

$$Res \equiv \int_{\text{shell}} J_1 \, \frac{(r_b)^2 (\boldsymbol{r} - \boldsymbol{r}_b')}{|\boldsymbol{r} - \boldsymbol{r}_b'|^3} \sin \theta' d\theta' d\varphi' + \int_{\text{II}} J_2 \, \frac{a_{\text{I}}(\boldsymbol{r} - \boldsymbol{r}')}{|\boldsymbol{r} - \boldsymbol{r}'|^3} d^3 \boldsymbol{r}', \tag{19}$$

where

$$J_{1} \equiv \frac{G}{(a_{\rm I})} \int_{0}^{t} a_{\rm I}(\delta\rho)_{\rm I} dt,$$

$$J_{2} \equiv \frac{G}{(a_{\rm I})^{2}} \int_{0}^{t} \left(\frac{(\delta\rho)_{\rm II}}{(\delta\rho)_{\rm I}} - 1\right) \bar{\rho}_{\rm I} a_{\rm I}^{2} \delta_{\rm I} dt.$$
(20)

The first term (A) on the right-hand side of equation (18) represents the peculiar velocity induced in the model that is homogeneous in the entire region with $H_0^{\rm I}$ and $\Omega_0^{\rm I}$. The second term (*Res*) gives the contribution of the local void to the peculiar velocity, which comes from the inhomogeneous matter distribution. In the homogeneous case without Res, the above expression accords with the usual one (e.g. Strauss & Willick (1995)). The peculiar velocity of the Local Group $\boldsymbol{v}_{\rm G}$ is given by the velocity at $\boldsymbol{r} = \boldsymbol{r}_{\rm LG}$ as $\boldsymbol{v}_{\rm G} = \boldsymbol{v}(\boldsymbol{r}_{\rm LG})$.

The integrands in A and *Res* are proportional to $|\mathbf{r} - \mathbf{r}'|^{-2}$, and the main contributions in the integrations A and *Res* come from the region of distances ~ 40 Mpc and > 200 Mpc, respectively. Accordingly, *Res* may be smaller by the factor 5^{-2} than A, so that the rough estimate for the peculiar velocity is given by A and *Res* is negligible compared with A.

Under the assumption that the small-scale density perturbations in the luminous matter is $\beta \times$ those in the dark matter and β is ≈ 1 , $v(\mathbf{r})$ is calculated from equation (18) with Res = 0 by investigating the inhomogeneity of spatial distributions in galactic redshift surveys. The values of $v(\mathbf{r}_{\text{LG}})$ were derived by Rowan-Robinson et al. (2000) using the PSCz redshift survey including 15,459 galaxies. When we average their values (in Table 1 of their paper) over the interval of $|\mathbf{r}| = (200 - 300)h^{-1}$ Mpc, we obtain the expression in equation (2).

3. Doppler velocity due to the CMB anisotropy for an off-center comoving observer

In this section we consider the CMB anisotropy that is measured by an off-center observer in the inner region. For this purpose we must investigate the behavior of light rays in our model with a local void, by solving null-geodesic equations and extracting the multipole components. The detail is described in the previous paper (Tomita 2000b). Here we pay attention to light rays that are emitted at the recombination epoch in the outer region and reached the observer at the present epoch in the inner region. The redshift $(z_{\rm rec}^{\rm II})$ corresponding to the recombination epoch depends on the direction ϕ between the light rays and $\overrightarrow{\rm CO}$. The value of $z_{\rm rec}^{\rm II}$ is numerically calculated for $0 < \phi < \pi$, and the dipole moment is derived. When $z_{\rm rec}$ (= $z_{\rm rec}^{\rm II}(\phi)$) is given, the temperature $T(\phi)$ of the cosmic background radiation is proportional to $1/(1 + z_{\rm rec})$, and the dipole moment D and quadrupole moment Q are defined as

$$D \equiv \left| \int_{0}^{\pi} \int_{0}^{2\pi} (1+z_{\rm rec})^{-1} Y_{10} \sin \phi d\phi d\varphi \right| / \langle (1+z_{\rm rec})^{-1} \rangle,$$

= $2\pi \left| \int_{0}^{\pi} (1+z_{\rm rec})^{-1} Y_{10} \sin \phi d\phi \right| / \langle (1+z_{\rm rec})^{-1} \rangle,$ (21)

and

$$Q \equiv \left| \int_{0}^{\pi} \int_{0}^{2\pi} (1+z_{\rm rec})^{-1} Y_{20} \sin \phi d\phi d\varphi \right| / \langle (1+z_{\rm rec})^{-1} \rangle,$$

= $2\pi \left| \int_{0}^{\pi} (1+z_{\rm rec})^{-1} Y_{20} \sin \phi d\phi \right| / \langle (1+z_{\rm rec})^{-1} \rangle,$ (22)

where $\langle \rangle$ means the average value taken over the whole sky, and

$$Y_{10}(\phi) = \sqrt{\frac{3}{4\pi}} \cos \phi,$$

$$Y_{20}(\phi) = \sqrt{\frac{5}{4\pi}} \left(\frac{3}{2} \cos^2 \phi - \frac{1}{2}\right).$$
(23)

The Doppler velocity v_d corresponding to D is given by

$$v_d \equiv c[(3/4\pi)^{1/2}D].$$
 (24)

This velocity v_d is a function of $R \ (\equiv a_0^{\mathrm{I}}\chi_0^{\mathrm{I}}), z_b, (H_0^{\mathrm{I}}, \Omega_0^{\mathrm{I}})$ and $(H_0^{\mathrm{II}}, \Omega_0^{\mathrm{II}})$. Here we fix H_0^{I} as $h^{\mathrm{I}} = 0.7$, and fix z_b as $z_b = 0.067$, which corresponds to the case $r_b (\equiv a_b^{\mathrm{I}}\chi_b^{\mathrm{I}}) = 200(h^{\mathrm{I}})^{-1}$ Mpc. In the 6th and 7th columns of Table 1, D and v_d for $R = 10(h^{\mathrm{I}})^{-1}$ Mpc are displayed for various model parameters of the background models. First seven models are spatially flat in the outer region, and next three models are open. In the inner region all models are open. In the last column of Table 1 the values of $R \ (\equiv R_{170})$ corresponding to $v_d = 170 \text{ km s}^{-1}$ are shown. It is found from this table that the distance R must be in the interval $(10 - 20)(h^{\mathrm{I}})^{-1}$ Mpc.

On the other hand, we assume that the Doppler velocity vector v_d gives the difference between $v_{\rm G}$ and $v_{\rm CMB}$. From the comparison between $v_{\rm G}$ in equation (2) and $v_{\rm CMB}$ in equation (1), then the estimated v_d is given by equation (3). It is to be noticed that this velocity is in the direction of the Galactic center.

The above derivation of D and v_d is quite the same as that in Tomita (2000b), but it was found that the expressions for D, Q and v_d in Table 1 of the previous paper have some mistakes with respect to a factor 2π , and we should read $D(\times 10^4), Q(\times 10^4)$ and v_d (km s⁻¹) as $D(\times 10^4/2\pi), Q(\times 10^4/2\pi)$ and v_d (km s⁻¹/2 π) in Table 1 of Tomita (2000b).

// Table 1 //

4. Large-scale bulk flows

At each point in the inner region we have a peculiar velocity v(r) given by equation (18) that is caused by the inhomogeneous matter distribution. Averaging the velocity over scales ~ r_{bulk} longer than $50h^{-1}$ Mpc, we obtain the bulk flow velocity $v_b(r_{\text{bulk}})$. Branchini et al. (1999) derived the bulk flow velocity for IRAS galaxies in PSCz redshift survey including 11206 galaxies, measured out to $200h^{-1}$ Mpc. Its average value is $400 - 500 \text{ km s}^{-1}$ for $R_{\text{bulk}} (\equiv a_0 r_{\text{bulk}}) = 60h^{-1}$ Mpc and $350 - 450 \text{ km s}^{-1}$ for $R_{\text{bulk}} = 80h^{-1}$ Mpc. The amplitude and direction of bulk flows are sensitive to the measured scales of the matter distribution. If the scales are extended to more than $300h^{-1}$ Mpc, the average values of the velocities may increase over the above ones.

In our inhomogeneous models with a local void, we have not only the above bulk flow velocity $\boldsymbol{v}_b(\boldsymbol{r})$ corresponding to the peculiar velocity in the region V^I, but also the velocity $(\boldsymbol{v}_{\rm I})$ of the comoving observer (in V^I) relative to the rest frame in V^{II} which is the CMB frame.

The latter velocity \mathbf{v}_{I} is equal to $(H_0^{\mathrm{I}} - H_0^{\mathrm{II}})R_g$ in the direction $\overrightarrow{\mathrm{CG}}$ from the center to each galaxy G, where R_g is the distance between C and G. This velocity \mathbf{v}_{I} is expressed as the sum of the radial component \mathbf{v}_r (in the direction $\overrightarrow{\mathrm{OG}}$) and the constant component \mathbf{v}_o (in the direction $\overrightarrow{\mathrm{CO}}$), where $|\mathbf{v}_o| = (H_0^{\mathrm{I}} - H_0^{\mathrm{II}})R_o$ and R_o is the distance between C and O, i.e., $\mathbf{v}_{\mathrm{I}} = \mathbf{v}_r + \mathbf{v}_o$. The constancy of this component was shown in the previous paper (Tomita 2000b, 2002). The constant component \mathbf{v}_o behaves as a peculiar velocity, while the radial component \mathbf{v}_r can be regarded as part of the Hubble motion from the viewpoint of the observer O. Accordingly, the total observed bulk flow velocity $\mathbf{V}_{\mathrm{bulk}}$ is practically expressed using $\mathbf{v}_b(\mathbf{r}_{\mathrm{bulk}})$ and \mathbf{v}_o as

$$\boldsymbol{V}_{\text{bulk}} = \boldsymbol{v}_b(\boldsymbol{r}_{\text{bulk}}) + \boldsymbol{v}_o. \tag{25}$$

Here let us estimate this velocity, tentatively using one of the values of Branchini for $v_b(r_{\text{bulk}})$ on a scale of $60h^{-1}$ Mpc. Assuming the PSCz M1 method we adopt

$$\boldsymbol{v}_b(\boldsymbol{r}_{\text{bulk}}) = 450 \ \boldsymbol{n}_{\text{G}} \ \text{km s}^{-1}, \tag{26}$$

whose direction is approximately taken to be equal to that of $\boldsymbol{v}_{\rm G}$ derived by Rowan-Robinson et al. (2000). We consider the case $\Omega_0^{\rm I} = 0.3$, $\Omega_0^{\rm II} = 1.0$ and $\lambda_0^{\rm I} = \lambda_0^{\rm II} = 0$ and $h_0^{\rm II}/h_0^{\rm I} = 0.82$, which is the best case for representing the SN data including the data of z = 1.7 (Tomita 2002). If we take $R_o = 10/h_0^{\rm I}$ as in Table 1, we have

$$v_o = 180 \ n_d \ \mathrm{km \ s^{-1}}.$$
 (27)

Then we obtain from equation (25)

$$V_{\text{bulk}} = 378.3 \ n_{\text{bulk}} \ \text{km s}^{-1},$$
 (28)

where $n_{\text{bulk}} \equiv (0.135, -0.906, 0.402)$ is a directional unit vector, so that the total bulk flow velocity has the amplitude 378 km s⁻¹, which is 62% of v_{CMB} , but its direction is comparatively close to that of v_{CMB} in equation (1).

5. Concluding remarks

In this paper we first derived the expression for peculiar velocities $v_b(\mathbf{r})$ in the inner region, which are caused gravitationally by linear matter density perturbations. Next we derived the Doppler velocity $v_d(\mathbf{r})$ corresponding to the component of the CMB anisotropy caused by a local void. Using these results we obtained some constraints to our observer's position O relative to the center C in the local void, under the assumption that v_d gives the difference between v_{CMB} and $v_{\text{G}} (\equiv v(\mathbf{r}_{\text{LG}}))$ for the Local Group. It was found that the distance CO must be $(10 - 20)/h^{\text{I}}$ Mpc, depending on the parameters of the background models, and $\overrightarrow{\text{CO}}$ is in the direction of the Galactic center.

Moreover, we estimated the amplitude and the direction of the bulk flow velocities on scales of $\sim 60h^{-1}$ Mpc, as the sum of $\boldsymbol{v}_b(\boldsymbol{r})$ and the additional flow velocity velocity \boldsymbol{v}_o in the direction of \overrightarrow{CO} which is specified so as to give the difference between the inner and outer Hubble constants.

In our previous treatments (Tomita 2000b, 2002) we considered only the additional velocity to explain the origin of the bulk flow velocities obtained by Hudson et al. (1999) and Willick (1999). However, when we consider the peculiar velocity at the same time, the amplitude of the resultant bulk flow velocity is constrained by the observation so as to be much smaller than their flow velocities ($\approx 700 \text{ km/s}$), but the directions may be roughly consistent.

In §5 we neglected *Res* for the estimation of $v_{\rm G}$, but because *Res* is of the order of 4% of $v_{\rm G}$, it may give to $v_{\rm G}$ an error ~ 25 km s⁻¹, so that it may bring an error ~ 14% to the difference of the velocities ~ 170 km s⁻¹, or the distance CO (~ 10 h^{-1} Mpc).

This work was supported by Grant-in Aid for Scientific Research 12440063 from the Ministry of Education, Science, Sports and Culture, Japan.

REFERENCES

- Benson, A.J., Frenk, C.S., Baugh, C.M., Cole, S. and Lacey, C.G. 2001, MNRAS, 327, 1041
- Branchini, E. et al. 1999, MNRAS, 308, 1
- Courbin, F., Magain, P., Keeton, C.R., et al. 1997, A&A, 324, L1
- Fassnacht, C.D., Pearson, T.J., et al. 1999, ApJ, 527, 498
- Folkes, S., Ronen, S., Price, I., Lahav, O., et al. 1999, MNRAS, 308, 459
- Freedman, W.L., Madore, B.F., Gibson, B.K., et al. 2001, ApJ, 553, 47
- Hudson, M. J., Smith, R. J., Lucey, J. R., Schlegel D. J. and Davies, R. L. 1999, ApJ, L79

- Juszkiewicz, R., Vittorio, N., Wyse R.F.G., 1990, ApJ, 349, 408
- Kauffmann, G., Colberg, J.M., Diaferi, A. and White, S.D.M. 1999, MNRAS, 303, 188
- Keeton, C.R. and Kochanek, C.S. ApJ, 487, 42
- Kochanek, C.S. 2002a, astro-ph/0204043
- Kochanek, C.S. 2002b, ApJ, 578, 25
- Kochanek, C.S. 2002c, astro-ph/0206006
- Lahav, O. 1987, MNRAS, 225, 213
- Lange, A.E. et al. 2001, Phys. Rev. D, D63, 042001
- Lawrence, A. 1999, MNRAS, 308, 897
- Marinoni, C., Monaco, P., Giuricin, G. and Costantini, B. 1999, ApJ, 521, 50
- Marzke, R.O., da Costa, L.N., Pellegrini, P.S., Willmer, C.N.A. and Geller, M.J. 1998, ApJ, 503, 617
- Meiksin, A. and Davis M. 1986, AJ, 91, 191
- Perlmutter et al. 1999, ApJ, 517, 565
- Plionis M. and Kolokotronis, V. 1998, ApJ, 500,1
- Pryke, C. et al. 2002, ApJ, 568, 46
- Reese, E.D. et al. 2000, ApJ, 533, 38
- Riess, A.G. et al. 1999, AJ, 118, 2668
- Riess, A.G. et al. 1998, AJ, 116, 1009
- Riess, A.G. et al. 2001, ApJ, 560, 49
- Rowan-Robinson, M. et al. 1990, MNRAS, 247, 1
- Rowan-Robinson, M. et al. 2000, MNRAS, 314, 375
- Sakai, S. et al. 2000, ApJ, 529, 698
- Scaramella, R., Vettolani, G. & Zamorani, G. 1994, ApJ, 422, 1
- Schmidt, B.P. et al. 1998, ApJ,507,46
- Schmoldt, I. et al. 1999, MNRAS, 304, 893

- Sommerville, R.S. et al. 2001, MNRAS, 320, 289
- Stomper, R. et al. 2001, ApJ, 561, L7
- Strauss M.A. and Willick J.A. 1995, Phys. Rep. 261, 271
- Strauss M.A. et al. 1992, ApJ, 397, 395
- Tada, M. and Futamase, T. 2001, Prog. Theor. Phys. 104, 971
- Tomita K. 2000, ApJ, 529, 26
- Tomita K. 2000, ApJ, 529, 38
- Tomita K. 2001, MNRAS, 326, 287
- Tomita K. 2001, Prog. Theor. Phys., 106, 929, astro-ph/0104141
- Tomita K. 2002, Prog. Theor. Phys., 108, 103, astro-ph/0203125
- Willick, J. A. 1999, ApJ, 522, 647
- Williams, L.L.A. 2000, AJ, 119, 439
- Yahil, A., Walker, D., and Rowan- Robinson, M. 1986, ApJ, 301, L1
- Zehavi, I. Riess, A.G., Kirshner, R.P. and Dekel, A. 1998, ApJ, 503, 483
- Zucca, E., et al. 1997, A&A, 326, 477

This preprint was prepared with the AAS ${\rm IAT}_{\rm E}{\rm X}$ macros v5.0.





Fig. 1.— A simple model with a spherical boundary. A light ray is received by an observer O being near the center C, and z_b and z_{rec} are the redshifts at the boundary and at the recombination epoch, respectively, depending on angle ϕ .

Table 1: Dipole moment D and the velocity v_d in the case $z_b = 0.067$ and $h^{\rm I} = 0.7$.

Ω_0^1	Ω_0^{Π}	λ_0^1	λ_0^{Π}	h^{11}/h^1	$D(\times 10^{4})$	$v_d \; (\mathrm{km/sec})$	$R_{170}(\times h^{\mathrm{I}}) \mathrm{Mpc}$
0.3	1.0	0.0	0.0	0.82	11.76	172.8	9.8^{1}
0.4	1.0	0.0	0.0	0.82	9.59	141.0	12.1^{1}
0.5	1.0	0.0	0.0	0.82	7.66	112.6	15.1^{1}
0.6	1.0	0.0	0.0	0.82	5.90	86.7	19.6^{1}
0.3	0.6	0.205	0.4	0.80	7.47	109.9	15.5^{1}
0.3	0.6	0.303	0.4	0.87	6.72	98.9	17.2^{1}
0.4	0.6	0.269	0.4	0.82	5.31	78.0	21.8^{1}
0.3	0.6	0.0	0.0	0.80	6.83	100.4	16.9^{2}
0.3	0.6	0.0	0.0	0.87	6.53	95.9	17.7^{2}
0.4	0.6	0.0	0.0	0.82	4.44	65.2	26.1^2

¹The outer space is flat.

 2 The outer space is open.