

Spectroscopy of Planetary Nebulae in Sextans A and Sextans B

Laura Magrini¹, Pierre Leisy^{2,3}, Romano L. M. Corradi³, Mario Perinotto¹, Antonio Mampaso², and José Vilchez⁴

¹ Dipartimento di Astronomia e Scienza dello Spazio, Università di Firenze, Italy

² Instituto de Astrofísica de Canarias, Tenerife, Canarias, Spain

³ Isaac Newton Group of Telescopes, La Palma, Canarias, Spain

⁴ Instituto de Astrofísica de Andalucía, Granada, Spain

Abstract. Sextans A and Sextans B are two dIr galaxies situated in the outskirts of the Local Group (LG), in which both PNe and H II regions have been detected (Jacoby & Lesser [5], Magrini et al. [6], [7]). We present spectroscopic observations of PNe and HII regions in these two galaxies (obtained with the VLT) Preliminary results about PNe physico-chemical properties are presented.

1 The galaxies: Sextans A and Sextans B

Sextans A and Sextans B are both dwarf Irregular galaxies (Ir V and Ir IV-V morphological types, respectively, cf. van den Bergh [12], hereafter vdB00) with approximately the same V luminosity, belonging to the outer fringes of the Local Group. Their distances from the barycenter of the LG are quite similar (1.60 and 1.72 Mpc, see vdB00) and their separation on the sky is relatively small (~ 10 degree), which corresponds to about 280 kpc. Moreover their velocity difference is only 23 ± 6 km s⁻¹. These results suggest the hypothesis of a common formation of these two galaxies, probably together with NGC 3109 and the Antlia galaxy. Considering the mean distance of the four galaxies from the barycenter of the LG, 1.7 Mpc, this sub-group is located beyond the zero velocity surface of the LG (cf. vdB00) and thus it can be considered the nearest external group of galaxies. As said by Mateo ([9]) “No two Local Group dwarfs have the same star-formation history”. This is true also when comparing very similar galaxies. In fact, in spite of a probable common formation, Sextans A and Sextans B show a different star formation history as indicated by the different amounts of stars in the various evolutionary phases, and also reflected in the different number of PNe and HII regions observed. Sextans A shows a large amount of old stellar population, whereas it has a modest intermediate (4-10 Gyrs ago) and recent star formation (1-4 Gyrs ago). On the other hand, Sextans B exhibits a very strong recent star formation (1-4 Gyrs ago), together with a very old population (cf. Mateo’s ([9] review about the star formation history of the LG dwarfs).

In this context, PNe represent an useful tool to trace the chemical enrichment history of galaxies, moreover when compared with chemical abundance derived

from H II regions, because they are tracers of stellar population in a wide range of ages, from old to intermediate age.

2 Observations and data reduction

PNe (1 in Sextans A and 5 in Sextans B) and H II regions (10 and 9, respectively) have been observed in December 2003 with VLT (ESO, Paranal) equipped with FORS2 spectrograph in multi-object spectroscopy (MOS) observing mode. The FORS2 spectrograph was used with two setup: the 300 V and 300 I gratings providing a dispersion of $3.0 \text{ \AA}/\text{pixel}$. The total resulting spectral range, from $\sim 3200 \text{ \AA}$ to $\sim 10000 \text{ \AA}$, includes basic lines needed for the determination of chemical abundances. For each galaxy, a total of six exposures were taken, three with the 300 V grating (total exposure time 5400 s) and three using the 300 I one (3600 s). A spectrophotometric standard star, GD 108, was observed once during each night. The data were reduced using the IRAF LONGSLIT package and complemented with MIDAS (MOS and LONG packages). The spectra are presented in Figs. 1 and 2. Identified emission lines are marked with the name of the corresponding ion and/or wavelength.




Fig. 1. VLT spectra of the planetary nebula in Sextans A

Table 1. Chemical abundances of Sextans A and Sextans B planetary nebulae derived with the ICFs method. Metal abundances are expressed in $12+\log(X/H)$.

Name	He/H	N/H	O/H	Ne/N	Ar/H	S/H
SexA-PN	0.085±0.01	8.45±0.1	8.0±0.1	6.7±0.2	5.2±0.3	5.8±0.1
SexB-PN1	0.082±0.02	-	8.0±0.2	7.0±0.3	-	-
SexB-PN2	0.05±0.03	-	7.65±0.3	6.7±0.3	-	-
SexB-PN3	0.083±0.01	5.4†	7.5±0.2	6.9±0.3	5.5±0.3	-
SexB-PN4	0.084±0.02	5.5†	8.0±0.3	7.0±0.3	-	5.4‡
SexB-PN5	0.118±0.02	8.0±0.3	8.2±0.1	7.65±0.2	-	-

Note: † this is the ionic N^+ abundance. ‡ this is the ionic S^{2+} abundance. In both † and ‡ cases, ICF(N) and ICF(S), respectively, were not computed because [O II] lines were not detected.

Table 2. Some nebular and stellar parameters of Sextans A and Sextans B planetary nebulae. † Derived with the CLOUDY model. T_* marked with : are upper limits to the stellar effective temperature obtained when the He II 4686 Å was not measurable.

Name	N_e	$T_{e[O III]}$	$T_{e[N II]}$	Radius(pc)†	T_* †	$\log(L_*/L_\odot)$ †	$c\beta$
SexA-PN	2700	21000	13400	0.13	190000	3.8	0.22
SexB-PN1	-	12900	-	0.08	50000:	3.2	0.11
SexB-PN2	-	17300	-	0.02	73000:	3.3	-
SexB-PN3	-	17800	-	0.03	74000:	3.5	0.11
SexB-PN4	-	13300	-	0.03	70000:	3.1	-
SexB-PN5	-	12700	-	0.01	84000	3.4	0.14

this definition, the Sextans A PN is a Type I PN, while the Sextans B PNe are non-Type I. Using the stellar luminosity and temperature derived with the CLOUDY model, we compared the location of the six PNe in the $\log T_*$ - $\log L_*$ space with the H-burning evolutionary tracks from Vassiliadis & Wood ([13]). We found a more massive progenitor for the Sextans A PN ($\sim 2.5M_\odot$) than those of Sextans B PNe ($\sim 1-1.5M_\odot$), in agreement with its Type I nature.

Another interesting aspect from Table 1 is the large range of O/H, ~ 0.7 dex, of Sextans B PNe in spite of their similar evolutionary age. Since oxygen is not processed by PN progenitors, this represents a sign of the dishomogeneity of the interstellar medium at the time of the formation of the PN progenitors. We plan to compare this dishomogeneity with the spatial distribution of H II regions chemical abundances.

4 The metallicity-luminosity relationship using PNe

In Fig. 3, the relationship of V-luminosity vs. oxygen abundance from PNe in the LG is shown. Galaxies are identified with different symbols according to their morphological type. A different behaviour of star forming (irregulars and




Fig. 3. Relationship of V-luminosity vs. oxygen abundance from PNe in the LG. PN abundances: Sextans A and Sextans B from this work, other galaxies data present in literature.

spirals) and non-star forming (spheroidals) galaxies was already noted by several authors (cf. Mateo [9]): spheroidal galaxies are generally metal richer than irregular galaxies with the same V luminosity. There are several hypotheses about this behaviour (cf. Pilyugin [11]), but a satisfactory understanding is still missing. The uniform way to derive chemical abundances with PNe, which are present in galaxies of every morphological type, will help to understand the origins of this fundamental relation.

References

1. Ambartsumian, V. A., 1932, Poulkovo Obs. Circ 4, 8
2. Ferland, G. J., Korista, K. T., Verner, D. A., Ferguson, J. W., Kingdon, J. B., Verner, E. M., 1998, PASP, 110, 761
3. Gurzadyan, G. A., 1988, ApSS, 149, 343
4. Kingsburgh, R. L. & Barlow, M. J., 1994, MNRAS, 271, 257
5. Jacoby, G. H. & Lesser, M. P., 1981, AJ, 86, 185
6. Magrini, L., Corradi, R. L. M., Walton, N. A., et al., 2002, A&A, 386, 869
7. Magrini, L., Corradi, R. L. M., Greimel, R., et al., 2003, A&A, 407, 51
8. Magrini, L., Perinotto, M., Mampaso, A., Corradi, R. L. M., 2004, A&A in press, astro-ph/0406642
9. Mateo, M. L., 1998, ARA&A, 36, 435
10. Perinotto, M., Magrini, L., Mampaso, A., Corradi, R. L. M., 2004, to appear in the Proceedings of the ESO-workshop “Planetary Nebulae beyond the Milky Way”
11. Pilyugin, L. S., 2001, A&A, 374, 412
12. van den Bergh, S., 2000 in *The Galaxies of the Local Group*, Cambridge University Press
13. Vassiliadis, E. & Wood, P. R., 1994, ApJS, 92, 125