Mass and angular momentum loss during RLOF in Algols

W. Van Rensbergen, C. De Loore & D. Vanbeveren

Astrophysical Institute, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

Abstract. We present a set of evolutionary computations for binaries with a B-type primary at birth. Some liberal computations including loss of mass and angular momentum during binary evolution are added to an extensive grid of conservative calculations. Our computations are compared statistically to the observed distributions of orbital periods and mass ratios of Algols. Conservative Roche Lobe Over Flow (RLOF) reproduces the observed distribution of orbital periods decently but fails to explain the observed mass ratios in the range \in [0.4-1]. In order to obtain a better fit the binaries have to lose a significant amount of matter, without transferring too much angular momentum.

1. Introduction

Eggleton (2000) introduced the denomination "liberal" to make a distinction between binary evolution with mass and angular momentum loss and the conservative case where no mass and consequently no angular momentum leave the system. Refsdal et al. (1974) showed that the binary AS Eri is the result of liberal binary evolution; the amounts of mass and angular momentum lost by the system are however uncertain. Sarna (1993) showed that only 60 % of the mass lost by the loser of β Per was captured by the gainer and that 30 % of the initial angular momentum was lost during RLOF. Hence it is clear that the liberal evolutionary scenario is important for binary evolution calculations. However the amount of mass and angular momentum that has to be removed from interacting systems is far from obvious.

With the Brussels simultaneous evolution code (a description is given in "The Brightest Binaries", Vanbeveren et al. (1998)) we calculated a representative grid of conservative evolution of binaries with a B type primary at birth. Application of the criterion of Peters (2001) allows then to determine for each of the evolutionary sequences the beginning and ending of various Algol-stages. This criterion states that in the semi-detached system:

- The less massive star fills its Roche lobe (RL)
- The most massive star does not fill its RL and is still on the main sequence
- The less massive star is the coolest, the faintest and the largest

The grid allows to determine an expected distribution of orbital periods and mass ratios of Algols. These can be compared to the well established observed distributions (Budding et al. (2004)).

Considering the examples mentioned above (Refsdal et al. (1974), Sarna (1993)) it may be expected that the match between the observations and the

 $\mathbf{2}$

conservative results are far from satisfactory. Therefore we add liberal calculations to our conservative library. The detailed evolutionary tracks can be found in http://www.vub.ac.be/astrofys/

2. Details of the conservative calculation

Our grid contains binaries with sufficiently small initial periods to lead to Case A RLOF (RLOF A): i.e. during H core burning of the donor (initially primary star that becomes the less massive after Algol-ignition). The track across the HRD is traced for every system in our grid. A binary lives its era of "Algolism" (De Loore and Van Rensbergen (2004)) when it obeys the criterion of Peters (2001). Every binary shows its Algol A (Algol during H core burning) aspect for some time during RLOF A. The drastical change of the mass ratio and the orbital period during this process can be followed in detail for every system in our grid. It happens frequently that RLOF A is succeeded by Case B RLOF (RLOF B): i.e. during H shell burning of the donor. Systems that have sufficiently large initial orbital periods so that RLOF A does not occur will also show a short living Algol B (Algol during H shell burning) appearance. These systems have been considered in a previous paper (Van Rensbergen (2003)). Figure 1 shows a typical example of a case AB. The $(7+4.2)M_{\odot}$ binary with an initial period of 2.5 d starts as Algol A with q=1 and P=2.06 d. It ends this Algol A stage after some 20 million years with q=0.27 and P=6.87 d. From these initial values the system remains an Algol B for some 1.5 million years during H shell burning of the donor. The system eventually evolves into a long periodic (P>50 d) Algol B with $q \approx 0.1$. The grid contains some 240 more evolutionary tracks.



Figure 1. Conservative evolution of a $(7+4.2)M_{\odot}$ binary with an initial period of 2.5 d. The system remains Algol A for ≈ 20 million years before it is Algol B for ≈ 1.5 million years.

3. Conservative simululation

From our grid of conservative calculations we made a simulation of the distribution of orbital periods and mass ratios of Algols using a Monte Carlo algorithm. As starting conditions we selected:

- The primaries IMF of Salpeter (1955): $\zeta(M) \div M^{-2.35}$
- an initial distrubution of orbital periods from Popova et al. (1982): $\Pi(\mathbf{P}) \div \frac{1}{P}$

• an initial mass ratio distribution as derived by Van Rensbergen (2001) from the non-evolved systems of the catalogue of Spectroscopic Binaries shown on http://sb9.astro.ulb.ac.be/

The distribution of the initial mass ratio
$$q = \frac{M_2^0}{M_1^0}$$
 obeys relation (1):

 $\Psi(q) \div (1+q)^{-\alpha}$; $\alpha = 3.37$ for early B & 1.47 for late B primaries (1)

Figure 2 shows the observed distribution of orbital periods of Algols. Here we find more A than B-cases, because the Algol A-phase goes on for a fraction of the nuclear time scale, whereas the Algol B-phase lasts only for a fraction of the much shorter Kelvin-Helmholtz time scale. RLOF A produces Algols that follow the observed distribution rather well. Since the periods of the Algol B cases peak towards the long periods, a contribution of a few % of B cases to the Algol population will mimic the observed period distribution well.



Figure 2. Observed distribution of the orbital periods of Algols compared to conservative evolution. Cases B produce an excessive amount of long periods. Cases A follow the observed distribution better. There is no doubt that there are far more Algol A than Algol B cases.

Figure 3 shows the observed mass ratio distribution of Algols. The mass ratio q is now defined as the observers do: $q = \frac{M_{donor}}{M_{gainer}}$.

Van Rensbergen et al.

The mass ratios of the Algol B cases peak towards the smallest mass ratios, whereas RLOF A produces a majority of Algols in the q-bin [0.2-0.4]. As a consequence an admixture of cases A and B will never reproduce well the observed mass ratio distribution in the q-bins [0.4-1]. We may conclude that liberal binary evolution is needed to describe the mass ratio distribution of Algols.



Figure 3. Observed distribution of mass ratios of Algols compared to conservative evolution. Cases B produce more than 80% Algols with the smallest mass ratio $q \in [0-0.2]$. Cases A produce most of their Algols with $q \in [0.2-0.4]$. The fact that almost 80% Algols are observed with $q \in [0.4-1]$ excludes conservative evolution as the major channel through which Algols can be formed.

4. Confining the liberal model

Mass loss is defined by a parameter β describing the fraction of the mass lost by the loser that is accreted by the gainer:

$$\dot{M}_{gainer} = -\beta \ \dot{M}_{donor} \quad with \quad 0 \leq \beta \leq 1$$

$$\tag{2}$$

Conservative evolution is described with $\beta = 1$, whereas the liberal case uses values of $\beta < 1$ which are not known beforehand. Conservative evolution implies that no angular momentum can be lost by the system, whereas the amount of loss of angular momentum in the liberal case is also a free parameter if no physics restricts the assumptions. It is clear that realistic hydrodynamical calculations should learn us the appropriate choice of the amounts of mass and angular momentum which are lost by a binary at any moment of the RLOF process. In the mean time we have performed a number of liberal evolutionary calculations and compared the results with the observations.

Our calculations reveal that the time dependent parameter $\beta(t)$ should sufficiently often differ drastically from 1.

For a given value of β , the amount of angular momentum lost by the system is defined by the position of the site where matter is recoiled from the system. Soberman et al. (1997) have argued that matter can be trapped in a Keplerian ring after transit across the second Lagrangian point L_2 . The radius of the ring is η times the semi major axis of the orbit. This Keplerian ring passes outside L_2 which is located at $\eta \approx 1.25$. This yields a minimum value of $\eta \approx 1.25$. Hydrodynamical calculations of Lubow and Shu (1975) locate the ring at $\eta \approx$ 3. This yields a maximum value of $\eta \approx 3$. A Keplerian ring located at $\eta \approx 2.25$ takes away as much angular momentum as the co-rotating point L_2 would do. A value of $\eta \approx 2.25$ is thus a fair value in the interval [1.25-3] that can be used to calculate the change of the orbital period of a binary as a consequence of loss of mass and angular momentum through a ring which rotates with Keplerian velocity around the center of mass of the system (Soberman et al. (1997)).

Our calculations reveal that if the system loses angular momentum across points located near $\eta \approx 2.25$ the orbital periods shrink drastically. Most binaries become mergers before they show Algolism. The obtained distribution of orbital periods of Algols is completely shifted towards the shortest periods. Since this conclusion conflicts with observations, the mass that leaves the system carries rather the angular momentum of points located near $\eta \approx 0$ than near $\eta \approx 2.25$.



Figure 4. Liberal evolution with a lot of mass loss (β =0.5) and a little loss of angular momentum leaves the distribution of orbital periods of Algols almost unaltered. This figure shows the result for Algols issued from RLOF A and a 7 M_{\odot} primary at birth.

For all these (statistically sound but physically not well understood) reasons we performed a number of calculations with constant $\beta = 0.5$ and angular momentum lost at the edge of the gainer. Until now only the representative cases with a $7M_{\odot}$ primary at birth and initial periods leading to RLOF A have been performed. Figure 4 shows that the assumption of mass loss through a point near $\eta = 0$ does not alter the conservative distribution of the orbital periods of Algols. A conclusion that meets the observations since the conservative orbital period distribution matched the observations fairly well. Figure 5 shows that our liberal assumption ($\beta=0.5$, $\eta \approx 0$) deviates radically from the conservative case. The q-bins [0.4-1] which were not populated in the conservative scenario are now populated properly as required by the observations (see figure 3).



Figure 5. Liberal evolution with a lot of mass loss ($\beta = 0.5$) and a little loss of angular momentum changes the mass ratio distribution of Algols drastically. The observed mass ratios in the q-interval [0.4-1] which were not produced by conservative evolution are created by this liberal model. This plot shows the result for Algols issued from RLOF A and a 7M_{\odot} primary at birth.

References

- Budding, E., Erdem, A., Cicek, C., Bulut, I., Soydugan, E., Soydugan, V., Bakis, V., Demircan, O. A&A, 217, 263-268, 2004.
- De Loore, C., Van Rensbergen, W. in Zdenek Kopal's Binary Star Legacy ed. Drechsel, H.. Space Sci.Rev., accepted.
- Eggleton, P. New Astron. Rev., 44, 111-117, 2000.
- Lubow, S., Shu, F. ApJ, 198, 383-405, 1975.
- Peters, G. Ap&SS, 264, 79-94, 2001.
- Popova, E., Tutukov, A., Yungelson, L Ap&SS, 88, 55-80, 1982.
- Refsdal,S., Roth,M., Weigert, A. A&A, 36, 113-122, 1974.
- Salpeter, E. ApJ, 121, 161-167, 1955.
- Sarna, M. MNRAS, 262, 534-542, 1993.
- Soberman, G., Phinney, E., Van den Heuvel, E. A&A, 327, 620-635, 1997.
- Vanbeveren, D., Van Rensbergen, W., De Loore, C. The Brightest Binaries. Ap&SS Library, 232, 1998.
- Van Rensbergen, W. In "The Influence of Binaries on Stellar population Studies" (ed. Vanbeveren, D.), Ap&SS Library, 264, 21-35, 2001.
- Van Rensbergen, W. In "Stellar Astrophysics" (eds. Cheng,K., Leung,K., Li,T.), Ap&SS Library, 298, 117-125, 2003.