

Extreme Properties Of GRB 061007: A Highly Energetic Or A Highly Collimated Burst?

P. Schady¹, M.De Pasquale¹, M.J. Page¹, L.Vetere³, S.B. Pandey¹, X.Y. Wang^{3,5}, J. Cummings², B. Zhang⁴, S. Zane¹, A. Breeveld¹, D.N. Burrows³, N. Gehrels², C. Gronwall³, S. Hunsberger³, C. Markwardt², K.O. Mason^{1,6}, P. Mészáros^{3,7}, S.R. Oates¹, C. Pagani³, T.S. Poole¹, P.W.A. Roming³, P.J. Smith¹ and D.E. Vanden Berk³

¹ *The UCL Mullard Space Science Laboratory, Holmbury St Mary, Dorking, Surrey, RH5 6NT, UK.*

² *NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA.*

³ *Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802, USA.*

⁴ *Department of Physics, University of Nevada, Las Vegas, NV 89154, USA.*

⁵ *Department of Astronomy, Nanjing University, Nanjing 210093, China.*

⁶ *The Particle Physics and Astronomy Research Council, Polaris House, North Star Avenue, Swindon, Wiltshire, SN2 1SZ, UK.*

⁷ *Department of Physics, Pennsylvania State University, University Park, PA 16802, USA.*

Received:

ABSTRACT

GRB 061007 is the brightest gamma-ray burst (GRB) to be detected by *Swift* and is accompanied by an exceptionally luminous afterglow that had a *V*-band magnitude < 11.1 at 80 s after the prompt emission. From the start of the *Swift* observations the afterglow decayed as a power law with a slope of $\alpha_X = 1.66 \pm 0.01$ in the X-ray and $\alpha_{opt} = 1.64 \pm 0.01$ in the UV/optical, up to the point that it was no longer detected above background in the optical or X-ray bands. The brightness of this GRB and the similarity in the decay rate of the X-ray, optical and γ -ray emission from 100 s after the trigger distinguish this burst from others and present a challenge to the fireball model. The lack of a cooling or jet break in the afterglow up to $\sim 10^5$ s constrains any model that can produce the large luminosity observed in GRB 061007, which we found to require either an excessively large kinetic energy or highly collimated outflow. Analysis of the multi-wavelength spectral and high-resolution temporal data taken with *Swift* suggest an early time jet-break to be a more plausible scenario. This must have occurred within 80 s of the prompt emission, which places an upper limit on the jet opening angle of $\theta_j = 0.8^\circ$. Such a highly collimated outflow resolves the energy budget problem presented in a spherical emission model, reducing the isotropic equivalent energy of this burst to $E_\gamma^{corr} = 10^{50}$ ergs; consistent with other GRBs.

Key words: gamma-rays: bursts - gamma-ray: observations - ISM: jets and outflows

1 INTRODUCTION

Gamma-ray bursts (GRBs) are by far the most energetic and luminous astrophysical phenomena in the Universe, releasing isotropic equivalent energies of the order of $10^{48} - 10^{54}$ ergs in γ -rays alone on timescales of $10^{-3} - 10^3$ s. The favoured mechanism to produce such a large amount of energy on such a short timescale is the creation of a black hole, either from a merger of two compact objects or the gravitational core collapse of a massive star. There appear to be two distinct populations of GRBs, distinguished by the duration of their prompt emission and

their spectral properties. The most robust model for long, soft GRBs, which typically have prompt emissions lasting > 5 s, is the collapsar model (Woosley 1993), in which a GRB results from the energy released during the death of a massive star. Evidence supporting this model includes the large number of long GRBs located in regions of high star-formation rate (e.g. Tanvir et al. 2004; Fruchter et al. 2006), and the association of several nearby GRBs with supernovae, e.g. GRB 980425/SN1998bw (Kulkarni et al. 1998), GRB 030329/SN2003dh (Hjorth et al. 2003) and GRB 060218/SN2006aj (Campana et al. 2006).

There is evidence to suggest that collimated outflows reduce the amount of energy emitted by a GRB from isotropic emission by ~ 2 orders of magnitude, where the beam-corrected energy is clustered at $10^{50} - 10^{51}$ ergs (Frail et al. 2001; Bloom, Frail & Kulkarni 2003). Bipolar jets are a natural consequence of accretion onto a black hole and are observed in many high-energy astrophysical phenomena (e.g. AGN, pulsars, micro-quasars), where the ejecta take the easiest exit path, along the rotational axis.

A signature of highly relativistic jets is an achromatic break in the broadband afterglow light curves, caused by the expansion of the radiation beaming angle as the bulk Lorentz factor, Γ , decreases. Once the Lorentz factor has decreased such that $\theta_j = 1/\Gamma$, where θ_j is the jet opening angle, the rate of decay increases, resulting in a break in the light curve.

Swift is a rapid-response mission with multi-wavelength capabilities designed specifically to study the early time evolution of GRBs (Gehrels et al. 2004). The spacecraft is equipped with three telescopes; the Burst Alert Telescope (BAT; Barthelmy et al. 2005), which detects γ -rays from ~ 10 up to several hundred keV, the X-Ray Telescope (XRT; Burrows et al. 2005) which covers the 0.3–10 keV energy range, and the Ultra-Violet and Optical Telescope (UVOT; Roming et al. 2005), which can observe in six colours and a broadband white band filter, covering the wavelength range from 1600 Å to 6000 Å.

Swift's slewing capabilities provide GRB afterglow light curves from ~ 60 s after the trigger, and well-sampled X-ray and optical light curves have revealed far fewer achromatic breaks than had been expected prior to launch (Ghirlanda, Ghisellini & Lazzati 2004; O'Brian et al. 2006). The jet opening angle, θ_j , increases with t_j as $\theta_j \propto t_j^{3/8}$, and thus the lack of observed jet breaks may be the result of large average jet opening angles.

GRB 061007 is an example of a burst that showed no evidence of a break in the X-ray or UV/optical afterglow from 80 s after the prompt emission, until it was no longer detected above background, at $\sim 10^6$ s. Prompt observations by ground based observatories measured the redshift of this burst to be $z=1.26$ (Osip, Chen & Prochaska 2006; Jakobsson et al. 2006). At this redshift, GRB 061007 had one of the highest isotropic equivalent energies ever seen, releasing $\sim 10^{54}$ ergs in γ -radiation alone, with an optical afterglow comparable to that of GRB 990123 (Akerlof et al. 1999), with $V < 11.1$ mag at $T + 71.7$ s.

In this paper we describe in detail the γ -ray, X-ray and optical observations taken by *Swift*. From the high resolution early time data we constrain the type of environment in which the burst occurred and determine the conditions that distinguish this burst from others. In section 2 we describe the data reduction and analysis techniques used followed by a discussion on the physical parameters and emission mechanisms involved that could reproduce our observations in section 3.1–3.2.2. We also analysis the properties of the circumburst medium of GRB 061007 in section 3.4 using the afterglow spectral energy distribution (SED), and a summary of our results is given in section 4. Throughout the paper all errors quoted are 1σ unless otherwise specified and the temporal decay rates, α , and spectral indices, β , are defined such that $F_t \propto t^{-\alpha}$ and $F_\nu \propto \nu^{-\beta}$, respectively. Luminosities and the energy release are calculated assuming




Figure 1. Prompt emission light curve of GRB 061007 shown in the four BAT energy bands.

a standard cosmology model with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\lambda = 0.7$.

2 OBSERVATIONS AND ANALYSIS

2.1 BAT

The BAT on board *Swift* triggered on GRB 061007 on 7th October, 2006 at 10:08:08 UT, resulting in an immediate slew to point the narrow field instruments in the direction of the burst. From here on we shall refer to the time of the trigger as T . The refined BAT position for GRB 061007 is RA = $03^h 05^m 11.8^s$, Dec = $-50^\circ 29' 47.4''$ (J2000) with an uncertainty (systematic and statistical) of $0.9'$ at 90% containment (Markwardt et al. 2006).

The time-interval over which 90% of the 15–150 keV γ -radiation is emitted is $T_{90} = 75 \pm 5$ s. The BAT light curve consists of three large and distinct peaks (see Fig. 1) with weak emission detected out to ~ 150 s after the trigger. The time lag between the peak emission in the 50–100 keV and the 15–25 keV energy bands was 16_{-6}^{+8} ms, and between the 100–350 keV and 25–50 keV bands the lag was 21_{-5}^{+7} ms (private communication with J. Norris).

2.2 XRT

The first XRT observation was a centroiding image taken 80.4 s after the BAT trigger, and a bright uncatalogued source was found at RA = $03^h05^m19.5^s$, Dec = $-50^\circ30'01.9''$, with a 90% containment radius of $3.5''$ (Vetere et al. 2006).

At 86.6 s after the BAT trigger the XRT began taking data in Window Timing mode, in which data are stored with a 1.7 ms time resolution and 1-dimensional imaging. At $T + 2000$ s the count rate fell to below ~ 1 count s^{-1} and the automated sequence switched to Photon Counting (PC) mode, in which the time resolution is 2.5 ms and full imaging and spectroscopic resolution are available. The X-ray afterglow was detected for 10^6 s after the BAT trigger, before it fell below the detection threshold. During the first 2200 s of observations the count rate of the burst was high enough to cause pile-up in both WT and PC mode data. To account for this effect, the WT data were extracted in a rectangular 40×20 -pixel region with a 9×20 -pixel region excluded from its centre. The size of the exclusion region was determined following the procedure illustrated in Romano et al. (2006a). The standard correction for pile-up was applied to the PC data, as described by Vaughan et al. (2006). Due to a large amount of pile-up in the first 30 s of WT observations, these data were not used in our spectral analysis.

The 0.3–10 keV light curve can be represented well by a power law with decay index $\alpha = 1.66 \pm 0.01$ from $T + 80$ s until at least $T + 10^6$ s, when it reached the detectability threshold with XRT (see Fig. 2). We also determined the earliest times that a cooling break or jet break could fit the data with a goodness of fit within 3σ of the best-fit power law model. To do this we applied a broken power law fit modelled on the cooling break, for which $\alpha_2 = \alpha_1 + 0.25$, and on a jet break, where $\alpha_2 = \alpha_1 + 0.75$. In the former model $t_{\nu_c} > 10^5$ s before the change in χ^2 from a power law fit was more than 3σ level, and in the latter model this corresponded to at $t_j > 2.65 \times 10^5$ s.

A power law fit to the WT spectrum, where the Galactic column density is fixed at $N_H = 2.13 \times 10^{20}$ cm^{-2} (Dickey & Lockman 1990), gives a best-fit host galaxy column density at a redshift of $z = 1.26$ of $(5.1 \pm 0.3) \times 10^{21}$ cm^{-2} for a spectral index $\beta_X = 0.99 \pm 0.02$.

2.3 UVOT

The UVOT began observing GRB 061007 71.7 s after the BAT trigger, at which point it took a 9 s settling exposure in the V -filter. This was followed by a 100 s exposure in the white light filter ($\lambda \sim 1600 - 6500$ Å) and a 400 s exposure in the V -band. After this the automated sequence rotated twice through the UVOT filters taking a series of short exposures ($V, UVM2, UVW1, U, B, \text{white}, UVW2$) (10 s for B and white filters and 20 s for the rest). The central wavelengths to the optical and UV filters are 5460 Å, 4370 Å, 3460 Å, 2600 Å, 2260 Å and 2030 Å for the $V, B, U, UVM1, UVM2$ and $UVW2$, respectively. A further 400 s V -band exposure and 100 s white filter were taken, followed by another series of rapid rotations around the filter wheel up to 2260 s after the burst. All observations up to this time were taken in event mode, which has a time resolution of 11 ms, after which time image mode exposures were taken. The re-

finned position of the afterglow is RA = $03^h05^m19.6^s$, Dec = $-50^\circ30'02.4''$ to a certainty of $0.5''$ (J2000). The Galactic extinction along this line of sight is $E(B - V) = 0.021$ (Schlegel, Finkbeiner & Davis 1998).

For all short exposures (20 s) and exposures taken later than $T + 2260$ s photometric measurements were obtained from the UVOT data using a circular source extraction region with a $6''$ radius for the V, B and U optical filters and a $12''$ radius for the three UVOT ultra-violet filters to remain compatible with the current effective area and zeropoint calibrations¹. The background was measured in a source-free region near to the target using an extraction radius of $20''$.

The first V -band settling exposure and first white-band exposure were affected by coincidence loss to a degree that is outside the photometrically calibrated range of UVOT; for a V -band settling exposure, complete saturation within a standard $6''$ aperture implies that this afterglow is brighter than $V = 11.1$ for the whole of the 9 s exposure. During the first white band exposure the source is so bright that relative photometry can be obtained using a very large aperture, which includes the wings of the point spread function, where coincidence loss is not a problem. The exposure was divided into 5 s bins, and source counts were obtained from an extraction region of radius $\sim 26''$. To calibrate this photometry, observations of the white dwarf standards GD50 and HZ2 were taken with UVOT and the white filter an October 18th 2006. From these observations, we determined the relationship between observed count rate within the $26''$ aperture and corrected count rate within a standard $6''$ UVOT source aperture, which we assume to be linear. To apply this calibration to GRB 061007 we apply constant offset to this relation to account for the different background count rate between the GRB 061007 observations and the white dwarf standard count rate observations. This offset was chosen to match the mean white-band count rate in the saturated exposure to the white filter light curve extrapolated from later times; we caution that calibration of this first white exposure is crude and thus the apparent steepening of the light curve towards the end of the exposure may not be real. However, the overall behaviour of a steadily decreasing count rate over the course of the exposure is not in doubt. For the first 400 s V -band exposure a higher resolution light curve was created by splitting it into 4 s bins, and the second into 20 s bins, consistent with the rest of the exposures during that time.

We created a single UV/optical light curve from all the UVOT filters in order to get the best measurement of the optical temporal decay. To do this the light curve in each filter was individually fitted to find the corresponding normalisation, and this was then used to renormalised each light curve to the V -band. A fit to the combined UV and optical light curve, from $T + 85$ s up to $T + 2.5 \times 10^4$ s, is best fit with a power law decay with temporal index $\alpha_{opt} = 1.64 \pm 0.01$. The combined light curve shows no apparent colour evolution, as can be seen in Fig. 2.

The combined UVOT light curve is shown next to the 0.3–10 keV and 15–350 keV light curve in Fig. 2. The magnitude along the left axis corresponds to the V -band, and the count rate along the right axis applies to the X-ray and γ -ray light curves.

¹ <http://heasarc.gsfc.nasa.gov/docs/heasarc/caldb/swift/docs/uvot/>




Figure 2. GRB afterglow light curve shown in all six UVOT lenticular filters, in the X-ray (0.3–10 keV) and in the γ -ray (15–350 keV). All light curves have been shifted vertically with respect to the V band filter for clarity. The B, U, UVW1, UVM2, UVW2 and white filter have all been normalised to the V band light curve. The left hand axis corresponds to the V-band light curve, and the count rate axis on the right applies to the X-ray and γ -ray light curves. The best fit decay indices to the combined UVOT light curve, the X-ray light curve and the BAT light curve are $\alpha_{UVOT} = 1.64 \pm 0.01$, $\alpha_{XRT} = 1.66 \pm 0.01$ and $\alpha_{BAT} = 1.54 \pm 0.12$, respectively.

2.4 Observations By Other Facilities

The prompt emission from GRB 061007 also triggered Konus-Wind and Suzaku, and the afterglow was monitored by ground based telescopes, which carried out both spectroscopic and photometric observations. GRB 061007 was measured to be at $z=1.26$ (Osip et al. 2006; Jakobsson et al. 2006), for which the rest-frame isotropic energy release and maximum luminosity in the 20 keV–10 MeV is $E_{iso} \sim 1.0 \times 10^{54}$ erg, and $L_{iso} \sim 1.8 \times 10^{53}$ erg s^{-1} (Golenetskii et al. 2006). The optical afterglow was also detected by ROTSE at 26.4 s after the BAT trigger at a magnitude of 13.6 mag in an unfiltered exposure (Rykoff & Rujopakarn 2006), and the Faulkes Telescope South measured an R-band magnitude of $R = 10.15 \pm 0.34$ 142 s after the trigger (Bersier et al. 2006), consistent with the UVOT observations and indicative of a significant brightening during the first two minutes. The *Swift* observations from $T+80$ s show that by this time the afterglow had entered a constant power law decay (see Fig. 2). Radio observations were also performed 1–1.24 days after the prompt emission, which resulted in no detection of a radio source with a flux measurement of $1 \pm 36 \mu\text{Jy}$ at 19 GHz (van der Horst & Rol 2006).

2.5 Multi-Wavelength Light Curve Analysis

At $\sim T + 100$ s, after the last flare in the BAT light curve (Fig. 1), the BAT light curve appears to decay smoothly and monotonically. A power law fit to the BAT light curve from $\sim T + 100$ s provides an acceptable fit ($\chi^2 = 32$ for 29 dof) with a best-fit decay index of $\alpha_{BAT} = -1.54 \pm 0.12$, which is consistent with the decay observed in both the UV/optical and X-ray light curves. Furthermore, a power law spectral fit to this last part of the BAT light curve, beyond $\sim T + 100$ s, gives a best fit energy spectral index $\beta_{BAT}(> T + 100 \text{ s}) = -0.80 \pm 0.09$, in fairly good agreement with the spectral indices determined for the X-ray and optical afterglows. This supports a scenario in which the γ -emission at this epoch is generated by the same radiation mechanism as the X-ray and UV/optical afterglow.

2.6 The Spectral Energy Distribution

Using the full range of data available with *Swift* a SED was produced using optical, X-ray and γ -ray data, which were normalised to an instantaneous epoch at 600 s after the BAT trigger, at which time there are data available from all three instruments, and good colour information for UVOT. For each of the UVOT lenticular filters the tool UVOT2PHA (v1.1) was used to produce spectral files compatible with XSPEC (v12.2.1), and the count rate in each band was normalised




Figure 3. The combined γ -ray, X-ray and UV/optical spectral energy distribution of GRB 061007 at $T + 600$ s (see text) with best-fit models for each corresponding dust extinction curve shown; SMC (dashed), LMC (solid) and Galactic (dotted). Open circles are data points at wavelength $\lambda < 1215$ Å in the rest frame, and therefore not used in the spectral fitting. The rest-frame positions of the Galactic absorption feature at 2175 Å the beginning of the Lyman- α forest at 1215 Å and the Lyman- α break at 912.5 Å are indicated.

to the estimated value at $T + 600$ s determined from the best-fit models to the light curves. The XRT spectrum was produced from WT mode data in the 0.5–10 keV energy range, omitting the first 30 s of data, which suffered from pile-up. This was scaled to the corresponding count rate at $T + 600$ s. For the BAT spectrum, data were taken from the time interval $T + 100$ s – $T + 500$ s, during which the BAT light curve decayed at a rate consistent with X-ray and UV/optical afterglows and had a similar spectrum. Beyond 500 s the γ -ray emission became indistinguishable from the background. This spectrum was normalised to $T + 600$ s in the same way as the XRT spectrum. The SED is shown in Fig. 3.

The SED was fit with a power law emission spectrum and two dust and gas components to model the Galactic and host galaxy photoelectric absorption and dust extinction. The column density and reddening in the first absorption system were fixed at the Galactic values (see sections 2.2 and 2.3). The second photoelectric absorption system was set to the redshift of the GRB, and the neutral hydrogen column density in the host galaxy was determined assuming solar abundances. The dependence of dust extinction on wavelength in the GRB host galaxy was modelled on three

extinction laws, taken from observations of the Milky Way (MW), the Large Magellanic Cloud (LMC) and in the Small Magellanic Cloud (SMC). The greatest differences observed in these extinction laws are the amount of far UV (FUV) extinction, which is greatest in the SMC and least in the MW, and the strength of the 2175 Å absorption feature, which is most prominent in the MW and negligible in the SMC. To model these extinction laws the parameterisations from Pei (1992) were used, where $R_V = A_V/E(B - V) = 3.08, 2.93,$ and 3.16 for the Galactic, SMC and LMC extinction laws respectively.

At a redshift of $z = 1.26$ the beginning of the Lyman- α forest is redshifted to an observer-frame wavelength of ~ 2580 Å, which falls squarely on the UVW1 filter, the reddest of the UV filters. The unknown equivalent width of Lyman- α absorption from the host galaxy of the GRB and the variation of the Ly- α forest along different lines of sight makes it difficult to model this additional source of absorption in the SED. We, therefore, exclude the three data points affected by Lyman- α from our spectral analysis. The results from our spectral analysis are summarised in Table 1. The fitted value of the spectral index depends on the amount of optical/UV extinction, which in turn is affected

Table 1. Power law spectral fit results to the spectral energy distribution of GRB 061007

Spectral Model	Host galaxy N_H^1 (10^{21} cm $^{-2}$)	Rest-Frame visual extinction, A_V	$\beta_{o,x}$	χ^2 (dof)	Null-hypothesis probability
SMC	$3.66^{+0.29}_{-0.28}$	0.39 ± 0.01	0.90 ± 0.005	398 (308)	4×10^{-4}
LMC	$5.30^{+0.33}_{-0.32}$	0.66 ± 0.02	0.98 ± 0.007	347 (308)	0.063
MW	$7.62^{+0.81}_{-0.77}$	1.13 ± 0.10	$1.13^{+0.04}_{-0.03}$	1042 (308)	0.000

¹ Equivalent hydrogen column density assuming solar abundances at the GRB host galaxy.

by the dust extinction model used. Similarly, the amount of soft X-ray absorption fitted to the model will affect the best-fit spectral index. Therefore, although the X-ray absorption and optical/UV extinction are independent components in the fit, they are coupled because they both depend on the spectral index.

A fit to the optical, X-ray and γ -ray data, with an LMC extinction law for the GRB host galaxy, gives an acceptable fit, with $\chi^2 = 347$ for 308 degrees of freedom (dof), implying a null-hypothesis probability of $p = 0.063$. Models with an SMC and MW extinction law at the host galaxy were rejected with 99.96% confidence and 100% confidence, respectively.

3 DISCUSSION

GRB 061007 is one of the brightest bursts to be observed by *Swift*, with an isotropic energy released in γ -rays that is second only to GRB 050904 (Tagliaferri et al. 2005) and an optical flux comparable to that observed in GRB 990123 (Akerlof et al. 1999). The similarity of the afterglow of GRB 061007 in the flux decay of γ -ray, X-ray and UV/optical bands, and the lack of features in X-ray and optical light curves make this GRB highly unusual. Optical afterglows typically decay as power laws, although usually with decay indices ~ 1.1 (Zeh, Klose & Kann 2006). Indeed, prior to the launch of *Swift* GRB 061007 could have been categorised a dark burst. In most GRB X-ray afterglows, the light curve is characterised by several discrete segments, where the canonical shape is made up of three, and sometimes four, power law segments with different decay indices (Nousek et al. 2006; Zhang et al. 2006a).

The lack of a break in the light curves provides a constraint on the frequency of the characteristic synchrotron frequency, ν_m and the peak flux density, $F_{\nu, \max}$. From the response to the white filter it can be seen that around half the counts correspond to the *V* and *B* filter, and the lack of a break in the first white filter exposure, therefore, indicates that at $T + 80$ s ν_m was no bluer than the *B*-band filter. This corresponds to an upper limit on the characteristic synchrotron frequency $\nu_m < 7.5 \times 10^{14}$ Hz. From the UV/optical data we also determine that at $T + 80$ s the flux density in the *V*-band, corresponding to $\nu = 5.5 \times 10^{14}$ Hz, is $F_\nu = 730$ mJy, after correcting for absorption at the host galaxy.

With these constraints in place we investigate two possibilities that could explain the temporal and spectral properties of GRB 061007, where the fundamental difference is in the effect of collimation on our observations. In the first model we assume that a jet break does not occur before

$T + 2.65 \times 10^5$ s, based on the constraints we obtained in section 2.2, and the observed emission is, therefore, indistinguishable from isotropic emission. We then look into a model in which the jet break occurs before our first X-ray and UV/optical observations at $T + 80$ s such that assumption of isotropic emission no longer applies.

3.1 Spherical Expansion Model

The relation between α and β for synchrotron emission at different intervals in the synchrotron spectrum are defined by a set of closure relations, both for a constant density medium (Sari, Piran & Narayan 1998) and a wind density profile (Chevalier & Li 1999). Using these we find our observations to be in closest agreement with a constant density circumburst medium, where the UV/optical and X-ray bands are in the same slow-cooling regime ($\nu_m < \nu < \nu_c$). In this case, the temporal and spectral flux decay indices are related as $\alpha = 3\beta/2$. For a value of $\beta_X = 0.99 \pm 0.02$, we expect $\alpha = 1.49 \pm 0.03$, which is shallower than the observed decay rate of $\alpha = 1.65 \pm 0.01$. However, this deviation from the relation is small, and could be accommodated if the density profile is slightly deviated from an absolute constant medium. In this regime $\beta = (p - 1)/2$, where p is the electron energy index. Thus, for $\beta_X = 0.99 \pm 0.02$ the electron energy index is in the range $2.94 < p < 3.02$. We therefore take $p = 3.0$. These relations hold as long as the characteristic synchrotron frequency, ν_m , lies below the optical band, i.e. $\nu_m < 7.5 \times 10^{14}$ Hz, at $T + 80$ s, as discussed previously. There is also the additional condition that the cooling frequency, ν_c , must lie above the peak X-ray frequency up to at least $T + 10^5$ s, before which a cooling break would be detected in our X-ray observations (see section 2.2). This corresponds to the requirement that $\nu_c > 6 \times 10^{17}$ Hz at $T + 10^5$ s.

For a constant density medium in which synchrotron radiation is the dominating emission mechanism, the characteristic synchrotron frequency, the cooling frequency and the peak density flux are given by the following equations from Zhang et al. (2006b)

$$\nu_m = 3.3 \times 10^{15} \text{ Hz} \left(\frac{p-2}{p-1} \right)^2 (1+z)^{1/2} \epsilon_B^{1/2} \epsilon_e^2 E_{k,52}^{1/2} t_d^{-3/2} \quad (1)$$

$$\nu_c = 6.3 \times 10^{12} \text{ Hz} (1+z)^{-1/2} (1+Y)^{-2} \epsilon_B^{-3/2} E_{k,52}^{-1/2} n^{-1} t_d^{-1/2} \quad (2)$$

$$F_{\nu, \max} = 16 \text{ mJy} (1+z) D_{28}^{-2} \epsilon_B^{1/2} E_{k,52} \Omega^{1/2}, \quad (3)$$

where ϵ_B is the fraction of energy in the circumburst magnetic field, ϵ_e is the fraction of energy in electrons, $E_{k,52}$ is the kinetic energy in the GRB in units of 10^{52} , t_d is the time since the prompt emission in units of days, n is the density

of the surrounding circumburst medium, Y is the Compton parameter, and D_{28} is the luminosity distance in units of 10^{28} cm.

For a spectral index $\beta = 1$, νF_ν is constant for $\nu_m < \nu < \nu_c$. At $T + 80$ s $F_\nu = 730$ mJy at $\nu = 5.5 \times 10^{14}$ Hz, and therefore $\nu_m F_{\nu, \max} = 730 \times (5.5 \times 10^{14})$. By using this together with Eq. 1 and 3 we obtain:

$$n = \frac{3.4 \times 10^{-6}}{\epsilon_B^2 \epsilon_e^4 E_{k,52}^3} \quad (4)$$

In Eq. 2 we apply the condition that at 10^5 s after the prompt emission ν_c must still be greater than 6×10^{17} Hz, and substituting in Eq. 4 gives

$$(1 + Y)^{-2} \epsilon_B^{1/2} \epsilon_e^4 E_{k,52}^{5/2} > 0.52 \quad (5)$$

In the case of the characteristic synchrotron frequency we know that $\nu_m < 7.5 \times 10^{14}$ Hz, and this reduces Eq. 1 to

$$\epsilon_B^{1/2} \epsilon_e^2 E_{k,52}^{1/2} < 1.7 \times 10^{-5}, \quad (6)$$

This, and the inequality given in Eq. 5 provide an upper limit to the value of ϵ_B in terms of $E_{k,52}^3$, such that

$$\epsilon_B < 3.1 \times 10^{-19} (1 + Y)^{-4} E_{k,52}^3 \quad (7)$$

The above equation bounds us to extreme values of $E_{k,52}$ and ϵ_B . For $E_{k,52} = 10^3$, $\epsilon_B < 3.1 \times 10^{-10} (1 + Y)^{-4}$, which is an unrealistically low value of ϵ_B . Even a value as small as $\epsilon_B = 10^{-6}$ requires a huge amount of kinetic energy to meet the condition that $E_{k,52} > 1.5 \times 10^4 (1 + Y)^{4/3}$. The amount of kinetic energy assumed is an isotropic equivalent value which would, thus, be smaller if it were corrected for collimation. A 3σ lower limit of $t_j = 2.65 \times 10^5$ s can be imposed as the earliest time that a jet break could have occurred. t_j is the time of the jet break measured from the onset of the afterglow, which we take to be equal to T . A kinetic energy $E_{k,52} = 1.5 \times 10^4 (1 + Y)^{4/3}$ and $\epsilon_B = 10^{-6}$ would require $\epsilon_e = 0.012$ and $n = 49 \text{ cm}^{-3}$ for the conditions given above to be satisfied. From this we estimate the jet opening angle using

$$\theta_j = 0.2 \left(\frac{t_{j,d}}{1+z} \right)^{3/8} \left(\frac{n}{E_{k,52}} \right)^{1/8}, \quad (8)$$

(Panaitescu & Kumar 2000), where $t_{j,d}$ is in units of days. For the physical parameters assumed, a jet opening angle for GRB 061007 would need to be $\theta_j > 6.3^\circ$ to produce a jet break later than $T + 2.65 \times 10^5$ s. This would provide a beam-corrected kinetic energy of $E_k^{corr} > 9 \times 10^{53}$ ergs, which is three orders of magnitude greater than usual bursts (Freedman & Waxman 2001). The energetics required by this model are unreasonably large and, therefore, an unlikely solution to explain the properties of GRB 061007.

3.2 Highly Collimated Models

The large amount of energy required to power a model with isotropic emission leads us to investigate a scenario in which the emission from the GRB is highly collimated, where we still assume the circumburst environment to be of constant density. A very narrow jet opening angle could reduce the energy required to explain our observations by several orders of magnitude, where t_j must be prior to our first afterglow observations in order to account for the lack of a break in

the X-ray and UV/optical light curves at later times. If we assume the afterglow begins at the time of trigger, this corresponds to a time $t_j < 80$ s. Within the premise that the outflow is highly collimated we consider two cases in which the cooling frequency is either above or below the observing frequencies. In these two regimes p and β are related by $\beta = (p - 1)/2$ and $\beta = p/2$, respectively, corresponding to $p \sim 3.0$ or $p \sim 2.0$. After the jet break the flux decays as $F_t \propto t^{-p}$ (Sari, Piran & Halpern 1999), where p is either 2 or 3, which is steeper than the observed decay rate of $\alpha \sim 1.65$. However, this does not account for continual energy injection, either from the central engine itself, or late time refreshed shocks that slow down the decay of the afterglow. The shallow decay phase observed in many XRT light curves is interpreted to be the result of energy injection, and although this phase typically lasts for 10^4 s, there are also examples of GRBs with evidence of energy injection for prolonged periods of time, and with very smooth decay slopes (e.g. Romano et al. 2006; Huang, Cheng & Gao 2006). Taking this into account, an explanation whereby continual energy injection maintains the afterglow and slows down its decay over a long period of time is not unreasonable.

3.2.1 $\nu_m < \nu < \nu_c$

For a laterally expanding jet with constant density circumburst medium and $\nu_m < \nu < \nu_c$, the relation between the temporal decay of the GRB afterglow, α , and the spectral index, β , post jet-break is given by

$$\alpha = (1 + 2\beta) - \frac{2}{3}(1 - q)(\beta + 2), \quad (9)$$

(Panaitescu et al. 2006) where q is the energy injection parameter such that $L = L_0(t/t_0)^{-q}$, (Zhang & Mészáros 2001). For our observed temporal and spectral indices, $q = 0.3$ and $p = 3$, as previously. Prior to the jet break Eq. 1-3 still apply, but in order take into account the effect of energy injection, the kinetic energy requires an additional factor of $(t/t_j)^{(1-q)}$. The time of jet break and the jet opening angle are related as $\theta_j \propto t_j^{3/8}$. To keep θ_j as large as our observations will allow we take $t_j = 80$ s. At $T + 80$ s the additional factor of $(t/t_j)^{(1-q)}$ is unity and there is no change in Eq. 1-3. Thus at $T + 80$ s, Eq. 4 and Eq. 6 stand. For Eq. 2 the constraint that $\nu_c > 6 \times 10^{17}$ at $T + 10^5$ s means that the evolution of ν_c post jet-break has to be taken into account. If ν_{c_j} is the frequency of ν_c at $T + 80$ s,

$$\nu_{c_j} = 4.05 \times 10^{19} \epsilon_B^{1/2} \epsilon_e^4 E_{k,52}^{5/2} \quad (10)$$

However, we are concerned with where ν_c lies at $T + 10^5$ s. After a jet break $\nu_c \propto \gamma^{-4} t^{-2}$, where $\gamma \propto t^{-(2+q)/6}$ is the Lorentz factor of the electrons in the circumburst environment (Panaitescu et al. 2006). The cooling frequency after $T + 80$ s is given by

$$\nu_c = \nu_{c_j} \left(\frac{\gamma}{\gamma_j} \right)^{-4} \left(\frac{t}{t_j} \right)^{-2} = \nu_{c_j} \left(\frac{t}{t_j} \right)^{\frac{2}{3}(q-1)}, \quad (11)$$

where γ_j is the electron Lorentz factor at $T + 80$ s. Using the condition that $\nu_c > 6 \times 10^{17}$ Hz at $T + 10^5$ s, for $q = 0.3$ Eq. 11 reduces to

$$(1 + Y)^{-2} \epsilon_B^{1/2} \epsilon_e^4 E_{k,52}^{5/2} > 0.41 \quad (12)$$

In the same way that we determined the inequality given in Eq. 7, we combine the above equation with Eq. 6 and determine that

$$\epsilon_B < 5.0 \times 10^{-19} (1+Y)^{-4} E_{k,52}^3 \quad (13)$$

For $E_{k,52} = 5 \times 10^4$, $\epsilon_B < 6.2 \times 10^{-5} (1+Y)^{-4}$, and if we take this upper limit as the value of ϵ_B , $\epsilon_e = 3.1 \times 10^{-3} (1+Y)^{1/2}$. Although the assumed kinetic energy is large, this corresponds to the isotropic equivalent value and the beam-corrected kinetic energy is a few orders of magnitude smaller than this. For $p = 3$, the inverse Compton parameter Y is proportional to $[\eta(\epsilon_e/\epsilon_B)]^{1/2}$, where $\eta = \min[1, (\nu_m/\nu_c)^{1/2}]$, and for these values of ϵ_e and ϵ_B , $Y < 0.3$. For values of $E_{k,52} = 5 \times 10^4$, $\epsilon_B = 6.2 \times 10^{-5}$, and $\epsilon_e = 3.1 \times 10^{-3}$ the particle density in the circumburst environment is $n = 0.08 \text{ cm}^{-3}$, which, using Eq. 8 gives an opening angle of $\theta_j = 2.0 \times 10^{-3} \text{ rad}$. This corresponds to $\theta_j = 0.11^\circ$, which gives a beam-corrected kinetic energy $E_k^{corr} = 10^{51}$ and γ -ray energy $E_\gamma^{corr} = 2 \times 10^{48}$. The jet opening angle that we estimate assumes the hydrodynamics as described by Panaitescu & Kumar (2000), and variations on the hydrodynamics, such as that discussed in Sari et al. (1999), result in a different coefficient in Eq. 8. However, these differences are of order unity, and do not affect the validity of the model discussed.

This model is, therefore, able to satisfy our observations without the requirement of an excessively large kinetic energy. However, the opening angle is a factor of at least 30 narrower than previously observed (Bloom et al. 2003) and the inferred γ -ray energy is extremely low for a long GRB, prompting us to consider other options.

3.2.2 $\nu > \max[\nu_m, \nu_c]$

Keeping in place the condition that $t_j < 80 \text{ s}$, we investigate a model in which the X-ray and UV/optical bands are above ν_m and ν_c . In this model our observational constraints on the values of ν_m , ν_c and $F_{\nu, \max}$ only apply at $T + 80 \text{ s}$. Since we assume a jet break to occur at $t_j = 80 \text{ s}$, the factor of $(t/t_j)^{(1-q)}$ in the kinetic energy is unity, and q is therefore not of importance in our calculations. For ν_c above the observing band, $\beta = p/2$, and therefore $p = 2$, in which case the dependence on the minimum Lorentz factor of the shock accelerated electrons γ_m changes from

$$\gamma_m = \frac{(p-2)}{(p-1)} \epsilon_e \frac{m_p}{m_e} \Gamma \quad (14)$$

to

$$\gamma_m = \ln\left(\frac{\epsilon_m}{\epsilon_M}\right) \epsilon_e \frac{m_p}{m_e} \Gamma, \quad (15)$$

where ϵ_m and ϵ_M are the minimum and maximum energy of the shock accelerated electrons, and Γ is the bulk Lorentz factor in the shocked medium (Pe'er & Zhang 2006). $\nu_m \propto \gamma_m$, and therefore the expression for ν_m given in Eq. 1 has to be modified such that $[(p-2)/(p-1)]^2$ is replaced by $[\ln(\epsilon_m/\epsilon_M)]^2$. The value of $\ln(\epsilon_M/\epsilon_m)$ is not well determined due to the unknown magnetic field in the upstream of the GRB shock, although it generally ranges between 5 and 10 (Li & Waxman 2006). We therefore take $\ln(\epsilon_M/\epsilon_m) = 7$, such that Eq. 1 becomes

$$\nu_m = 6.7 \times 10^{13} \text{ Hz} (1+z)^{1/2} \epsilon_B^{1/2} \epsilon_e^2 E_{k,52}^{1/2} t_d^{-3/2} \quad (16)$$

We assume the same jet break time as before ($t_j = 80 \text{ s}$), whereby Eq. 2 and 16 reduce to

$$\epsilon_B^{-3/2} E_{k,52}^{-1/2} n^{-1} (1+Y)^{-2} < 5.45 \quad (17)$$

and

$$\epsilon_B^{1/2} \epsilon_e^2 E_{k,52}^{1/2} < 2.1 \times 10^{-4} \quad (18)$$

In the regime where $\nu > \max[\nu_m, \nu_c]$,

$$F_\nu = F_{\nu, \max} \nu_c^{1/2} \nu_m^{(p-1)/2} \nu^{-p/2} \quad (19)$$

By taking the measured value of $F_\nu = 730$ at $\nu = 5.5 \times 10^{14} \text{ Hz}$, and using Eq. 3, 17 and 18, the expression for F_ν reduces to

$$\epsilon_e E_{k,52} (1+Y)^{-1} = 3.65 \quad (20)$$

From these equalities we then determine that a kinetic energy of $E_{k,52} = 10^3$ would require $\epsilon_e = 3.7 \times 10^{-3} (1+Y)$, $\epsilon_B < 0.25 (1+Y)^{-4}$ and $n > 0.05 (1+Y)^4$. For a particle density of $n = 10^4 \text{ cm}^{-3}$ in the GRB local environment, which is a typical density for a molecular cloud (Reichart & Price 2002), the jet opening angle would be $\theta_j = 0.8^\circ$, a factor of a few smaller than that determined for some other GRBs (e.g. GRB 980519; Nicastro et al. 1999). This gives a beam-corrected kinetic energy of $E_{k,52}^{corr} = 9.7 \times 10^{50}$.

This model provides an alternative scenario in which a highly collimated jet is the cause for the large flux observed in GRB 061007, but where a larger jet opening angle and more standard beam-corrected γ -ray can satisfy our observations, with respect to the previous model.

3.3 Synchrotron Self-Compton Emission

For completeness we look into a scenario in which synchrotron self-Compton (SSC) emission contributes to the emission. In the case where the X-ray emission is dominated by SSC radiation, and the UV/optical emission is produced by synchrotron radiation, both observed bands would need to be in the fast-cooling regime, with $\nu_{opt} > \nu_c > \nu_m$ and $\nu_X > \nu_c^{IC} > \nu_m^{IC}$, in order for the decay slope and spectral index in the X-ray and UV/optical emission to be consistent. In this case the spectral index is given by $\beta = p/2$, which would give a value of $p = 2.0$. However, the rate of synchrotron optical afterglow is expected to be $-(2-3p)/4$, which is inconsistent with our observations of $\alpha = 1.65 \pm 0.01$.

3.4 The Circumburst Medium

From our spectral analysis we determine the dust extinction in the circumburst medium to be most consistent with a wavelength dependence similar to that of the LMC. This is consistent with previous studies of GRB environments (e.g. Fruchter et al. 2006, Kann, Klose & Zeh 2006; Schady et al. 2006), which also find GRB hosts to be irregular galaxies, and is also in agreement with the collapsar model, which requires a sub-solar metallicity progenitor star, with an upper limit of $Z_\odot \lesssim 0.3$ (e.g. Hirschi, Meynet & Maeder 2005; Woosley & Heger 2006).

The best-fit parameters for the X-ray column density and dust extinction local to the GRB are $N_H = (5.3 \pm 0.3) \times 10^{21} \text{ cm}^{-2}$ and $A_V = 0.66 \pm 0.02$, and this gives a gas-to-dust

ratio of $N_H/A_V = (8.0 \pm 0.32) \times 10^{21}$. However, N_H was determined assuming solar abundances, which are unlikely to be correct for an irregular, Magellanic-type galaxy. Correcting the X-ray column measurement to a metallicity of 1/3 solar, appropriate for an ISM which is similar to that of the LMC, we find that the gas-to-dust ratio is a factor of four larger than that measured in the SMC or LMC. The gas-to-dust ratio measured in the local environment of GRB 061007 is also larger than the average of those determined for a set of previous *Swift* GRBs (Schady et al. 2006) by a factor of ~ 1.5 . This could be the result of dust destruction in the environment of the GRB.

The lack of evolution observed in the soft X-ray absorption later than ~ 80 s after the prompt emission implies that the burst must have completely ionised the gas in its immediate environment within this time scale. Given the large column density that is measured from the SED we, therefore, infer that the absorbing medium we observe is far enough from the burst that it was not completely ionised in the first few 100 s of seconds after the initial explosion. The probability of there being an absorption system in the line of sight from an unrelated, intervening galaxy that is of sufficient optical depth to contribute a measurable amount of soft X-ray absorption is very small (O’Flaherty & Jakobsen 1997; Campana et al. 2006), leading us to conclude that the absorbing material measured in the spectrum of GRB 061007 is local to the burst. From the results in Perna & Lazzati (2002) we estimate the absorbing medium to lie at approximately ~ 10 pc from the GRB. The large value of N_H is comparable to that of giant molecular clouds, from which we infer that GRB 061007 was embedded in a dense, star-forming region, or giant molecular cloud, in common with many other GRBs (e.g. Dai & Lu 1999; Wang et al. 2000; Reichart & Price 2002).

3.5 Comparing Between Models

It is interesting to note that with $E_{\gamma,iso} = 10^{54}$ ergs in the 10 keV–20 MeV energy range, and a rest-frame peak energy of $E_p = 902$ keV, GRB 061007 satisfies the Amati relation, given by $E_p \propto E_{\gamma,iso}^{0.5}$ (Amati et al. 2002). It also satisfies a correlation between the peak isotropic luminosity, $L_{p,iso}$, and spectral lag in the prompt emission, τ_{lag} , whereby $L_{p,iso} \propto \tau_{lag}^{-1.1}$. Both these correlations are satisfied by a large fraction of long GRBs (e.g. Amati 2006; Gehrels et al. 2006), suggesting that GRB 061007 has the same class of progenitor as other long GRBs.

The consistency observed in the spectral and temporal afterglow behaviour in the X-ray and UV/optical bands, and in the late time γ -ray band, implies that the emission observed comes from the same spectral segment. If we take the afterglow to be in the slow cooling regime and assume a jet break to occur at $t_j > 2.65 \times 10^5$ s a large amount of kinetic energy, of at least $E_k^{corr} = 9 \times 10^{53}$ is required. The implausibly large amount of energy required by this model makes it an unsatisfactory scenario to explain the properties of GRB 061007.

We also investigate a second scenario in which a highly collimated outflow produces a jet break before $T + 80$ s. This requires a jet angle no larger than 0.8° , which satisfies the observational requirements with a lower energy budget and more reasonable set of parameters. The upper limit on

the jet opening angle of 0.8° is a factor of 4 smaller than that previously determined in GRBs (Berger, Kulkarni & Frail 2003), although still physically possible. Furthermore, the need to observe the jet break in order to measure θ_j introduce selection effects, and the smaller and larger end of the current jet opening angle distribution will be underestimated. For the very smallest of jet opening angles to be measured, very early time data of the afterglow is needed, preferentially covering several energy bands.

A consequence of a very early jet break time is the faster evolution of Γ into a non-relativistic phase, which will produce a break in the light curve as the decay rate slows down. However, $\Gamma \propto t^{-1/2}$, and therefore the GRB outflow will not enter a non-relativistic phase until $t_{nr} = t_j \theta_j^{-2}$. For $\theta_j < 0.8^\circ$, this corresponds to $t_{nr} > 4 \times 10^5$ s, which is not in conflict with our observations.

For a jet break earlier than $T + 80$ s our observations are valid for a spectral regime in which $\nu_m < \nu < \nu_c$, and $\nu_c < \nu$, where in the latter a larger jet opening angle is obtained by assuming a density $n = 10^4 \text{ cm}^{-3}$, consistent with that observed in molecular clouds. The dependence of the circumstellar particle density on ϵ_B , ϵ_e and $E_{k,52}$ in the former of these models does not allow for such high densities, for which we estimate $n = 0.08 \text{ cm}^{-3}$. Our spectral analysis on the afterglow resulted in a large column density, which supports a scenario in which the GRB is embedded within a dense molecular cloud or active star-forming region. This therefore favours a model in which $\nu_c < \nu$, where the number density is inferred to be larger.

In the case of a uniform jet, where the energy and Lorentz factor are constant across the jet, the afterglow after a jet break should decay as t^{-p} . In both the post jet-break models that we consider the predicted decay slope is steeper than the observed value of $\alpha = 1.65 \pm 0.01$. A possible explanation is continual energy injection, which would maintain the energy in the afterglow and slow down the rate of decay. The requirement for steady energy injection over the entire light curve is of some concern, but seems plausible on the basis of many other examples of energy injection in *Swift* afterglow observations. Evidence of prolonged energy injection in the form of plateaus in X-ray light curves is observed in a large fraction of *Swift* GRBs, although typically up to 10^3 – 10^4 s after the prompt emission. In the case of GRB 061007 a constant amount of injection is needed for at least an order of magnitude longer than this, during which no break is observed in the light curve. This is not to say that the central engine needs to be active for this length of time, but simply that it is active for the duration of the prompt emission and ejects shells with a large distribution in the Lorentz factor. A range in Γ from a few tens to ten (e.g. Rees & Meszaros 1998; Sari & Mészáros 2000) can provide a source of energy to the afterglow for a long time after the central engine has stopped through ongoing refreshed shocks. Nevertheless, an energy injection mechanism that remains extremely constant for such a long time requires very specific conditions in which the energy injection mechanism dominates the afterglow during the entire period of observations.

Alternatively, the slow decay rate of the afterglow in the context of post jet-break could also be the result of a jet-edge effect in which there is no sideways expansion. In such a case the afterglow decay index would steepen by 3/4 when

$\Gamma = 1/\theta_j$, and would continue to decay at this rate due to the lack of lateral expansion (Mészáros & Rees 1999). In such a scenario there is no need for energy injection, although additional conditions would be required to prevent the jet from expanding further, such as a high magnetic field within the jet.

A possibly more natural form of non-laterally expanding jet is a structured jet or outflow with an energy distribution that is a decreasing function of angular distance from the jet axis, θ_0 , and a narrow, high-energy core at the axis. A jet break is, then, observed when the edge of the uniform inner core becomes visible to the observer. In order for a jet break to occur before $T+80$ s the viewing angle, therefore, needs to be close to the jet axis, which is not that unreasonable given the brightness of GRB 061007. At the time of jet break the smoother distribution in energy, ϵ , compared to a uniform jet results in a smaller change in the decay slope that will be shallower than the 3/4 increase in decay index observed in a uniform jet break. For a power law distribution of E , where $E \propto \theta^{-k}$, the smaller the value of k the shallower the afterglow light curve will be after the jet-break. In the case of $\nu_m < \nu_{obs} < \nu_c$, where $p = 3$, a decay rate after a jet break with index $\alpha = 1.65$ will occur in structured jet with an E distribution index $k = 0.3$, and for $\nu_c < \nu_{obs}$ we require $k = 1.4$ (Panaitescu 2005).

In the context of post-jet break emission, a decay slope of $\alpha = 1.65$ can, therefore, be accounted for by a non-laterally expanding jet as well as continual energy injection through refreshed shocks. We, therefore, find an early jet break before $T + 80$ s can satisfy the conditions imposed by our observations without requiring a large energy budget and with a reasonable set of physical parameters.

4 CONCLUSIONS

In this paper we have presented multi-wavelength spectral and temporal data of GRB 061007 taken with *Swift*; an incredibly bright GRB with unusually smooth panchromatic decay. Using these data we explored the circumstances that could produce such a bright burst, and so determined the conditions that distinguish GRB 061007 from other, more standard GRBs. On many levels we find GRB 061007 to be typical of others, suggesting that the same progenitor model for most long GRBs is also applicable to GRB 061007. The surrounding dust has properties most consistent with an LMC-like environment, and the circumburst material is likely to be dense, suggesting that it is embedded in a large molecular cloud or a region of active star formation. Furthermore, its consistency with both the Amati and timelag-luminosity relations, which apply to a large fraction of long GRBs, indicate that the source of energy in GRB 061007 is no different than that of most other long GRBs.

The enormous brightness of GRB 061007 is, therefore, unlikely to be due to a different progenitor model, but the result of extreme values for a few defining characteristics. Two inherently different models can satisfy our observations of GRB 061007, where the main difference lies in the time of the jet break; either before or after our observations. Both require certain extreme conditions to account for the brightness of this burst, either in the form of a large energy budget, or in the collimation of ejecta into a very narrow jet open-

ing angle. However, given the brightness of GRB 061007, it should not be a surprise that some of defining properties are non-standard. The large amount of kinetic energy required in a late-time jet break model lead us to favour a model in which the GRB 061007 is produced by the concentration of energy into a highly narrow jet $< 0.8^\circ$; a condition that can more easily be met than the production of large amounts of energy.

The brightness of GRB 061007 across the electromagnetic spectrum makes it ideal to explore the conditions surrounding the production of GRBs, and its uniqueness provides further insight into the range in properties that GRBs have. The small opening angle that we determine for GRB 061007 has important implications on the true range in jet opening angles, and this could provide a clue as to the cause for the large fraction of *Swift* GRBs with no clear signs of jet breaks.

ACKNOWLEDGEMENTS

We gratefully acknowledge the contributions of all members of the *Swift* team. SZ thanks PPARC for support through an Advanced fellowship.

REFERENCES

- Akerlof, C., et al. 1999, *Nature*, 398, 400
- Amati, L., et al. 2002, *A&A*, 390, 81
- Amati, L. 2006, *MNRAS*, 372, 233
- Barthelmy, S. D., et al. 2005, *Space Science Reviews*, 120, 143
- Berger, E., Kulkarni, S. R., & Frail, D. A. 2003, *ApJ*, 590, 379
- Bersier, D., et al. 2006, *GCN*, 5709
- Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, *ApJ*, 594, 674
- Burrows, D. N., et al. 2005, *Space Science Reviews*, 120, 165
- Campana, S., et al. 2006, *A&A*, 449, 61
- Chevalier, R. A., & Li, Z.-Y. 1999, *ApJ*, 520, L29
- Dai, Z. G., & Lu, T. 1999, *ApJ*, 519, L155
- Dickey, J. M., & Lockman, F. J. 1990, *ARA&A*, 28, 215
- Frail, D. A., et al. 2001, *ApJ*, 562, L55
- Freedman, D. L., & Waxman, E. 2001, *ApJ*, 547, 922
- Fruchter, A. S., et al. 2006, *Nature*, 441, 463
- Gehrels, N., et al. 2004, *ApJ*, 611, 1005
- Gehrels, N., et al. 2006, *astro-ph/0610635*
- Ghirlanda, G., Ghisellini, G., & Lazzati, D. 2004, *ApJ*, 616, 331
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'Shin, V., Frederiks, D., & Cline, T. 2006, *GCN*, 5722
- Hirschi, R., Meynet, G., & Maeder, A. 2005, *A&A*, 443, 581
- Hjorth, J., et al. 2003, *Nature*, 423, 847
- Huang, Y. F., Cheng, K. S., & Gao, T. T. 2006, *ApJ*, 637, 873
- Jakobsson, P., Fynbo, J. P. U., Tanvir, N., & Rol, E. 2006, *GCN*, 5716
- Kann, D. A., Klose, S., & Zeh, A. 2006, *ApJ*, 641, 993
- Kulkarni, S. R., et al. 1998, *Nature*, 395, 663

- Li, Z., & Waxman, E. 2006, astro-ph/0603427
Markwardt, C., et al., 2006, GCN, 5713
Meszaros, P., & Rees, M. J. 1997, ApJ, 476, 232
Mészáros, P., & Rees, M. J. 1999, MNRAS, 306, L39
Nicastrò, L., et al. 1999, A&A supp., 138, 437
Nousek, J. A., et al. 2006, ApJ, 642, 389
O'Brien, P. T., et al. 2006, ApJ, 647, 1213
O'Flaherty, K. S., & Jakobsen, P. 1997, ApJ, 479, 673
Osip, D., Chen, H. W., & Prochaska, J. X. 2006, GCN, 5715
Panaitescu, A., & Kumar, P. 2000, ApJ, 543, 66
Panaitescu, A. 2005, MNRAS, 363, 1409
Panaitescu, A., Mészáros, P., Gehrels, N., Burrows, D., & Nousek, J. 2006, MNRAS, 366, 1357
Pe'er, A., & Zhang, B. 2006, astro-ph/0605641
Pei, Y. C. 1992, ApJ, 395, 130
Perna, R., & Lazzati, D. 2002, ApJ, 580, 261
Rees, M. J., & Meszaros, P. 1998, ApJ, 496, L1
Reichart, D. E., & Price, P. A. 2002, ApJ, 565, 174
Romano, P., et al. 2006a, A&A, 456, 917
Romano, P., et al. 2006b, A&A, 450, 59
Roming, P. W. A., et al., 2005, Space Sci. Rev, 120, 95
Rykoff, E. S., & Rujopakarn, W., 2006, GCN, 5706
Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
Sari, R., Piran, T., & Halpern, J. P. 1999, ApJ, 519, L17
Sari, R., & Mészáros, P. 2000, ApJ, 535, L33
Schady, P., et al., 2006, submitted to MNRAS
Tagliaferri, G., et al. 2005, A&A, 443, L1
Tanvir, N. R., et al. 2004, Astronomical Society of the Pacific Conference Series, 312, 275
van der Horst, A. J., & Rol, E. 2006, GCN, 5720
Vaughan, S., et al. 2006, ApJ, 638, 920
Vetere, L., Pagani, C., Burrows, D. N., & Schady, P. 2006, GCN, 5714
Wang, X. Y., Dai, Z. G., & Lu, T. 2000, MNRAS, 317, 170
Woosley, S. E. 1993, ApJ, 405, 273
Woosley, S. E., & Heger, A. 2006, ApJ, 637, 914
Zeh, A., Kloze, S., & Kann, D. A. 2006, ApJ, 637, 889
Zhang, B., & Mészáros, P. 2001, ApJ, 552, L35
Zhang, B., Fan, Y. Z., Dyks, J., Kobayashi, S., Mészáros, P., Burrows, D. N., Nousek, J. A., & Gehrels, N. 2006a, ApJ, 642, 354
Zhang, B., et al. 2006b, ApJ, in press (astro-ph/0610177)