

TABLE 1. X-ray Galaxies and AGN Behind the Magellanic Clouds

X-ray Source ^a RX J	X-ray Position (J2000.0)			X-ray Count Rate ^b (cts s ⁻¹)	V	Redshift	Finding Chart ^c	Notes
	R.A.	Dec.						
0005.3–7427 = SMC 1	00 ^h 05 ^m 19.5 ^s	–74°26′42″		0.0315 ± 0.0058	16.3	0.1316	×	resolved galaxy
0033.3–6915 = SMC 70	00 33 21.5	–69 15 13		0.020 ± 0.003	15.5	0.0975	×	cluster A2789; 1.
0119.5–7301 = SMC 66	01 19 30.3	–73 01 28		0.006 ± 0.002	15.8	0.0656	×	cluster; 2.
0135.4–7048	01 35 23.5	–70 48 29		0.010 ± 0.005	17.9	0.0647	×	cluster?; 3.
0136.4–7105 = SMC 68	01 36 25.6	–71 05 25		0.0078 ± 0.0020	19.1	0.4598	×	
0454.2–6643	04 54 10.6	–66 43 17		0.0082 ± 0.0023	18.2	0.2279	×	
0503.1–6634 = CAL F	05 03 05.1	–66 33 47		0.0511 ± 0.0077	16.8	0.064	P1	
0517.3–7044 = CAL 21	05 17 17.1	–70 44 03		0.0063 ± 0.0011	18.2	0.1688	P2	
0524.0–7011 = CAL 32	05 24 02.4	–70 11 06		0.0338 ± 0.0052	17.7	0.151	P1	
0531.5–7130 = CAL 46	05 31 32.5	–71 29 48		0.0140 ± 0.0040	19.2	0.2214	P1	
0532.0–6920	05 31 59.2	–69 19 52		0.0050 ± 0.0011	18.9	0.1495	P1	
0534.0–7145	05 34 00.5	–71 45 28		0.0124 ± 0.0018	13.8	0.0242	×	large galaxy; 4.
0534.8–6739	05 34 45.7	–67 38 57		0.0045 ± 0.0011	17.8	0.0720	P2	‘star’ 2; 5.
0547.8–6745	05 47 45.4	–67 45 03		0.0097 ± 0.0021	20.5	0.3905	P2	radio source; 6.
0550.5–7110	05 50 32.1	–71 09 56		0.0024 ± 0.0007	19.9	0.4429	×	
0550.6–6637 = CAL 91	05 50 33.3	–66 36 59		-----	17.0	0.076	pub.	7.

^aSMC sources: Wang & Wu 1992; CAL sources, alternately called LHG: Long et al. 1981

^b*ROSAT*-HRI count rate in 25'' diameter aperture; unless otherwise specified, these are point sources

^cfinding charts: ×: this paper; P1: Schmidtke et al. 1994; P2: Cowley et al. 1997

Notes:

1. extended X-ray source; count rate in 72'' cell; associated with a cluster of galaxies; V mag measured in 40'' aperture; z for central cD gal
2. extended X-ray source; count rate in 72'' cell; optical galaxy V mag measured in 28'' aperture; double nucleus (see text)
3. extended X-ray source; count rate in 72'' cell; galaxy cluster?; z for brightest galaxy within the X-ray contours
4. X-ray point source in center of very large, optical galaxy (Up 053448–7147.3, S0(r))
V mag measured in 80'' aperture; spectrum has narrow [O II], [O III] emission
5. “note added in proof” in Paper 2 identifies this source with ‘star’ 2
6. compact radio source MDM 100 (Marx et al. 1997)
7. *Einstein* count rate and finding chart (Cowley et al. 1984); spectrum described (Cristiani et al. 1984)

TABLE 2. Grens Observations

Galaxy	Plate No.	Exposure (min.)	Prism Angle	1950 Coordinates of Center	
				R.A.	Dec.
NGC 45	2482	90	90°	00 ^h 11 ^m 37 ^s	-23°37' 00''
M 31 - E ^a	2481	60	0	00 33 58	+40 51 14
M 31 - C29	2466	60	245	00 47 53	+39 51 51
NGC 185	1499	90	0	00 36 13	+48 03 49
NGC 253	2483	90	90	00 45 10	-25 33 39
IC 1613	2476	90	90	01 02 23	+01 51 53
M 33 - A	1496	75	270	01 31 11	+30 39 11
	2477	90	180		
M 33 - B	1501	90	270	01 32 47	+30 15 10
	2484	90	180		
M 33 - C	1489	55	0	01 31 05	+30 39 11
	2471	90	270		
NGC 2366	2478	90	300	07 23 42	+69 19 05
NGC 2403	2485	90	300	07 32 14	+65 42 32
NGC 6822	1484	60	0	19 42 10	-14 55 56
	1498	90	190		
	2463	155	157		
	2474	90	90		

^a Close to C12 (Sargent et al. 1977)

TABLE 3. Quasar Candidates Around Nearby Galaxies

Quasar Candidates	I.D.	J2000 Coordinates	m_J	Class	Emission Lines ^a & Comments ^b
<u>NGC 45 Quasars:</u>					
0012.3–2305	15	00 ^h 12 ^m 19 ^s .0 –23°05′33″	21.0	4	5200; red; high z?
0012.3–2331	31	00 12 23.7 –23 31 29	20.8	4	poor focus; W of pair
0012.8–2308	F	00 12 51.5 –23 08 13	19.9	2	4320,3400; z = 1.8
0013.2–2302	C	00 13 13.3 –23 02 46	17.5	5	dA
0013.4–2324	25	00 13 25.6 –23 24 40	21.0	4	emission? short spectrum
0013.4–2304	G	00 13 27.5 –23 04 45	21.0	2	4100,5200
0013.5–2332	N	00 13 35.2 –23 32 32	20.0	2	4120
0013.7–2306	H	00 13 44.0 –23 06 18	19.1	2	4200,5100
0013.7–2259	D	00 13 46.5 –22 59 39	21.3	1	3840,4900; z = 2.2
0013.8–2328	O	00 13 51.4 –23 28 34	21.1	4	
0013.9–2319	M	00 13 54.2 –23 19 32	21.2	4	
0013.9–2316	J	00 13 56.2 –23 16 00	21.0	3	3875
0013.9–2327	P	00 13 58.5 –23 27 00	20.8	4	3980
0014.0–2250	B	00 14 00.4 –22 50 17	19.7	4	
0014.1–2309	12	00 14 08.8 –23 09 35	18.5	4	blue object; emission from NGC 45?
0014.2–2328	28	00 14 13.2 –23 28 34	20.4	4	overlap; W of pair
0014.4–2259	E	00 14 29.2 –22 59 02	21.2	1	3840,4890; z = 2.2
0014.5–2303	8	00 14 31.4 –23 03 18	21.1	4	
0014.7–2303	I	00 14 44.6 –23 03 12	20.0	3	3750
0014.8–2317	S	00 14 50.6 –23 17 49	21.4	4	
0014.9–2309	K	00 14 56.2 –23 09 36	20.2	4	4100
0014.9–2327	V	00 14 57.5 –23 27 39	20.0	3	3400,3850,4300; z = 1.8
0014.4–2321	R	00 14 26.7 –23 21 19	20.9	4	3600
0015.0–2320	U	00 15 04.2 –23 20 37	20.2	4	4060
0015.2–2312	T	00 15 14.7 –23 12 37	18.4	3	4500,3900,5200
0015.3–2306	L	00 15 23.4 –23 06 30	21.2	4	
<u>M 31 – Field E Quasars:</u>					
0035.3+4107	F	00 ^h 35 ^m 18 ^s .8 +41°07′02″	16.5	5	
0035.7+4046	A	00 35 46.2 +40 46 19	18.9	2	4960,4025; z = 1.6?
0035.7+4134	5	00 35 46.6 +41 34 02	18.5	3	
0035.8+4053	C	00 35 49.7 +40 53 57	16.9	5	
0035.9+4056	E	00 35 54.4 +40 56 17	20.9	3	3920,5210
0035.9+4112	K	00 35 57.8 +41 12 21	20.3	2	3550
0036.3+4117	O	00 36 19.2 +41 17 48	20.0	3	4000
0036.4+4125	T	00 36 24.9 +41 25 37	21.1	4	4140
0036.4+4130	U	00 36 26.7 +41 30 09	17.8	5	dA
0036.4+4105	J	00 36 29.6 +41 05 02	20.2	1	4120,5170; z = 2.4
0036.6+4134	8	00 36 40.1 +41 34 35	20.7	4	4500
0036.9+4120	S	00 36 55.1 +41 20 25	20.5	4	4800,4100
0036.9+4101	I	00 36 54.1 +41 01 57	20.6	4	4140
0036.9+4119	R	00 36 59.2 +41 19 25	20.5	1	4760,3960
0037.6+4053	H	00 37 41.5 +40 53 47	20.5	4	
0037.6+4049	G	00 37 41.5 +40 49 55	20.5	3	3780,4200
0037.8+4103	M	00 37 53.4 +41 03 00	20.3	4	4000
0037.9+4104	N	00 37 56.7 +41 04 54	20.3	4	3840
0038.4+4106	Q	00 38 27.0 +41 06 50	17.2	5	dA

TABLE 3. (continued)

Quasar Candidates	I.D.	J2000 Coordinates	m_J	Class	Emission Lines ^a & Comments ^b
<u>M 31 – Field C29 Quasars:</u>					
0048.8+4009	O	00 ^h 48 ^m 52 ^s .7 +40°09′18″	17.9	4	4040
0048.9+4005	V	00 48 58.4 +40 05 13	18.5	4	white dwarf?
0049.5+4006	U	00 49 33.2 +40 06 13	18.9	2	3750,4630; $z = 1.4$
0049.6+3952	H	00 49 41.0 +39 52 50	20.4	2	4160
0050.1+3958	I	00 50 07.8 +39 58 09	19.4	3	4000; very weak line(s)
0050.2+3941	P	00 50 13.7 +39 41 24	19.8	4	4030,3540
0050.3+4006	L	00 50 20.4 +40 06 22	21.0	4	4040; very weak lines
0051.1+4017	N	00 51 08.4 +40 17 38	20.5	4	
0051.2+3954	F	00 51 14.7 +39 54 55	19.0	2	4590,5160
0051.3+3947	C	00 51 21.1 +39 47 13	20.4	4	5120,3950
0051.4+3950	D	00 51 25.2 +39 50 06	19.3	1	4080,5160; $z = 2.4$
0051.5+3955	R	00 51 31.3 +39 55 49	20.4	4	
0051.8+3942	2	00 51 52.8 +39 42 50	18.1	4	
0052.1+3951	Q	00 52 07.2 +39 51 58	20.6	4	
0052.2+4019	14	00 52 13.9 +40 19 11	20.9	4	
0052.2+4012	J	00 52 15.7 +40 12 52	20.2	2	4530
0052.3+3944	B	00 52 23.3 +39 44 40	20.2	3	4740,3950; $z = 2.9?$
<u>NGC 185 Quasars:</u>					
0037.4+4803	D	00 ^h 37 ^m 27 ^s .5 +48°03′29″	18.0	3	3725,4925; white dwarf?
0039.6+4817	H	00 39 40.7 +48 17 58	19.0	2	3600,4600
0039.8+4802	G	00 39 48.4 +48 02 57	19.4	2	3750
0040.9+4820	K	00 40 56.9 +48 20 18	19.7	4	
<u>NGC 253 Quasars:</u>					
0045.7–2515	H	00 ^h 45 ^m 43 ^s .0 –25°15′35″	19.9	4	4980; *
0045.9–2503	V	00 45 59.6 –25 03 56	21.2	3	3880
0046.1–2503	W	00 46 11.8 –25 03 32	19.8	3	5000
0046.3–2528	N	00 46 22.4 –25 28 20	20.9	1	4550; $z = 2.7?$
0046.3–2532	R	00 46 22.1 –25 32 37	20.8	4	4060
0046.5–2457	U	00 46 32.6 –24 57 20	21.3	3	5100
0046.5–2518	Y	00 46 30.9 –25 18 15	21.2	3	3900
0046.6–2529	O	00 46 36.6 –25 29 28	19.9	3	3460,5120
0046.6–2508	E	00 46 39.3 –25 08 33	21.0	1	4140,4800,5220; $z = 2.4$
0046.6–2500	B	00 46 41.8 –25 00 27	20.0	3	4300
0046.7–2519	I	00 46 43.6 –25 19 34	21.2	2	4010
0046.7–2521	J	00 46 47.5 –25 21 51	20.9	4	3830; brighter on grems plate
0046.9–2531	A2	00 46 55.6 –25 31 22	21.0	4	5100?
0048.1–2459	C	00 48 08.7 –24 59 11	20.7	4	3750
0048.1–2535	S	00 48 08.0 –25 35 07	19.8	1	4330; *
0048.3–2523	K	00 48 23.0 –25 23 53	20.8	4	4100?; overlap?
0048.3–2510	F	00 48 23.2 –25 10 10	21.3	4	4170
0048.4–2528	P	00 48 27.0 –25 28 58	18.8	3	3920
0048.7–2512	X	00 48 44.7 –25 12 06	21.0	4	4300
0048.7–2529	Q	00 48 44.1 –25 29 40	20.7	3	3860
0049.2–2525	M	00 49 17.2 –25 25 57	19.6	4	
0049.2–2524	L	00 49 17.9 –25 24 42	20.6	3	3800

TABLE 3. (continued)

Quasar Candidates	I.D.	J2000 Coordinates	m_J	Class	Emission Lines ^a & Comments ^{ba}
<u>IC 1613 Quasars:</u>					
0102.9+0154	V	01 ^h 02 ^m 57 ^s .5 +01°54'49''	19.4	4	4750
0103.1+0202	P	01 03 08.5 +02 02 18	18.3	2	3530,4450; *
0103.2+0204	O	01 03 12.0 +02 04 14	20.2	2	4420,5100
0103.5+0150	W	01 03 30.8 +01 50 41	19.1	4	3700
0103.6+0212	J	01 03 36.9 +02 12 32	21.2	3	4000
0103.6+0212	K	01 03 39.2 +02 12 48	21.2	3	3900
0103.8+0224	B	01 03 52.5 +02 24 00	21.3	3	*
0104.1+0224	C	01 04 06.9 +02 24 57	20.4	3	4050
0104.2+0209	L	01 04 14.9 +02 09 28	21.1	4	4000
0104.3+0220	I	01 04 20.0 +02 20 11	20.6	2	4100,5050; $z = 1.6$
0104.7+0222	E	01 04 44.0 +02 22 16	21.3	2	3900
0104.7+0225	D	01 04 47.7 +02 25 02	21.2	1	4750,4020; $z = 2.9$
0104.9+0152	Y	01 04 59.6 +01 52 20	19.8	4	3900
0105.0+0201	S	01 05 05.3 +02 01 08	21.3	4	
0105.6+0158	A2	01 05 40.0 +01 58 14	20.7	4	
0105.7+0211	M	01 05 43.7 +02 11 31	21.5	4	4100
0105.7+0152	E2	01 05 44.1 +01 52 10	21.4	4	5050; $z = 3.2?$
0105.9+0214	N	01 05 54.2 +02 14 06	21.0	4	3760,4680
0106.0+0200	U	01 06 00.3 +02 00 30	21.0	3	4670
0106.0+0220	G	01 06 01.1 +02 20 53	20.7	3	3900
0106.3+0220	H	01 06 22.3 +02 20 50	20.3	3	4030
0106.4+0158	C2	01 06 27.1 +01 58 46	21.3	4	3625,4140
0106.7+0211	F2	01 06 42.3 +02 11 27	19.5	4	3925?
<u>M 33 Quasars:</u>					
0132.1+3022	A89	01 ^h 32 ^m 10 ^s .0 +30°22'57''	16.7	5	dA
0132.6+3107	J77	01 32 40.2 +31 07 50	18.7	4	overlap?
0134.0+3118	I77	01 34 00.5 +31 18 09	19.8	1	4025,5090; $z = 2.3$
0134.6+3114	C77	01 34 37.8 +31 14 48	20.1	2	4150,5200
0135.7+3035	D84	01 35 43.6 +30 35 35	20.6	4	4085
0135.3+3027	E84	01 35 23.1 +30 27 26	20.7	3	3800?
<u>NGC 2366 Quasars:</u>					
0726.2+6909	F	07 ^h 26 ^m 17 ^s .8 +69°09'22''	20.4	2	3820,4800; $z = 2.1$
0726.4+6856	S	07 26 26.2 +68 56 26	20.4	4	4080
0726.5+6857	P	07 26 30.8 +68 57 51	17.2	4	*
0729.3+6857	D	07 29 21.3 +68 57 45	18.2	1	3880
0729.7+6901	G	07 29 43.2 +69 01 19	20.4	2	3750
0729.7+6914	X	07 29 44.4 +69 14 19	20.5	4	3700
0729.7+6855	U	07 29 42.5 +68 55 51	19.4	5	
0729.9+6904	V	07 29 54.7 +69 04 31	21.1	3	4100
0730.6+6851	B	07 30 38.9 +68 51 51	20.5	2	4330,5000
0730.9+6920	L	07 30 59.3 +69 20 21	20.4	3	3970
0732.2+6905	I	07 32 15.5 +69 05 10	20.6	3	3600
0732.8+6909	W	07 32 50.7 +69 09 33	16.3	5	dA white dwarf

TABLE 3. (concluded)

Quasar Candidates	I.D.	J2000 Coordinates	m_J	Class	Emission Lines ^a & Comments ^b
<u>NGC 2403 Quasars:</u>					
0734.2+6548	P	07 ^h 34 ^m 13 ^s .8 +65°48'49''	19.2	4	3920
0734.6+6536	I	07 34 37.9 +65 36 17	20.9	2	4400
0734.8+6555	S	07 34 51.2 +65 55 38	19.2	3	4150; *
0735.1+6548	O	07 35 08.0 +65 48 19	19.2	2	4100,3550; *
0735.1+6553	R	07 35 09.2 +65 53 05	19.4	3	3500; *
0735.5+6535	K	07 35 31.5 +65 35 06	20.3	4	3850?
0735.7+6520	E	07 35 45.2 +65 20 36	20.8	2	5050
0737.2+6557	V	07 37 16.4 +65 57 39	20.3	4	5200
0737.5+6508	D	07 37 35.6 +65 08 25	18.4	2	3900,5000; *
0737.8+6516	F	07 37 52.5 +65 16 36	20.4	4	4000,4950
0738.3+6536	N	07 38 23.7 +65 36 40	19.9	2	4320; *
0739.2+6528	M	07 39 13.8 +65 28 11	19.7	4	3800
0740.4+6534	Q	07 40 24.0 +65 34 19	18.5	4	*
0740.4+6539	U	07 40 24.7 +65 39 57	19.4	2	4400; *
<u>NGC 6822 Quasars:</u>					
1943.9-1502	Q	19 ^h 43 ^m 58 ^s .9 -15°02'49''	19.9	2	5210; $z = 3.3?$
1944.4-1436	D	19 44 27.4 -14 36 43	19.3	5	
1944.6-1435	E	19 44 39.8 -14 35 17	20.1	4	
1944.8-1511	R	19 44 48.9 -15 11 35	19.7	4	5100,4020
1945.0-1441	L	19 45 01.0 -14 41 35	19.9	3	5225
1945.0-1439	J	19 45 02.8 -14 39 13	18.9	4	white dwarf?
1945.0-1457	U	19 45 04.7 -14 57 31	19.0	6	H II?
1945.1-1436	G	19 45 06.0 -14 36 40	18.1	3	
1945.0-1452	E98	19 45 00.3 -14 52 11	20.3	4	
1945.4-1448	B	19 45 26.6 -14 48 58	18.3	4	
1945.7-1429	A	19 45 47.1 -14 29 40	18.4	4	white dwarf?
1945.8-1437	I	19 45 53.0 -14 37 18	17.6	4	

^a estimated wavelengths given in order of decreasing intensity

^b tabulated redshifts estimated from gres spectra; * indicates note at end of table

Notes:

NGC 253 - H: identical to MS 00433-2531; $z = 0.861$ (Garilli et al. 1992)

NGC 253 - S: $z = 2.521$ (Clowes & Savage 1983)

IC 1613 - P: $z = 1.909$ (Chaffee et al. 1991)

IC 1613 -B: very faint; 2 lines, but wavelength scale uncertain

NGC 2366 - P: $z = 0.111$; [O II], H β , [O III], H α

NGC 2403 - S: $z = 1.76$; Ly α , [Si IV], C IV

NGC 2403 - O: $z = 2.53$; Ly α , Ly β , C IV

NGC 2403 - R: $z = 1.924$; Ly α , [N V], [Si IV], C IV, C III]; identical to 0730+65W1;
 $z = 1.927$ (Tytler & Fan 1992)

NGC 2403 - D: $z = 2.24$; Ly α , [Si IV], C IV

NGC 2403 - N: $z = 2.58$; Ly α , [Si IV], C IV

NGC 2403 - Q: quasar with one unidentified line at 4190Å

NGC 2403 - U: $z = 2.81$; Ly α , Ly β , [N V]

Probes for Nearby Galaxies

David Crampton^{1,2} and G. Gussie^{2,4}

Dominion Astrophysical Observatory, National Research Council of Canada,
Victoria, B.C. V8X 4M6, Canada

A.P. Cowley^{1,2,3} and P.C. Schmidtke^{1,3}

Department of Physics & Astronomy, Arizona State University, Tempe, AZ, 85287-1504

ABSTRACT

Data are given for sixteen extragalactic objects (predominantly AGN) behind the Magellanic Clouds and for 146 quasar candidates behind the nearby galaxies NGC 45, 185, 253, 2366, 2403 and 6822, IC 1613, M31 and M33. The Magellanic Cloud objects were identified by their X-ray emission, and precise optical and X-ray positions and optical photometry and spectra are presented for all of these. The quasar candidates surrounding the other nearby galaxies were identified through a CFHT slitless spectral survey. Although redshifts for only eight of these candidates have been obtained, previous observations indicate that the majority are likely to be quasars. A subsample of 49 of the brighter objects could confidently be used, in addition to the Magellanic Cloud sources, as probes of the gas in nearby galaxies for rotation curve studies, for studies of their halos, for comparison with higher redshift QSO absorption lines, or as references for proper motion studies.

Subject headings: Galaxies: Active – Galaxies: Quasars: General – X-rays: Galaxies

1. INTRODUCTION

The UV spectroscopic capabilities of HST allow nearby galaxies to be probed via absorption lines in the same way that QSO absorption line systems probe galaxies at high redshift. Although

¹Visiting Astronomers, Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation

²Visiting Astronomers, Canada-France-Hawaii Telescope, operated by the National Research Council of Canada, the Centre de la Recherche Scientifique de France, and the University of Hawaii

³Guest Observer, Multiple Mirror Telescope Observatory, operated jointly by the Smithsonian Observatory and the University of Arizona

⁴Student assistant, 1985. Present address: Dept. of Physics, U. of Tasmania, Hobart, Australia

recent studies have shown that Mg II absorption lines at $z < 1$ appear to be associated with luminous, massive galaxies (Steidel, Dickinson & Persson 1994), there is still a debate about the nature of the absorbing material. For example, it is unclear whether the absorption occurs in an extended disk, halo or even satellite galaxies. Studying absorption lines in AGN and QSOs behind nearby galaxies allow these galaxies to be probed along multiple sightlines, thereby yielding information on how the absorption correlates with various parameters of the foreground galaxy. Accurate measurements of velocities of the interstellar material can also be used to extend rotation curves to larger galactocentric distances, thereby providing improved estimates of the mass distributions. Furthermore, the background quasars can be used for precise proper motion studies of the foreground objects. Unfortunately, however, most surveys for quasars have avoided directions towards nearby galaxies. Monk et al. (1986) give a list of the brightest ($m < 17.5$) quasars located within ~ 200 kpc of nearby galaxies. In this paper we report results from an X-ray survey of the Magellanic Clouds and an optical slitless-spectra survey of nearby northern hemisphere galaxies.

For several years we have been undertaking a census of the X-ray sources in the Magellanic Clouds, first with data from *Einstein* Observatory (e.g., Cowley et al. 1984) and, more recently, with *ROSAT* (Schmidtke et al. 1994, hereafter Paper I; Cowley et al. 1997, hereafter Paper II). Several background galaxies, AGN, and QSOs were detected in addition to X-ray bright objects within the Clouds. Some of these were reported in Papers I and II, but the identifications and redshifts of most have only recently been determined. In this paper we bring together the data for all of these objects. We give photometry and redshifts for sixteen X-ray selected extragalactic objects, five in the SMC field and eleven in the vicinity of the LMC. We also present finding charts for those that are not already in the literature.

In the course of another project, searches for quasar candidates were carried out on a series of CFHT gress plates of nearby galaxies. These plates are ideally suited for the detection of QSO candidates (e.g., Crampton, Schade & Cowley 1985). Although we have not been able to confirm the identifications or measure the redshifts for most of these candidates, our previous high success rate in identifying QSOs (Crampton, Cowley & Hartwick 1987) demonstrates that most of them can confidently be expected to be AGN. For this reason we present these data now since, apart from the faintest low quality identifications, most will be quasars and hence useful as probes regardless of their redshift. However, a low S/N spectrum confirming the QSO identification is recommended before investing large amounts of telescope time for, say, absorption line studies.

2. X-RAY SELECTED AGN BEHIND THE MAGELLANIC CLOUDS

Nearly 200 point X-ray sources in the direction of the Magellanic Clouds were detected by *Einstein* Observatory (Long, Helfand, Grabelsky 1981; Wang et al. 1991; Seward & Mitchell 1981, Wang & Wu 1992; Cowley et al. 1984). Details of recent *ROSAT* X-ray observations of many of these Magellanic Cloud sources are given in Papers I and II. Identifications of the optical

counterparts have been carried out through extensive photometric and spectroscopic observations at CTIO. This program reveals that many of these sources are associated with foreground stars or background extragalactic objects (e.g. Papers I and II). The optical data which we have obtained are described below.

Our *ROSAT*-HRI survey of the Magellanic Clouds was aimed at obtaining improved positions of *Einstein* sources which had not been already identified to enable detection of new optical counterparts. Therefore, our material does not cover the entire fields of the LMC and SMC and thus does not comprise a complete sample of AGN behind these galaxies. The objects which are included in our X-ray sample are listed in Table 1 together with their *ROSAT* positions and count rates. We have carefully investigated the positional accuracy delivered by the *ROSAT* detectors (see discussion in Papers I and II) and the positions reported here are accurate to about $\pm 5''$, making the search for optical counterparts relatively straightforward. Figures 1 and 2 show the distribution of X-ray bright extragalactic objects we have found near these two galaxies. Finding charts are given in Figures 3 and 4 for those which are not already in the literature. The X-ray positions are indicated by a ‘+’ and the optical counterpart is marked with a dash. Three of the X-ray sources are spatially extended, and for those we have overlaid their X-ray contours on the optical finding charts. Each of these appears to be associated with a cluster of galaxies.

2.1. Photometry

All the photometric data were obtained from CCD photometry carried out at CTIO during observing runs between 1992 and 1996 using the 0.9-m telescope. The *V* magnitudes presented in Table 1 are based on aperture photometry, calibrated using observations of Landolt (1992) standard stars. The accuracy of the magnitudes is about ± 0.02 mag. A few special cases are mentioned below where the object was extended or complicated in some way. The astrometry of each CCD frame used for the finding charts has been tied to the coordinate system of the *HST Guide Star Catalogue* (Lasker et al. 1990) by measuring the positions of ~ 6 – 8 stars on the digitized *GSC*.

2.2. Spectroscopy

Our highest resolution AGN spectra were obtained with the CTIO 4-m telescope in November 1996 with the KPGL1 grating and Loral 3K detector. These spectra cover the wavelength range 3700–6700Å and have a resolution of ~ 1.0 Å per pixel. With a $1''.5$ slit, corresponding to three pixels, the spectral resolution is 3Å. The spectra of the optical counterparts of CAL 21 and RX J0532.0–6920 were taken with the CTIO ARGUS fiber system in December 1995. These spectra cover the range 3650–5800Å with a resolution of 1.8Å per pixel. The spectrum of RX J0547.8–6745 has a wavelength range of 3720–5850Å and also has a resolution of 1.8Å per pixel.

One-dimensional spectra were extracted and processed following standard techniques with IRAF to yield wavelength-calibrated spectra. The spectra are shown in Figure 5, shifted to restframe wavelengths according to their redshifts (Table 1).

Some of the earlier spectra were taken with the SIT vidicon detector on the CTIO 4-m RC spectrograph. These are not shown in Figure 5, but the objects observed are included in Table 1.

2.3. Individual Sources

Some of the newly identified individual sources deserve comment beyond just listing them in Table 1. Of the ones previously published (see references in Table 1), we point out that RX J0534.8–6739 was only listed as a “note added in proof” by Cowley et al. (1997) so it would be very easy to overlook it in that paper. The AGN is “star” 2 in the finding chart for this source in Paper II.

RX J0005.3–7427 (SMC 1): This X-ray point source falls nearly on the galaxy shown in the finding chart in Fig. 3. Its spectrum, shown in Fig. 5, indicates the object is a Seyfert I galaxy with a redshift of $z = 0.1316$. The optical image shows the galaxy is just resolved.

RX J0033.3–6915 (SMC 70): The X-ray contours for this source are very extended, as shown in Fig. 3. The X-ray position falls on a large cD galaxy in the center of the cluster Abell 2789. Thus, the X-rays appear to result from hot gas in the cluster. The redshift of the cD galaxy is $z = 0.0975$.

RX J0119.5–7301 (SMC 66): The extended X-ray contours suggest that this source also arises from hot gas, but this is a previously unknown cluster. Many galaxies are visible in the field (see finding chart in Fig. 3). A spectrum of the bright cD galaxy to the west of the central X-ray contour shows it to have a redshift of $z = 0.0658$. The optical image of this galaxy has three parts, consisting of a foreground star and two non-stellar condensations. The magnitude given in Table 1 refers to the brighter (south-western) of the two non-stellar parts.

RX J0135.4–7048: This weak, extended source also appears to be associated with an unknown cluster of galaxies (visible on our original image, but not easily seen in Figure 3). The center of the extended X-ray contours is not coincident with any bright galaxy, but we have observed the nearest (bright) one, which is south-east of the third X-ray contour (as marked on the finding chart in Fig. 3). Its spectrum gives a redshift of $z = 0.0647$. The identification of this X-ray source with the cluster, and whether the galaxy we observed is a member, should be verified by obtaining redshifts of some of the fainter galaxies.

RX J0136.4–7105 (SMC 68), RX J0454.2–6643, and RX J0550.5–7110: These three sources are all associated with AGN, as shown by their redshifts in Table 1 and their spectra in Fig. 5.

RX J0534.0–7145: This X-ray point source appears to be located in or near to the nucleus of the very large optical galaxy, Up 053448–7147.3, which is listed as an S0(r). The V magnitude was measured in an $80''$ aperture. The optical spectrum shows narrow [O II] and [O III] emission but otherwise is relatively normal. It thus appears to be one of the narrow-emission-line galaxies that comprise about $\sim 15\%$ of the extragalactic X-ray galaxy population (e.g., Griffiths et al. 1996). The large extent of the optical disk can be seen in Fig. 4.

RX J0547.8–6745: This point X-ray source is identified with an AGN with redshift $z = 0.3905$. It was also recently found to be a compact radio source, MDM 100 (Marx et al. 1997).

3. QSOS AND QSO CANDIDATES AROUND NEARBY GALAXIES

Quasars with $z < 3.4$ and $B < 21$ can be easily recognized from CFHT blue greys images. The blue greys, a grating-prism-lens combination designed by E.H. Richardson, produces spectra with a dispersion of $945\text{\AA} \text{ mm}^{-1}$ and a wavelength range of $3500\text{--}5300\text{\AA}$ when recorded on IIIaJ emulsion. The greys plates cover a $55' \times 55'$ field, although some parts of this field may be vignetted by the guide probe. Greys exposures were obtained of eight fields centered on northern nearby galaxies and two “halo” M31 fields, centered on field C29 of Sargent et al. (1977) and a field to the SW, primarily to study objects in the galaxies themselves. In some cases, observations with different greys orientations were taken to alleviate problems of overlapping images. A list of the plates is given in Table 2. Unfortunately, the seeing was not very good ($> 1''$) during most of these observations.

As in previous surveys, four visual searches for objects with a UV excess and/or emission lines were made of each plate by at least two of the authors (in this case, by DC, GG and APC). Objects satisfying these criteria but which were likely to be associated with the nearby galaxy were ignored in this survey. Subsequently, PDS scans were made of the spectra of all candidates with a 50 micron square aperture. These spectra were converted to intensity versus wavelength with a software package written by Graham Hill. As in previous quasar surveys (e.g., Crampton, Schade & Cowley 1985), the candidates were assigned a class based on these tracings and a final visual inspection of the original image. Class 1 candidates are certain quasars with strong emission lines, class 4 objects show UV excess with no definite spectral features, and classes 2 and 3 are intermediate between these extremes. Recognizable white dwarfs are included as class 5 since they might otherwise be selected as quasar candidates on the basis of their colors. Emission-line galaxies with no significant spatial extension or “extragalactic H II regions” are assigned class 6. Spectroscopic follow-up observations with the MMT of candidates with $m < 20.5$ indicate that, on average, 100% of class 1 objects are quasars, 93% of class 2, 76% of class 3 and 42% of class 4 candidates are quasars. Further details of the observational and identification procedures are given by Crampton, Schade & Cowley (1985). Due to the low galactic latitude of many of these fields, and to internal absorption in the galaxies themselves, fewer candidates than typical were

identified in these fields.

Positions and magnitudes of all candidates were measured from glass copies of the Palomar Sky Survey O plates using the method and software described by Stetson (1979). Scans of 100×100 $10\mu\text{m}$ square pixel boxes were made of each candidate with the PDS and the positions were related to nearby SAO stars. Subsequently, the new SKYCAT software was used in conjunction with the digitized Palomar Observatory Sky Survey to double-check the coordinates and charts and, in some cases, to correct for errors. The resulting positions are accurate to $\sim 2''$ and magnitudes to ~ 0.3 mag.

A list of all candidates is given in Table 3. The first column gives the candidate name derived from truncated 2000 coordinates in the form HHMM.M+DDMM. An internal identification symbol is given in the second column, followed by the 2000 coordinates, the magnitude as estimated from the POSS plates, and the class or certainty of the identification. In the notes column we give: (1) the estimated wavelengths of any emission features visible on the gress spectra, listed in order of decreasing intensity (2) redshifts, listed to one digit accuracy if they were estimated from the gress observations, (3) other comments or remarks (for explanation of the abbreviations and any measured redshifts, see the Notes to the table). Rather than give a table listing projected distances from the centers of the nearby galaxies, the locations of the candidates (marked with their ID as given in Table 3) are shown in Figures 6 – 15 so that their distances relative to the optical extent of the galaxies are obvious.

Spectra of nine of the candidates in NGC 2366 and NGC 2403 were obtained in rather poor weather conditions with the MMT in 1987 February with the photon-counting spectrograph. The spectra cover the $3000\text{--}8000\text{\AA}$ region with a resolution of $\sim 7\text{\AA}$. Identified features and redshifts for these candidates are given in the notes to Table 3.

4. SUMMARY

Sixteen galaxies, clusters, and AGN behind the Large and Small Magellanic Clouds have been identified through their X-ray emission. Redshifts of most of these indicate that they are relatively nearby, with only three having $z > 0.3$. Quasar candidates in ten fields in the direction of nearby galaxies have been identified on the basis of their colors and emission lines. Of these, twenty-two have magnitudes brighter than $m=19$ and thus are excellent targets for high spectral resolution studies with HST or 10-m class telescopes. Forty-nine candidates with magnitudes brighter than $m=20.5$ and classes 1 – 3 have an extremely high probability of being quasars, based on similar surveys.

We thank Dr. Martha Hazen of the Harvard College Observatory who located the photographs of the Magellanic Clouds and kindly sent them to us, and Dr. Sidney van den Bergh for permission to use his plate of M 33. We also thank Y. Yuan for her careful checking of the material in Table

3 and for assistance with the diagrams. The excellent new SKYCAT tool developed jointly by ESO and the CADC was extremely useful in verifying and confirming positions and magnitudes of all our objects. We thank D. Durand for his support in its use and help in installing additional catalogs. A.P.C. and P.C.S. gratefully acknowledge support from NSF for this work.

REFERENCES

- Chaffee, F.H. et al. 1991, *AJ*, 102, 461
- Clowes, R.G. & Savage, A. 1983, *MN*, 204, 365
- Cowley, A.P., Crampton, D., Hutchings, J.B., Helfand, D.J., Hamilton, T.T., Thorstensen, J.R., & Charles, P.A. 1984, *ApJ*, 286, 196
- Cowley, A.P., Schmidtke, P.C., McGrath, T.K., Ponder, A.L., Fertig, M.R., Hutchings, J.B., & Crampton, D. 1997, *PASP*, 109, 21 (Paper II)
- Crampton, D., Cowley, A.P., & Hartwick, F.D.A. 1987, *ApJ*, 314, 129
- Crampton, D., Schade, D., & Cowley, A.P. 1985, *AJ*, 90, 987
- Cristiani, S. & Tarenghi, M. 1984, *A&A*, 132, 351
- Garilli, B., Bottini, D., Maccagni, D., Vettolani, G., & Maccacaro, T. 1992, *AJ*, 104
- Griffiths, R.E., Della Ceca, R., Georgantopoulos, I., Boyle, B.J., Stewart, G.C., Shanks, T. & Fruscione, A. 1996, *MNRAS*, 281, 71
- Landolt, A.U. 1992, *AJ*, 104, 340
- Lasker, B.M., Sturch, C.R., McLean, B.J., Russell, J.L, Jenkner, H., and Shara, M.M. 1990, *AJ*, 99, 2019
- Long, K.S., Helfand, D.J., & Grabelsky, D.A. 1981, *ApJ*, 248, 925
- Marx, M., Dickey, J.M., & Mebold, U. 1997, *A&A*, in press
- Monk, A.S., Penston, M.V., Pettini, M., & Blades, J.C. 1986, *MN*, 222, 787
- Sargent, W.L.W., Kowal, C.T., Hartwick, F.D.A., & van den Bergh, S. 1977, *AJ*, 82, 947
- Schmidtke, P.C., Cowley, A.P., Frattare, L.M., McGrath, T.K., Hutchings, J.B., & Crampton, D. 1994, *PASP*, 106, 843 (Paper I)
- Seward, F.D. & Mitchell, M. 1981, *ApJ*, 243, 736
- Steidel, C.C., Dickinson, M., & Persson, E. 1994, *ApJ*, 437, L75
- Stetson, P.B. 1979, *AJ*, 84, 1056
- Tytler, D. & Fan, X.-M. 1992, *ApJS*, 79, 1
- Wang, Q., Hamilton, T., Helfand, D.J., & Wu, X. 1991, *ApJ*, 374, 475
- Wang, Q. & Wu, X. 1992, *ApJS*, 78, 391

Fig. 1.— The Large Magellanic Cloud, showing the distribution of X-ray galaxies and AGN discussed in this paper. The photograph was taken with the 8-inch Bache photographic doublet at the Boyden Station of the Harvard College Observatory in Arequipa, Peru on 1897 Nov. 25. This 6-hour and 5-minute exposure was taken using a blue-sensitive emulsion. (Harvard plate no. B20849, reproduced with the permission of Harvard College Observatory)

Fig. 2.— The Small Magellanic Cloud, showing the distribution of X-ray galaxies and AGN discussed in this paper. The photograph was taken with the 10-inch Metcalf photographic triplet at the Boyden Station of the Harvard College Observatory in Bloemfontein, South Africa on 1950 Oct. 4. This 45-minute exposure was taken using a blue sensitive emulsion. (Harvard plate no. MF39311, reproduced with the permission of Harvard College Observatory)

Fig. 3.— Finding charts for the Magellanic Cloud X-ray sources RX J0005.3–7427 (SMC 1), RX J0033.3–6915 (SMC 70), RX J0119.5–7301 (SMC 66), RX J0135.4–7048, RX J0136.4–7105 (SMC 68), and RX J0454.2–6643. Note that the finding charts have different scales. For the extended X-ray sources (SMC 70, SMC 66, and RX J0135.4–7048) the X-ray contours have been overlaid on the optical images. These three are associated with clusters of galaxies. For the point X-ray sources, the X-ray position is shown by the + and the optical identification is marked by a dash.

Fig. 4.— Finding charts for RX J0534.0–7145 and RX J0550.5–7110. The X-ray contours are shown for RX J0534.0–7145, although it is a point source. The large, central galaxy is Up 053448–7147.3, S0(r). For RX J0550.5–7110 the X-ray position is shown by a + and the associated AGN is marked by a dash. Note that the two charts have different scales.

Fig. 5.— Spectra of galaxies and AGN associated with X-ray sources behind the Magellanic Clouds. The spectra have been ‘de-redshifted’ and gaussian smoothed for display purposes. They were then normalized to a peak continuum intensity of one and arbitrarily shifted so that they could be presented in order of increasing redshift from bottom to top. Rest wavelengths of some prominent features are indicated. The sources are: (1) RX J0119.5–7301, the central galaxy in a cluster, (2) RX J0534.0–7145, (3) RX J0033.3–6915, (4) RX J0135.4–7048, (5) 0534.8–6739, Sy I galaxy, (6) RX J0005.3–7427, Sy I galaxy, (7) RX J0517.3–7044, (8) RX J0532.0–6920, (9) RX J0454.2–6643, (10) RX J0531.5–7130, (11) RX J0547.8–6745, (12) RX J0550.5–7110, and (13) RX J0136.4–7105.

Fig. 6.— NGC 45, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5′.

Fig. 7.— M 31 – Field E, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5′.

Fig. 8.— M 31 – Field C29 (Sargent et al. 1977), showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 9.— NGC 185, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 10.— NGC 253, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 11.— IC 1613, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 12.— M 33, showing the distribution of quasar candidates and white dwarfs from all three fields listed in Tables 2 and 3. The image is reproduced from a Palomar 5-m 103aO plate taken by S. van den Bergh. North is to the top and East to the left, and the bar indicates 5'.

Fig. 13.— NGC 2366, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 14.— NGC 2403, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.

Fig. 15.— NGC 6822, showing the distribution of quasar candidates and white dwarfs. The image is taken from the Sky Survey print. North is to the top and East to the left, and the bar indicates 5'.