

This figure "fig1\_1.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig2\_1.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig1\_2.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig2\_2.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig1\_3.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig3.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig1\_4.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig4.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>



This figure "fig5.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig6.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig7.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig8.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

This figure "fig9.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9906468v1>

# A SURVEY FOR INFALL MOTIONS TOWARD STARLESS CORES. I. CS (2 – 1) AND N<sub>2</sub>H<sup>+</sup> (1 – 0) OBSERVATIONS

Chang Won Lee<sup>1,2</sup>, Philip C. Myers<sup>1</sup>, & Mario Tafalla<sup>1,3</sup>

<sup>1</sup>Harvard-Smithsonian Center for Astrophysics,  
60 Garden Street, MS 42, Cambridge, MA 02138, USA

<sup>2</sup>Taeduk Radio Astronomy Observatory, Korea Astronomy Observatory,  
36-1 Hwaam Dong, Yusung Ku, Taejon 305-348, Korea

<sup>3</sup>Observatorio Astronómico Nacional,  
Campus Universitario, Apartado 1143, E-28800, Alcalá de Henares (Madrid), Spain  
E-mail: cwl@hae.issa.re.kr, pmyers@cfa.harvard.edu, tafalla@oan.es

## ABSTRACT

We present the first results of a survey of 220 starless cores selected primarily by their optical obscuration (Lee & Myers 1999) and observed in CS(2–1), N<sub>2</sub>H<sup>+</sup> (1–0), and C<sup>18</sup>O (1–0) using the NEROC Haystack 37-m telescope. We have detected 163 out of 196 sources observed in CS, 72 out of 142 in N<sub>2</sub>H<sup>+</sup>, and 30 out of 30 in C<sup>18</sup>O. In total, 69 sources were detected in both CS and N<sub>2</sub>H<sup>+</sup>.

The isolated component of the N<sub>2</sub>H<sup>+</sup> (1–0) spectrum (F<sub>1</sub>F = 01 – 12) usually shows a weak symmetric profile which is optically thin. In contrast, a significant fraction of the CS spectra show non-Gaussian shapes, which we interpret as arising from a combination of self absorption due to lower excitation gas in the core front and kinematics in the core.

The distribution of the normalized velocity difference ( $\delta V_{CS}$ ) between the CS and N<sub>2</sub>H<sup>+</sup> peaks appears significantly skewed to the blue ( $\delta V_{CS} < 0$ ), as was found in a similar study of dense cores with embedded young stellar objects (YSOs) (Mardones et al. 1997). The incidence of sources with blue asymmetry tends to increase as the total optical depth or the integrated intensity of the N<sub>2</sub>H<sup>+</sup> line increases. This overabundance of “blue” sources over “red” sources suggests that inward motions are a significant feature of starless cores. We identify 7 strong infall candidates and 10 probable infall candidates. Their typical inward speeds are sub-sonic, approximately 0.04 – 0.1 km s<sup>-1</sup>, so they contain ‘thermal’ infall motions, unlike the faster inward

speeds associated with most YSOs (Mardones et al. 1997). We discuss the importance of the choice of a consistent set of line frequencies when using the velocity shift between an optically thick and a thin line as a tracer of infall, and show how the results of the survey depend on that frequency choice.

*Subject headings:* ISM: Globules; ISM: Kinematics and Dynamics; Stars: Formation

## 1. Introduction

A “starless” core is a dense core with no embedded IRAS source and no associated T Tauri star, implying that it is in the earliest identifiable stage of star formation (Shu, Adams, & Lizano 1987; Ward-Thompson et al. 1994; Lee & Myers 1999, hereafter LM99). Low mass stars are believed to form through the gravitational collapse of such a core when its central density becomes high enough (Shu 1977).

The last decade has seen remarkable progress on the understanding of low mass star formation. However, most studies of infall phenomena have been focussed on Young Stellar Objects (YSOs) or cores with embedded YSOs (Zhou et al. 1993; Gregersen et al. 1997; Mardones et al. 1997 hereafter M97). Most of our understanding of the physics of the pre-stellar stage is still entirely theoretical (e.g., Shu, Adams, & Lizano 1987, Ciolek & Mouschovias 1995). A systematic observation of the starless cores has been carried out in continuum (Ward-Thompson et al. 1994, 1999), but not yet in spectral lines sensitive to inward motions. Thorough observational studies of starless cores in molecular lines are required to understand initial conditions for star formation and evolution of star-forming cores.

The physical and kinematical state of starless cores is still unclear. Using sub-millimeter data of 21 starless cores, Ward-Thompson et al. (1994) suggested that starless cores are currently in the ambipolar diffusion phase prior to protostellar collapse. However, combining observations of optically thick and thin lines, Tafalla et al. (1998) and Williams et al. (1999) have shown that even a starless core like L1544 can present many symptoms of advanced evolution toward star formation. These authors found that the large area with infall asymmetry in L1544 is inconsistent with the ‘inside-out’ collapse model (Shu 1977), and that the large inward speed (up to  $0.1 \text{ km s}^{-1}$ ) is also too fast to result from ambipolar diffusion. Therefore L1544 was interpreted as having some type of inward motions in a manner not explained by any present ‘standard’ star formation model. We still do not know whether L1544 is an exceptional case or it represents a phase that every core experiences in the early course of star formation.

Systematic and extensive observational studies of starless cores may allow us to find good answers for the above questions. Recently, Lee & Myers (1999) have presented a new catalogue of dense cores optically selected by using the STScI Digitized Sky Survey, where 306 starless cores are identified. We have made extensive multi-molecular line surveys toward starless cores, mainly based on the list of starless cores in the LM99 catalogue which provides the largest available database to study starless cores with a statistical approach. Here we report the first result of one such survey using optically thick [CS (2–1)] and thin [N<sub>2</sub>H<sup>+</sup> (1–0) and C<sup>18</sup>O (1–0)] tracers, as the first step to study the physical and kinematical status of starless cores. Our main goals in this paper are to quantify the incidence of infall motions toward starless cores and to find infall candidates for further study in the future.

In §2 we explain how we selected and observed the starless cores, and reduced the molecular line data. Results of the qualitative and quantitative analysis of the molecular line survey data are



presented in §3. Implications of our statistics on starless cores, identification of infall candidates, kinematics of the suggested infall candidates, and future work are discussed in §4; we summarize our results in §5.

## 2. Observations

We have conducted single-pointing observations for 220 starless cores in the lines of CS (2–1), N<sub>2</sub>H<sup>+</sup> (1–0), and C<sup>18</sup>O (1–0). Most source positions were chosen from the catalogue of optically selected dense cores by LM99, to which we added 3 new positions (L1622A-2, L183B, and L1155C-2) and slightly modified 17 within about 1′ – 2′ using our recent FCRAO maps (Lee, Myers, & Tafalla 1999) so that they coincide with molecular peaks. Nine of the 220 positions were adopted from Benson & Myers (1989), although 7 of them are listed in LM99 catalogue with similar coordinates because the LM99 catalogue was not available at the beginning of our survey. Tables 1 and 2 list the names and positions of the observed sources, together with the position references. Including the observations of all sources with similar positions to those in the LM99 catalogue, we have observed about 93% (212) of the 227 sources in the catalogue which are observable from Haystack ( $\delta \gtrsim -25^\circ$ ).

In order to understand the optical depth and kinematical characteristics of the spectra, it is necessary to observe both optically thick and thin molecular lines. In this way, we can distinguish profile asymmetries that arise from self absorption from those due to multiple components in the emitting gas. We have chosen CS (2–1) as our thick tracer and N<sub>2</sub>H<sup>+</sup> (1–0) as our thin tracer because they have been already successfully used in identifying infall signatures in YSOs (e.g., M97). The C<sup>34</sup>S isotope would be the ideal thin partner for CS, as it probes the same physical properties (temperature, density, chemistry, and size scale) as the main isotope. However, it is usually too weak to determine its peak velocity with a good signal-to-noise (S/N) ratio, and for this reason, we have chosen N<sub>2</sub>H<sup>+</sup>. The N<sub>2</sub>H<sup>+</sup> (1–0) line at 93 GHz consists of 7 hyperfine components (Caselli, Myers, & Thaddeus 1995; see also Fig. 5), allowing us to determine the peak velocity with about twice the precision of a single line with the same S/N ratio. The isolated component (F<sub>1</sub>F = 01 – 12) is usually optically thin (M97), and does not show self-absorption features for most sources. The C<sup>18</sup>O (1–0) line was used as a substitute optically thin tracer for several sources not detected in N<sub>2</sub>H<sup>+</sup> (1–0).

For the quantitative study of spectral line asymmetries, it is necessary to use accurate and consistent frequencies for the different molecular lines. Despite the high accuracy of laboratory microwave spectroscopy, the frequency uncertainties in standard line catalogs (e.g, Lovas 1992, Pickett et al. 1998) are of the order of tens of kHz, and therefore close to some of the observed line velocity differences (few tenths of km s<sup>−1</sup>, see Section 4). This makes the choice of the line frequencies potentially critical in our statistical study of velocity differences between the peaks of the thick and the thin tracers, and a careful consideration of the frequency values is necessary. For this reason we have adopted the frequencies recommended by Tafalla et al. (1998), which seem to

constitute a consistent set: 97980.950 MHz for CS (2–1) from Pickett et al. (1998), 109782.182 MHz for C<sup>18</sup>O (1–0) from Lovas & Krupenie (1974), and 93176.2651 MHz for the N<sub>2</sub>H<sup>+</sup> (1–0) “isolated” component (F<sub>1</sub>F = 01 – 12) from Caselli et al. (1995). In the Appendix, we describe in detail how and why this set of frequencies was chosen.

Our observations were made using the NERO Haystack 37m telescope<sup>1</sup> in 1997 February, March, and December, and in 1998 January, March, and May. Two receivers were used simultaneously to obtain spectra in the left and right circular polarizations, and the signals were later combined. The back-end was an autocorrelator with 17.8 MHz bandwidth and 2.17 kHz (0.007 km s<sup>-1</sup>) spectral resolution. The spectra were obtained in frequency-switching mode with a frequency throw of  $\pm$  a quarter of the bandwidth –  $\pm$ 4.425 MHz. The full width at half maximum (FWHM) of the telescope beam is about 27'' at 86 GHz (Barvainis et al. 1993), and the telescope main beam efficiencies are  $18 \pm 3\%$  at 86 GHz and  $16.9 \pm 2\%$  at 115 GHz (Ball et al. 1993). The total integration time for each spectrum was usually between 30 and 120 minutes, depending on the source brightness, in order to achieve a S/N ratio better than  $\sim 8$ . Pointing was checked by using mostly Venus, Saturn, and Jupiter whenever the region of the observing source was changed. The usual pointing error was within about 6''.

### 3. Results

#### 3.1. Detection Statistics

In total, 220 starless cores have been observed in the CS (2–1) or N<sub>2</sub>H<sup>+</sup> (1–0) lines. With a detection criterion of  $T_A^* \gtrsim 5\sigma$ , 163 out of 196 sources observed in CS (2–1) were detected, and 72 out of 142 sources were detected in N<sub>2</sub>H<sup>+</sup> (1–0). The ranges of line intensities ( $T_A^*$ ) and widths (FWHMs) are approximately 0.05 – 0.4 K and 0.3 – 0.9 km s<sup>-1</sup> for CS in about 80% of the detected sources, and 0.05 – 0.4 K and 0.2 – 0.55 km s<sup>-1</sup> for the N<sub>2</sub>H<sup>+</sup> isolated component in about 90% of the detected sources. The peak intensity and FWHM of the CS line for the detected sources have mean and standard deviation of about  $0.28 \pm 0.17$  K and  $0.68 \pm 0.29$  km s<sup>-1</sup>, and the corresponding values for N<sub>2</sub>H<sup>+</sup> are  $0.19 \pm 0.11$  K and  $0.35 \pm 0.14$  km s<sup>-1</sup>. Table 1 lists the source coordinates and velocities for the cores which have been observed in CS (2–1) or N<sub>2</sub>H<sup>+</sup> (1–0), but detected only in one line or not detected in either line.

As a result of our survey, 69 starless cores were detected in both CS (2–1) and N<sub>2</sub>H<sup>+</sup> (1–0), and these sources are suitable for a comparison between the CS and N<sub>2</sub>H<sup>+</sup> line velocities. Table 2 lists their coordinates, line velocities, N<sub>2</sub>H<sup>+</sup> line width and optical depth, and the line asymmetry parameter  $\delta V_{CS}$  which will be explained in §3.4.

---

<sup>1</sup>Radio astronomy observations at the Haystack Observatory of the Northeast Radio Observatory Corporation are supported by a grant from National Science Foundation

The CS peak velocities and their  $1\sigma$  uncertainties in Tables 1 and 2 were determined from Gaussian fits. CS spectra consisting of two components or of a brighter component with a shoulder were fitted only for the brighter component by masking the fainter part or the shoulder in order to obtain the velocity of the peak component of the spectrum. The values of LSR velocity, FWHM, and their uncertainties for  $\text{N}_2\text{H}^+$  (1–0) were obtained by fitting seven Gaussian hyperfine components with the line parameters (velocity offsets and relative intensities) as determined by Caselli et al. (1995), using the Gaussian hyperfine structure (hfs) fit routine in the ‘CLASS’ reduction program (Buisson et al. 1994).

In Figure 1, we show 43 pairs of CS (2–1) and  $\text{N}_2\text{H}^+$  (1–0) (isolated component) spectra for sources with strong detections, and in Fig. 2 we present similar pairs for sources whose isolated components were barely detected or not seen. In this second group, the six less-isolated hyperfine components of the  $\text{N}_2\text{H}^+$  line are usually bright enough to determine the peak velocity of the  $\text{N}_2\text{H}^+$  line with useful precision, and the velocity is indicated in Fig. 1 and Fig. 2 by a dashed vertical line. We also present, in Fig. 3, spectra of 9 sources that show interesting non-Gaussian features in the CS line, while the  $\text{N}_2\text{H}^+$  line was not observed or not detected.

We observed 30 sources in  $\text{C}^{18}\text{O}$  (1–0) and all were detected. The  $\text{C}^{18}\text{O}$  spectra for 5 sources which have not been detected in the  $\text{N}_2\text{H}^+$  (1–0) line are shown in Fig. 3. These  $\text{C}^{18}\text{O}$  lines help to examine possible velocity shifts in the CS spectra.

### 3.2. Analysis of the CS Lines

As illustrated in Figs. 1, 2, and 3, a substantial fraction of the CS(2–1) spectra show non-Gaussian features, with some type of asymmetry like two peaks or a peak with shoulder. This is in contrast with what is seen in  $\text{N}_2\text{H}^+$ , where the isolated component appears as a single Gaussian feature, narrower than the equivalent CS line. This suggests that the CS asymmetries may be related to the higher optical depth of this line, and that they probably arise from self absorption by lower excitation foreground gas. Furthermore, we notice that the spectral shapes of the CS(2–1) spectra vary from source to source, indicating that the characteristics of the absorbing gas vary strongly from one object to the other.

We classify the CS spectra in 6 different groups depending on their shape (Fig.4). The line characteristics and typical examples for each group are as follows;

1. Two peaks with the blue peak brighter than the red (Fig. 4a) - e.g., L1521F, L183B, L1689B, L694-2, and L1445<sup>1</sup>

---

<sup>1</sup>It is difficult to decide from our data whether in this source the double peak originates from self absorption because the  $\text{N}_2\text{H}^+$  line was not detected. Our recent NRAO-12m survey (Lee, Myers, & Plume 1999) shows that the peak velocity of another optically thin tracer,  $\text{DCO}^+$ (2-1), lies between the two CS(2–1) peaks

2. Blue peak and red shoulder (Fig. 4b) - e.g., L1355, L1498, L1551S-2, TMC2, B18-3, CB23, L1622A-2, L1696B, L158, L234E-1, L492, L1155C-1, and L1197.

3. Single symmetric peak (Fig. 4c) - e.g., L1333, L1400A, L204C-2, L1517A, L1517B, L462-1, and L1063.

4. Red peak and blue shoulder (Fig. 4d) - e.g., L1521E

5. Two peaks with the red peak brighter than the blue peak (Fig. 4e) - e.g., CB246-2

6. Two peaks with similar brightness (Fig. 4f) - e.g., L1521B, L1517C-1, L1544, L204C, and L429-1

### 3.3. Analysis of the $N_2H^+$ Lines

The  $N_2H^+$  (1–0) line consists of 7 hyperfine components of different optical depths, being the isolated component ( $F_1F = 01 - 12$ ) usually optically thin (e.g., Caselli et al. 1995). Fig. 5 shows four types of  $N_2H^+$  spectra seen in our survey. The overwhelming majority ( $\sim 53$ ) of the detected  $N_2H^+$  lines have Gaussian shape, as shown in Fig. 5a for L183. A few sources, however, have double-peaked spectra, even in the isolated component, as in L492 (Fig. 5d). For  $\sim 15$  sources, the isolated component is too weak to be detected, while the other six components of the spectrum are fairly bright (e.g., Fig. 5b). For these sources, the six less-isolated components play an important role in determining the peak velocity of  $N_2H^+$  (1–0) line. The spectral shapes of several sources are sometimes different from component to component. We find that 6 sources (L1521F, L1551S-2, L1544, L63, L429-1, and L492) have double-peaked lines in the several hyperfine components (e.g., Fig5-c, and d). Our recent surveys in  $H^{13}CO^+$  (1 – 0) (Lee, Myers, & Park 1999) and  $DCO^+$  (2 – 1) (Lee, Myers, & Plume 1999) show that all these sources present single Gaussian components in other thin tracers [ $H^{13}CO^+$  (1 – 0) and  $DCO^+$  (2 – 1)], indicating that the two  $N_2H^+$  peaks arise not from overlapping components, but from self-absorption due to high optical depth. It is noted that L1521F and L1544 show a blue asymmetry in the main component ( $F_1F = 23 - 12$ ) of the  $N_2H^+$  line.

### 3.4. Distribution of the Velocity Differences between CS (2–1) and $N_2H^+$ (1–0)

In a previous section we have classified qualitatively the CS line profiles into 6 groups relying on their spectral shapes. Now we quantify this by estimating the amount by which the CS (2–1) spectrum is blue- or red-shifted with respect to the  $N_2H^+$  (1–0) line. To do this, we follow M97 and use the normalized velocity difference  $\delta V = (V_{CS} - V_{N_2H^+})/\Delta V_{N_2H^+}$ , where  $V_{CS}$  and  $V_{N_2H^+}$  are the peak velocities of the CS and  $N_2H^+$  lines, and  $\Delta V_{N_2H^+}$  is the FWHM of  $N_2H^+$  (1–0). The calculated values of  $\delta V_{CS}$  for the 69 sources in our survey with both CS and  $N_2H$  detections are listed in Table 2. However, for further analysis, we drop 2 sources (L1544 and L429-1) whose

intensity ratio between the blue component ( $T_b$ ) and the red component ( $T_r$ ) of the CS spectra is not significantly different from unity [ $\frac{T_b}{T_r} \lesssim 1 + \sigma_{(\frac{T_b}{T_r})}$ ], so any choice of a brighter component is ambiguous.

In Fig. 6 we show two histograms of the distribution of  $\delta V_{CS}$  in our sample. The left panel represents the distribution for the set of frequencies recommended by Tafalla et al. (1998), which corresponds to the  $N_2H^+$  (1–0) frequency determined by Caselli et al. (1995) and the CS (2–1) frequency recommended by Pickett et al. (1998). The right panel represents the same distribution when we use Lovas (1992) recommended value for CS (2–1) and keep the same value for  $N_2H^+$ . In the Appendix we discuss in detail why the first frequency set should be preferred, although here we will discuss both options in parallel. Probably the strongest conclusion from Fig. 6 is how crucial (and urgent) are high-precision frequency measurements for future infall searches.

Adopting the first (and preferred) set of frequencies, the  $\delta V_{CS}$  distribution is clearly skewed to the blue side ( $\delta V < 0$ ), with a mean value ( $\pm$  standard error of the mean) for the 67 sources of  $-0.24 \pm 0.04$ . This number, very close to the  $-0.28 \pm 0.10$  found by M97 for a set of Class 0 sources with broader lines, suggests that inward motions play an important role in the kinematics of starless cores (see below). A statistical t-test with unknown standard deviation shows that the probability of drawing the observed  $\delta V_{CS}$  distribution from a zero mean t-distribution is less than 5%, so the skewness is highly significant.

In Table 3, we have marked a ‘B’ for the 20 sources with a significant blue asymmetry ( $\delta V_{CS} \leq -5 \sigma_{\delta V_{CS}}$ ), and a ‘R’ for the 3 sources with significant red asymmetry ( $\delta V_{CS} \geq 5 \sigma_{\delta V_{CS}}$ ).

If we adopt the second set of frequencies, however, the distribution appears symmetric, with a mean indistinguishable from zero, and a high probability (about 80% in the t-test) of arising from a symmetric population. Until higher precision CS (2–1) frequencies are available, no further progress seems possible in determining the statistics of  $\delta V_{CS}$  with more accuracy.

A more robust result, as it holds for the two possible frequency choices, is the systematic decrease of  $\delta V_{CS}$  as a function of the  $N_2H^+$  (1–0) total optical depth  $\tau_{N_2H^+}$ . To study this, we have measured  $\tau_{N_2H^+}$  using the hyperfine structure (hfs) fitting routine in CLASS (Buisson et al. 1994) with the relative frequencies of the components as determined by Caselli et al. (1995), and the derived values are indicated in Table 2. The error in  $\tau_{N_2H^+}$  can become unacceptably large for very high or very low optical depths, or if the  $N_2H^+$  (1–0) spectrum is too noisy, and for that reason, we have discarded 13 sources for most of which the estimated  $\tau_{N_2H^+}$  error is larger than the measurement (these sources have an ‘\*’ before their  $\tau_{N_2H^+}$  values in Table 2).

Fig. 7 illustrates the distribution of  $\delta V_{CS}$  as a function  $\tau_{N_2H^+}$  for our preferred set of frequencies (see Appendix), although the same trend is found if the Lovas (1992) value is adopted. Fig. 7 shows that there is a strong tendency for  $\delta V_{CS}$  to become more negative with increasing total optical depth, and all 6 sources with  $\tau_{N_2H^+}$  larger than 11 have large negative  $\delta V_{CS}$ . A similar trend is also seen in Fig. 8 which shows  $\delta V_{CS}$  as a function of the total integrated intensity

of the  $\text{N}_2\text{H}^+$  line – a measure of the  $\text{N}_2\text{H}^+$  column density. The incidence of sources with blue asymmetry ( $\delta V < 0$ ) clearly increases with increasing  $\text{N}_2\text{H}^+$  integrated intensity.

## 4. Discussion

### 4.1. Implication of the velocity difference distribution

In the last section we have shown that if we adopt our preferred set of frequencies, the statistics of the velocity difference between  $\text{CS}(2-1)$  and  $\text{N}_2\text{H}^+(1-0)$  is significantly skewed to the blue, and that even if we adopt the Lovas (1992) frequency for CS, the objects with large  $\tau_{\text{N}_2\text{H}^+}$  systematically have negative  $\delta V_{\text{CS}}$ . Here we discuss the possible kinematic origin for this trends.

Bipolar outflows can strongly affect the shape of molecular lines and give rise to a strong asymmetric profiles even in dense gas tracers used in our survey (e.g., M97). Our sample, however, was selected to contain starless cores, and therefore should have no molecular outflows. To test this assumption, we have examined all spectra looking for high velocity wings. Among 163 CS spectra, only two sources (L183B and L429-1) show small wing components, in agreement with our expectation that our sources are truly starless and have no detectable signs of star formation. We therefore conclude that outflow contamination is not affecting our statistics of the velocity difference distribution.

Core rotation can also produce blue and red spectral asymmetry at either side of the rotational axis (Adelson & Leung 1988, Zhou 1995). Our recent mapping survey in  $\text{CS}(2-1)$  and  $\text{N}_2\text{H}^+(1-0)$  (Lee, Myers, & Tafalla 1999), however, shows that the incidence of sources with a well-defined shift from blue to red asymmetry is also small, suggesting that core rotation does not affect our statistics of velocity difference distribution either.

Finally, inward motions (infall) will also give rise to blue asymmetry in spectral profiles. Line excitation in a centrally concentrated core will tend to decrease outwards (unless compensated by a strong temperature gradient), so the foreground gas will appear in absorption in optically thick lines. If inward motions prevail, the foreground absorbing material will be red shifted (approaching the background), so the self absorption will preferentially affect the red side of the line and the resulting profile will have a brighter blue peak (e.g., Leung & Brown 1977). Inward motions (if spherical) have the property of systematically producing blue asymmetry in the spectra, in contrast with bipolar outflow and rotation, which produce both blue and red asymmetries in different parts of the core. This suggests that even if outflow and rotation were present, when studying a large sample of objects like ours, their effect will cancel out, and if infall is present, it will dominate the statistics of spectral asymmetries (see also M97). We therefore conclude that the most likely explanation for the overabundance of blue asymmetric starless cores in our sample (if our frequency choice is correct), and the overabundance of blue asymmetric cores at the highest

$\text{N}_2\text{H}^+$  (1 – 0) optical depths (independent of the frequency choice) is caused by the presence of inward motions in the gas.

We emphasize that the above conclusion refers to a statistical property of the sample as derived from single-pointing observations, and it does not guarantee that each source with blue asymmetric profile has necessary inward motions over a significant range of gas density and spatial extent. Spatial mapping of the spectral asymmetry is still necessary to verify that a given potential infall candidate is a truly infalling core.

#### 4.2. Infall Candidates

One important objective of this paper is identifying infall candidates for further, more detailed study. To carry out this identification, we analyze the following three spectral properties: (1) value of  $\delta V_{CS}$ , (2) intensity ratio of the blue to the red component in double-peaked CS spectra, and (3) intensity ratio of the blue to the red component in double-peaked  $\text{N}_2\text{H}^+$  spectra. The result of this analysis is summarized in Table 3 (for our frequency choice, see Appendix).

Property (1) is indicated as ‘B’, ‘N’, or ‘R’ in the second column of Table 3 depending on the value of  $\delta V_{CS}$  (‘B’ for  $\delta V_{CS} \leq -5\sigma_{V_{CS}}$ , ‘N’ for  $|\delta V_{CS}| < 5\sigma_{V_{CS}}$ , and ‘R’ for  $\delta V_{CS} \geq 5\sigma_{V_{CS}}$ ). To be a good infall candidate, a source should have a ‘B’ in this column.

Property (2) can be also expressed as ‘B’, ‘N’, or ‘R’ according to the ratio  $T_b/T_r$  (‘B’ for  $T_b/T_r > 1 + \sigma$ , ‘N’ for  $1 - \sigma \leq T_b/T_r \leq 1 + \sigma$ , and ‘R’ for  $T_b/T_r < 1 - \sigma$ ), where the values of  $T_b$  and  $T_r$  were derived from Gaussian fits to each component of the CS spectra with double peaks or single peak with distinct shoulder. To fulfill our criterion of being a good infall candidate, the ratio should be greater than 1.0 within a  $1\sigma$  error. This criterion allows us include sources whose  $\delta V_{CS}$  are poorly determined due to weak  $\text{N}_2\text{H}^+$  emission, but whose CS spectra show an apparently self-absorbed infall asymmetry (e.g., L1689B).

Property (3) can also help identifying infall candidates, because the  $\text{N}_2\text{H}^+$  hyperfine components trace higher density gas (a few  $10^5 \text{ cm}^{-3}$ ) than CS (2–1) (a few  $10^4 \text{ cm}^{-3}$ ), when their different optical depths are taken into account. Like with property (2), we have marked ‘B’, ‘N’, or ‘R’ those sources satisfying  $T_b/T_r > 1 + \sigma$ ,  $1 - \sigma \leq T_b/T_r \leq 1 + \sigma$ , or  $T_b/T_r < 1 - \sigma$ , respectively.

The above properties used for selecting infall candidates are similar to those used by M97 and Gregersen et al. (1997). Given their multiplicity, we can define different quality degrees for the candidates. Thus, we define as a “strong” infall candidate a source that is blue (B) according to at least 2 criteria, and is not red (R) according to any. A “probable” infall candidate is defined as one which is blue (B) according to one criterion and is not red (R) according to any. With these rules, we find 7 “strong” infall candidates – **L1355, L1498, L1521F, L183B, L158, L694-2, and L1155C-1**, and 10 “probable” infall candidates – **L1524-4, L1445, TMC2, B18-3, CB23, L1544, L1622A-2, L1689B, L234E-1, and L492**.

For the list of “probable” infall candidates, we have dropped several sources that are ‘B’ according to property (1) due to a very narrow  $\text{N}_2\text{H}^+$  line, but that do not show any pronounced asymmetry in the spectral profiles. These sources are L1400K, TMC1, L183, L1696A, L922-2, L1155C-2, and L1234. Two sources in our infall-candidate list, L1544 and L1498, are previously known and well studied infall candidates (see Tafalla et al. 1998, Williams et al. 1999 for L1544; Kuiper, Langer, & Velusamy 1996, Willacy, Langer, & Velusamy 1998 for L1498), but the additional 15 sources in our list are new infall candidates. In contrast, we have found 1 strong outflow candidate (CB246-2) and one probable outflow candidates (L63), according to the equivalent rules as those described before (L1049-1 is marked with “R” according to the spectral property (1), but it does not show any pronounced asymmetry in the CS spectrum, so we have excluded it from the category of “probable” outflow candidate).

All the above analysis has been done using our preferred set of frequencies (see Appendix). If the Lovas (1992) CS (2–1) frequency is used instead, the number of infall candidates decreases, but all 7 ‘strong’ infall candidates still remain ‘strong’. Five of the 10 ‘probable’ infall candidates, however, become ‘N,’ while 5 other sources (L1445, TMC2, L1544, L1689B, & L492) remain ‘probable’.

### 4.3. Infall Kinematics of The Starless Cores

Typical infall speeds for our candidates can be approximately calculated by taking half of the velocity dispersion ( $= \frac{1}{2}\sigma_v = \frac{1}{2} \frac{\Delta V_{\text{N}_2\text{H}^+}}{(8 \ln 2)^{\frac{1}{2}}}$ ) of the optically thin  $\text{N}_2\text{H}^+$  line (e.g., Leung & Brown 1977, Myers et al. 1996). The infall speeds of all sources are found to be approximately between  $0.04 - 0.1 \text{ km s}^{-1}$ .

For each core, we have calculated the line FWHM for thermal motions [ $\Delta V_T = (8 \ln 2 kT/m)^{1/2}$ , where  $k$  is the Boltzmann constant,  $T$  the kinetic temperature, and  $m$  the mean molecular weight (2.3 amu)], and the line FWHM for non-thermal motions [ $\Delta V_{NT} = (\Delta V_{\text{N}_2\text{H}^+}^2 - \frac{8 \ln 2 kT}{m_{\text{N}_2\text{H}^+}})^{\frac{1}{2}}$ , where  $m_{\text{N}_2\text{H}^+}$  is the molecular weight of  $\text{N}_2\text{H}^+$  (29 amu)] using the observed  $\Delta V_{\text{N}_2\text{H}^+}$  and assuming  $T = 10 \text{ K}$ . Taking the ratio of the two ( $\Delta V_{NT}/\Delta V_T$ ), we find that it is always smaller than 1 except for one source (L492; ratio equal to 1.13), meaning that most of the starless infall candidates are in state of ‘thermal’ infall. This is contrast to the result of M97, who found a prevalence of turbulent infall motions when studying a sample of cores with embedded YSOs. We note that this difference arises from the broader lines in the M97 sample, and not from their  $\delta V_{CS}$  value, which is very similar to that found here for starless cores.



#### 4.4. Future Work

Though our statistical analysis of the distribution of velocity differences strongly suggests that inward motions dominate the kinematics in starless cores, it is still premature to state that the blue asymmetry in each of our candidates is purely from inward motions, and it is not contaminated by other elements like outflow or rotation. In addition, even if the motions we observe are truly inward, it can not be decided yet whether they arise from ‘inside-out’ gravitational collapse (Shu 1977), or from contraction due to ambipolar diffusion (Mestel & Spitzer 1956) or any other process [e.g., pressure-driven motions (Myers & Lazarian 1998)].

A more comprehensive picture of the kinematics of starless cores requires further observations of additional transitions [e.g., CS (3 – 2)], different gas tracers (e.g., ions such as HCO<sup>+</sup>), and 2-dimensional mapping in both lines and continuum. Work in the above directions is currently in progress.

### 5. Conclusions

We have carried out a survey of 220 starless cores by observing them with the Haystack 37-m telescope in the optically thick CS (2–1) line, and the optically thin N<sub>2</sub>H<sup>+</sup> (1–0) and C<sup>18</sup>O (1–0) lines. The main goal of this survey has been to study the presence and frequency of inward motions in cores which have not yet formed stars.

The sources for these observations were selected primarily from the catalogue of optically selected cores by LM99, although a few new sources were added and several positions were slightly changed. Out of 227 sources in the LM99 catalogue observable from the latitude of the Haystack Observatory, about 93% (212) have been observed.

As a result of the survey, 163 out of 196 sources observed were detected in CS, 72 out of 142 were detected in N<sub>2</sub>H<sup>+</sup>, and 30 out of 30 were detected in C<sup>18</sup>O. In total, 69 sources were detected in both CS and N<sub>2</sub>H<sup>+</sup>, and these are the sources used for our analysis of inward motions.

The main conclusions of this analysis are:

1. The CS lines change in shape strongly from one source to the other, and very frequently present asymmetric features. From a comparison with the optically thin lines of N<sub>2</sub>H<sup>+</sup> (1–0) and C<sup>18</sup>O (1–0), which are usually symmetric and Gaussian, we conclude that most of the CS asymmetries arise from a combination of optical depth effects and kinematics. We classify the CS spectra according to their shape in the 6 following groups: (1) two peaks, with blue peak brighter than red, (2) blue peak and red shoulder, (3) single symmetric peak, (4) red peak and blue shoulder, (5) two peaks, with red peak brighter than blue peak, and (6) two peaks with similar brightness.

2. The isolated component of N<sub>2</sub>H<sup>+</sup> (1–0) (F<sub>1</sub>F = 01 – 12) is found to be optically thin in

most sources. For 6 sources, however, several hyperfine components of the  $\text{N}_2\text{H}^+$  line show what appear to be self absorption features that differ from component to component. L1521F and L1544 show blue asymmetry in the main component of the  $\text{N}_2\text{H}^+$  line ( $F_1F = 23 - 12$ ).

3. We find consistent indications that inward motions are a significant feature in the starless cores of our large sample.

First, we find that the distribution of the normalized velocity difference  $\delta V_{CS}$  between the CS and  $\text{N}_2\text{H}^+$  lines for our sample is significantly skewed to the blue side ( $\delta V_{CS} < 0$ ). The statistics of the average value of  $\delta V_{CS}$  and of the student's t-test are similar to those found for cores with embedded Class 0 sources (Mardones et al. 1997). There is also an excess of sources (20) with  $\delta V_{CS} \leq -5 \sigma_{\delta V_{CS}}$  over (3) sources with  $\delta V_{CS} \geq 5 \sigma_{\delta V_{CS}}$ . In addition, the incidence of the sources with infall asymmetry tends to increase as the total optical depth or the integrated intensity of the  $\text{N}_2\text{H}^+$  line increases.

Second, among sources with self-absorbed features in the CS line, there is an excess of sources (10) for which the intensity ratio between the blue and the red peak is larger than 1.0, while there is just one source where the ratio is less than 1.0.

5. Based on three spectral properties – the value of  $\delta V_{CS}$ , the intensity ratio of the blue to the red peak in two-peaked CS spectra, and the intensity ratio of the blue to the red peak in two-peaked main hyperfine components of  $\text{N}_2\text{H}^+$  (1–0), we identify 7 *strong infall candidates* – **L1355, L1498, L1521F, L183B, L158, L694-2, and L1155C-1**, and 10 *probable infall candidates* – **L1524-4, L1445, TMC2, B18-3, CB23, L1544, L1622A-2, L1689B, L234E-1, and L492**. All sources but two (L1498 and L1544) are newly found from this survey.

6. If the motions we observe are true infall, their typical speed is about  $0.04 - 0.1 \text{ km s}^{-1}$ . This implies infall in starless cores is subsonic and so in thermal motion, in contrast with the turbulent infall speeds found by M97 in cores with embedded YSOs.

7. Our identification of strong infall candidates and the increase in infall asymmetry with line optical depth are independent of the choice of CS and  $\text{N}_2\text{H}^+$  line frequencies. The statistical distribution of infall asymmetry, however, is crucially dependent on these frequencies. Higher precision, laboratory measurements of the line frequencies of these and other dense gas tracers are urgently needed for future progress in infall research.

This survey would have been impossible without the dedicated support of the staff of Haystack Observatory, among whom we would like to give special thanks to John Ball, Phil Shute, Joseph Carter, Kevin Duvet, and Mike Titus. Especially we would like to thank Phil Shute for his considerable help through our 2 years of observations. We are also grateful to the Director, Joe Salah for his outstanding support of our program. We wish to thank Tyler Bourke for his help in the beginning of the survey and Diego Mardones for many useful discussions. C.W.L acknowledges the financial support by 1996 Overseas Postdoctoral Support Program of

Korea Research Foundation and partial support by Star Project Grant 98-2-400-01 of the Korea Astronomy Observatory. C.W.L also thanks the Harvard-Smithsonian Center for Astrophysics for support while working on this project. M.T acknowledges partial support from Spanish DGES grant PB96-104. This research was supported by NASA Origins of Solar System Program, Grant NAGW-3401.

### A. Selection of the Line Frequencies

Tafalla et al. (1998) discussed how the narrow lines observed towards the starless core L1544 could be used to check the consistency of the different sets of recommended line frequencies, and they proposed a set of values that seemed self-consistent when checked against observations of that core. The results of the L1544 analysis from Tafalla et al. (1998) were independent from the exact frequency values, so the frequency discussion was stated without further illustration. The work presented in the present paper, however, depends sensitively on the exact frequencies adopted for the lines (see Fig. 6), so it seems necessary to provide further details on how the frequency comparison was done.

To minimize systematic errors, it is convenient to use a set of observations done with the same telescope using a similar spatial resolution and sampling, so for our discussion we will use only FCRAO QUARRY data (a similar analysis using IRAM 30m data leads to the same conclusions). To increase the S/N of the spectra, necessary for the weak lines, we average the spectra over a FCRAO QUARRY footprint ( $\sim 5' \times 4.2'$  area), and show the results in Fig. 9. The first panel of this figure shows the result of using the frequencies recommended by Lovas (1992) [109782.160 MHz for  $C^{18}O$  ( $1-0$ ), 96412.982 MHz for  $C^{34}S$  ( $2-1$ ), and 97980.968 MHz for  $CS$  ( $2-1$ )]. As Fig. 9-(a) clearly shows, unless the velocity shifts between the different tracers are real, which is very unlikely given their similar line widths and spatial distributions, the frequencies are inconsistent.

Next (Fig. 9-(b)), we use the set of frequencies recommended by Lovas & Krupenie (1974), one of the original references for the  $CS$  and  $CO$  frequencies in Lovas (1992) catalogue. These values are 109782.182 MHz for  $C^{18}O$  ( $1-0$ ), 96412.953 MHz for  $C^{34}S$  ( $2-1$ ), and 97981.007 MHz for  $CS$  ( $2-1$ ) and produce an excellent agreement between the  $C^{18}O$  and  $C^{34}S$  spectra. However, there is still a clear discrepancy with  $CS$  ( $2-1$ ), which seems slightly displaced to the red.

To correct the  $CS$  discrepancy, Tafalla et al. (1998) chose 97980.950 MHz for the  $CS$  ( $2-1$ ) frequency, which is the value that Pickett et al. (1998) recommend, and as shown in Fig. 9-(c), produces a good agreement with the other lines.

As a further check, Fig. 9-(d) presents a comparison of the  $C^{18}O$  ( $1-0$ ) and  $C^{34}S$  ( $2-1$ ) lines with the isolated  $N_2H^+$  ( $1-0$ ) component ( $F_1F = 01-12$ ), whose frequency was determined with high accuracy by Caselli et al. (1995) (93176.2651 MHz, a value we have adopted for this work). As the figure shows, there is an excellent agreement between the three lines, reinforcing our belief that the frequency selection is correct.

One possible point of concern is the fact that L1544 shows infall asymmetry, and that this may in some way bias the adopted frequency to be higher than the real value. If this was the case, however, one would expect that the bias would produce in non infalling objects an opposite signature, and would give rise to a false overabundance of outward motions in a statistical study like ours. As shown in Fig. 6, the opposite is the case, making us to believe that the infall nature of L1544 is immaterial for the frequency determination.

Finally, we examine the consistency of our frequencies by comparing the  $\text{C}^{18}\text{O}$  (1–0) peak velocities with those of  $\text{N}_2\text{H}^+$  (1–0) for 16 sources whose  $\text{C}^{18}\text{O}$  (1–0) spectra are Gaussian and have good S/N. The difference ( $V_{\text{N}_2\text{H}^+} - V_{\text{C}^{18}\text{O}}$ ) is negligible (about  $-0.02 \pm 0.057 \text{ km s}^{-1}$ ), again in agreement with our expectations.

In summary, the frequencies recommended here [97980.950 MHz for the CS (2–1), 93176.2651 MHz for the  $\text{N}_2\text{H}^+$  (1–0) isolated component, and 109782.182 MHz for the  $\text{C}^{18}\text{O}$  (1–0)] give the best agreement among the lines observed toward L1544. We note, however, that our comparison is a relative one, so it is possible the the absolute frequency values may be slightly off. If this is the case, all frequencies would have to be changed accordingly, and the results in this paper would not be affected.

## REFERENCES

- Adelson, L.M. & Leung, C.M. 1988, MNRAS, 235, 349
- Ball, J.A., Carter, J.C., Shute, P.A., Philips, A.E.E., Whalen, R.D. & Johnson, H.N. 1993, in An Introduction to the Haystack Observatory
- Barvainis, R., Ball, J.A., Ingalls, R.P. & Salah, E.J. 1993, PASP, 105, 1334
- Beichman, C. A., Myers P.C., Emerson, J. P., Harris, S., Mathieu, R., Benson, P.J., & Jennings, R. E., 1986 ApJ, 307, 337
- Benson P.J., & Myers P.C. 1989, ApJS, 71, 89
- Buisson, G., Desbats, L., Duvert, G., Forveille, T., Gras, R., Guilloteau, S., Lucas, R., & Valiron, P. 1994, CLASS Manual (Grenoble: IRAM)
- Ciolek, G.E., & Mouschovias, T.Ch. 1995, ApJ, 454, 194
- Caselli, P., Myers, P.C., & Thaddeus, P. 1995, ApJ, 455, L77
- Gregersen, E.M., Evans, N. J., Zhou, S., & Choi, M. 1997, ApJ, 484, 256
- Kuiper, B.H., Langer, W.D., & Velusamy, T. 1996, ApJ, 468, 761
- Lee, C. W., & Myers, P.C. 1999, ApJS, in press (LM99)
- Lee, C. W., Myers, P.C., & Park, Y.S. 1999, in preparation
- Lee, C. W., Myers, P.C., & Plume, L. 1999, in preparation
- Lee, C. W., Myers, P.C., & Tafalla, M. 1999, in preparation
- Leung, C.M., & Brown, R.B. 1977, ApJ, 214, L73
- Lovas, F.J. 1992, J. Phys. Chem. Ref. Data, 21, 181
- Lovas, F.J. & Krupenie, P.H. 1974, J. Phys. Chem. Ref. Data, 3, 25
- Mardones, D., Myers, P.C., Tafalla, M., Wilner, D.J., Bachiller, R., & Garay, G. 1997, ApJ, 489, 719 (M97)
- Mestel, L., & Spitzer, L. 1956, MNRAS, 116, 505
- Myers, P.C., Mardones, D., Tafalla, M., Williams, J.P., & Wilner, D.J. ApJ, 1996, 465, 133
- Myers, P.C., & Lazarian, A. 1998 ApJ, 507, L157
- Pickett, H. M., Poynter, R. L., Cohen, E.A., Delitsky, M.L., Pearson, J.C., & Muller, H.S.P. 1998, J. Quant. Spectrosc. & Rad. Transfer 60, 883
- Shu, F.H. 1977, ApJ, 214, 488
- Shu, F.H., Adams, F.C., & Lizano, S. 1987, ARAA, 25, 23
- Tafalla, M., Mardones, D., Myers, P.C., Caselli, P., Bachiller, R., & Benson, P.J. 1998, ApJ, 504, 900
- Ward-Thompson, D., Scott, P.F., Hills, R.E., & André P. 1994, MNRAS, 268, 276

Ward-Thompson, D., Motte, F., & André, P. 1999, MNRAS, in press.

Willacy, K., Langer, W.D., & Velusamy, T. 1998, ApJ, 507, L171

Williams, J.P., Myers, P.C., Wilner, D.J., & Di Francesco, J. 1999, ApJ, 513, L61

Zhou, S., Evans, N.J., Kompe, C., & Walmsley, C. M. 1993, ApJ, 404, 232

Zhou, S. 1995, ApJ, 442, 685.

## FIGURE CAPTIONS

**Fig. 1.** — CS (2–1) and  $N_2H^+$  (1–0) (isolated component) line spectra for 43 sources with strong detections. The dashed line indicates the velocity of  $N_2H^+$  obtained from the Gaussian hfs fit of its 7 hyperfine components.

**Fig. 2.** — CS (2–1) and  $N_2H^+$  (1–0) (isolated component) spectra for 21 sources with weak or no detection of the  $N_2H^+$  isolated component, but for which the other  $N_2H^+$  components are bright enough to determine a velocity from the hyperfine structure fit.

**Fig. 3.** — Spectra for 9 selected sources with non Gaussian CS lines for which  $N_2H^+$  was not detected or not observed. For 5 sources,  $C^{18}O$  (1–0) spectra are also presented, and their central velocities are indicated by dashed lines.

**Fig. 4.** — Different types of CS spectra - (a) Two peaks, with blue peak brighter than red, (b) Blue peak and red shoulder, (c) Single symmetric peak, (d) Red peak and blue shoulder, (e) Two peaks, with red peak brighter than blue peak, and (f) Two peaks with similar brightness. Dashed lines are the Gaussian fit velocities of the optically thin  $N_2H^+$  (1–0) or  $C^{18}O$  (1–0).

**Fig. 5.** — Representative spectra of  $N_2H^+$  (1–0) - (a) bright line with no self absorption features in any of the hyperfine components, (b) isolated component barely detectable while the other hyperfine components are bright enough to determine the LSR velocity of the  $N_2H^+$  line, (c) symmetric isolated component but possible self absorption in the other hyperfine components, (d) self-absorption in the isolated component as well as in the other components. The dotted lines in each panel mark the velocities of the 7 components of the  $N_2H^+$  obtained from a Gaussian hyperfine structure (hfs) fit using CLASS (Buisson et al. 1994) with the velocity offsets determined by Caselli et al. (1995). A Gaussian hfs fit spectrum is overlaid on the observed spectrum of L204C-2 in panel (b).

**Fig. 6.** — Histograms of the normalized velocity difference ( $\delta V_{CS}$ ) between  $V_{CS(2-1)}$  and  $V_{N_2H^+(1-0)}$  for a sample of 67 objects with detection in both tracers. The left panel represents the distribution using our preferred set of frequencies (see Appendix for a full discussion on frequencies), and shows a significant excess of blue shifted sources ( $\delta V < 0$ ). The right panel shows the histogram we would get if the CS (2–1) frequency from Lovas (1992) is used and the  $N_2H^+$  (1–0) frequency is not changed.

**Fig. 7.** — Distribution of  $\delta V_{CS}$  as a function of total optical depth ( $\tau_{N_2H^+}$ ) of the  $N_2H^+$  (1–0) line. The  $1\sigma$  error bars drawn in the Figure are the means of the errors in  $\tau_{N_2H^+}$  and  $\delta V_{CS}$ , respectively.

**Fig. 8.** — Distribution of  $\delta V_{CS}$  as a function of total integrated intensity of the  $N_2H^+$  hyperfine components.

**Fig. 9.** — Comparison of  $\text{C}^{18}\text{O}$  ( $1 - 0$ ) (thick solid spectra),  $\text{CS}$  ( $2 - 1$ ) (thin solid spectra),  $\text{C}^{34}\text{S}$  ( $2 - 1$ ) (dotted spectra), and  $\text{N}_2\text{H}^+$  ( $1 - 0$ ) (dashed spectrum) towards L1544 (average over a FCRAO QUARRY footprint, see text). (a) Lovas (1992) frequencies for  $\text{CS}$  ( $2 - 1$ ),  $\text{C}^{34}\text{S}$  ( $2 - 1$ ), and  $\text{C}^{18}\text{O}$  ( $1 - 0$ ), (b) Lovas & Krupenie (1974) frequencies for  $\text{CS}$  ( $2 - 1$ ),  $\text{C}^{34}\text{S}$  ( $2 - 1$ ), and  $\text{C}^{18}\text{O}$  ( $1 - 0$ ), (c) Pickett et al. (1998) frequency  $\text{CS}$  ( $2 - 1$ ), and Lovas & Krupenie (1974) frequencies for  $\text{C}^{34}\text{S}$  ( $2 - 1$ ) and  $\text{C}^{18}\text{O}$  ( $1 - 0$ ), and (d) Caselli et al. (1995) frequency for  $\text{N}_2\text{H}^+$  ( $1 - 0$ ), Lovas & Krupenie (1974) frequencies for  $\text{C}^{18}\text{O}$  ( $1 - 0$ ) and  $\text{C}^{34}\text{S}$  ( $2 - 1$ ). The intensities of all the spectral lines are scaled up to be the same as the intensity of  $\text{C}^{18}\text{O}$ .



TABLE 1  
 STARLESS CORES OBSERVED IN CS (2-1), N<sub>2</sub>H<sup>+</sup> (1-0), AND C<sup>18</sup>O (1-0) <sup>a</sup>

| Source <sup>b</sup> | RA 1950<br>h m s | DEC 1950<br>° ' " | V <sub>CS</sub><br>km s <sup>-1</sup> | V <sub>N<sub>2</sub>H<sup>+</sup></sub><br>km s <sup>-1</sup> | V <sub>C<sup>18</sup>O</sub><br>km s <sup>-1</sup> |
|---------------------|------------------|-------------------|---------------------------------------|---|--|
| L1275               | 00 00 48.2       | 67 02 05          | -6.73±0.04                            | ND  | ...  |
| L1291               | 00 39 47.6       | 61 44 35          | 5.56±0.02                             | ND  | ...  |
| L1358-2             | 02 51 55.1       | 69 12 03          | -3.24±0.03                            | ND  | ...  |
| B5-2                | 03 45 15.5       | 32 45 07          | 9.81±0.04                             | ...   | ...  |
| L1495-2             | 04 14 51.2       | 28 30 40          | 8.06±0.03                             | ...   | ...  |
| L1506B              | 04 16 03.9       | 25 13 01          | 7.53±0.01                             | ND  | ...  |
| L1521B*             | 04 21 08.5       | 26 30 00          | 6.57±0.02                             | ND  | 6.48 ±0.01   |
| B213-7              | 04 22 12.8       | 26 26 10          | 6.42±0.03                             | ND  | ...  |
| L1400A-2            | 04 25 05.8       | 54 39 40          | 3.52±0.05                             | ND  | ...  |
| L1400F              | 04 25 37.0       | 54 07 23          | 1.66±0.08                             | ND  | 1.87 ±0.11   |
| L1521E              | 04 26 09.6       | 26 07 46          | 6.86±0.01                             | ND  | 6.86 ±0.01   |
| L1445               | 04 28 28.8       | 46 31 01          | 5.12±0.01                             | ND  | ...  |
| L1551S-1            | 04 28 53.9       | 18 01 14          | 6.71±0.02                             | ND  | ...  |
| B18-2               | 04 29 34.4       | 24 45 46          | 6.17±0.02                             | ...   | ...  |
| L1536-2             | 04 29 50.7       | 22 53 29          | ND                                    | ...   | ...  |
| TMC2-3              | 04 29 57.6       | 24 11 26          | 6.52±0.02                             | ND  | ...  |
| L1536-3             | 04 30 01.7       | 22 47 41          | 5.69±0.10                             | ...   | ...  |
| L1536-1             | 04 30 19.7       | 22 36 51          | 5.60±0.03                             | ...   | ...  |
| L1527A-2            | 04 35 54.0       | 26 15 38          | 5.76±0.06                             | ...   | ...  |
| L1507-2             | 04 39 53.4       | 29 38 25          | 6.26±0.01                             | ND  | ...  |
| L1426               | 04 43 17.5       | 52 59 03          | 4.09±0.04                             | ND  | ...  |
| L1517C-1**          | 04 51 35.9       | 30 30 50          | 5.45±0.01                             | ND  | 5.55 ±0.01   |
| L1517B-2            | 04 51 56.4       | 30 38 03          | 5.82±0.02                             | ND  | ...  |
| L1544-3             | 05 00 38.6       | 25 14 18          | 7.12±0.05                             | ...   | ...  |
| L1544-2             | 05 01 39.1       | 25 07 30          | 6.87±0.01                             | ND  | ...  |
| L1523               | 05 03 02.7       | 31 37 30          | 7.03±0.02                             | ...   | ...  |
| CB28                | 05 03 51.2       | -04 00 20         | 8.74±0.01                             | ND  | ...  |
| L1621-2             | 05 52 49.8       | 02 08 07          | 1.25±0.02                             | ND  | ...  |
| L1621-1             | 05 53 23.0       | 02 17 39          | 1.82±0.01                             | ...   | ...  |
| CB37                | 05 57 29.9       | 31 39 25          | 0.92±0.09                             | ...   | ...  |
| L1574               | 06 05 08.9       | 18 28 41          | 0.22±0.03                             | ...   | ...  |
| L1633               | 06 22 04.0       | 03 23 53          | 8.93±0.03                             | ND  | ...  |
| L1696B-2            | 16 26 09.4       | -24 06 43         | 3.18±0.02                             | ND  | ...  |
| L1704-2             | 16 28 09.3       | -23 34 29         | ...                                   | ND  | ...  |
| L1757-1             | 16 29 01.9       | -19 35 24         | 1.45±0.07                             | ...   | ...  |
| L1757-2             | 16 29 08.5       | -19 42 05         | ...                                   | ND  | ...  |
| L1689A              | 16 29 10.4       | -24 56 03         | ...                                   | ND  | ...  |
| L1709C-2            | 16 30 18.0       | -23 40 01         | ...                                   | ND  | ...  |
| L1709C              | 16 30 52.6       | -23 33 54         | 2.50±0.02                             | ND  | 2.57 ±0.02   |
| L121                | 16 36 41.0       | -13 59 21         | ND                                    | ...   | ...  |
| L1782-1             | 16 39 13.2       | -19 36 22         | 5.34±0.01                             | ...   | ...  |
| L158-2              | 16 44 02.6       | -13 53 26         | 3.87±0.05                             | ...   | ...  |
| L204C-1             | 16 44 30.6       | -12 14 14         | 4.20±0.02                             | ND  | ...  |
| L204C**             | 16 44 44.2       | -12 04 27         | 4.89±0.02                             | ND  | 4.13 ±0.01   |
| L191-1              | 16 44 59.6       | -12 34 43         | 3.91±0.02                             | ND  | ...  |
| L204B*              | 16 45 00.0       | -11 50 00         | 3.74±0.02                             | ND  | 3.70 ±0.01   |
| L204F**             | 16 45 01.3       | -11 50 50         | 3.79±0.02                             | ND  | 3.69 ±0.01   |
| L191-2              | 16 45 01.7       | -12 47 38         | 3.32±0.03                             | ...   | ...  |
| L255-2              | 16 45 30.9       | -09 58 47         | 3.51±0.04                             | ...   | ...  |
| L234E3              | 16 45 49.0       | -10 45 45         | 3.02±0.02                             | ND  | ...  |
| L63-4               | 16 46 49.1       | -17 49 47         | ND                                    | ...   | ...  |
| L63-3               | 16 47 10.5       | -17 53 51         | 5.69±0.10                             | ...   | ...  |
| L63-2               | 16 47 11.2       | -17 59 52         | 5.86±0.03                             | ND  | ...  |
| L141-2              | 16 48 03.1       | -15 16 54         | 3.62±0.02                             | ND  | ...  |
| L141-3              | 16 48 11.1       | -15 12 04         | 3.33±0.01                             | ND  | ...  |
| L129-2              | 16 51 57.6       | -16 21 42         | 5.79±0.02                             | ND  | ...  |
| L129-1              | 16 52 18.4       | -16 17 10         | 5.80±0.01                             | ND  | ...  |
| L122                | 16 52 22.4       | -16 42 03         | 6.18±0.01                             | ND  | ...  |
| CB68-2              | 16 54 00.0       | -15 57 11         | 4.99±0.02                             | ...   | ...  |
| L111                | 17 11 59.5       | -20 27 15         | ...                                   | ND  | ...  |
| L173                | 17 14 25.4       | -18 27 45         | ...                                   | ND  | ...  |
| L392                | 17 50 24.3       | -08 26 43         | ND                                    | ...   | ...  |
| L421                | 17 51 38.9       | -05 46 02         | 7.43±0.06                             | ...   | ...  |
| L460                | 17 55 24.4       | -03 46 24         | 6.82±0.02                             | ...   | ...  |
| L432                | 17 55 46.2       | -05 43 45         | 7.37±0.03                             | ND  | ...  |
| L468                | 17 57 38.0       | -03 30 03         | ND                                    | ...   | ...  |
| L422                | 18 09 20.0       | -08 04 46         | 3.65±0.04                             | ...   | ...  |
| L438                | 18 11 23.8       | -07 10 01         | 4.61±0.03                             | ...   | ...  |
| L436                | 18 11 53.0       | -07 23 49         | 4.68±0.03                             | ...   | ...  |
| L490                | 18 12 42.5       | -03 47 11         | 8.09±0.04                             | ...   | ...  |
| CB130-2             | 18 13 37.5       | -02 24 28         | 7.17±0.04                             | ND  | ...  |

TABLE 1—*Continued*

| Source <sup>b</sup> | RA 1950<br>h m s | DEC 1950<br>° ′ ″ | $V_{CS}$<br>km s <sup>-1</sup> | $V_{N_2H^+}$<br>km s <sup>-1</sup> | $V_{C^{18}O}$<br>km s <sup>-1</sup> |
|---------------------|------------------|-------------------|--------------------------------|------------------------------------|-------------------------------------|
| CB130-1             | 18 13 39.2       | -02 34 06         | ...                            | 7.65 ±0.028                        | ...                                 |
| CB130-3             | 18 13 41.5       | -02 17 47         | ...                            | 7.20 ±0.005                        | ...                                 |
| L328                | 18 14 05.3       | -18 03 10         | ...                            | 6.62 ±0.006                        | ...                                 |
| L429-2              | 18 14 32.8       | -08 19 02         | ...                            | ND                                 | ...                                 |
| L429-3              | 18 14 50.8       | -08 15 17         | ND                             | ...                                | ...                                 |
| L429-4              | 18 15 05.2       | -08 10 07         | ND                             | ...                                | ...                                 |
| L483-2              | 18 15 26.5       | -04 34 24         | ...                            | ND                                 | ...                                 |
| L466                | 18 17 04.5       | -06 07 36         | ...                            | ND                                 | ...                                 |
| L539-1              | 18 21 05.7       | -01 04 28         | ND                             | ...                                | ...                                 |
| L539-2              | 18 21 46.5       | -01 01 05         | 8.71±0.04                      | ...                                | ...                                 |
| L588-2              | 18 32 47.0       | -00 43 53         | 10.79±0.03                     | ...                                | ...                                 |
| L588-3              | 18 33 18.2       | -00 33 25         | ...                            | ND                                 | ...                                 |
| L495                | 18 36 32.6       | -06 41 06         | ...                            | ND                                 | ...                                 |
| L648-2              | 18 37 20.5       | 12 47 05          | 5.22±0.02                      | ND                                 | ...                                 |
| L530D-3             | 18 46 48.0       | -05 03 12         | 3.32±0.05                      | ...                                | ...                                 |
| L530D-1             | 18 47 20.5       | -04 51 55         | 3.48±0.07                      | ...                                | ...                                 |
| L530D-2             | 18 47 22.1       | -04 43 25         | ND                             | ...                                | ...                                 |
| L549                | 18 59 29.8       | -05 23 08         | ...                            | ND                                 | ...                                 |
| L709                | 19 11 37.3       | 16 22 01          | 5.91±0.02                      | ...                                | ...                                 |
| L763                | 19 16 45.9       | 23 21 45          | 11.18±0.06                     | ...                                | ...                                 |
| L730-1              | 19 17 40.5       | 19 51 09          | 8.24±0.03                      | ND                                 | ...                                 |
| L655                | 19 17 47.2       | 08 23 33          | 7.97±0.02                      | ...                                | ...                                 |
| L730-2              | 19 18 12.4       | 19 56 53          | 8.10±0.05                      | ...                                | ...                                 |
| L673-4              | 19 18 47.1       | 11 20 11          | 6.67±0.02                      | ND                                 | ...                                 |
| L673-5              | 19 18 53.6       | 11 27 16          | 6.77±0.04                      | ...                                | ...                                 |
| L673-6              | 19 19 06.2       | 11 30 39          | 6.67±0.04                      | ...                                | ...                                 |
| L673-8              | 19 19 15.4       | 11 12 20          | 7.09±0.04                      | ...                                | ...                                 |
| L673-10             | 19 19 32.5       | 11 14 21          | ...                            | ND                                 | ...                                 |
| L677-1              | 19 19 50.0       | 11 27 19          | ...                            | ND                                 | ...                                 |
| L677-2              | 19 19 52.8       | 11 30 46          | ...                            | ND                                 | ...                                 |
| L677-3              | 19 20 07.5       | 11 15 30          | ...                            | ND                                 | ...                                 |
| L677-4              | 19 20 22.8       | 11 16 57          | ...                            | ND                                 | ...                                 |
| L774-1              | 19 20 44.9       | 23 19 21          | 11.27±0.05                     | ...                                | ...                                 |
| L774-2              | 19 21 34.2       | 23 17 18          | ND                             | ...                                | ...                                 |
| L769-4              | 19 22 47.0       | 22 56 36          | ...                            | ND                                 | ...                                 |
| L698                | 19 23 37.5       | 12 59 48          | ...                            | ND                                 | ...                                 |
| L778-2              | 19 25 22.0       | 23 39 45          | ND                             | ND                                 | ...                                 |
| L778-3              | 19 25 54.8       | 23 37 14          | ND                             | ND                                 | ...                                 |
| L778-4              | 19 26 02.6       | 23 51 12          | ...                            | ND                                 | ...                                 |
| L694-1              | 19 38 06.5       | 10 43 33          | ...                            | ND                                 | ...                                 |
| L694-3              | 19 38 54.0       | 10 43 42          | ND                             | ...                                | ...                                 |
| L758                | 19 44 12.8       | 18 54 47          | ND                             | ...                                | ...                                 |
| L880                | 20 11 49.4       | 40 03 59          | ND                             | ...                                | ...                                 |
| L917                | 20 38 16.6       | 43 58 37          | 3.10±0.03                      | ...                                | ...                                 |
| L918                | 20 39 02.5       | 43 54 57          | 1.35±0.06                      | ...                                | ...                                 |
| L1147**             | 20 39 59.9       | 67 10 10          | 2.78±0.02                      | ND                                 | ...                                 |
| L896-2              | 20 40 26.3       | 39 36 01          | -1.95±0.10                     | ...                                | ...                                 |
| L896-1              | 20 40 28.8       | 39 42 22          | ND                             | ...                                | ...                                 |
| L1155D              | 20 43 17.9       | 67 26 12          | 2.53±0.02                      | ND                                 | ...                                 |
| L915                | 20 43 23.0       | 42 58 52          | -0.80±0.10                     | ...                                | ...                                 |
| L1082C-3            | 20 47 59.4       | 60 00 36          | -2.90±0.01                     | ND                                 | ...                                 |
| L981-2              | 20 58 38.0       | 50 05 09          | -0.24±0.05                     | ...                                | ...                                 |
| L971-1              | 20 59 08.1       | 49 17 21          | -0.41±0.07                     | ...                                | ...                                 |
| L971-2              | 20 59 15.8       | 49 07 34          | -0.48±0.04                     | ...                                | ...                                 |
| L1172A-2            | 21 01 10.8       | 67 34 34          | 2.82±0.03                      | ND                                 | ...                                 |
| L1172A-3            | 21 03 22.6       | 67 29 06          | 2.89±0.02                      | ND                                 | ...                                 |
| L1125-1             | 21 13 07.8       | 61 32 26          | 0.99±0.05                      | ...                                | ...                                 |
| L944-1              | 21 14 42.6       | 42 59 47          | ND                             | ...                                | ...                                 |
| L944-3              | 21 16 59.7       | 43 12 57          | ND                             | ...                                | ...                                 |
| L952-1              | 21 18 08.5       | 43 18 07          | ND                             | ...                                | ...                                 |
| L952-2              | 21 18 25.0       | 43 19 59          | ND                             | ...                                | ...                                 |
| L953                | 21 19 31.8       | 43 08 46          | ND                             | ...                                | ...                                 |
| L1021               | 21 20 34.8       | 50 50 17          | 2.78±0.10                      | ...                                | ...                                 |
| L1014-1             | 21 22 20.6       | 49 56 52          | 4.02±0.07                      | ...                                | ...                                 |
| L1014-2             | 21 22 22.2       | 49 46 10          | ND                             | ...                                | ...                                 |
| L1099-1             | 21 34 27.9       | 57 13 18          | -8.27±0.02                     | ND                                 | ...                                 |
| L1111               | 21 38 53.6       | 57 34 24          | ND                             | ...                                | ...                                 |
| L1103-1             | 21 39 49.8       | 56 23 32          | ND                             | ...                                | ...                                 |
| L1103-2             | 21 40 41.0       | 56 30 07          | 4.44±0.03                      | ...                                | ...                                 |
| L1113-2             | 21 42 45.4       | 56 57 33          | ND                             | ...                                | ...                                 |
| L1113-1             | 21 42 56.7       | 56 56 25          | ND                             | ...                                | ...                                 |

TABLE 1—*Continued*

| Source <sup>b</sup> | RA 1950<br><i>h m s</i> | DEC 1950<br><i>° ' "</i> | $V_{CS}$<br>km s <sup>-1</sup> | $V_{N_2H^+}$<br>km s <sup>-1</sup> | $V_{C^{18}O}$<br>km s <sup>-1</sup> |
|---------------------|-------------------------|--------------------------|--------------------------------|------------------------------------|-------------------------------------|
| L1139-1             | 21 53 14.7              | 58 20 59                 | -2.32±0.04                     | ...                                | ...                                 |
| L1166               | 22 03 46.7              | 59 19 07                 | ND                             | ...                                | ...                                 |
| L1235B**            | 22 15 04.1              | 73 07 30                 | -4.19±0.02                     | ND                                 | -4.00 ±0.01                         |
| L1195               | 22 25 18.1              | 61 01 06                 | ND                             | ...                                | ...                                 |
| L1185               | 22 27 27.6              | 58 53 30                 | ND                             | ...                                | ...                                 |
| L1229               | 23 12 16.8              | 61 46 47                 | ND                             | ...                                | ...                                 |
| L1263               | 23 54 54.2              | 64 32 27                 | ND                             | ...                                | ...                                 |
| L1271-1             | 23 57 32.6              | 67 05 07                 | ND                             | ...                                | ...                                 |
| L1271-3             | 23 59 15.3              | 67 01 48                 | ND                             | ...                                | ...                                 |

<sup>a</sup>This lists velocity information for sources which have been observed in CS (2-1) and N<sub>2</sub>H<sup>+</sup> (1-0), but detected in only one line or not detected in either line. The C<sup>18</sup>O (1-0) line data which have been complementarily obtained for several sources are also listed in this table. 'ND' and '...' mean 'No Detection' and 'Not Observed', respectively.

<sup>b</sup>Position of the source marked with '\*' behind the source name is from Benson & Myers (1989), '\*\*' from the FCRAO map (Lee, Myers, & Tafalla 1999), and none from LM99 catalogue.



TABLE 3  
SPECTRAL PROPERTIES OF STARLESS CORES

| Source   | $\delta V_{CS}^a$ | $T_b/T_r^b$<br>in CS | $T_b/T_r^b$ in<br>$N_2H^+$ main component | Source   | $\delta V_{CS}^a$ | $T_b/T_r^b$<br>in CS | $T_b/T_r^b$ in<br>$N_2H^+$ main component |
|----------|-------------------|----------------------|---|----------|-------------------|----------------------|---|
| L1333    | N                 | ...                  | ...                                       | L1696A   | B                 | ...                  | ...                                       |
| L1355    | B                 | B (1.53±0.17)        | ...                                       | L1696B   | N                 | ...                  | ...                                       |
| B5-1     | N                 | ...                  | ...                                       | L1704-1  | N                 | ...                  | ...                                       |
| L1498    | B                 | B (2.03±0.17)        | ...                                       | L1709B-2 | N                 | ...                  | ...                                       |
| L1399-1  | N                 | ...                  | ...                                       | L1689B   | N                 | B (1.46±0.22)        | ...                                       |
| L1521B-2 | N                 | ...                  | ...                                       | L1782-2  | N                 | ...                  | ...                                       |
| L1399-2  | N                 | ...                  | ...                                       | L158     | B                 | B (1.37±0.19)        | ...                                       |
| L1521B   | ...               | N (0.91±0.19)        | ...                                       | L204C    | ...               | N (0.84±0.20)        | ...                                       |
| L1399-3  | N                 | ...                  | ...                                       | L204C-2  | N                 | ...                  | ...                                       |
| B217-2   | N                 | ...                  | ...                                       | L234E-1  | B                 | ...                  | ...                                       |
| L1521F   | B                 | B (1.26±0.16)        | B(1.13±0.12)                              | L63      | R                 | ...                  | ...                                       |
| L1521-2  | N                 | ...                  | ...                                       | L462-1   | N                 | ...                  | ...                                       |
| L1400K   | B                 | ...                  | ...                                       | L462-2   | N                 | ...                  | ...                                       |
| L1400A   | N                 | ...                  | ...                                       | L492     | B                 | ...                  | N(1.15±0.16)                              |
| L1524-4  | B                 | ...                  | ...                                       | L429-1   | ...               | N (0.79±0.24)        | N(1.43±0.45)                              |
| L1551S-2 | N                 | ...                  | N(0.94±0.19)                              | L648-1   | N                 | ...                  | ...                                       |
| L1445    | ...               | B (1.29±0.20)        | ...                                       | L673-7   | N                 | ...                  | ...                                       |
| TMC2     | B                 | ...                  | ...                                       | L694-2   | B                 | B (1.70±0.20)        | ...                                       |
| B18-3    | N                 | B (1.24±0.20)        | ...                                       | L1041-2  | N                 | ...                  | ...                                       |
| B18-5    | N                 | ...                  | ...                                       | L922-1   | N                 | ...                  | ...                                       |
| CB22     | N                 | ...                  | ...                                       | L922-2   | B                 | ...                  | ...                                       |
| TMC1     | B                 | ...                  | ...                                       | L1049-1  | R                 | ...                  | ...                                       |
| L1507A-1 | N                 | ...                  | ...                                       | L1049-2  | N                 | ...                  | ...                                       |
| CB23     | B                 | ...                  | ...                                       | L1148    | N                 | ...                  | ...                                       |
| L1517C-1 | ...               | N (0.96±0.16)        | ...                                       | L1155C-2 | B                 | ...                  | ...                                       |
| L1517A   | N                 | ...                  | ...                                       | L1155C-1 | B                 | B (2.59±0.19)        | ...                                       |
| L1517B   | N                 | ...                  | ...                                       | L1155E   | N                 | ...                  | ...                                       |
| L1512-1  | N                 | ...                  | ...                                       | L1082C-2 | N                 | ...                  | ...                                       |
| L1544    | ...               | N (1.13±0.20)        | B(1.47±0.22)                              | L981-1   | N                 | ...                  | ...                                       |
| L1552    | N                 | ...                  | ...                                       | L1063    | N                 | ...                  | ...                                       |
| L1582A   | N                 | ...                  | ...                                       | L944-2   | N                 | ...                  | ...                                       |
| L1622A-2 | B                 | ...                  | ...                                       | L1251A-2 | N                 | ...                  | ...                                       |
| L1622A-1 | N                 | ...                  | ...                                       | L1197    | N                 | ...                  | ...                                       |
| CB45     | N                 | ...                  | ...                                       | L1251    | N                 | ...                  | ...                                       |
| L134A    | N                 | ...                  | ...                                       | L1234    | B                 | ...                  | ...                                       |
| L183B    | B                 | B (1.46±0.22)        | ...                                       | CB246-2  | R                 | R (0.83±0.12)        | ...                                       |
| L183     | B                 | ...                  | ...                                       |          |                   |                      | ...                                       |

<sup>a</sup>‘B’ and ‘R’ are given for sources with blue or red asymmetry of  $|\delta V_{CS}| \geq 5 \sigma_{\delta V_{CS}}$ , and ‘N’ for sources with neither asymmetry of  $|\delta V_{CS}| \leq 5 \sigma_{\delta V_{CS}}$

<sup>b</sup>The ratio of intensities of blue component ( $T_b$ ) and red component ( $T_r$ ) from the Gaussian fit after masking the other fainter component. ‘B’, ‘N’, or ‘R’ are designated for the sources of  $T_b/T_r > 1 + \sigma$ ,  $1 - \sigma \leq T_b/T_r \leq 1 + \sigma$ , or  $T_b/T_r < 1 - \sigma$ , respectively.