

Discovery of a ~ 1 Hz quasi-periodic oscillation in the low-mass X-ray binary 4U 1746–37

Peter G. Jonker¹, Michiel van der Klis¹, Jeroen Homan¹, Rudy Wijnands^{1,2}, Jan van
Paradijs^{1,3}, Mariano Méndez^{1,4}, Erik Kuulkers^{5,6}, Eric C. Ford¹

Subject headings: accretion, accretion disks — stars: individual (4U 1746–37)
— stars: neutron — X-rays: stars

Received _____; accepted _____

¹Astronomical Institute “Anton Pannekoek”, University of Amsterdam, and Center for High-Energy Astrophysics, Kruislaan 403, 1098 SJ Amsterdam; peterj@astro.uva.nl, michiel@astro.uva.nl, homan@astro.uva.nl, rudy@astro.uva.nl, jvp@astro.uva.nl, mariano@astro.uva.nl, ecford@astro.uva.nl

²MIT, Center for Space Research, Cambridge, MA 02139

³University of Alabama, Huntsville

⁴Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque S/N, 1900 La Plata, Argentina

⁵Space Research Organization Netherlands, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands; e.kuulkers@sron.nl

⁶Astronomical Institute, Utrecht University, P.O. Box 80000, 3507 TA Utrecht, The Netherlands

ABSTRACT

We have discovered a ~ 1 Hz quasi-periodic oscillation (QPO) in the persistent X-ray emission and during type I X-ray bursts of the globular cluster source, dipper and low-mass X-ray binary (LMXB) 4U 1746–37. The QPO properties resemble those of QPOs found recently in the LMXB dippers 4U 1323–62, and EXO 0748–676, which makes 4U 1746–37 the third source known to exhibit this type of QPOs. We present evidence for X-ray spectral changes in this source similar to those observed in LMXBs referred to as atoll sources. We detect two states, a low intensity and spectrally hard state, and a higher intensity and spectrally soft state. This may explain the different spectral characteristics reported for 4U 1746–37 earlier. The high intensity state resembles the banana branch state of atoll sources. The QPOs are only seen in the low intensity state, and are absent when the source is in the banana branch. This strongly suggests that either the accretion disk or an extended central source change shape between the low intensity state and the banana branch.

Twelve bursts were detected, of which 5 took place while the source was on the banana branch and 7 when the source was in the low intensity state. The bursts occurring on the banana branch had an e-folding time ~ 3 times longer than those which occurred in the low intensity state.

Whereas previously detected dips showed only a decrease in count rate of $\sim 15\%$, we found in one observation a dip in which the count rate dropped from ~ 200 counts per second to ~ 20 counts per second. This dip lasted only ~ 250 seconds, during which clear spectral hardening occurred. This is the first time strong evidence for spectral changes during a dip are reported for this source.

1. Introduction

The low-mass X-ray binary (LMXB) 4U 1746–37 shows type I X-ray bursts (Li & Clark 1977) and dips (Parmar et al. 1989) which recur on the 5.7 hours (Sansom et al. 1993) orbital period. The source is located in the globular cluster NGC 6441. Deutsch et al. (1998) reported the probable detection of the optical counterpart. The dips in dip sources such as 4U 1746–37 are thought to be caused by periodic obscuration of the central source by a structure formed in an interplay between the accretion stream from the companion towards the disk and the disk itself (White & Swank 1982; White, Nagase, & Parmar 1995; Frank, King, & Lasota 1987). This structure moves through the line of sight fully or partially obscuring the central source once per binary cycle. Often, the X-ray spectral properties change during the dips. Spectral hardening occurs in 4U 1323–62 (Parmar et al. 1989), and spectral softening in deep dips seen in 4U 1624–49 (Church & Balucińska-Church 1995). In 4U 1746–37 no energy dependence was found so far (Sansom et al. 1993). The type I X-ray bursts are observed to have peak luminosities around the Eddington luminosity for the estimated distance to the globular cluster (10.7 kpc; Djorgovski 1993). Combined with the fact that the ratio of the X-ray to optical luminosities $L_x/L_{\text{opt}} \sim 10^3$, this makes it likely that outside the dips the central source is viewed directly (Parmar et al. 1999).

Recently, in two dipping LMXBs, 4U 1323–62 (Jonker et al. 1999a) and EXO 0748–676 (Homan et al. 1999), quasi-periodic oscillations (QPOs) with a typical frequency of 1 Hz were discovered. In EXO 0748–676 these frequencies vary between 0.58–2.44 Hz on timescales ranging from several days to weeks (Homan et al. 1999). Observations with intervals of several days to weeks revealed similar frequency shifts in 4U 1323–62 (Jonker et al. 1999b). These QPOs are strong, with fractional rms amplitudes of $\sim 10\%$, which are the same in the persistent emission, during dips, and during type I X-ray bursts. The rms amplitude of the QPOs does not, or only weakly, depend on the photon energy, unlike

that of various other types of QPO observed from LMXBs (van der Klis 1995, 1998). From these properties, and the fact that the ~ 1 Hz QPOs are observed in high inclination systems, Jonker et al. (1999a) suggested that partial obscuration of the central source by a near-opaque or gray medium in or on the disk covering and uncovering the source quasi-periodically at the observed QPO frequency could explain the QPO properties. The most natural location for such a medium would be at the radius where the orbital frequency in the disk is equal to the QPO frequency, which corresponds to a radius of $\sim 2 \times 10^8$ cm for a $1.4M_{\odot}$ neutron star (Jonker et al. 1999a).

On the basis of the timing and spectral properties of LMXBs Hasinger & van der Klis (1989) defined the atoll and the Z sources. The atoll sources show at least two states, named after the associated structures in the X-ray color-color diagram (CD): the island and the banana branch. The timing and spectral properties of the sources in these states are different. In the island state strong band limited noise is present in the power spectra. When the source moves up the banana branch the fractional rms amplitude of this noise component decreases and simultaneously the rms amplitude of a very low frequency power law noise (VLFN) component increases. The Z sources have higher luminosities and trace out a Z track in the CD. The timing properties are correlated with the position along this track (for a complete overview of the timing and spectral properties of the atoll and Z sources, see van der Klis 1995).

In this Letter, we report the discovery of a ~ 1 Hz QPO in the source 4U 1746–37. We also show that the spectral characteristics and some of the timing properties of this source are reminiscent of those of an atoll source.

2. Observations and analysis

We analysed 9 observations of 4U 1746–37 obtained with the proportional counter array (PCA) on board the *RXTE* satellite (Bradt, Rothschild, & Swank 1993), in 1996 on October 25, 27, 31, and in 1998 on June 5, June 6, August 3, November 7, and November 22. The total amount of good data analysed was ~ 129 ksec. A log of the observations can be found in Table 1. Data were obtained over an energy range of 2–60 keV simultaneously with time resolutions of 0.125 s, 16 s, and 1 μ s in 1, 129, and 255 energy bins; the Standard 1, Standard 2, and GoodXenon data modes, respectively.

The average persistent emission count rates varied from 66 counts per second (observation 3) to nearly 800 counts per second (observation 9; see Table 1). All count rates reported are background subtracted and for five PCA detectors, unless otherwise stated. We observed shallow dips in the lightcurve showing a decrease in count rate of only $\sim 15\%$, and one deep dip with evidence for spectral hardening (see Fig. 1, and Section 3). During the last part of observation 9 a solar flare occurred; we excluded these data from our analysis.

Since the dips were often difficult to distinguish from the persistent emission (Sansom et al. 1993; Parmar et al. 1999), we divided the data into just two categories; non-burst (which combines persistent emission and dip data) and burst data.

For the selection of the burst data the start of a burst was characterized by the sharp increase in flux and simultaneous decrease in hardness. The burst e-folding time was determined by fitting an exponential to the burst decay in the 2–60 keV lightcurve of the Standard 1 data. The end of the burst was chosen to be three times the e-folding time after the onset (see also Jonker et al. 1999a).

Using the Good Xenon data, we calculated power spectra of non-burst data segments

Table 1. Log of the observations. The 5 detector average 2–60 keV persistent emission count rate of each observation is given in column 6.

Number	Observation ID	Date	Start time (UTC)	Amount of good data (ksec.)	Persistent emission (c/s)
1	10112-01-01-00	October 25 1996	00:10:55	~22	73
2	10112-01-01-01	October 27 1996	03:25:07	~17	73
3	10112-01-01-02	October 31 1996	08:26:19	~21	66
4	30701-11-01-00	June 5 1998	00:09:55	~6	365
5	30701-11-01-01	June 5 1998	14:09:03	~15	415
6	30701-11-01-02	June 6 1998	07:45:43	~3	414
7	30701-11-02-00	August 3 1998	03:15:03	~25	296
8	30701-11-03-01	November 7 1998	08:19:37	~7	728
9	30701-11-04-00	November 22 1998	11:18:33	~13	785

of 64 s length and of burst data segments of 16 s length in the energy bands 2–60 keV, 2–6.4 keV, and 6.4–60 keV, with a Nyquist frequency of 512 Hz. We also calculated power spectra of segments of 16 s length with a Nyquist frequency of 2048 Hz in the 2–60 keV band.

The power spectra within each energy band were added and averaged for each observation, separately for burst and non-burst data. To search for burst oscillations we also calculated power spectra of a length of 1 s with a Nyquist frequency of 2048 Hz.

We constructed a CD from all non-burst Standard 2 data, using four detectors (0, 1, 2, and 3). The hard color in this diagram is defined as the logarithm of the ratio between the count rates in the 9.7–16.0 keV band and the count rates in the 6.0–9.7 keV band, and the soft color as the logarithm of the ratio between the count rates in the 3.5–6.0 keV band and the count rates in the 2–3.5 keV band

3. Results

In the non-burst data of October 25 1996 (observation 1), we discovered a QPO in the 2–60 keV band with a frequency, FWHM, and rms amplitude of 1.04 ± 0.03 Hz, 0.36 ± 0.09 Hz, and $7.7\% \pm 0.7\%$ (6σ), respectively (see Fig. 2). In the observations of October 27 (observation 2) and October 31 (observation 3) 1996, we observed QPOs with similar frequencies, FWHMs and rms amplitudes, albeit at a lower significance level (see Table 2).

We measured the QPO properties in the 2–6.4 keV and 6.4–60 keV bands. Upper limits on the presence of the QPO were derived fixing the values of the FWHM and frequency obtained in the 2–60 keV energy band, except for observation 2. For this observation we used the values obtained in the 2–6.4 keV band, since the QPO was not detected at a significance higher than 3σ in the 2–60 keV band (see Table 2). Only for observation 1 the QPO was detected at a significance higher than 3σ in both the 2–6.4 keV and the 6.4–20

keV energy band. Within the errors the rms amplitude of the QPO in the two energy bands was the same (see Table 2).

The QPO was also detected when all 16 s segment power spectra (9 in total) of the decay of the three bursts of observation 1 were combined, at a similar fractional rms amplitude ($7.2\% \pm 0.8\%$). The averaged count rate was ~ 720 counts per second, i.e., about ten times that in the persistent emission. The frequency increased significantly with respect to the frequency of the QPO in the persistent emission to $1.39_{-0.10}^{+0.05}$ Hz. The FWHM of the QPO was consistent with being the same as in the persistent emission. In the bursts of observations 2 and 3 no significant QPO was detected, with upper limits on the rms amplitude of 7% to 8%.

We fitted a power law to the noise component evident in Fig. 2. Its rms amplitude integrated over 0.0156–1 Hz varied from $7.1\% \pm 0.3\%$ to $5.9\% \pm 0.4\%$ and to $9.3\% \pm 0.4\%$, from observation 1–3, while the power law index was consistent with being constant at 0.5 ± 0.3 . In the observations 4–9, the noise amplitude was in the range of 1.6%–5.7%. Here, the fractional rms amplitude increased with increasing intensities (see Table 1), except for observation 8 where it was only 1.6%. The power law indices ranged from 1.7 ± 0.4 to 2.1 ± 0.4 . The rms amplitude of this power law component is slightly higher in the 6.4–20 than in the 2–6.4 keV band. The 6.4–20/2–6.4 keV spectral hardness did not significantly depend on the presence or absence of the QPO.

Combining all non-burst data into a CD, we observed a pattern which resembles the atoll shape (Hasinger & van der Klis 1989), (Fig. 3). Two distinct parts can be distinguished in the CD, a hard (low intensity, upper part of the diagram) island-like state, and a softer (higher intensity, lower part of the diagram) banana branch. The ~ 1 Hz oscillations were only found when the source was in the upper (hard) part of the diagram (made up of observations 1–3). Upper limits on the presence of a QPO in the range of 0.5–2.5 Hz of

1%–2% were derived for the banana part of the diagram (made up of observations 4–9).

We observed a total of 12 type I bursts. One burst in the second observation and two bursts in the third observation showed secondary bursts, occurring several hundred seconds after the first burst and with a 3–4 times lower peak flux (see Lewin, van Paradijs, & Taam 1995). The burst e-folding times ranged from 7.5 to 11.3 seconds for bursts observed in observations 1–3, and from 7.5 to 14.5 for the secondary bursts. The burst e-folding time of 4 of the 5 bursts observed in observation 8 and 9 ranged from 27.7–30.1 seconds, while one burst in observation 9 which occurred ~ 600 seconds after a burst with a 2.3 higher peak flux had an e-folding time of only 3.0 seconds (the peak flux of this very short burst was similar to that of the bursts in the low intensity state). Upper limits on burst oscillations in the 100–1000 Hz frequency range during the peak and decay of the bursts adding the different bursts of typically $\sim 15\%$ were derived.

One of the dips observed in observation 7 was different from the others; it showed a simultaneous sharp increase in hard color of $\sim 28\%$, and drop in count rate from ~ 200 counts per second to ~ 21 counts per second. The dip was quite short (~ 250 s), whereas previous dips were shallow ($\sim 15\%$), and exhibited no clear spectral hardening (see Fig. 1). Due to the uncertainties in the orbital ephemeris, we could not relate the exact time of this deep dip to the dips in the other observations.

Upper limits on the presence of kHz QPOs in the frequency range of 100–1000 Hz were derived using a fixed FWHM of 50 Hz. The upper limits were in the range of 3.4%–6.5%.

4. Discussion

We discovered a QPO in 4U 1746–37 with a frequency of ~ 1 Hz, and an rms amplitude of $\sim 7.5\%$. The QPO was also observed during type I X-ray bursts where the count rate

was a factor of 10 higher, with similar fractional rms amplitudes. The QPO in 4U 1746–37 was only observed when the source intensity was low. The frequency of the QPO, the weak energy dependence of its rms amplitude, and the ratio Q between the QPO frequency and its FWHM are similar to the frequencies, the rms amplitude energy dependence, and Q -values (3–4) of the QPOs discovered recently in the persistent emission, dips, and during the type I X-ray bursts in the dippers and LMXBs 4U 1323–62 (Jonker et al. 1999a) and in EXO 0748–676 at low intensity (Homan et al. 1999). The similarities in QPO properties and the fact that 4U 1746–37 is a dipper suggest that the ~ 1 Hz QPO is a general property of high-inclination LMXBs at low intensities, and has a common origin in these systems.

We found for the first time that 4U 1746–37 showed spectral characteristics of an atoll source; the timing properties of observations 4–9 were consistent with banana branch behavior reported for atoll sources (van der Klis 1995). Although no band limited noise was detected, upper limits (1–512 Hz) in the low intensity (hard) state were 20%–28%, whereas they were 2.5%–5.5% when the source was on the apparent banana branch. So, with respect to the band limited noise the timing behavior of the source is consistent with that of an atoll source. Also the fact that on the banana branch the rms amplitude of the power law noise component increased as the source moved up in count rate is consistent with the behavior of the VLFN on the banana branch of an atoll source. However, the high (5.9%–7.1%) rms amplitude and low power law index (0.5) of the power law noise component when the source is at low intensities is not consistent with island-state VLFN. We note that this could be due to effects observable only at high inclinations, similar to those which cause the dips and the ~ 1 Hz QPO. In EXO 0748–676 strong power law noise (6%–11%) with indices 0.5–0.8 was also present. This noise had no or weak dependence on photon energy, just as the ~ 1 Hz QPOs (Homan et al. 1999), suggesting a similar origin for both phenomena. In the one observation of EXO 0748–676 where no QPO was found the rms amplitude of the VLFN was lower ($\sim 4\%$) and the power law index was ~ 1 . The

spectral changes we report in 4U 1746–37 related to changes in the position along the atoll, probably reflect the changing spectral characteristics reported before for this source (eg. see Parmar et al. 1999).

Homan et al. (1999) found that the QPO in EXO 0748–676 was present in all observations except one, where the persistent emission count rate was ~ 2.5 times higher than in the other observations. The fact that the QPO is not detected at higher source count rates (and inferred mass accretion rates) may be due to changes in the accretion geometry with \dot{M} . We expect that if we observe 4U 1323–62 at higher \dot{M} , the QPO may disappear as well. In 4U 1746–37 the QPO also disappeared when the count rate increased. The rms amplitude dropped by at least a factor of five when the source moved from the low intensity state to the banana branch. This can be accounted for if the size of the central source increased by a factor of 2–3 in radius, decreasing the modulated fraction of the X-rays below our threshold of 1%–2%. If this is the explanation the dip fraction should decrease, making the dips more shallow. Since in 4U 1746–37 the dips were shallow and the source count rates were low, we were not able to check this prediction for this source. Another possibility is that the structure in or on the disk responsible for the ~ 1 Hz QPO disappeared or decreased in size, or the optical depth of the gray medium changed.

We found that the burst e-folding time is shorter (~ 10 s) when the source is in the low intensity state than when the source is on the higher intensity banana branch (~ 30 s). This is opposite to what was found by van Paradijs, Penninx, & Lewin (1988) for several sources. Van der Klis et al. (1990) found for 4U 1636–53 that the burst duration was longer (> 20 s) when the source was in the island part of the CD than when the source was in the banana branch (< 10 s). Either the state where we observed the short burst e-folding times is not the product of an island state and high inclination effects as we propose, but a state at a higher mass accretion rate than the banana branch mass accretion rate, or another

parameter rather than \dot{M} is also affecting the burst duration.

The frequency of the QPO was 0.35 Hz higher during the type I X-ray bursts than during the persistent emission of observation 1. Similar increases in the QPO frequency during bursts were also observed in EXO 0748–676 by Homan et al. (1999).

This work was supported by the Netherlands Organization for Scientific Research (NWO) under contract number 614-51-002, and by NWO Spinoza grant 08-0 to E.P.J.van den Heuvel. This research has made use of data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. MM is a fellow of the Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina.




Fig. 1.— A: Hardness curve (ratio between the count rates in the 9.7–16 keV and the 2–9.7 keV bands) of the last part of observation 7 (time zero corresponds to August 3 14:15:03 UTC). B: The simultaneous 2–60 keV, background-subtracted light curve. There is clear spectral hardening during the dip apparent in the lower panel. No dead time corrections were applied (the deadtime fraction was $<1.5\%$). Since the photon-photon deadtime is completely negligible both in and outside the dip, the deadtime does not affect the hardening during the dip.




Fig. 2.— Normalized (van der Klis 1988) power spectrum of observation 1. The Poisson noise has been subtracted. The solid line represents the best fit to the data using two components in the fit: a Lorentzian at ~ 1 Hz (the QPO), and a power law.

Table 2. The fractional rms amplitudes, frequencies and FWHMs of the QPO in the observations where 4U 1746–37 was observed in the low intensity state. If no QPO at a significance level higher than 3σ was detected, the 95% confidence upper limit is given.

The FWHM and the frequency were measured in the 2–60 keV band, except for observation 2, where they were measured in the 2–6.4 keV band.

Observation	rms amplitude (2–60 keV)	rms amplitude (2–6.4 keV)	rms amplitude (6.4–20 keV)	ν_{QPO} (Hz)	FWHM (Hz)
1	$7.7\% \pm 0.7\%$	$7.8\% \pm 0.9\%$	$6.7\%^{+2.0\%}_{-1.1\%}$	1.04 ± 0.03	0.36 ± 0.09
2	$< 7\%$	$6.9\%^{+1.2\%}_{-0.9\%}$	$< 6.8\%$	1.59 ± 0.05	0.4 ± 0.2
3	$7.5\%^{+1.4\%}_{-1.1\%}$	$< 7.4\%$	$< 8.7\%$	1.01 ± 0.06	0.25 ± 0.13




Fig. 3.— Color-color diagram of 4U 1746–37. The hard color is defined as the log of the 9.7–16.0 keV / 6.0–9.7 keV count rate ratio, the soft color as the log of the 3.5–6.0 keV / 2–3.5 keV ratio. No dead-time corrections have been applied (the deadtime fraction was $<2\%$). The data are background subtracted and bursts were removed. The data points are 64 s and 128 s averages for the banana branch (lower part) and for the presumed (see text) island state (upper part), respectively. Typical error bars are shown for data in the hard (island) state and for data on the banana branch.

REFERENCES

- Bradt, H.V., Rothschild, R.E., Swank, J.H., 1993, A&AS, 97, 355
- Church, M.J., Balucińska-Church, M. 1995, A&A, 300, 441
- Deutsch, E.W., Anderson, S.F., margon, B., & Downes, R. 1998, ApJ, 493, 775
- Djorgovski, S., 1993, In: Djorgovski S., Meylen G., (eds.) ASP, conf. ser. 50, Structure and Dynamics of Globular Clusters. ASP, San Francisco, p. 373
- Frank, J., King, A.R., Lasota, J.-P. 1987, A&A, 178, 137
- Hasinger, G., & van der Klis, M. 1989, A&A, 225, 79
- Homan, J., Jonker, P.G., Wijnands, R., van der Klis, M., van Paradijs, J. 1999, ApJ, 516, L91
- Jonker, P.G., van der Klis, M., Wijnands, R. 1999a, ApJ, 511, L41
- Jonker, P.G., van der Klis, M., Homan, J., Wijnands, R. 1999b, in prep
- Lewin, W.H.G., van Paradijs, J., Taam, R.E. 1995, pag. 175 in X-Ray Binaries, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van der Heuvel (Cambridge: Cambridge Univ. Press)
- Li, F.K., & Clark, G.W. 1977, IAU Circ. 3095
- Parmar, A.N., et al. 1989, A&A, 222, 96
- Parmar, A.N., et al. 1999, submitted
- Sansom, A., Dotani, T., Asai, K., Lehto, H.J. 1993, MNRAS, 262, 429

- van der Klis, M. Timing neutron stars, eds. H. Ögelman, E.P.J. van den Heuvel, NATO ASI Series C, Vol. 262, p. 27–70, 1988
- van der Klis, M., Hasinger, G., Damen, E., Penninx, W., van Paradijs, J., & Lewin, W.H.G. 1990, ApJ, 360, L19
- van der Klis, M., 1995 pag. 252 in X-Ray Binaries, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van der Heuvel (Cambridge: Cambridge Univ. Press)
- van der Klis, M., 1998, astro-ph/9812395
- van Paradijs, J., Penninx, W., & Lewin, W.H.G., 1988, MNRAS, 233, 437
- White, N.E., and Swank, J.H., 1982, ApJ 253, L61
- White, N.E., Nagase, F., Parmar, A.N. 1995, pag. 1 in X-Ray Binaries, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van der Heuvel (Cambridge: Cambridge Univ. Press)