

# Ab initio results for the electronic structure of $C_{50}Cl_{10}$ .

Steven W.D. Bailey<sup>1</sup> and Colin J. Lambert<sup>1</sup>

<sup>1</sup> Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK

In this paper we use ab initio density functional theory (DFT) to calculate the electronic properties of  $C_{50}Cl_{10}$ . In comparison with the unstable  $C_{50}$  which has a small  $t_{lu}(LUMO) - h_u(HOMO)$  energy gap and a high total free energy compared with  $C_{60}$ , the belt of chlorine atoms stabilize the  $C_{50}Cl_{10}$  fullerene by increasing the energy gap to approximately that of  $C_{60}$  and lowering the total free energy. We also examine the effects of inter-cage separation on the band structure for infinite periodic  $C_{50}Cl_{10}$  chains where a high degree of dispersion is found to persist for separations beyond the predicted  $C_{60} - C_{60}$  distance of closest approach.

A new example of the fullerene family,  $C_{50}Cl_{10}$  has recently been produced as a stable solid [1]. The fullerenes smaller than the  $C_{60}$  buckyball are very difficult to produce in a stable form as the structure breaks the isolated pentagon rule (IPR) [2]. In  $C_{50}$  there are five sets of adjacent pentagons and this together with the high curvature of the molecular surface means it is highly unstable [3–5]. Xie et al have overcome this instability by attaching chlorine atoms around the  $C_{50}$  equator as seen in Fig. 1(a). In what follows, we examine the electronic properties and stability of  $C_{50}Cl_{10}$ , along with the electronic properties of chains of  $C_{50}Cl_{10}$  molecules. To model  $C_{50}Cl_{10}$  we use a double zeta basis set with a cutoff radius of 150 Ry to describe the carbon and Chlorine atoms within the SIESTA [6] implementation of DFT, where nonlocal norm-conserving pseudopotentials describe the core electrons and a linear combination of atomic orbitals the valence electrons. The relaxed, isolated molecule is shown in Fig. 1(a). In the case of a chain we consider molecules aligned along the z axis as shown in Fig. 1(b).




FIG. 1. Relaxed atomic coordinates of  $C_{50}Cl_{10}$ . The chlorine atoms form a fan around the equator of the  $C_{50}$  molecule (a). The proposed chain of  $C_{50}Cl_{10}$  in the z direction (b)

**TABLE 1.** compares the Fermi energy  $E_F$ (eV), the total free energy  $E_{tot}$ (eV) and the  $t_{lu}(LUMO) - h_u(HOMO)$  energy gap  $E_{\Delta}$ (eV) between the  $C_{60}$ ,  $C_{50}$  molecules and  $C_{50}Cl_{10}$  as a linear chain with varying inter-cage separations in (Å), up to the isolated molecule.

**TABLE 1.**

Case	a	b	c	d	e	f
	$C_{60}$ mol	$C_{50}$ mol	$C_{50}Cl_{10}2.5\text{Å}$	$C_{50}Cl_{10}2.9\text{Å}$	$C_{50}Cl_{10}3.4\text{Å}$	$C_{50}Cl_{10}$ mol
$E_F$ (eV)	-4.61	-7.04	-5.41	-5.41	-5.47	-5.67
$E_{tot}$ (eV)	-9201	-7670	-11751	-11751	-11751	-11750
$E_{\Delta}$ (eV)	1.80	0.16	—	—	—	1.80

The table shows that the unstable  $C_{50}$  has a high  $E_{tot}$  of  $-7670\text{eV}$  in comparison to that of  $-9201\text{eV}$  for the  $C_{60}$  molecule but that the belt of ten Cl atoms stabilizes the  $C_{50}Cl_{10}$  to produce a lower  $E_{tot}$  of  $-11751\text{eV}$ . This is reflected in the result for the  $t_{lu}(LUMO) - h_u(HOMO)$  energy gap, ( $E_{\Delta}$ ), where the energy gap of  $0.16\text{eV}$  for  $C_{50}$  opens to  $1.80\text{eV}$ , matching that of  $C_{60}$ , for the  $C_{50}Cl_{10}$  molecule. It is noted that the value of  $E_{\Delta} = 1.80\text{eV}$  for  $C_{60}$  is in very good agreement with the experimental values of  $E_{\Delta}$  between  $(1.6 - 1.8)\text{eV}$  [7]

Fig. 2 shows the energy levels of the isolated molecules in cases (a,b and f) and the electronic band structures of  $C_{50}Cl_{10}$  chains with the different inter-molecule separations in cases (c - e), of Table 1. In each case the energy is measured from the  $E_F$  for each relaxed structure studied. In cases(c - e) the band structure is shown for a wavevector  $kz$  in the z direction as defined in Fig. 1. The  $C_{60} - C_{60}$  distance of closest approach is predicted to be approximately

3.0Å [8] therefore the range of cage separations for cases (c - e) were selected to cover this interval and beyond. The band structures for the infinite periodic chains of  $C_{50}Cl_{10}$  for cases (c - e) show the dispersion of the  $t_{lu}(LUMO)$  and  $h_u(HOMO)$  bands persists up to and beyond a 3.4Å separation and that the  $C_{50}Cl_{10}$  chain is an insulator. In conclusion we have used DFT calculations to investigate the electronic properties of the newly produced, solid  $C_{50}Cl_{10}$  fullerene. We find that the belt of chlorine atoms around the equator of the  $C_{50}$  stabilize the structure by dramatically lowering the total free energy of  $C_{50}Cl_{10}$  and increasing the  $h_u(HOMO) - t_{lu}(LUMO)$  gap to match that of  $C_{60}$ . The  $t_{lu}(LUMO)$  and  $h_u(HOMO)$  bands show a marked dispersion even for separations of the  $C_{50}Cl_{10}$  cages in excess of the predicted closest  $C_{60} - C_{60}$  approach of 3.0Å. This points to some interesting properties arising from carbon nanotube peapods such as (12, 12)@ $C_{50}Cl_{10}$ .




FIG. 2. For the structures (a - f) defined in Table 1, the energy levels of the isolated molecules are seen in cases (a,b and f) and the electronic band structures for wavevectors  $kz$  in the  $z$  direction as defined in Fig. 1, in cases (c - e).

- 
- [1] Su - Yuan Xie et al, Science. **304**, 5671 (2004).
  - [2] K. M. Kadish and R. S. Ruoff, Fullerene: Chemistry, Physics and Technology (Wiley, new York, 2002).
  - [3] T. Guo et al., Science **257**, 1661 (1992).
  - [4] H. W. Kroto, Nature **329**, 529 (1987).
  - [5] J. R. Heath, Nature **393**, 730 (1998).
  - [6] S. D. S-Portal, P. Ordejon, E. Artacho, J. Soler Int. J. Quantum Chem. **65**, 453 (1997).
  - [7] M.S. Dresselhaus, G. Dresselhaus and P. C. Eklund, Science of Fullerenes and Carbon Nanotubes (Academic, San Diego, 1996).
  - [8] A. Rochefort, Phys. rev. B, **67**, 115401, (2003).