## Theory of single-particle properties of the Hubbard model

Y.M. Vilk, and A.-M.S. Tremblay

Département de Physique and Centre de Recherche en Physique du Solide,

Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1

(24 January 1995)

It is shown that it is possible to quantitatively explain quantum Monte Carlo results for the Green's function of the two-dimensional Hubbard model in the weak to intermediate coupling regime. The analytic approach includes vertex corrections in a paramagnon-like self-energy. All parameters are determined self-consistently. This approach clearly shows that in two dimensions Fermi-liquid quasiparticles disappear in the paramagnetic state when the antiferromagnetic correlation length becomes larger than the electronic thermal de Broglie wavelength.

For more than thirty years, the Hubbard model has served as a theoretical basis for a wide range of interacting electron phenomena, from itinerant electron magnetism to, more recently, high-temperature superconductivity. In the latter context, one of the key issues has been the possibility of non-Fermi liquid behavior in the single-particle properties. These properties have been intensively studied in the weak to intermediate coupling regime using paramagnon-type theories. [1] [2] Such approaches are physically attractive since they describe the effect of low-lying collective modes on single-particle properties in a manner similar to that of phonons. However in these theories one faces the difficulty that, contrary to the case of phonons, vertex corrections are not small (no Migdal theorem) and have to be taken into account.

In this letter, we introduce a simple approach that has the straightforward physical interpretation of paramagnon theories and uses no adjustable parameter. It does include vertex corrections, and achieves, for both two-particle and single-particle properties, better quantitative agreement with Monte Carlo simulations than previous theories [3] [4] [5]. Our analytical approach allows us to unambiguously interpret the Monte Carlo data and to show that in two dimensions spin fluctuations destroy the Fermi liquid in the paramagnetic state when the antiferromagnetic correlation length becomes larger than the thermal de Broglie wavelength of electrons. This corresponds to the disappearance of the quasiparticle peak and the appearance of a pseudogap, an issue that has been controversial mostly due to finite-size effects in Monte Carlo [6].

We consider the one-band Hubbard model on the square lattice with unit lattice spacing, on-site repulsion U and nearest-neighbor hopping t. We work in units where the lattice spacing is unity,  $k_B = 1$ ,  $\hbar = 1$  and t = 1. The theory has a simple structure that we now explain physically, postponing the formal derivation momentarily.

The calculation proceeds in two steps: we first obtain spin and charge susceptibilities, then we inject them in the self-energy calculation. In the calculation of susceptibilities we make the approximation that spin and charge susceptibilities  $\chi_{sp}$ ,  $\chi_{ch}$  are given by RPA-like forms but with two different effective interactions  $U_{sp}$  and  $U_{ch}$  which are then determined self-consistently. The necessity to have two different effective interactions for spin and for charge is dictated by the Pauli exclusion principle  $\langle n_{\sigma}^2 \rangle = \langle n_{\sigma} \rangle$  which implies that both  $\chi_{sp}$  and  $\chi_{ch}$  are related to only one local pair correlation function  $\langle n_{\uparrow}n_{\downarrow} \rangle$ . Indeed, using the Fluctuation-Dissipation theorem in Matsubara formalism and the Pauli principle one can write:

$$\frac{1}{\beta N} \sum_{q} \chi_{ch,sp}(q) = \frac{1}{\beta N} \sum_{q} \frac{\chi_0(q)}{1 + \frac{(-1)^{\ell}}{2} U_{ch,sp} \chi_0(q)} = n + 2 (-1)^{\ell} \langle n_{\uparrow} n_{\downarrow} \rangle - (1 - \ell) n^2 \tag{1}$$

where  $\ell = 0$  for charge (ch),  $\ell = 1$  for spin (sp),  $\beta \equiv 1/T$ ,  $n = \langle n_{\uparrow} \rangle + \langle n_{\downarrow} \rangle$ ,  $q = (\mathbf{q}, iq_n)$  with  $\mathbf{q}$  the wave vectors of an N site lattice,  $iq_n$  Matsubara frequencies and  $\chi_0(q)$  the susceptibility for non-interacting electrons. The value of  $\langle n_{\uparrow}n_{\downarrow} \rangle$  may be obtained self-consistently [7] by adding to the above set of equations the relation  $U_{sp} = g_{\uparrow\downarrow}(0) U$  with  $g_{\uparrow\downarrow}(0) \equiv \langle n_{\uparrow}n_{\downarrow} \rangle / \langle n_{\downarrow} \rangle \langle n_{\uparrow} \rangle$ .

As shown in Ref. [7], the above procedure reproduces both Kanamori-Brueckner screening as well as the effect of Mermin-Wagner thermal fluctuations, giving a phase transition only at zero-temperature in two-dimensions. There is however a crossover temperature  $T_X$  below which the magnetic correlation length  $\xi$  grows exponentially. Quantitative agreement with Monte Carlo simulations is obtained [7] for all fillings and temperatures in the weak to intermediate coupling regime U < 8.

We now turn to the discussion of the single-particle properties. In order, to be consistent with the two-particle correlation functions, the self-energy  $\Sigma_{\sigma}(k)$  must satisfy the sum rule

$$\lim_{\tau \to 0^{-}} \frac{1}{\beta N} \sum_{k} \Sigma_{\sigma} \left( k \right) G_{\sigma} \left( k \right) e^{-ik_{n}\tau} = U \left\langle n_{\uparrow} n_{\downarrow} \right\rangle, \tag{2}$$

which follows from the definition of  $\Sigma_{\sigma}(k)$  [8]. Here, we encounter the same key quantity  $\langle n_{\uparrow}n_{\downarrow}\rangle$  that appears in the sum rule for the susceptibilities Eq. (1). We find, using the formal approach described below, the following expression for  $\Sigma_{\sigma}(k)$ 

$$\Sigma_{\sigma}(k) = Un_{-\sigma} + \frac{U}{4} \frac{T}{N} \sum_{q} \left[ U_{sp} \chi_{sp}(q) + U_{ch} \chi_{ch}(q) \right] G^{0}_{\sigma}(k+q),$$
(3)

which satisfies Eq. (2) with  $G_{\sigma}$  replaced by  $G_{\sigma}^{0}$  on the left-hand side. This self-energy expression (3) is physically appealing since, as expected from general skeleton diagrams, one of the vertices is the bare one U, while the other vertex is dressed and given by  $U_{sp}$  or  $U_{ch}$  depending on the type of fluctuation being exchanged. Eq.(3) already gives good agreement with Monte Carlo data but the accuracy can be improved even further by requiring that the consistency condition (2) be satisfied with  $G_{\sigma}$  instead of  $G_{\sigma}^{0}$ . To do so we replace  $U_{sp}$  and  $U_{ch}$  on the right-hand side of (3) by  $\alpha U_{sp}$  and  $\alpha U_{ch}$  with  $\alpha$  determined self-consistently by Eq.(2). For U < 4, we have  $\alpha < 1.15$ . This concludes the description of the structure of our theory. We now turn to the formal derivation.

Formal aspects: We use the notation  $1 = (\mathbf{R}_1, \tau_1)$ , and an overbar  $(\overline{1})$  to indicate a space-time or spin variable which is integrated or summed over. Using the Matsubara Green's function  $G_{\sigma}(1,3)$  in the presence of an external potential  $\phi_{\sigma}(2)$ , one can write the exact equations for the generalized susceptibilities  $\chi_{\sigma\sigma'}(1,3;2) \equiv \delta G_{\sigma}(1,3) / \delta \phi_{\sigma'}(2)$  and the self-energy  $\Sigma_{\sigma}(1,2)$  as [8]:

$$\chi_{\sigma,\sigma'}\left(1,3;2\right) = G_{\sigma}\left(1,2\right)G_{\sigma}\left(2,3\right)\delta_{\sigma,\sigma'} + G_{\sigma}\left(1,\overline{6}\right)G_{\sigma}\left(\overline{7},3\right)\Gamma^{ir}_{\sigma,\overline{\sigma}}\left(\overline{6},\overline{7};\overline{4},\overline{5}\right)\chi_{\overline{\sigma},\sigma'}\left(\overline{4},\overline{5};2\right) \tag{4}$$

$$\Sigma_{\sigma}(1,2) = Un_{-\sigma}\delta(1-2) - UG_{\sigma}(1,\overline{6})\Gamma_{\sigma\bar{\sigma}}^{ir}(\overline{6},2;\overline{4},\overline{5})\chi_{\bar{\sigma},-\sigma}(\overline{4},\overline{5};1).$$

$$\tag{5}$$

where  $\Gamma_{\sigma,\sigma'}^{ir} = \delta \Sigma_{\sigma} / \delta G_{\sigma'}$  are irreducible vertices. Our approximation for spin and charge susceptibilities and for selfenergy Eq.(3) is obtained from these exact equations by using as initial guess on the right-hand side of *both* equations a self-energy  $\Sigma_{\sigma}^{(0)}$  and irreducible vertices  $\Gamma_{\sigma\bar{\sigma}}^{(0)ir}$  which are functional derivatives of

$$\Phi\left[G\right] = \frac{1}{4} (U_{ch} - U_{sp}) \widetilde{n}_{\overline{\sigma}} \left(\overline{1}\right) \widetilde{n}_{\overline{\sigma}} \left(\overline{1}\right) + \frac{1}{4} (U_{ch} + U_{sp}) \widetilde{n}_{\overline{\sigma}} \left(\overline{1}\right) \widetilde{n}_{-\overline{\sigma}} (\overline{1})$$

$$\tag{6}$$

where  $\tilde{n}_{\sigma}(1) \equiv G_{\sigma}(1, 1^+)$ . The initial self-energy  $\Sigma_{\sigma}^{(0)} = \delta \Phi / \delta G_{\sigma}|_{\phi=0}$  is a constant and hence can be absorbed in a chemical potential shift so that all Green's functions on the right-hand side of Eqs. (4) and (5) are bare ones  $G_{\sigma}^0(1, 2)$ . The corresponding particle-hole irreducible vertices  $\Gamma^{(0)ir} = \delta^2 \Phi / (\delta G^2)|_{\phi=0}$  are given in Fourier space by two constants

$$\Gamma_{\sigma;-\sigma}^{(0)ir} = (U_{ch} + U_{sp})/2 \quad ; \quad \Gamma_{\sigma,\sigma}^{(0)ir} = (U_{ch} - U_{sp})/2 \tag{7}$$

which are chosen so that the susceptibilities are consistent with the correct double-occupancy  $\langle n_{\uparrow}n_{\downarrow}\rangle$  and with the exclusion principle  $\langle n_{\sigma}^2 \rangle = \langle n_{\sigma} \rangle$ , as described above.

Important advantages of our iterative procedure are that: a) The approximation is conserving since  $\Sigma_{\sigma}^{(0)}$  and  $\Gamma_{\sigma,\sigma'}^{(0)ir}$  are functional derivatives of a single functional  $\Phi$  [9]. b) The following identity

$$\Sigma_{\sigma}\left(1,\overline{3}\right)G_{\sigma}\left(\overline{3},2\right) = -U\left\langle T_{\tau}\left[n_{-\sigma}\left(1\right)\psi_{\sigma}\left(1\right)\psi_{\sigma}^{+}\left(2\right)\right]\right\rangle \tag{8}$$

is satisfied with  $\Sigma_{\sigma}^{(1)}$  and  $G_{\sigma}^{(0)}$  and it can be used as a convergence check, namely  $\Sigma_{\sigma}^{(1)}(1,\overline{3}) G_{\sigma}^{(1)}(\overline{3},1^+) \approx \Sigma_{\sigma}^{(1)}(1,\overline{3}) G_{\sigma}^{(0)}(\overline{3},1^+)$ . For U < 4, this convergence criterion is satisfied at worse within 15%. The sum rule (2) follows from the identity (8) when  $2 = 1^+$ .

Comparisons with other theories and with Quantum Monte Carlo data: Fig.1(a) shows  $G(\mathbf{k},\tau)$  for filling n = 0.875, temperature T = 0.25 and U = 4 for the wave vector on the  $8 \times 8$  lattice which is closest to the Fermi surface, namely  $(\pi, 0)$ . For these parameters, size effects are negligible. Our theory is in agreement with Monte Carlo data and with the parquet approach but in this regime second-order perturbation theory for the self-energy gives the same result. This surprising performance of perturbation theory (see also [10]) is a consequence of compensation between the renormalized vertices and susceptibilities  $(U_{sp} < U, \chi_{sp}(q) > \chi_0(q); U_{ch} > U, \chi_{ch}(q) < \chi_0(q))$ . Half-filling n = 1 is an ideal situation for numerical studies of low-energy phenomena since some of the allowed wave vectors on finite lattices lie exactly on the Fermi surface. Fig.1(b) shows  $G(\mathbf{k}_F, \tau)$  for  $\mathbf{k}_F = (\pi, 0)$  in a regime where the antiferromagnetic correlation length is growing exponentially. The Green's function is strongly renormalized since the non-interacting value is -0.5, independent of  $\tau$ . Our theory shows better agreement with Monte-Carlo than previous approaches. Despite the very large correlation length, finite-size effects for the Matsubara Green's function  $G(\mathbf{k}_F, \tau)$ are not large and almost completely disappear when the system size becomes larger than the thermal de Broglie wavelength of the electrons  $(\xi_{th} \equiv \langle v_F \rangle / \pi T)$  as we explain below.

Our most dramatic numerical results addressing the issue of the Fermi liquid are shown in Fig. 2 where we plot

$$\tilde{z}(T) = -2G(\mathbf{k}_F, \beta/2) = \int \frac{d\omega}{2\pi} \frac{A(\mathbf{k}_F, \omega)}{\cosh\left(\beta\omega/2\right)}.$$
(9)

This quantity  $\tilde{z}(T)$  is an average of the single-particle spectral weight  $A(\mathbf{k}_F, \omega)$  within  $T \equiv 1/\beta$  around the Fermi level  $(\omega = 0)$  and it is a generalization of the usual zero-temperature quasiparticle renormalization factor  $z \equiv 1/(1-\partial \Sigma/\partial \omega)$ . For non-interacting particles  $\tilde{z}(T)$  is unity. For a normal Fermi liquid it becomes equal to a constant less than unity as the temperature decreases since the width of the quasiparticle peak scales as  $T^2$  and hence lies within T of the Fermi level. The quantity  $\tilde{z}(T)$  is the best estimate of z one can get from Monte Carlo data for  $G(\mathbf{k}, \tau)$ . Moreover  $\tilde{z}(T)$  gives an estimate of  $A(\mathbf{k}_F, \omega)$  around the Fermi surface even when the Fermi liquid does not exist.

One can clearly see from Fig. 2 that while second-order perturbation theory exhibits typical Fermi-liquid behavior for  $\tilde{z}(T)$ , both Monte Carlo data and a numerical evaluation of our expression for the self-energy lead to a rapid fall-off of  $\tilde{z}(T)$  below  $T_X$  (for U = 4,  $T_X \approx 0.2$  [7]). The decrease of  $\tilde{z}(T)$  in this regime is approximately  $\tilde{z} \sim T^2$ which clearly suggests non-Fermi liquid behavior. The physical origin of this effect is that the quasi-particles of the *two-dimensional* paramagnetic state become overdamped when the energy scale associated with the proximity to antiferromagnetism  $\delta U \equiv U_{mf,c} - U_{sp}$  ( $U_{mf,c} \equiv 2/\chi_0(\mathbf{Q}, 0)$ ) becomes exponentially small or, more precisely, when the two-particle antiferromagnetic correlation length  $\xi$  becomes larger than the single-particle thermal de Broglie wavelength  $\xi_{th} \equiv v_F/\pi T$ .

Pseudogap: While size effects and statistical errors make continuation of the Monte Carlo data to real frequencies particularly difficult, in our approach we can make this continuation analytically to show that the above effect corresponds to a pseudogap. We first explain the behavior  $\tilde{z} \sim T^2$  then do real frequency analysis.

Since the spin susceptibility  $\chi_{sp}(\mathbf{q},0)$  below  $T_X$  is almost singular at the antiferromagnetic wave vector  $\mathbf{Q} = (\pi,\pi)$ , the main contribution to  $\Sigma$  in Eq. (3) comes from  $iq_n = 0$  and wave vectors  $(\mathbf{q} - \mathbf{Q})^2 \leq \xi^{-2}$  near  $\mathbf{Q}$ . Approximating  $\chi_{sp}(\mathbf{q},0)$  in Eqs. (1) and (3) by its asymptotic form  $\chi_{sp}(\mathbf{q},0) \approx 2 \left[ U_{sp} \xi_0^2 (\xi^{-2} + (\mathbf{q} - \mathbf{Q})^2) \right]^{-1}$  where  $\xi_0^2 \equiv \frac{-1}{2\chi_0(Q)} \frac{\partial^2 \chi_0(Q)}{\partial q_x^2}$  and  $\xi \equiv \xi_0 (U_{sp}/\delta U)^{1/2}$ , the integrals over  $\mathbf{q}$  can be done to obtain the asymptotic results

$$\xi \sim \exp\left(\pi \tilde{\sigma}^2 \xi_0^2 \frac{U_{sp}}{T}\right) \tag{10}$$

$$\Sigma\left(\mathbf{k}_{F}, ik_{n}\right) = \frac{U}{2} - i \frac{UT}{8\pi\xi_{0}^{2}\sqrt{k_{n}^{2} - v_{F}^{2}\xi^{-2}}} \ln \frac{k_{n} + \sqrt{k_{n}^{2} - v_{F}^{2}\xi^{-2}}}{k_{n} - \sqrt{k_{n}^{2} - v_{F}^{2}\xi^{-2}}} + \mathcal{R}.$$
(11)

Here  $\mathcal{R}$  is the regular part which remains finite as  $T \to 0$  and  $\tilde{\sigma}^2 \equiv n - 2\langle n_{\uparrow}n_{\downarrow}\rangle - C < 1$  is the righthand side of (1) minus corrections C that come from the sum over non-zero Matsubara frequencies (quantum effects) and from  $(\mathbf{q} - \mathbf{Q})^2 \gg \xi^{-2}$ . The corresponding value of  $\tilde{z}(T)$  can be written as the alternating series  $-2G(\mathbf{k}_F, \beta/2) = -4T \sum_{n=1}^{\infty} (-1)^n / (ik_n + \mu - \Sigma(\mathbf{k}_F, ik_n))$ . We can understand qualitatively both the temperature and size dependence of the Monte Carlo data in Fig. 2 by using the first term of this series along with Eqs. (10) and (11)

$$\tilde{z}(T) \sim \frac{T^2}{\tilde{\sigma}^2 U U_{sp}} \sqrt{1 - \frac{\xi_{th}^2}{\xi^2}}.$$
(12)

On the infinite lattice,  $\xi$  starts growing exponentially below  $T_X$ , quickly becoming much larger than  $\xi_{th}$ . This implies  $\tilde{z}(T) \simeq T^2$ . On finite lattices Eq.(1) gives  $\xi \sim \sqrt{N}$  below  $T_X$  which explains the size effect observed in Monte Carlo *i.e.* smaller  $\tilde{z}$  for smaller size N,  $(\xi_{th}(T_X) \sim 5$  for Fig. 2).

The analytic continuation of  $\Sigma(\mathbf{k}_F, ik_n)$  in Eq. (11) is

$$\Sigma^{R}(\mathbf{k}_{F},\omega) = \frac{U}{2} + \frac{UT}{8\pi\xi_{0}^{2}\sqrt{\omega^{2} + v_{F}^{2}\xi^{-2}}} \ln \left| \frac{\omega + \sqrt{\omega^{2} + v_{F}^{2}\xi^{-2}}}{\omega - \sqrt{\omega^{2} + v_{F}^{2}\xi^{-2}}} \right| - i\frac{UT}{8\xi_{0}^{2}\sqrt{\omega^{2} + v_{F}^{2}\xi^{-2}}} + \mathcal{R}.$$
(13)

At small frequencies the regular part  $\mathcal{R}$  can be neglected so that exactly at the Fermi surface ( $\omega = 0$ ) the imaginary part of the self-energy for  $\xi > \xi_{th}$  increases exponentially when the temperature decreases,  $\Sigma''(\mathbf{k}_F, \omega) \sim U\xi/(\xi_{th}\xi_0^2) \sim$  $\exp(\pi \tilde{\sigma}^2 \xi_0^2 U_{sp}/T)$  [11]. This corresponds to a pseudogap: Instead of a quasiparticle peak, the spectral weight has a minimum at the Fermi level and two symmetrically located maxima away from it. Clearly this is not a Fermi liquid, yet the symmetry of the system remains unbroken at any finite T. By contrast [12] with 3D, in 2D this effect exists in a wide temperature range  $T < T_X$ . These conclusions persist slightly away from half-filling. In particular, we do not find a quasiparticle peak in the pseudogap close to half-filling when  $\xi \gg \xi_{th}$ . This is different from the results inferred from a phenomenological zero-temperature calculation [2] ( $\xi_{th} = \infty$ ) which physically corresponds to  $1 \ll \xi \ll \xi_{th}$ .

In photoemission experiments on real quasi two-dimensional materials we predict a rapid decrease of the spectral weight and density of states at the Fermi level in a wide temperature range, from  $T_X$  to the Néel temperature  $T_N$   $(T_X - T_N \sim 10^2 K)$ .

We thank Liang Chen, R. Côté and A.E. Ruckenstein for numerous enlightening discussions. We acknowledge financial support from the Natural Sciences and Engineering Research Council of Canada (NSERC), the Fonds pour la formation de Chercheurs et l'aide à la Recherche from the Government of Québec (FCAR), and (A.-M.S.T) the Canadian Institute of Advanced Research (CIAR) and the Killam foundation.

- [1] N.F. Berk, and J.R. Schrieffer, Phys. Rev. Lett. 17, 433 (1966). For a review, P.C.E. Stamp, J. Phys. F, 15, 1829 (1985).
- [2] A.P. Kampf, and J.R. Schrieffer, Phys. Rev. B 41, 6399 (1990); Phys. Rev. B 42, 7967 (1990).
- [3] N. E. Bickers and D. J. Scalapino, Ann. Phys. (N. Y.) 193, 206 (1989).
- [4] N. E. Bickers, Int. J. of Mod. Phys. B 5, 253 (1991).
- [5] N. E. Bickers and S. R. White, Phys. Rev. B 43, 8044 (1991).
- [6] S. R. White, Phys. Rev. B 46, 5679 (1992); M. Vekić and S. R. White, Phys. Rev. B 46, 5679 (1992)
- [7] Y.M. Vilk, Liang Chen and A.-M.S. Tremblay, Phys. Rev. B 49, 13 267 (1994).
- [8] L. P. Kadanoff and G. Baym, Quantum Statistical Mechanics (Benjamin, Menlo Park, 1962)
- [9] Gordon Baym, Phys. Rev 127, 1391 (1962).
- [10] J. Galán, J.A. Vergés, and A.Martin-Rodero, Phys. Rev. B 48, 13 654 (1993).
- [11] It can be shown that for  $T > T_X$ ,  $\xi \sim 1$  the regular part  $\mathcal{R}$  dominates in equation (13) and one recovers standard behavior with the maximum in the spectral weight at the Fermi level.
- [12] In the isotropic 3D case  $\Sigma''(\mathbf{k}_F, \omega) \sim (\ln \xi) / \xi_{th}$  and hence the pseudogap exists only in a very narrow temperature range.

Fig. 1. Comparison of our results for  $G(\mathbf{k}, \tau)$  with Monte Carlo data, FLEX, parquet, and second-order pertubation theory, all on  $8 \times 8$  mesh. Monte Carlo data and results for FLEX and parquet are from Ref. [5]. a) n = 0.875, T = 0.25; b) n = 1, T = 0.17.

Fig. 2. Temperature dependence of the generalized renormalization factor  $\tilde{z}$  defined in Eq.(9). Lines are results of our calculations for infinite lattice and  $16 \times 16$  mesh. Symbols are Monte Carlo data from Ref. [6].