Accepted for publication in Physical Review D UM-P 014-2001 RCHEP 001-2001

## TeV-scale electron Compton scattering in the Randall-Sundrum scenario.

S.R.Choudhury<sup>1</sup>

International Centre for Theoretical Physics, Trieste, Italy and School of Physics, University of Melbourne, Parkville, Victoria 3108, Australia A.S.Cornell<sup>2</sup>, and G.C.Joshi<sup>3</sup> School of Physics, University of Melbourne, Parkville, Victoria 3108, Australia

## Abstract

The spin-2 graviton excitations in the Randall-Sundrum gravity model provides a *t*-channel contribution to electron Compton scattering which competes favourably with the standard QED contributions. The phenomenological implications of these contributions to the unpolarized and polarized cross-sections are evaluated.

The Standard model (SM) has so far been very successful for explaining most experimental observations relevant to the model. However, it has one very ugly feature namely the hierarchy problem, which refers to the enormous difference in the order of magnitude of the Planck scale ( $M_{pl} \sim 10^{19}$ GeV) and the weak scale (~ 1TeV). Recent ideas that have been proposed to solve the problem introduce extra compact spatial dimensions. In the first idea proposed in this direction, given by Arkani-Hamed, Dimopoulos and Dvali (ADD)[1], spacetime was a direct product of a four dimensional Minkowski spacetime and a compact *n*-manifold. While gravity was free to propagate through the extra *n*-spatial dimensions as well, SM fields were restricted to the four-dimensional spacetime. The Planck-scale in (4+*n*)-dimensional world could be much much smaller than  $M_{pl}$  - even as low as the TeV scale. The weakness of gravity for distances  $\gg$  compact dimension arises not because of a fundamental weakness of gravitational interaction but because of SM-fields being restricted to lie on a brane of the entire (4+*n*)-dimensional spacetime.

Phenomenologically, the ADD-model implies the existence of a large number of 'massive' gravitons coupled with (four-dimensional) Planck strength to SMparticles, with masses ranging from near zero to some high TeV scale  $M_s$ .

 $<sup>^1 {</sup>m src} @ {
m ducos.ernet.in}$ 

 $<sup>^2</sup>a. cornell@tauon.ph.unimelb.edu.au$ 

 $<sup>^3</sup>$ joshi@physics.unimelb.edu.au

An alternative scenario considered first by Randall and Sundrum (RS)[2], the extra dimension is a single  $S^1/Z_2$  orbifold with 3-branes residing at the boundaries of the spacetime. With suitably tuned brane tensions and a bulk cosmological constant, RS obtain a non-factorizable metric

$$ds^{2} = e^{-2kr_{c}|\phi|}\eta_{\mu\nu}dx^{\mu}dx^{\nu} - r_{c}^{2}d\phi^{2}$$
<sup>(1)</sup>

as a solution of five-dimensional Einstein equations. Here  $\phi \epsilon [-\pi, \pi]$  is the fifth coordinate with  $\phi$  and  $-\phi$  identified.  $r_c$  is the radius of the fifth dimension and k is a parameter  $\sim M_{pl}$ . Choosing  $kr_c \sim 12$ , this model also solves the hierarchy problem. Phenomenologically, the RS-model gives rise to a discrete series of 'heavy' gravitons of increasing mass which couple with SM-particles with weak interaction strength (referred to now on as KK).

Phenomenological consequences of these excited 'gravitons' would naturally be felt at TeV-scale processes. In situations where the normal SM contributions are absent or small for some reason the KK-contributions are obviously the ones that are best suited for testing the theory. One such vertex is the KK- $\gamma\gamma$  vertex which exists in this theory, whereas the corresponding  $Z\gamma\gamma$  or  $H\gamma\gamma$  vertices are absent [3]. The existence of the KK- $\gamma\gamma$  vertex would imply a *t*-channel pole contribution to the Compton-scattering process  $\gamma + e \rightarrow \gamma + e$ . The SMcontribution with s- and u-channel electron pole will have comparable strength and hence the KK-contribution will show up in the cross-section of this process. The feasibility of performing the Compton process using backscattering laser beams in the NLC has been discussed by Davoudias [4], who has also calculated the effect of ADD-type excitations on the process. Phenomenologically, the RSmodel has differences from the ADD and it will be useful to examine Compton scattering in the RS-model as well. In this note we carry out the analysis not only to see the nature of changes in the cross-section compared to SM, but also to see if the process at high energies can distinguish between the two gravity models.

We consider the process

$$\gamma(k_1) + e(p_1) \to \gamma(k_2) + e(p_2)$$

and neglect the electron mass. The invariant amplitude for this process is denoted by  $M_{\lambda_1,\lambda_2,\lambda_3,\lambda_4}(s,t,u)$  where s,t,u are the usual Mandelstam variables and  $\lambda_1, \lambda_2, \lambda_3, \lambda_4$  denote respectively the helicities of initial photon, initial electron, final photon and final electron. The interaction of the KK-excitations  $h_{\alpha\beta}^{(n)}$ with SM-fields is described by the interaction Lagrangian [5];

$$\mathcal{L} = -\frac{1}{\Lambda_{\pi}} \cdot T^{\alpha\beta} h^{(n)}_{\alpha\beta} \tag{2}$$

where  $T^{\alpha\beta}$  is the SM-energy momentum tensor. The constant  $\Lambda_{\pi}$  is related to the parameters of RS metric and the Planck scale by

$$\Lambda_{\pi} = M_{pl} \cdot e^{-kr_c\pi}.$$
(3)



Figure 1: Diagrams for high energy Compton scattering, (a) and (b) are the usual QED contributions, whereas (c) represents the contribution from exchange of a KK-graviton

For purposes of phenomenological parametrization it is more convenient to choose the mass of the lowest KK-resonance  $m_1$  and  $\left(\frac{k}{M_{pl}}\right)$  as independent parameters with

$$\Lambda_{\pi}^{-1} = \left(\frac{k}{M_{pl}}\right) \cdot \left(\frac{x_1}{m_1}\right),$$

where  $x_1$  is the first zero of  $J_1(x)$ . For our numerical evaluation, we shall fix  $m_1$  at a reasonable value of 600GeV and vary  $\left(\frac{k}{M_{pl}}\right)$  over a range of values within the acceptable limits [5].

With the Lagrangian, equation (2), added to the standard Q.E.D. Lagrangian, the amplitudes for the diagrams of figure (1) work out to be

$$M_{++++}(s,t,u) = M_{----}(s,t,u) = \frac{8\pi\alpha\sqrt{s}}{\sqrt{-u}} - \sum_{i=1}^{12} \frac{2s^{\frac{3}{2}}\sqrt{-u}}{\Lambda_{\pi}^{2}(m_{i}^{2}-t)}$$
(4)

$$M_{+-+-}(s,t,u) = M_{-+-+}(s,t,u) = \frac{8\pi\alpha\sqrt{-u}}{\sqrt{s}} + \sum_{i=1}^{12} \frac{2u\sqrt{-su}}{\Lambda_{\pi}^2(m_i^2 - t)}$$
(5)

with all other amplitudes vanishing. We have retained in the summation in the right hand side of equation (4) and (5) the first 12 resonaces, having checked that the terms left out contribute less than 1%. In equations (4) and (5),  $m_i$ 's are the masses of the KK-resonance of the spin-2 graviton in the RS model, given by

$$m_i = \left(\frac{k}{M_{pl}}\right)\Lambda_{\pi} \cdot x_i$$

where the  $x_i$ 's are the zeroes of the Bessel function  $J_1(x)$ . Thus, for a given  $m_1$ , which we have taken to be 600 Gev., the  $m_i$ 's are simply in proportion to the  $x_i$ 's. In the limit of vanishing electron-mass, the contribution of the KK-exchange graphs like the QED-contribution conserve both electron and photon

helicities.

The differential cross-section for the two different initial electron polarizations j  $(j = \pm 1)$  are then given by

$$\left(\frac{d\sigma}{d\Omega}\right)_{j} = \frac{1}{64\pi^{2}s} \cdot \left|M_{+j+j}\right|^{2}.$$
(6)

The  $\gamma$ -beams are produced by backscattering of laser beams off electron beams; it is thus more appropriate to express the polarized cross-sections in terms of the helicities of the original laser beam  $(p_l)$  and the helicity of the electron beam  $(p_e)$  from which it was backscattered [4]. Since the backscattered photon carries only a fraction x of the energy of the electron beam, the Mandelstam variables entering the amplitudes M will be (xs, xt, xu) instead of (s, t, u). The fraction x theoretically has the maximum value

$$x_{max} = \frac{z}{1+z} \quad , \quad z = \frac{4E_e E_l}{m_e^2} \tag{7}$$

where  $E_l$  and  $E_e$  respectively are the energies of the laser beam and the electron beam from which it is backscattered. From equation (7), the value of  $x_{max}$ approaches unity as z tends towards infinity with  $E_l$  increasing indefinitely. However as  $E_l$  increases,  $e^+e^-$  are produced more and more because of the interaction of the laser beam with the backscattered photons [6]. The luminosity of the beam therefore erodes rapidly. An optimal value of z is thus finite and is given by [6]

$$z_{opt} = 2(1+\sqrt{2})$$
 (8)

which is the value we would use in equation (7). For a given value of x,  $p_l$ , and  $p_e$ , the photon number density  $f(x, p_e, p_l)$  and the average helicity  $\xi_2(x, p_e, p_l)$  are given by [4]:

$$f(x, p_e, p_l) = N^{-1} \left[ \frac{1}{1-x} + (1-x) - 4r(1-r) - rp_l p_e z(2r-1)(2-x) \right]$$
(9)

$$\xi_2(x, p_e, p_l) = \frac{1}{Nf(x)} \left\{ p_e \left[ \frac{x}{1-x} + x(2r-1)^2 \right] - p_l(2r-1) \left( 1 - x + \frac{1}{1-x} \right) \right\} (10)$$

where  $r = \frac{x}{z(1-x)}$  and

$$N = \ln(1+z) \cdot \left(1 - \frac{4}{z} - \frac{8}{z^2}\right) + \frac{1}{2} + \frac{8}{z} - \frac{1}{2(z+1)^2} + p_e p_l \cdot \left[\ln(1+z)\left(1 + \frac{2}{z}\right) - \frac{5}{2} + \frac{1}{1+z} - \frac{1}{2(z+1)^2}\right]$$
(11)

For a given  $\xi_2$  and a polarization  $p_{e_2}$  the electron beam undergoing Compton scattering, the cross-section can be written in terms of the projection operators

$$P_{ij} = \frac{1}{4} (1 + (-1)^{i} \xi_{2}) \cdot (1 + (-1)^{j} p_{e_{2}}) \text{ as}$$

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^{2}} \int \frac{f(x) dx}{xs} \left[ P_{++} |M_{++++}(xs, xt)|^{2} + P_{+-} |M_{+-+-}(xs, xt)|^{2} + P_{-+} |M_{-+-+}(xs, xt)|^{2} + P_{-+} |M_{----}(xs, xt)|^{2} \right]$$

$$= \frac{1}{128\pi^{2}} \int \frac{f(x) dx}{xs} \left[ \left( \frac{1 + \xi_{2} p_{e_{2}}}{2} \right) |M_{++++}(xs, xt)|^{2} + \left( \frac{1 - \xi_{2} p_{e_{2}}}{2} \right) |M_{+-+-}(xs, xt)|^{2} \right]$$
(12)

where we have made use of parity invariance. For computational purposes, we restrict the x integration in equation (12) to  $x\epsilon[0.1, x_{max}]$  and for computing the total cross-sections from equation (12), we made the cut  $\theta_{CM}\epsilon[\frac{\pi}{6}, \frac{5\pi}{6}]$  as has been done in [4]. Figures 2-4 are our results for the unpolarized and polarized cross-sections for a range of values of the parameter  $(\frac{k}{M_{pl}})$ . For comparison, the corresponding SM-values and the values based on ADD-scenarios as calculated in [4] for n=4 and  $M_s = 2$  TeV are also given.

The cross-sections in the present case as a function of s show an interesting pattern. At low s, the cross-section remains below the SM value, with the difference increasing with the parameter  $k/M_{pl}$ . This is because in the present case both the SM and the graviton exchange contributions are real and of opposite signs. The graviton contribution which increases with increasing  $k/M_{pl}$  thus reduces the amplitude and lowers the cross-section. With increasing energy, the graviton contribution starts dominating and for  $k/M_{pl}$  above 0.05 starts dominating the amplitude from  $\sqrt{s} = 1$ TeV resulting in a monotonic behaviour of the cross-section with the parameter  $k/M_{pl}$  in that range. In the case where the graviton enters as a direct channel resonance [5], the situation is different. There the SM contribution and the graviton contribution add up constructively throughout resulting in a monotonic dependence of the cross-section with the parameter  $k/M_{pl}$ .

Our results show that for values of the parameter  $k/M_{pl}$  equal to 0.10, the cross-sections are much higher than for SM as for SM with the inclusion of ADD grvitons, as  $\sqrt{s}$  exceed 1 TeV. However, for a lower value of  $k/M_{pl}=0.05$ , the deviation from SM does not occur till  $\sqrt{s} = 1.5$ TeV. For even lower values of  $k/M_{pl}$  deviations from SM would occur at even higher value of s. As discussed in reference 5, arguments based on heterotic string theory favour  $k/M_{pl} = 0.01$  and our results show that for such low values of the parameter, differences with SM would be insignificant all the way up to  $\sqrt{s} = 2$  TeV. This is similar to the direct observation signal of resonant peaks in the reaction  $e^+e^- \rightarrow \mu^+\mu^-[5]$ , where for  $k/M_{pl} = 0.01$  the resonant peak is about ten times the background; for a lower  $k/M_{pl} = 0.01$ , the peak practically disappears.

The polarized cross-section results show that the cross-section for the (+, -, +) combination of  $(p_{e1}, p_{l1}, p_{l2})$  dominates over the other combinations in the TeV range. This is expected since the energy distribution of the backscattered photons is strongly peaked for the polarization combination  $p_{e1}, p_{l1} = -1$  [7].

This, combined with the helicity dependence of the amplitudes (12) makes the (+, -, +) cross-sections the largest one at energies in the TeV range.

The signature of the presence of spin-2 graviton exchange is different in the present case than the one for  $e^+e^- \rightarrow \mu^+\mu^-$  [5]. The angular dependence of the cross-section, once the KK-term starts dominating is peaked in the forward direction unlike the pure SM case, where the differential cross-section is peaked in the backward direction. This feature of course is shared also in the ADD scenario. However, for high values of  $k/M_{pl}$  where the cross-sections in the RS scenario are much higher than in the ADD scenario, the pattern of the forward peaking in the two cases is different. Wheras in the ADD case, the differential cross-section in the forward direction to the one at  $\theta = \frac{\pi}{2}$  is a factor around 3, in the RS-case, it is over 10. Hopefully, this feature may help discriminate between the two models if high energy cross-sections reveal the presence of spin-2 graviton exchange in the *t*-channel.

In terms of significance of the cross-section experimentally, the feature of the total cross-section that to begin with at the low energy side, the cross-section should be lower than the SM value and then should start increasing, ultimately becoming greater than the SM value should be the most interesting one. The exact values where the changeover occurs of course depends very much on the parameter  $k/M_{pl}$ . Formally, we can calculate a  $\chi^2$  defined as

$$\chi^2 = L \frac{(\sigma(SM + RS) - \sigma(SM))^2}{\sigma(SM)}$$
(13)

where L is the integrated luminosity over the period of observation and the  $\sigma$ 's refer to cross-sections in the present model and in SM. The value of  $\chi^2$  is of course a function of s and assuming a value of  $L = 100(fb)^{-1}$ , we get the values as listed in table I. Noting that a 95% confidence level implies a value of  $\chi^2$  around 2.7, it seems from the table that it will be possible to see a signature of the RS-graviton resonances in electron compton scattering in the TeV range if the value of the parameter is not too low.

In conclusion, experimental study of electron Compton scattering in the TeV range which is feasible with laser backscattering of electron beams in the NLC would be very useful for revealing the direct coupling to  $\gamma\gamma$  of spin-2 Kaluza-Klein type excitations present in Weak scale Quantum gravity theories. This feature will not be shared in other scenarios like supersymmetry extensions of SM for new Physics at the TeV scale.

S.R.Choudhury thanks Prof.B.McKellar and the School of Physics, University of Melbourne for their hospitality.

## References

 N.Arkani-Hamed, S.Dimopoulos and G.Dvali, Phys.Lett. B429, 263 (1998);
 I.Antoniadis, N.Arkani-Hamed, S.Dimopoulos and G.Dvali, Phys.Lett. B436, 257 (1998); N.Arkani-Hamed, S.Dimopoulos and J.March-Russell, hep-th/9809124.



Figure 2: Unpolarized cross-sections for  $k/M_{pl}=0.01$ , 0.05 and 0.10 together with the SM and SM+ADD values. The ADD contribution corresponds to the choice n=4,  $M_s = 2$ TeV, and  $\omega = +1$ .

- [2] L.Randall and R.Sundrum, Phys.Rev.Lett., 83, 3370 (1999);
   Phys.Rev.Lett., 83, 4690 (1999).
- [3] Tao Han, J.D.Lykken and R.J.Zhang, Phys.Rev. D59, 105006 (1999).
- [4] H.Davoudiasl, Phys.Rev. D61, 044018 (2000).
- [5] H.Davoudiasl, J.L.Hewett and T.G.Rizzo, Phys.Rev.Lett., 84, 2080 (2000).
- [6] V.Telnov, Nucl.Instr. and Meth. A294, 72 (1990).
- [7] Ginzburg et. al., Nucl. Instr. and Meth. A205, 47 (1983); A219, 5 (1984).



Figure 3: The total polarized cross-sections as a function of energy: (a) and (b) correspond to the RS model with  $k/M_{pl}=0.05$  and 0.10 respectively, (c) corresponds to SM and SM+ADD value (evaluated with n=4,  $M_s=2$ TeV and  $\omega=+1$ ).



Figure 4: Unpolarized differential cross-section at  $\sqrt{s} = 1.5$ TeV in the RS model for  $k/M_{pl} = 0.10$  (a) and 0.05 (b). For comparison, the SM value as well as value for SM+ADD (evaluated with n=4,  $M_s = 2$ TeV and  $\omega = +1$ ) are shown.



Figure 5: Polarized differential cross-section for the dominant combination  $p_{e1}, p_{l1}, p_{e2}$  (+, -, +) at  $\sqrt{s} = 1.5$ TeV for  $k/M_{pl}=0.10$  (a) and 0.05 (b). For comparison SM and SM+ADD values (evaluated for parameters as in figure 4) are also shown.

$k/M_{pl}$	$\sqrt{s}=0.5 \text{ TeV}$	$\sqrt{s}$ =1.0 TeV
0.01	$10^{-3}$	0.04
0.05	1.64	8.41
0.10	16.01	326

Table 1: Table showing  $\chi^2$  values for two values of s, calculated through equation (13) for the cross sections evaluated in the present calculation including SM and RS-model KK exchange contribution.