

Low energy hadronic contribution to the QED vacuum polarization

H. Burkhardt*

CERN, AB Departement, CH-1211 Geneva 23, Switzerland

B. Pietrzyk

Laboratoire de Physique des Particules LAPP, IN2P3-CNRS, F-74941 Annecy-le-Vieux Cedex, France †

(Dated: June 30, 2005)

Recent improvements in the low energy e^+e^- annihilation data and their influence on the determination of the hadronic contribution to the running of the QED fine structure constant at m_Z are discussed. Using CMD-2 and KLOE measurements in the ρ region we obtain $\Delta\alpha_{\text{had}}^{(5)}(s) = 0.02758 \pm 0.00035$ at $s = m_Z^2$.

PACS numbers: 13.85.Lg, 13.66.Jn, 12.15.Lk

In the year 2001, we published an updated evaluation of the hadronic contribution to the running of the QED fine structure constant [1], based on a dispersion integral using a parametrization of the measured cross section of $e^+e^- \rightarrow$ hadrons. We obtained a hadronic contribution of $\Delta\alpha_{\text{had}}^{(5)}(s) = 0.02761 \pm 0.00036$ at $s = m_Z^2$.

Our parametrization in the c.m.s. energy region of the ρ , the contribution of the $\pi^+\pi^-$ final state from threshold to 1.8 GeV, was based on a pion form factor parametrization obtained by the CMD-2 Collaboration which used results of their measurements in the c.m.s. energy region between 0.61 and 0.96 GeV at the VEPP-2M collider [2]. The overall uncertainty of the ρ region integral, including the statistical uncertainty, was 2.3% (that of Γ_{ee} in [2]) in our analysis.

Since then, the CMD-2 collaboration improved the treatment of radiative corrections twice. An intermediate improvement has appeared in the published document [3] and an additional improvement has become available in 2004 [4]. We have concluded that the most recent CMD-2 results imply only a small change in the estimate of the hadronic contribution [5].

Recently, the KLOE collaboration [6] has measured the cross section of $e^+e^- \rightarrow \pi^+\pi^-$ with high statistical accuracy in small energy bins using the “radiative return” from the ϕ resonance to the ρ in the $\pi^+\pi^-$ mass range between 0.59 and 0.97 GeV.

We have been repeatedly asked to update our previous analysis and to comment on and quantify the influence of recent low energy measurements by KLOE and CMD-2 on our results. We find that the actual change turns out to be very small. Since the change is very small we have decided to submit this work as a brief report. To the extent that this report is an update of a previously published article, the choice is not to unnecessarily repeat the discussion for energy regions which did not change.

In our 2001 analysis, we used the parametrization of the pion form factor obtained by the CMD-2 collabora-

tion. The contribution of the new results on $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ is now obtained by direct integration between measured KLOE and CMD-2 data points separately. For CMD-2 we use the “bare” cross section and for KLOE the pion form factor data. These are quantities, in which the vacuum polarization corrections have been removed. The small ρ contribution from lower and higher energies, not covered by new data, is evaluated as previously using the CMD-2 parametrization of the pion form factor. We treat the systematic uncertainties as fully correlated between different c.m.s. energies within the CMD-2 experiment. For the integration of the KLOE data, we constructed a covariance matrix based on the statistical covariance matrix with the addition of fully correlated systematic uncertainties as provided by the KLOE collaboration.

The results obtained from the dispersion integration of the KLOE and CMD-2 data at m_Z^2 are in good agreement with each other. The systematic uncertainty of the CMD-2 integration (0.6%) is smaller than the corresponding uncertainty of the KLOE integration (1.4%). On the other hand, the statistical uncertainty of the CMD-2 integration is slightly larger than the systematic one, while the statistical uncertainty of the KLOE integration is negligible. The integration results are combined as independent measurements in the evaluation of the ρ contribution to $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$.

We obtain a value of the hadronic contribution to the running of the QED fine structure constant of $\Delta\alpha_{\text{had}}^{(5)}(s) = 0.02758 \pm 0.00035$ at $s = m_Z^2$ corresponding to $1/\alpha^{(5)}(m_Z^2) = 128.940 \pm 0.048$. The value of the ρ contribution has changed from 0.00350 in [1] to 0.00347 and the relative uncertainty has decreased from 2.3% to 0.9%. The change of the uncertainty corresponds to the change of precision from the preliminary CMD-2 [2] data to the combination of published CMD-2 [4] and KLOE [6] data. The change of the value and the uncertainty of the hadronic contribution to the running of the QED fine structure constant at m_Z^2 is very small. In fact the ρ region contributes to less than 13% to the dispersion integral and is known to much better precision than many of the other energy domains as can be concluded from the Table I, which is the updated version from the Ref. [1].

*Electronic address: Helmut.Burkhardt@cern.ch

†Electronic address: pietrzyk@lapp.in2p3.fr




FIG. 1: R_{had} including resonances. Measurements are shown with statistical errors. The relative uncertainty assigned to our parametrization is shown as band and given with numbers at the bottom.

We note however, that the shape of the hadronic cross-sections measured by the KLOE and CMD-2 collaborations differ for some individual points by more than the systematic uncertainty would indicate [7]. There also appears to be a small, but systematic energy shift in

the observed cross sections between the KLOE and the CMD-2 data, which at present is not understood. The effect on the integrated cross sections which contribute to $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$ is however negligible.

The situation is different for the hadronic contribution to the anomalous magnetic moment of the muon $(g-2)_\mu$ [8, 9, 10, 11]. There the ρ region provides the dominant contribution. The dispersion integral involves a different kernel which gives more weight to lower energies and larger sensitivity on systematic energy shifts.

Fig. 1, which is the updated version from the Ref. [1], gives the summary of R_{had} measurements by different experiments and the current precision in different e^+e^- center-of-mass (cms) energy regions. R_{had} is the measured QED cross-section of the process $e^+e^- \rightarrow \text{hadrons}$, normalized to the QED cross-section for lepton-pair production. The uncertainty in the 1-2 GeV energy region is 15%. This region contributes to about 40% to the uncertainty on dispersion integral at m_Z^2 , as can be seen from

TABLE I: Contributions to $\Delta\alpha_{\text{had}}^{(5)}(m_Z^2)$

Range \sqrt{s} , GeV	$\Delta\alpha$	Relative error
ρ	0.00347	0.9 %
Narrow resonances	0.00184	3.1 %
1.05 – 2.0	0.00156	15 %
2.0 – 5.0	0.00381	5.9 %
5 – 7	0.00183	6 %
7 – 12	0.00304	1.4 %
> 12	0.01203	0.2 %
	0.02758	1.3 %

the Table I and Fig. 2 in the Ref. [1]. We would like to strongly encourage efforts to measure precisely R_{had} in this cms energy region.

We would like to thank S.I. Eidelman from the CMD-2 Collaboration and A. Denig and G. Venanzoni from the KLOE Collaboration for useful discussions.

[1] H. Burkhardt and B. Pietrzyk, Phys. Lett. **B513**, 46 (2001).
[2] R.R. Akhmetshin et al., CMD-2 Collaboration, hep-ex/9904027.
[3] R.R. Akhmetshin et al., CMD-2 Collaboration, Phys. Lett. **B527**, 161 (2002).
[4] R.R. Akhmetshin et al., CMD-2 Collaboration, Phys. Lett. **B578**, 285 (2004).
[5] B. Pietrzyk and H. Burkhardt, preprint LAPP-EXP 2003-07, contribution to the Int. Europhysics Conf. on HEP, Aachen, 2003.

[6] A. Aloisio et al., Phys. Lett. **B606**, 12 (2005).
[7] see, for example, talks of G. Venanzoni and F. Teubert at 32nd Int. Conf. on HEP, Beijing, 2004.
[8] M. Davier, S. Eidelman, A. Hoecker and Z. Zhang, Eur. Phys. J. **C31**, 503 (2003).
[9] F. Jegerlehner, Nucl. Phys. B (Proc. Suppl.) **126**, 325 (2004); **131**, 213 (2004).
[10] T. Teubner, Nucl. Phys. B (Proc. Suppl.) **131**, 201 (2004).
[11] M. Davier, Nucl. Phys. B (Proc. Suppl.) **144**, 250 (2005).