

DILEPTON PRODUCTION IN NON-EQUILIBRATED QUARK-GLUON PLASMA.

S.S.Singh¹ and Agam K. Jha^{1*}

Department of Physics, University of Delhi, Delhi - 110007,INDIA

A model of cut-off momentum distribution functions in quark-gluon plasma with finite baryon chemical potential is discussed. This produces a quark-gluon plasma signature in relativistic nuclear collisions with the specific structure of the dilepton spectrum in the transverse momentum region of (1-4) GeV and the dilepton production rate is found to be a strong decreasing function of chemical potential.

Quantum chromodynamics(QCD), the fundamental theory of strong interactions[1], predicts a phase transition from the state of hadronic constituents to a phase of deconfined quarks and gluon, so called Quark Gluon Plasma (QGP). Presumably the early universe was in this state up to about $10\mu s$ after the big bang. Today the core program of ultrarelativistic nucleus-nucleus collision is to study the properties of the strongly interacting matter at very high energy density and temperature so that it can create this deconfined state of matter in heavy ion collision experiments. This new state of matter emits a large number of particles. Among these particles, electromagnetic probes which are known as photons leave this plasma volume without further rescattering and it carries the whole information about the existence of the plasma. This produced virtual photons are subsequently decayed into dileptons such as l^-l^+ , $\mu^+\mu^-$ and they represent as a possible signature for the formation of QGP.

So far, many authors have studied the production of dilepton in a QGP at finite temperature and this temperature is related to the energy density ϵ given by the Stefan Boltzmann $\epsilon \sim T^4$. However experiments at AGS and SPS energies[2] indicate that there is a sizable amount of chemical potential and even at RHIC energies $\sqrt{s} \leq 200 A GeV$, this chemical potential μ may not be neglected and calculation using microscopic models[3, 4] limit that the colliding heavy ions may not be fully transparent. In this case the dilepton production rate is in (local) thermodynamic equilibrium, is a function of both temperature and quark chemical potential μ of the QGP which is recently studied by Dumitru et. al.[5]. Again, recent work by Hammon and coworkers[6] have indicated that the initial QGP system produced at the RHIC energies have finite baryon density. Mujumder et. al.[7] have discussed the dileptons from QGP produced at the RHIC energies at finite baryon density and Bass et. al. [8] have pointed out that the parton rescattering and fragmentation lead to a substantial increase in the net -baryon density at midrapidity region. Now we consider this influencing quark chemical potential for one dilepton production. As stated above the Stefan-Boltzmann relation $\epsilon \sim T^4$ is free from the relation $\frac{\mu}{T}$. We use EOS for the

plasma which relates ϵ, T and μ . We consider the grand partition function for fermion (particle and anti particle) and boson for finding EOS of relativistic hydrodynamical model. Thermodynamic potentials for fermion and boson are defined in rich baryon QGP as:

$$(T \ln z)_f = \left(\frac{g_f V}{12} \right) \times \left(\frac{7\pi^2}{30} T^4 + \mu^2 T^2 + \frac{1}{2\pi^2} \mu^4 \right) \quad (1)$$

$$(T \ln z)_b = \frac{g_b V \pi^2}{90} T^4 \quad (2)$$

where g_f and g_b are the respective degeneracies that is degree of freedom. The energy density and number density are obtained through this grand partition function. They are $n = \frac{1}{V} \partial(T \ln Z) / \partial \mu$, $\epsilon = T \partial(T \ln Z) / \partial T + \mu n$ and we can easily establish the equation of state (EOS) of the ideal gas $p = \frac{\epsilon}{3}$ but this QGP is far from the ideal gas at temperature as high as several times T_c . In this paper we study the dilepton production of the expanding hot baryon-rich QGP on the basis of cut off momentum distribution function which are used by Gorenstein et. al.[9] in the calculation of dilepton production in small invariant mass system in free baryonic system.

Dilepton production from QGP: There are a good number of research works in dilepton production into the quark-gluon plasmas. Whenever there is a new model for plasma evolution then its impact on dilepton and virtual photon production are assessed. The dominant reaction for thermal emission of dilepton pairs[10] is the Drell-Yan mechanism $q\bar{q} \rightarrow l^+l^-$ or $q(\bar{q})g \rightarrow q(\bar{q}) + l^+l^-$. But, we exclusively use $q\bar{q} \rightarrow l^+l^-$ reaction. The dilepton production rate $Y = \frac{dR}{d^4P}$ (i.e the number of dilepton produced per space time volume and four dimensional momentum space volume) is given by

$$\frac{dR}{d^4P} = \int \frac{d^3k_1}{(2\pi)^3} \frac{d^3k_2}{(2\pi)^3} f_q(k_1, T, \mu) f_{\bar{q}}(k_2, T, \mu) \times v_{q\bar{q}} \sigma_{q\bar{q}}(M^2) \delta^4(P - k_1 - k_2) \quad (3)$$

where,

$$f_q(k_1, T, \mu) = \frac{d_q \Theta(|k| - K_q)}{\exp\left(\frac{E-\mu}{T}\right) + 1}, f_{\bar{q}}(k_2, T, \mu) = \frac{d_{\bar{q}} \Theta(|k| - K_q)}{\exp\left(\frac{E+\mu}{T}\right) + 1} \quad (4)$$

*Electronic address: sssingh@physics.du.ac.in, akjha@physics.du.ac.in.




FIG. 1: The dilepton emission rate, $Y = \frac{dR}{d^4P}$, $\log(Y)$ at parameter $M = 1.0 \text{ GeV}$, $T = 0.25 \text{ GeV}$, $K_q = 0.6 \text{ GeV}$ for different values of μ

at equilibrium, these two distribution functions are equal

$$f_q(k_1, T, \mu = 0) = f_{\bar{q}}(k_2, T, \mu = 0) \quad (5)$$

and

$$f_g(k, T, \mu) = \frac{d_g \Theta(|k| - K_g)}{\exp\left(\frac{E_g}{T}\right) - 1} \quad (6)$$

where d_g and d_q are the degeneracy factor for gluon and quarks. $v_{q\bar{q}}$ is the relative velocity between quark and antiquark. p_μ is lepton pair four momentum. ($M^2 = p^\mu p_\mu$ invariant lepton mass). $\sigma_{q\bar{q} \rightarrow l\bar{l}}$ is the electromagnetic annihilation cross section. It is given:

$$\sigma_{q\bar{q} \rightarrow l\bar{l}}(M^2) = 16\pi\alpha^2 \sum_{f=1}^2 \left(\frac{e_f}{e}\right)^2 \frac{1}{M^2} \left(1 + \frac{2m^2}{M^2}\right) \left(1 - \frac{4m^2}{M^2}\right). \quad (7)$$

where $\left(\frac{e_f}{e}\right)^2 = \frac{5}{9}$ the electric charge of the quark in the units of the electron charge, $\alpha = \frac{1}{137}$ and m is the lepton mass but we consider the lepton mass is zero (ie $m = 0$). Substituting in the equation (3) using (4) and (6) we integrate over q and \bar{q} momentum and obtain the dilepton production rate as:

$$\frac{dR}{d^4P} = \frac{5\alpha^2}{18\pi^3} \frac{T\Theta(|E| - 2K_q)}{(\exp\left(\frac{E}{T}\right) - 1)} \ln(1 - \exp^{-(E+\mu)/T}) \sqrt{\frac{\pi MT}{2}} \quad (8)$$




FIG. 2: The dilepton emission rate, $Y = \frac{dR}{d^4P}$, $\log(Y)$ at parameter $M = 3.0 \text{ GeV}$, $T = 0.25 \text{ GeV}$, $K_q = 0.6 \text{ GeV}$ for different values of μ




FIG. 3: The dilepton emission rate, $Y = \frac{dR}{d^4P}$, $\log(Y)$ at parameter $P = 1.0 \text{ GeV}$, $T = 0.25 \text{ GeV}$, $K_q = 0.6 \text{ GeV}$ for different values of μ

Results and Conclusions: The results are shown in the figures. This result is consistent with the results produced by Ref[9] and that of Ref[11] for $\mu = 0$ which is free from the chemical potential. The cut off in the momentum distribution function is represented by the threshold factor $\Theta(E - 2K_q)$ in the dilepton production. Now, in




FIG. 4: The dilepton emission rate, $Y = \frac{dR}{d^4P}$, $\log(Y)$ at parameter $P = 3.0\text{GeV}$, $T = 0.25\text{GeV}$, $K_q = 0.6\text{GeV}$ for different values of μ

the figure 1, for small lepton pair mass $M = 1.0\text{GeV}$ and quark momentum cut off $K_q = 0.6\text{GeV}$ we obtain a sudden fall in the production rate of the dilepton and at $K_q = 0$ the rate extends upto the level mark for $\mu = 0$ which is shown by solid line. Again in the figure 2 we

increase the lepton pair mass M upto 3.0GeV then the cut-off value of quark momentum $K_q = 0.6\text{GeV}$ does not play any role and sudden fall shown by figure 1 disappear. It implies that the threshold factor $\Theta(E - 2K_q)$ plays role mainly in small invariant lepton pair mass M , not for large M . Similarly it produces the variation of the emission rate with the lepton pair mass in figure 3 for a specific transverse momentum $P_T = 1.0\text{GeV}$. From the analysis we again observed that cut-off in the momentum distribution function plays role mainly in the small invariant mass M and for the large value of M it is not much effective. We obtain this for the different values of $M = (1 - 4)\text{GeV}$ for one fixed value of $P_T = 1.0\text{GeV}$. Again we increase this transverse momentum P_T upto 3GeV and the production rate obtained by this fixed P_T value shows that at higher P_T the rate is highly suppressed corresponding to the value of chemical potential. Lastly we can conclude from this analysis that the rate of production is decreasing function of the increasing in the chemical potential in QGP which implies that in the rich baryon region the anti quark population is strongly suppressed for $\frac{\mu}{T} > 0$. The result is quiet similar with the results produced by Dimutru et. al.[5] for different values of $\mu/T = 0, 1, 2, 3$. So the effect of this evolution of the system at finite baryon density on lepton production is very important.

Acknowledgments

We are very thankful to Dr. R. Ramanathan for constructive suggestions and discussions.

-
- [1] F.Wilczek, hep-ph/003183; D.E.Kharzeev, J.Ranfeisen, nucl-th/0206073.
[2] Nagamiya S 1992 Nucl.Phys. A 544, 5c; Busza W 1984 Nucl. Phys. A 418, 635c, Hansen O 1992 Proc.of the 20th Int. Workshop on Gross Production of Nuclei and nuclear Excitations (Hirschegg, Austria) ed H.Ferdmeier.
[3] Gustafson G 1992 Proc.of the Workshop on Relativistic Heavy-Ion Physics at present and future Accelerators (Budapest).
[4] Mohring H J and Ranft J 1991 z.phys.C 52,643.
[5] Dumitru A, Rischke D H, Schonfed Th, Winkelmann L,

- Stocker H and Greiner W 1993 phys.Rev.Lett. 70 2860.
[6] Hammon et. al. 1999, Phys. Rev. C 61, 014901.
[7] A. Majumdar et. al., 2001 Phys. Rev. D 63, 114008.
[8] S. A. Bass, 2003 Phys. Rev. Lett. 91, 052302.
[9] M.I.Gorenstein and O.A.Mogilevsky 1984 phys. lett.B 228 121.
[10] P.V.Ruuskanen in Quark-Gluon Plasma edited by R.C Hwa world scientific, 1991 singapore.
[11] J. Cleymans, J.Fingberg and K.Redlich 1987 phys.Rev. D 35,2153.