

NRQCD Prediction for the Polarization of the J/ψ Produced from b -decay

Sean Fleming

Department of Physics, University of Wisconsin, Madison, WI 53706, USA

Presented at the Meeting of the Division of Particles and Fields, August 1996, in Minneapolis. The work presented here is based upon a recent paper^a done in collaboration with Oscar F. Hernández, Ivan Maksymyk, and H el ene Nadeau. The NRQCD predictions for the polarization of the J/ψ produced in $b \rightarrow J/\psi + X$, as well as the helicity-summed production rate are presented.

^a(hep-ph/9608413)

A rigorous theoretical framework within which quarkonium production can be studied is provided by the NRQCD factorization formalism¹. This approach is based on NRQCD, an effective field theory that that can be made equivalent to full QCD to any desired order in the heavy-quark velocity v . The NRQCD Lagrangian is

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\text{tr} G_{\mu\nu}G^{\mu\nu} + \sum \bar{\Psi}_\ell i\not{D}\Psi_\ell \\ & + \psi^\dagger \left(iD_0 + \frac{\mathbf{D}^2}{2m_Q} \right) \psi + \chi^\dagger \left(iD_0 - \frac{\mathbf{D}^2}{2m_Q} \right) \chi + \dots \end{aligned} \quad (1)$$

where $G_{\mu\nu}$ is the gluon field-strength tensor, Ψ_ℓ is the Dirac spinor field for a light quark, $D^\mu = \partial^\mu + igA^\mu$ is the gauge-covariant derivative (with g being the QCD coupling constant), m_Q is the heavy quark mass, and ψ and χ are 2-component fields that annihilate heavy quarks and create heavy antiquarks respectively. Color and spin indices on the fields have been suppressed. Eq. 1 describes a fully relativistic field theory for the light degrees of freedom coupled with a Schr odinger field theory for the heavy quarks and antiquarks. The relativistic effects of full QCD are reproduced through additional terms represented by the \dots in Eq. 1. In principle there are infinitely many terms. However, using NRQCD v -scaling rules it is possible to retain only those that contribute through a given order in v .

The NRQCD factorization formalism is a rigorous derivation within NRQCD of a factored form for quarkonium production and decay rates. A central result

is that inclusive quarkonium production cross sections must have the form

$$\sigma(A + B \rightarrow H + X) = \sum_n \frac{F_n}{m_Q^{d_n-4}} \langle \mathcal{O}_n^H \rangle, \quad (2)$$

where the \mathcal{O}_n^H represent NRQCD four-fermion production operators, with the index n labeling color and angular-momentum quantum numbers. The NRQCD matrix elements $\langle \mathcal{O}_n^H \rangle \equiv \langle 0 | \mathcal{O}_n^H | 0 \rangle$, parameterize the hadronization of a heavy-quark-antiquark pair with quantum numbers n into a quarkonium state H . They scale as the parameter v raised to a power. The short-distance coefficients F_n are obtainable in perturbation theory as a series in $\alpha_s(m_Q)$. Information on the order in v of the matrix elements, along with the dependence of the F_n on coupling constants, permits us to decide which terms must be retained in expressions for observables to reach a given level of accuracy.

Consider the specific case of J/ψ production. In many instances the most important NRQCD matrix elements are $\langle \mathcal{O}_1^\psi(3S_1) \rangle$, $\langle \mathcal{O}_8^\psi(3S_1) \rangle$, $\langle \mathcal{O}_8^\psi(1S_0) \rangle$, and $\langle \mathcal{O}_8^\psi(3P_J) \rangle$. Inclusive production of J/ψ from b -decay provides two measurable combinations of these matrix elements. The first one is the helicity-summed rate $\Gamma(b \rightarrow J/\psi + X)$. The second combination concerns the polarization parameter α appearing in the electromagnetic decay rate of J/ψ to lepton pairs:

$$\frac{d\Gamma}{d\cos\theta}(b \rightarrow J/\psi + \mu^+ \mu^-(\theta)) \propto 1 + \alpha \cos^2 \theta, \quad (3)$$

where the polar angle θ is defined in the J/ψ rest frame for which the z -axis is aligned with the direction of motion of the J/ψ in the lab.

The branching ratio for $b \rightarrow J/\psi + X$ is calculated using a matching procedure². One obtains³

$$BR(b \rightarrow J/\psi + X) = 0.002 \langle \mathcal{O}_1^\psi(3S_1) \rangle + 0.1 \langle \mathcal{O}_8^\psi(3S_1) \rangle + 0.2 \langle \mathcal{O}_8^\psi(1S_0) \rangle + 0.6 \frac{\langle \mathcal{O}_8^\psi(3P_0) \rangle}{m_c^2}. \quad (4)$$

Only the leading color-singlet piece and the leading color-octet pieces in the relativistic v^2 -expansion are considered. The above formula concurs with previous results⁴.

According to the NRQCD v -scaling rules, the color-octet matrix elements in Eq. 4 are all expected to be suppressed by v^4 with respect to $\langle \mathcal{O}_1^\psi(3S_1) \rangle$. However the short-distance coefficients in the color-octet terms are some 50 times larger than the color-singlet coefficient!

Beneke and Rothstein⁵ and Braaten and Chen⁶ have developed techniques for deriving the production rates of quarkonia with specified helicities. Applying their methods one obtains an expression for the polarization parameter³

$$\alpha = \frac{-0.39\langle\mathcal{O}_1^\psi(^3S_1)\rangle - 17\langle\mathcal{O}_8^\psi(^3S_1)\rangle + 52\langle\mathcal{O}_8^\psi(^3P_0)\rangle/m_c^2}{\langle\mathcal{O}_1^\psi(^3S_1)\rangle + 44\langle\mathcal{O}_8^\psi(^3S_1)\rangle + 61\langle\mathcal{O}_8^\psi(^1S_0)\rangle + 211\langle\mathcal{O}_8^\psi(^3P_0)\rangle/m_c^2}. \quad (5)$$

While experimental determinations of helicity-summed $BR(b \rightarrow J/\psi + X)$ have already been carried out⁷, a measurement of the polarization parameter α is not yet available. Anticipating the availability of this latter measurement, it is interesting to determine the range of α which is consistent with existing information on the matrix elements $\langle\mathcal{O}_1^\psi(^3S_1)\rangle$, $\langle\mathcal{O}_8^\psi(^3S_1)\rangle$, $\langle\mathcal{O}_8^\psi(^1S_0)\rangle$, and $\langle\mathcal{O}_8^\psi(^3P_0)\rangle/m_c^2$, and with various constraints on linear combinations of these quantities⁸. The *maximum* value for α is -0.09 , and the *minimum* value for α is -0.33 .

Unfortunately, due to the large number of poorly determined parameters the expected range of α as predicted by NRQCD is large. It may be that an accurate NRQCD prediction of the polarization parameter (beyond leading order) will not be possible for a long time. This possibility, however, does not degrade the importance of this calculation, since an experimental determination of α — in conjunction with these results — will most certainly serve to tighten the constraints on the possible values of the color-octet matrix elements. In fact, the NRQCD prediction for α is very sensitive to the values of the color-octet matrix elements. This offers the hope that a measurement of the polarization of J/ψ produced in $b \rightarrow J/\psi + X$ will be instrumental in determining $\langle\mathcal{O}_8^\psi(^3S_1)\rangle$, $\langle\mathcal{O}_8^\psi(^1S_0)\rangle$, and $\langle\mathcal{O}_8^\psi(^3P_0)\rangle$.

This work was supported by the U.S. D.O.E. under Grant No. DE-FG02-95ER40896, by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

References

1. G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. **D51**, 1125 (1995), and references therein.
2. S. Fleming and I. Maksymyk, Phys.Rev. **D54**, 3608 (1996).
3. S. Fleming, O. Hernández, I. Maksymyk, and H. Nadeau preprint (hep-ph/9608413); I. Maksymyk preprint UTTG-12-96, in preparation.
4. P. Ko, J. Lee, H.S. Song, Phys. Rev. **D53**, 1409 (1996).
5. M. Beneke and I.Z. Rothstein, Phys. Lett. **B372**, 157 (1996); Phys. Rev. **D54**, 2005 (1996).

6. E. Braaten and Y.-Q. Chen, Phys.Rev. **D54**, 3216 (1996).
7. R.M. Barnett et al., Phys. Rev. **D54**, 1 (1996).
8. P. Cho and A. K. Leibovich, Phys. Rev. **D53**, 6203 (1996); G.T. Bodwin, D.K. Sinclair and S. Kim, preprint (hep-lat/9605023); J. Amundson, S. Fleming, and I. Maksymyk, preprint (hep-ph/9601298).