

Tensor Products of Some Special Rings¹

Masoud Tousi^{ab} and Siamak Yassemi^{ac}

(a) *Institute for Studies in Theoretical Physics and Mathematics*

(b) *Department of Mathematics, Shahid Beheshti University*

(c) *Department of Mathematics, University of Tehran.*

Abstract

In this paper we solve a problem, originally raised by Grothendieck, on the properties, i.e. Complete intersection, Gorenstein, Cohen–Macaulay, that are conserved under tensor product of algebras over a field k .

1991 Mathematics subject classification. 13H10.

Key words and phrases. Regular; Complete intersection; Gorenstein; Cohen–Macaulay; flat homomorphism of rings.

0. Introduction

Throughout this note all rings and algebras considered in this paper are commutative with identity elements, and all ring homomorphisms are unital. Throughout, k stands for a field.

Among local rings there is a well-known chain:

Regular \Rightarrow Complete intersection \Rightarrow Gorenstein \Rightarrow Cohen–Macaulay.

These concepts are extended to non-local rings: for example a ring is regular if for all prime ideal \mathfrak{p} of R , $R_{\mathfrak{p}}$ is a regular local ring.

In this paper, we shall investigate if these properties are conserved under tensor product operations. It is well-known that the tensor product $R \otimes_A S$ of regular rings is not

¹This research was supported in part by a grant from IPM.

regular in general, even if we assume R and S are A -algebra and A is a field, see Remark 1.7. In [5], Watanabe, Ishikawa, Tachibana, and Otsuka, showed that under a suitable condition tensor products of regular rings are complete intersections. It is proved in [3], that the tensor product $R \otimes_A S$ of Cohen–Macaulay rings are again Cohen–Macaulay if we assume R is flat A -module and S is a finitely generated A -module, and in [5], it is shown that the same is true for Gorenstein rings. Recently, in [1], Bouchiba and Kabbaj showed that if R and S are k -algebras such that $R \otimes_k S$ is Noetherian then $R \otimes_k S$ is a Cohen–Macaulay ring if and only if R and S are Cohen–Macaulay rings.

In this paper we shall show that the same is true for complete intersection and Gorenstein rings. Also it is shown that $R \otimes_k S$ satisfies Serre’s condition (S_n) if and only if R and S satisfy (S_n) .

1. Main results

A Noetherian local ring R is a complete intersection (ring) if its completion \hat{R} is a residue class ring of a regular local ring S with respect to an ideal generated by an S -sequence. We say that a Noetherian ring is locally a complete intersection if all its localizations are complete intersections.

A Noetherian ring R satisfies Serre’s condition (S_n) if $\text{depth } R_{\mathfrak{p}} \geq \text{Min}\{n, \dim R_{\mathfrak{p}}\}$ for all prime ideal \mathfrak{p} of R . Also, a Noetherian ring R satisfies Serre’s normality condition (R_n) if $R_{\mathfrak{p}}$ is a regular local ring for all prime ideal \mathfrak{p} with $\dim R_{\mathfrak{p}} \leq n$.

The following Theorem is collected from [2; Remark 2.3.5, Corollary 3.3.15, Theorem 2.1.7, and Theorem 2.2.12]:

Theorem 1.1. Let $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ be a flat local homomorphism of Noetherian local rings. Then the following hold:

- (a) S is a complete intersection (resp. Gorenstein, Cohen–Macaulay) $\Leftrightarrow R$ and $S/\mathfrak{m}S$ are complete intersections (resp. Gorenstein, Cohen–Macaulay).
- (b1) If S is regular then R is regular.

(b2) If R and $S/\mathfrak{m}S$ are regular then S is regular. □

Corollary 1.2. Let $\varphi : R \rightarrow S$ be a flat homomorphism of Noetherian rings. Then the following hold:

- (a) If R and the fibers $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_R S$, $\mathfrak{p} \in \text{Spec}(R)$, are regular (resp. locally complete intersections, Gorenstein, Cohen–Macaulay) then S is regular (resp. locally complete intersection, Gorenstein, Cohen–Macaulay).
- (b) If S is locally complete intersection (resp. Gorenstein, Cohen–Macaulay) then the fibres $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_R S$, $\mathfrak{p} \in \text{Spec}(R)$, are locally complete intersections (resp. Gorenstein, Cohen–Macaulay)

Proof. (a): Let $\mathfrak{q} \in \text{Spec}(S)$. Set $\mathfrak{p} = \mathfrak{q} \cap R \in \text{Spec}(R)$. The induced homomorphism $\tilde{\varphi} : R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$ is flat and local. It is clear that $S_{\mathfrak{q}}/\mathfrak{p}R_{\mathfrak{p}}S_{\mathfrak{q}}$ is a localization of $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_R S$. Now the assertion follows from Theorem 1.1.

(b): Let $\mathfrak{p} \in \text{Spec}(R)$. Then $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_R S \cong S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}}$, where $S_{\mathfrak{p}} = T^{-1}S$ and $T = R - \mathfrak{p}$, and we have

$$\text{Spec}(S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}}) = \{\mathfrak{q}S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}} \mid \mathfrak{q} \in \text{Spec}(S), \mathfrak{q} \supseteq \mathfrak{p}S, \mathfrak{q} \cap (R - \mathfrak{p}) = \emptyset\}.$$

For $\mathfrak{q}S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}} \in \text{Spec}(S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}})$ we have to show that $(S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}})_{\mathfrak{q}S_{\mathfrak{p}}/\mathfrak{p}S_{\mathfrak{p}}} \cong S_{\mathfrak{q}}/\mathfrak{p}S_{\mathfrak{q}}$ is complete intersection (resp. Gorenstein, Cohen–Macaulay). Consider the induced flat local homomorphism $\tilde{\varphi} : R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$. Now the assertion follows from Theorem 1.1. □

Theorem 1.3. (See [2; Propositions 2.1.16 and 2.2.21]) Let $\varphi : R \rightarrow S$ be a flat homomorphism of Noetherian rings. Then the following hold:

- (a) Let $\mathfrak{q} \in \text{Spec}(S)$ and $\mathfrak{p} = \mathfrak{q} \cap R$. If $S_{\mathfrak{q}}$ satisfies (S_n) (resp. (R_n)) then $R_{\mathfrak{p}}$ satisfies (S_n) (resp. (R_n)).
- (b) If R and the fibers $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_R S$, $\mathfrak{p} \in \text{Spec}(R)$, satisfy (S_n) (resp. (R_n)) then S satisfies (S_n) (resp. (R_n)). □

Corollary 1.4. Let $\varphi : R \rightarrow S$ be a faithfully flat homomorphism of Noetherian rings. Then the following hold:

- (a) If S is regular (resp. locally complete intersection, Gorenstein, Cohen–Macaulay), then so is R .
- (b) If S satisfies (S_n) (resp. (R_n)), then so does R .

Proof. Let $\mathfrak{p} \in \text{Spec}(R)$. Since φ is faithfully flat there exists $\mathfrak{q} \in \text{Spec}(S)$ such that $\mathfrak{p} = \mathfrak{q} \cap R$. Consider the flat local homomorphism $\tilde{\varphi} : R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$ where $\tilde{\varphi}(r/s) = \varphi(r)/\varphi(s)$. Now the assertion follows from Theorems 1.1 and 1.3. \square

Proposition 1.5. Let k be a field, L and K be two extension fields of k . Suppose that $L \otimes_k K$ is Noetherian. Then the following hold:

- (a) $L \otimes_k K$ is locally complete intersection.
- (b) If k is perfect then $L \otimes_k K$ is regular.

Proof. (a): With the same method in the proof of [4; Theorem 2.2], we can assume that K is a finitely generated extension field of k (note that, in view of Theorem 1.1, [4; Lemma 2.1] is true with “Gorenstein ring” replaced by “complete intersection”). Now using [2; Proposition 2.1.11] we have that $L \otimes_k K$ is isomorphic to

$$A = T^{-1}(L[x_1, x_2, \dots, x_n]) / (f_1, f_2, \dots, f_m) T^{-1}(L[x_1, x_2, \dots, x_n]),$$

where T is a multiplicatively closed subset of $L[x_1, x_2, \dots, x_n]$ and f_1, f_2, \dots, f_m is a $T^{-1}(L[x_1, x_2, \dots, x_n])$ –sequence. Therefore A is locally complete intersection, cf. [2; Theorem 2.3.3(c)].

(b): The assertion follows from the note on page 49 of [4]. \square

Theorem 1.6. Let R and S be non-zero k –algebras such that $R \otimes_k S$ is Noetherian. Then the following hold:

- (a) $R \otimes_k S$ is locally complete intersection (resp. Gorenstein, Cohen–Macaulay) if and only if R and S are locally complete intersections (resp. Gorenstein, Cohen–Macaulay).
- (b) $R \otimes_k S$ satisfies (S_n) if and only if R and S satisfy (S_n) .
- (c) If $R \otimes_k S$ is regular then R and S are regular.
- (d) If $R \otimes_k S$ satisfies (R_n) then R and S satisfy (R_n) .
- (e) The converse of parts (c) and (d) hold if $\text{char}(k) = 0$ or $\text{char}(k) = p$ such that $k = \{a^p | a \in k\}$.

Proof. Consider two faithfully flat homomorphism:

$$\varphi : R \rightarrow R \otimes_k S \quad \text{and} \quad \psi : S \rightarrow R \otimes_k S$$

of Noetherian rings.

If $R \otimes_k S$ is regular (resp. locally complete intersection, Gorenstein, Cohen–Macaulay) then by Corollary 1.4 we have R and S are regular (resp. locally complete intersections, Gorenstein, Cohen–Macaulay). Also if $R \otimes_k S$ satisfies (S_n) (resp. (R_n)) then by Corollary 1.4, R and S satisfy (S_n) (resp. (R_n)).

Now let R and S be locally complete intersection (resp. Gorenstein, Cohen–Macaulay). By Corollary 1.2 it is enough to show that the fibres $(R \otimes_k S) \otimes_R R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \cong R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_k S$ over every prime ideal \mathfrak{p} of R is locally complete intersection (resp. Gorenstein, Cohen–Macaulay). Consider the flat homomorphism $\gamma : S \rightarrow R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_k S$. Using Corollary 1.2, it is enough to show that the fibres $(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_k S) \otimes_S S_{\mathfrak{q}}/\mathfrak{q}S_{\mathfrak{q}} \cong R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_k S_{\mathfrak{q}}/\mathfrak{q}S_{\mathfrak{q}}$ over every prime \mathfrak{q} of S is locally complete intersection (resp. Gorenstein, Cohen–Macaulay). But it is clear to see that $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \otimes_k S_{\mathfrak{q}}/\mathfrak{q}S_{\mathfrak{q}}$ is Noetherian, since it is a localization of $R/\mathfrak{p} \otimes_k S/\mathfrak{q} \cong R \otimes_k S/(\mathfrak{p} \otimes_k S + R \otimes_k \mathfrak{q})$, which is Noetherian. Now the assertion follows from Proposition 1.5.

If R and S satisfy (S_n) , with the same proof $R \otimes_k S$ satisfies (S_n) .

By using the Proposition 1.5 the proof of part (e) is the same. □

Remark 1.7. The converse of part (c) in Theorem 1.6 is not true. For example, let k be an imperfect field of characteristic 3, let $a \in k$ be an element with no cube root in k . Then

$K = k[x]/(x^3 - a)k[x]$ is a splitting field of $x^3 - a$ over k . Thus $K \otimes_k K \cong K[x]/(x^3 - a)K[x]$, which is not regular.

Acknowledgment. The authors would like to thank the referee for his/her comments.

References

- [1] Samir Bouchiba and Salah-Eddine Kabbaj, *Tensor products of Cohen-Macaulay rings: solution to a problem of Grothendieck*, J. Algebra **252** (2002), 65–73.
- [2] Winfried Bruns and Jürgen Herzog, *Cohen-Macaulay rings*, Cambridge University Press, Cambridge, 1993.
- [3] A. Grothendieck, *Éléments de géométrie algébrique* IV. Étude locale des schémas et des morphismes des schémas. II. Inst. Hautes Études Sci. Publ. Math., No. 24, 1965.
- [4] Rodney Y. Sharp, *Simplifications in the theory of tensor products of field extensions*. J. London Math. Soc., **15**, (1977), 48–50.
- [5] Kei-ichi Watanabe, Takeshi Ishikawa, Sadao Tachibana, and Kayo Otsuka, *On tensor products of Gorenstein rings*, J. Math. Kyoto Univ., **9** (1969), 413–423.