

Pair Correlation of the zeros of the Riemann zeta function in longer ranges

Tsz Ho Chan

November 20, 2018

Abstract

In this paper, we study a more general pair correlation function, $F_h(x, T)$, of the zeros of the Riemann zeta function. It provides information on the distribution of larger differences between the zeros.

1 Introduction

First of all, we assume the Riemann Hypothesis on the Riemann zeta function $\zeta(s)$ throughout this paper; $\rho = \frac{1}{2} + i\gamma$ denotes a non-trivial zero of the Riemann zeta function.

In the early 1970s, Hugh Montgomery considered the pair correlation function

$$F(x, T) = \sum_{0 < \gamma, \gamma' \leq T} x^{i(\gamma - \gamma')} w(\gamma - \gamma') \text{ with } w(u) = \frac{4}{4 + u^2}.$$

Here the sum is a double sum over the imaginary parts of the non-trivial zeros of $\zeta(s)$. He proved in [9] that, as $T \rightarrow \infty$,

$$F(x, T) \sim \frac{T}{2\pi} \log x + \frac{T}{2\pi x^2} \log^2 T$$

for $1 \leq x \leq T$ (actually he only proved for $1 \leq x \leq o(T)$ and the full range was done by Goldston [5]). He conjectured that

$$F(x, T) \sim \frac{T}{2\pi} \log T$$

for $T \leq x$ which is known as the Strong Pair Correlation Conjecture. From this, one has the (Weak) Pair Correlation Conjecture:

$$\sum_{\substack{0 < \gamma, \gamma' \leq T \\ 0 < \gamma - \gamma' \leq 2\pi\alpha / \log T}} 1 \sim \frac{T}{2\pi} \log T \int_0^\alpha 1 - \left(\frac{\sin \pi u}{\pi u} \right)^2 du.$$

which draws connections with random matrix theory.

The author studied these further in his thesis [1] (see also [2] and [3]) and derived more precise asymptotic formulas for $F(x, T)$ when x is in various ranges under the Twin Prime Conjecture **TPC** (see section 4). In the present paper, we generalize $F(x, T)$ further to

$$F_h(x, T) = \sum_{0 < \gamma, \gamma' \leq T} \cos((\gamma - \gamma' - h) \log x) w(\gamma - \gamma' - h).$$

Note that $F_h(x, T) = F_{-h}(x, T)$ and $F_0(x, T) = F(x, T)$. This leads to a better understanding of the distribution of larger differences between the zeros. Our main results are the following theorems: Here and throughout the paper, $\tilde{h} = |h| + 1$.

Theorem 1.1. For $1 \leq x \leq \frac{T}{\log T}$,

$$\begin{aligned} F_h(x, T) &= \frac{T}{2\pi} \left[\frac{4 \cos(h \log x)}{4 + h^2} \log x - \frac{8h \sin(h \log x)}{(4 + h^2)^2} \right] \\ &\quad + \frac{T}{2\pi x^2} \left[\left(\log \frac{T}{2\pi} \right)^2 - 2 \log \frac{T}{2\pi} \right] + O(x \log x) + O\left(\frac{\tilde{h}T}{x^{1/2-\epsilon}} \right). \end{aligned}$$

Theorem 1.2. Assume **TPC**. For $M \geq 3$ and $\frac{T}{\log^M T} \leq x$,

$$\begin{aligned} &F_h(x, T) \\ &= \frac{T}{\pi} \left[\frac{2 \cos(h \log x)}{4 + h^2} \log x - \frac{4h \sin(h \log x)}{(4 + h^2)^2} \right] \\ &\quad + \frac{T}{\pi} \int_1^\infty \left[-\frac{2 \cos(h \log x)}{4 + h^2} \frac{1}{y} - \frac{4f(y)}{y^2} \cos(h \log x) + G_1(y) + G_2(y) \right] \frac{\sin \frac{Ty}{x}}{\frac{Ty}{x}} dy \\ &\quad - \frac{x}{\pi} \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{3 \cos(h \log x)}{9 + h^2} + \frac{h \sin(h \log x)}{9 + h^2} \right] \\ &\quad - \frac{x}{\pi} \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{\cos(h \log x)}{1 + h^2} - \frac{h \sin(h \log x)}{1 + h^2} \right] \\ &\quad + \frac{T}{\pi} \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \int_0^1 y \cos\left(h \log \frac{kx}{y}\right) \frac{\sin \frac{Ty}{x}}{\frac{Ty}{x}} dy \\ &\quad + O\left(\tilde{h} \frac{x^{1+6\epsilon}}{T}\right) + O(\tilde{h} x^{1/2+7\epsilon}) + O\left(\tilde{h} \frac{x^2}{T^{2-2\epsilon}}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right). \end{aligned}$$

where $G_1(y)$ and $G_2(y)$ are defined in Lemma 4.2.

Theorem 1.3. Assume **TPC**. For $M \geq 3$ and $\frac{T}{\log^M T} \leq x \leq T$,

$$F_h(x, T) = \frac{T}{\pi} \left[\frac{2 \cos(h \log x)}{4 + h^2} \log x - \frac{4h \sin(h \log x)}{(4 + h^2)^2} \right] + O(\tilde{h}x) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right).$$

Theorem 1.4. Assume **TPC**. For $M \geq 3$ and $T \leq x \leq T^{2-29\epsilon}$,

$$F_h(x, T) = \frac{T}{2\pi} \log \frac{T}{2\pi e} \left[\frac{4 \cos(h \log x)}{4 + h^2} \right] + O\left(\tilde{h}T \left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right).$$

For real α , let $F_h(\alpha) := \left(\frac{T}{2\pi} \log T\right)^{-1} F_h(T^\alpha, T)$. Then $F_h(\alpha) = F_h(-\alpha)$. Based on the above theorems, one may make the following

Conjecture 1.1. For any arbitrary large A , as $T \rightarrow \infty$,

$$F_h(\alpha) = \begin{cases} (1 + o(1))T^{-2\alpha} \log T + \alpha \frac{4 \cos(h \log T^\alpha)}{4 + h^2} + o(1), & \text{if } 0 \leq \alpha \leq 1, \\ \frac{4 \cos(h \log T^\alpha)}{4 + h^2} + o(1), & \text{if } 1 \leq \alpha \leq A. \end{cases}$$

By convolving $F_h(\alpha)$ with an appropriate kernel $\hat{r}(\alpha)$,

$$\left(\frac{T}{2\pi} \log T\right)^{-1} \sum_{0 < \gamma, \gamma' \leq T} r\left((\gamma - \gamma' - h) \frac{\log T}{2\pi}\right) w(\gamma - \gamma' - h) = \int_{-\infty}^{+\infty} F_h(\alpha) \hat{r}(\alpha) d\alpha \quad (1)$$

where $\hat{r}(\alpha) = \int_{-\infty}^{\infty} r(u) e^{-2\pi i \alpha u} du$ for even $r(u)$ only. Conjecture 1.1 and (1) leads to

Conjecture 1.2. For fixed $\alpha > 0$,

$$\left(\frac{T}{2\pi} \log T\right)^{-1} \sum_{\substack{0 < \gamma \neq \gamma' \leq T \\ |\gamma - \gamma' - h| \leq 2\pi\alpha / \log T}} 1 \sim \int_{-\alpha + h \log T / (2\pi)}^{\alpha + h \log T / (2\pi)} 1 - \frac{4}{4 + h^2} \left(\frac{\sin \pi u}{\pi u}\right)^2 du.$$

Conjecture 1.3. For $0 < \alpha < \beta \ll \log T$,

$$\left(\frac{T}{2\pi} \log T\right)^{-1} \sum_{\substack{0 < \gamma \neq \gamma' \leq T \\ 2\pi\alpha / \log T \leq \gamma - \gamma' \leq 2\pi\beta / \log T}} 1 \sim \int_{\alpha}^{\beta} 1 - \frac{1}{1 + (\pi u / \log T)^2} \left(\frac{\sin \pi u}{\pi u}\right)^2 du.$$

2 Some Lemmas

Lemma 2.1.

$$\begin{aligned} 2 \sum_{\gamma} \frac{x^{i(\gamma-t)}}{1 + (t - \gamma)^2} &= -\frac{1}{x} \sum_{n \leq x} \frac{\Lambda(n)}{n^{-1/2+it}} - x \sum_{n > x} \frac{\Lambda(n)}{n^{3/2+it}} + \frac{x^{1/2-it}}{\frac{1}{2} + it} + \frac{x^{1/2-it}}{\frac{3}{2} - it} \\ &\quad + \frac{\log \tau}{x} + \frac{1}{x} \left[\frac{\zeta'}{\zeta} \left(\frac{3}{2} - it\right) - \log 2\pi \right] + O\left(\frac{1}{x\tau}\right) \end{aligned} \quad (2)$$

where the sum is over all the imaginary parts of the zeros of the Riemann zeta function, and $\tau = |t| + 2$. $\Lambda(n)$ is von Mangoldt's lambda function.

Proof: This is Lemma 2.2 in [2].

Write (2) as $\text{Left}(x, t) = \text{Right}(x, t)$. Let

$$\begin{aligned} P(x, T) &= \frac{1}{x} \sum_{n \leq x} \frac{\Lambda(n)}{n^{-1/2+it}} + x \sum_{n > x} \frac{\Lambda(n)}{n^{3/2+it}} - \frac{x^{1/2-it}}{\frac{1}{2} + it} - \frac{x^{1/2-it}}{\frac{3}{2} - it}, \\ Q(x, T) &= \frac{\log \tau}{x}, \\ R(x, T) &= \frac{1}{x} \left[\frac{\zeta'}{\zeta} \left(\frac{3}{2} - it \right) - \log 2\pi \right], \\ S(x, T) &= O\left(\frac{1}{x\tau}\right). \end{aligned}$$

Lemma 2.2.

$$\begin{aligned} &\int_0^T |\text{Left}(x, t) + \text{Left}(x, t-h)|^2 dt \\ &= 2\pi F(x, T) + 2\pi F(x, T-h) + 4\pi F_h(x, T) + O(\log^3 T) + O(h \log^2 h). \end{aligned}$$

Proof: This follows from page 188 of Montgomery [9] and the fact that $F(x, T) \ll T \log^2 T$.

Lemma 2.3. For $x \geq 1$,

$$\begin{aligned} &\int_0^T |\text{Right}(x, t) + \text{Right}(x, t-h)|^2 dt \\ &= \int_0^T |P(x, t) + P(x, t-h)|^2 dt + \frac{4T}{x^2} \left[\left(\log \frac{T}{2\pi} \right)^2 - 2 \log \frac{T}{2\pi} \right. \\ &\quad \left. + \left(\frac{1}{2} \sum_{n=1}^{\infty} \frac{\Lambda^2(n)(1 + \cos(h \log n))}{n^3} + 2 \right) \right] + O(\tilde{h} \log^2 T). \end{aligned}$$

Proof: This is similar to the proof of Theorem 3.1 in [2].

Lemma 2.4. For $x \geq 1$,

$$\begin{aligned} &4\pi F_h(x, T) \\ &= \int_0^T |P(x, t) + P(x, t-h)|^2 dt - \int_0^T |P(x, t)|^2 dt - \int_0^T |P(x, t-h)|^2 dt \\ &\quad + \frac{2T}{x^2} \left[\left(\log \frac{T}{2\pi} \right)^2 - 2 \log \frac{T}{2\pi} + \left(\sum_{n=1}^{\infty} \frac{\Lambda^2(n) \cos(h \log n)}{n^3} + 2 \right) \right] + O(\tilde{h} \log^3 T). \end{aligned}$$

Proof: It follows from Lemma 2.2 and Lemma 2.3 as well as their special cases when $h = 0$.

Lemma 2.5. For any sequence of complex numbers $\{a_n\}_{n=1}^{\infty}$ with $\sum_{n=1}^{\infty} n|a_n|^2 < \infty$,

$$\int_0^T \left| \sum_{n=1}^{\infty} a_n n^{-it} \right|^2 dt = \sum_{n=1}^{\infty} |a_n|^2 (T + O(n)).$$

Proof: This is Parseval's identity for Dirichlet series. See [10].

Lemma 2.6.

$$\begin{aligned}\sum_{n \leq x} \Lambda^2(n)n &= \frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + O(x^{1/2+\epsilon}). \\ \sum_{n > x} \frac{\Lambda^2(n)}{n^3} &= \frac{1}{2} \frac{\log x}{x^2} + \frac{1}{4x^2} + O\left(\frac{1}{x^{5/2-\epsilon}}\right).\end{aligned}$$

Proof: Use partial summation and the prime number theorem.

Lemma 2.7. *For any real a and b not both zero,*

$$\begin{aligned}\int e^{ax} \sin bx \, dx &= \frac{a}{a^2 + b^2} e^{ax} \sin bx - \frac{b}{a^2 + b^2} e^{ax} \cos bx. \\ \int e^{ax} \cos bx \, dx &= \frac{a}{a^2 + b^2} e^{ax} \cos bx + \frac{b}{a^2 + b^2} e^{ax} \sin bx. \\ \int x e^{ax} \sin bx \, dx &= \left[\frac{ax}{a^2 + b^2} - \frac{a^2 - b^2}{(a^2 + b^2)^2} \right] e^{ax} \sin bx \\ &\quad - \left[\frac{bx}{a^2 + b^2} - \frac{2ab}{(a^2 + b^2)^2} \right] e^{ax} \cos bx. \\ \int x e^{ax} \cos bx \, dx &= \left[\frac{ax}{a^2 + b^2} - \frac{a^2 - b^2}{(a^2 + b^2)^2} \right] e^{ax} \cos bx \\ &\quad + \left[\frac{bx}{a^2 + b^2} - \frac{2ab}{(a^2 + b^2)^2} \right] e^{ax} \sin bx.\end{aligned}$$

Proof: One can use $\int e^{(a+ib)x} dx$, $\int e^{(a-ib)x} dx$, $\int x e^{(a+ib)x} dx$ and $\int x e^{(a-ib)x} dx$ which are simple to compute.

Lemma 2.8.

$$\begin{aligned}\frac{1}{x^2} \sum_{n \leq x} \Lambda^2(n)n \cos(h \log n) &= \frac{2 \cos(h \log x)}{4 + h^2} \log x + \frac{h^2 - 4}{(4 + h^2)^2} \cos(h \log x) \\ &\quad + \frac{h \sin(h \log x)}{4 + h^2} \log x - \frac{4h}{(4 + h^2)^2} \sin(h \log x) \\ &\quad + O\left(\frac{\tilde{h}}{x^{1/2-\epsilon}}\right). \\ x^2 \sum_{n > x} \frac{\Lambda^2(n)}{n} \cos(h \log n) &= \frac{2 \cos(h \log x)}{4 + h^2} \log x - \frac{h^2 - 4}{(4 + h^2)^2} \cos(h \log x) \\ &\quad - \frac{h \sin(h \log x)}{4 + h^2} \log x - \frac{4h}{(4 + h^2)^2} \sin(h \log x) \\ &\quad + O\left(\frac{\tilde{h}}{x^{1/2-\epsilon}}\right).\end{aligned}$$

Proof: We shall prove the first one. The other one is very similar. Let $A(x) = \frac{1}{x^2} \sum_{n \leq x} \Lambda^2(n)n$. By integration by parts and Lemma 2.6,

$$\begin{aligned}
& \frac{1}{x^2} \sum_{n \leq x} \Lambda^2(n)n \cos(h \log n) \\
&= \frac{A(x)}{x^2} \cos(h \log x) + \frac{h}{x^2} \int_1^x A(u) \frac{\sin(h \log u)}{u} du \\
&= \left[\frac{1}{2} \log x - \frac{1}{4} \right] \cos(h \log x) + \frac{h}{x^2} \int_1^x \left[\frac{1}{2} \log u - \frac{1}{4} \right] u \sin(h \log u) du \\
&\quad + O\left(\frac{\tilde{h}}{x^{1/2-\epsilon}}\right) \\
&= \left[\frac{1}{2} \log x - \frac{1}{4} \right] \cos(h \log x) + \frac{h}{x^2} \left[\frac{1}{2} \int_0^{\log x} v e^{2v} \sin hv dv \right. \\
&\quad \left. - \frac{1}{4} \int_0^{\log x} e^{2v} \sin hv dv \right] + O\left(\frac{\tilde{h}}{x^{1/2-\epsilon}}\right)
\end{aligned}$$

which gives the desired result after applying Lemma 2.7 with $a = 2$ and $b = h$, and some algebra.

3 Proof of Theorem 1.1

First, note that

$$P(x, t) = \frac{1}{x^{1/2}} \left[\sum_{n \leq x} \Lambda(n) \left(\frac{x}{n}\right)^{-1/2+it} + \sum_{n > x} \Lambda(n) \left(\frac{x}{n}\right)^{3/2+it} \right] + O\left(\frac{x^{1/2}}{\tau}\right).$$

Thus,

$$\begin{aligned}
P(x, t) + P(x, t - h) &= \frac{1}{x^{1/2}} \left[\sum_{n \leq x} \Lambda(n) (1 + n^{ih}) \left(\frac{x}{n}\right)^{-1/2+it} \right. \\
&\quad \left. + \sum_{n > x} \Lambda(n) (1 + n^{ih}) \left(\frac{x}{n}\right)^{3/2+it} \right] + O\left(\frac{x^{1/2}}{\tau}\right)
\end{aligned}$$

So, the first integral in Lemma 2.4

$$\begin{aligned}
&= \frac{1}{x} \int_0^T \left| \sum_{n \leq x} \Lambda(n) (1 + n^{ih}) \left(\frac{x}{n}\right)^{-1/2+it} + \sum_{n > x} \Lambda(n) (1 + n^{ih}) \left(\frac{x}{n}\right)^{3/2+it} \right|^2 dt \\
&\quad + O\left(\left[\sum_{n \leq x} \Lambda(n) \left(\frac{x}{n}\right)^{-1/2} + \sum_{n > x} \Lambda(n) \left(\frac{x}{n}\right)^{3/2} \right] \int_0^T \frac{1}{\tau^2} dt \right) + O\left(\int_0^T \frac{x}{\tau^4} dt \right) \\
&= \frac{1}{x} \sum_{n \leq x} \Lambda^2(n) |1 + n^{ih}|^2 \left(\frac{x}{n}\right)^{-1} (T + O(n)) \\
&\quad + \frac{1}{x} \sum_{n > x} \Lambda^2(n) |1 + n^{ih}|^2 \left(\frac{x}{n}\right)^3 (T + O(n)) + O(x) \\
&= \frac{2T}{x^2} \sum_{n \leq x} \Lambda^2(n) n (1 + \cos(h \log n)) + 2Tx^2 \sum_{n > x} \frac{\Lambda^2(n)}{n^3} (1 + \cos(h \log n)) \\
&\quad + O(x \log x)
\end{aligned}$$

Similarly (or by setting $h = 0$), each of the second and third integral in Lemma 2.4

$$= \frac{T}{x^2} \sum_{n \leq x} \Lambda^2(n) n + Tx^2 \sum_{n > x} \frac{\Lambda^2(n)}{n^3} + O(x \log x)$$

Therefore,

$$\begin{aligned}
4\pi F_h(x, T) &= 2T \left[\frac{1}{x^2} \sum_{n \leq x} \Lambda^2(n) n \cos(h \log n) + x^2 \sum_{n > x} \frac{\Lambda^2(n)}{n^3} \cos(h \log n) \right] \\
&\quad + \frac{2T}{x^2} \left[\left(\log \frac{T}{2\pi} \right)^2 - 2 \log \frac{T}{2\pi} \right] \\
&\quad + O\left(\frac{T}{x^2} \right) + O(\tilde{h} \log^3 T) + O(x \log x) \\
&= 2T \left[\frac{4 \cos(h \log x)}{4 + h^2} \log x - \frac{8h \sin(h \log x)}{(4 + h^2)^2} \right] \\
&\quad + \frac{2T}{x^2} \left[\left(\log \frac{T}{2\pi} \right)^2 - 2 \log \frac{T}{2\pi} \right] + O(x \log x) + O\left(\frac{\tilde{h}T}{x^{1/2-\epsilon}} \right)
\end{aligned}$$

by Lemma 2.8. The theorem follows after dividing through by 4π .

4 Twin Prime Conjecture and smooth weight

We shall use a quantitative form of the Twin Prime Conjecture **TPC** as follow:
For any $\epsilon > 0$,

$$\sum_{n=1}^N \Lambda(n) \Lambda(n+d) = \mathfrak{S}(d)N + O(N^{1/2+\epsilon}) \text{ uniformly in } |d| \leq N.$$

$\mathfrak{S}(d) = 2 \prod_{p>2} (1 - \frac{1}{(p-1)^2}) \prod_{p|d, p>2} \frac{p-1}{p-2}$ if d is even, and $\mathfrak{S}(d) = 0$ if d is odd.

Let K and M be some large positive integers (K may depend on ϵ). Set $U = \log^M T$ and $\Delta = 1/(2^K U)$. We recall the smooth weight $\Psi_U(t)$ in [3] with:

1. support in $[-1/U, 1 + 1/U]$,
2. $0 \leq \Psi_U(t) \leq 1$,
3. $\Psi_U(t) = 1$ for $1/U \leq t \leq 1 - 1/U$,
4. $\Psi_U^{(j)}(t) \ll U^j$ for $j = 1, 2, \dots, K$.

This weight function satisfies the requirements in Goldston and Gonek [6]. One more thing to note is that

$$\operatorname{Re} \hat{\Psi}_U(y) = \frac{\sin 2\pi y}{2\pi y} \left(\frac{\sin 2\pi \Delta y}{2\pi \Delta y} \right)^{K+1}$$

where $\hat{f}(y) = \int_{-\infty}^{\infty} f(t)e(yt)dt$.

We also need to study

$$S_\alpha^h(y) := \sum_{k \leq y} \mathfrak{S}(k) k^\alpha \cos(h \log \frac{kx}{y}) - \int_0^y u^\alpha \cos(h \log \frac{ux}{y}) du \text{ for } \alpha \geq 0,$$

and

$$T_\alpha^h(y) := \sum_{k > y} \frac{\mathfrak{S}(k)}{k^\alpha} \cos(h \log \frac{kx}{y}) - \int_y^\infty \frac{1}{u^\alpha} \cos(h \log \frac{ux}{y}) du \text{ for } \alpha > 1.$$

Then from [4],

$$S_0(y) := S_0^0(y) = -\frac{1}{2} \log y + O((\log y)^{2/3}) = -\frac{1}{2} \log y + \epsilon(y). \quad (3)$$

By partial summation and Lemma 2.7,

$$\begin{aligned} S_\alpha^h(y) = & \epsilon(y) y^\alpha \cos(h \log x) - \frac{\alpha \cos(h \log x)}{2(\alpha^2 + h^2)} y^\alpha - \frac{h \sin(h \log x)}{2(\alpha^2 + h^2)} y^\alpha \\ & - \int_0^y \epsilon(u) u^{\alpha-1} \left[\alpha \cos(h \log \frac{ux}{y}) - h \sin(h \log \frac{ux}{y}) \right] du, \end{aligned} \quad (4)$$

and

$$\begin{aligned} T_\alpha^h(y) = & -\frac{\epsilon(y)}{y^\alpha} \cos(h \log x) - \frac{\alpha \cos(h \log x)}{2(\alpha^2 + h^2)} \frac{1}{y^\alpha} + \frac{h \sin(h \log x)}{2(\alpha^2 + h^2)} \frac{1}{y^\alpha} \\ & + \int_y^\infty \frac{\epsilon(u)}{u^{\alpha+1}} \left[\alpha \cos(h \log \frac{ux}{y}) + h \sin(h \log \frac{ux}{y}) \right] du. \end{aligned} \quad (5)$$

Let

$$f(y) := \int_0^y \epsilon(u) - \frac{B}{2} du$$

where $B = -C_0 - \log 2\pi$ and C_0 is Euler's constant. Note that

$$f(y) \ll y^{1/2+\epsilon} \quad (6)$$

(see Lemma 2.2 of [3]). From (4) and (5),

$$\begin{aligned} & S_2^h(y) \frac{1}{y^3} + T_2^h(y)y \\ &= -\frac{2 \cos(h \log x)}{4+h^2} \frac{1}{y} - \frac{1}{y^3} \int_0^y u \epsilon(u) \left[2 \cos\left(h \log \frac{ux}{y}\right) - h \sin\left(h \log \frac{ux}{y}\right) \right] du \\ & \quad + y \int_y^\infty \frac{\epsilon(u)}{u^3} \left[2 \cos\left(h \log \frac{ux}{y}\right) + h \sin\left(h \log \frac{ux}{y}\right) \right] du. \end{aligned} \quad (7)$$

Lemma 4.1.

$$\begin{aligned} I + J &= -\frac{1}{y^3} \int_0^y u \epsilon(u) \left[2 \cos\left(h \log \frac{ux}{y}\right) - h \sin\left(h \log \frac{ux}{y}\right) \right] du \\ & \quad + y \int_y^\infty \frac{\epsilon(u)}{u^3} \left[2 \cos\left(h \log \frac{ux}{y}\right) + h \sin\left(h \log \frac{ux}{y}\right) \right] du \\ &= -\frac{4f(y)}{y} \cos(h \log x) \\ & \quad + \frac{1}{y^3} \int_0^y f(u) \left[(2-h^2) \cos\left(h \log \frac{ux}{y}\right) - 3h \sin\left(h \log \frac{ux}{y}\right) \right] du \\ & \quad + y \int_y^\infty \frac{f(u)}{u^4} \left[(6-h^2) \cos\left(h \log \frac{ux}{y}\right) + 5h \sin\left(h \log \frac{ux}{y}\right) \right] du. \end{aligned}$$

Proof: I can be rewritten as

$$\begin{aligned} & -\frac{1}{y^3} \int_0^y u \left(\epsilon(u) - \frac{B}{2} \right) \left[2 \cos\left(h \log \frac{ux}{y}\right) - h \sin\left(h \log \frac{ux}{y}\right) \right] du \\ & -\frac{B}{2} \frac{1}{y^3} \int_0^y u \left[2 \cos\left(h \log \frac{ux}{y}\right) - h \sin\left(h \log \frac{ux}{y}\right) \right] du = -I_1 - I_2. \end{aligned}$$

By a substitution $v = \log \frac{ux}{y}$ and Lemma 2.7,

$$I_2 = \frac{B}{2} \frac{1}{y} \cos(h \log x). \quad (8)$$

By integration by parts and (6),

$$\begin{aligned} I_1 &= \frac{f(y)}{y^2} [2 \cos(h \log x) - h \sin(h \log x)] \\ & \quad - \frac{1}{y^3} \int_0^y f(u) \left[(2-h^2) \cos\left(h \log \frac{ux}{y}\right) - 3h \sin\left(h \log \frac{ux}{y}\right) \right] du. \end{aligned} \quad (9)$$

Similarly, J can be rewritten as

$$y \int_y^\infty \frac{\epsilon(u) - \frac{B}{2}}{u^3} \left[2 \cos \left(h \log \frac{ux}{y} \right) + h \sin \left(h \log \frac{ux}{y} \right) \right] du \\ + \frac{B}{2} y \int_y^\infty \frac{1}{u^3} \left[2 \cos \left(h \log \frac{ux}{y} \right) + h \sin \left(h \log \frac{ux}{y} \right) \right] du = J_1 + J_2$$

By a substitution $v = \log \frac{ux}{y}$ and Lemma 2.7,

$$J_2 = \frac{B}{2} \frac{1}{y} \cos(h \log x). \quad (10)$$

By integration by parts and (6),

$$J_1 = -\frac{f(y)}{y^2} [2 \cos(h \log x) + h \sin(h \log x)] \\ + y \int_y^\infty \frac{f(u)}{u^4} \left[(6 - h^2) \cos \left(h \log \frac{ux}{y} \right) + 5h \sin \left(h \log \frac{ux}{y} \right) \right] du. \quad (11)$$

(8), (9), (10) and (11) together gives the lemma.

Lemma 4.2.

$$S_2^h(y) \frac{1}{y^3} + T_2^h(y)y = -\frac{2 \cos(h \log x)}{4 + h^2} \frac{1}{y} - \frac{4f(y)}{y^2} \cos(h \log x) + G_1(y) + G_2(y)$$

where

$$G_1(y) = \frac{1}{y^3} \int_0^y f(u) \left[(2 - h^2) \cos \left(h \log \frac{ux}{y} \right) - 3h \sin \left(h \log \frac{ux}{y} \right) \right] du,$$

and

$$G_2(y) = y \int_y^\infty \frac{f(u)}{u^4} \left[(6 - h^2) \cos \left(h \log \frac{ux}{y} \right) + 5h \sin \left(h \log \frac{ux}{y} \right) \right] du.$$

Proof: Combine (7) and Lemma 4.1.

Lemma 4.3. For any integer $n \geq 1$,

$$\int_1^\infty \frac{1}{y^n} \operatorname{Re} \hat{\Psi}_U \left(\frac{T y}{2\pi x} \right) dy = \int_1^\infty \frac{1}{y^n} \frac{\sin \frac{T}{x} y}{\frac{T}{x} y} dy + O\left(\Delta \log \frac{1}{\Delta} \right).$$

When $n \neq 2$, the error term can be replaced by $O(\Delta)$.

Proof: This is Lemma 3.3 in [3].

Lemma 4.4. If $F(y) \ll y^{-3/2+\epsilon}$ for $y \geq 1$, then

$$\int_1^\infty F(y) \operatorname{Re} \hat{\Psi}_U \left(\frac{T y}{2\pi x} \right) dy = \int_1^\infty F(y) \frac{\sin \frac{T}{x} y}{\frac{T}{x} y} dy + O(\Delta).$$

Proof: This is Lemma 3.4 in [3].

5 Proof of Theorem 1.2

Throughout this section, we assume $\tau = T^{1-\epsilon} \leq T/\log^M T \leq x \leq T^{2-2\epsilon}$, $U = \log^M T$ for $M > 2$, $H^* = \tau^{-2}x^{2/(1-\epsilon)}$, and $\Psi_U(t)$ is defined as in the previous section. The implicit constants in the error terms may depend on ϵ , K and M .

Our method is that of Goldston and Gonek [6] and it is very similar to [3]. Let $s = \sigma + it$,

$$A_h(s) := \sum_{n \leq x} \frac{\Lambda(n)(1+n^{ih})}{n^s} \text{ and } A_h^*(s) := \sum_{n > x} \frac{\Lambda(n)(1+n^{ih})}{n^s},$$

$$A(s) := \frac{1}{2}A_0(s) \text{ and } A^*(s) := \frac{1}{2}A_0^*(s).$$

By Lemma 2.4, with slight modifications, one has

$$\begin{aligned} 4\pi F_h(x, T) &= \int_0^T \left| \frac{1}{x} \left(A_h\left(-\frac{1}{2} + it\right) - \int_1^x (1+u^{ih})u^{1/2-it} du \right) \right. \\ &\quad \left. + x \left(A_h^*\left(\frac{3}{2} + it\right) - \int_x^\infty (1+u^{ih})u^{-3/2-it} du \right) \right|^2 dt \\ &\quad - 2 \int_0^T \left| \frac{1}{x} \left(A\left(-\frac{1}{2} + it\right) - \int_1^x u^{1/2-it} du \right) \right. \\ &\quad \left. + x \left(A^*\left(\frac{3}{2} + it\right) - \int_x^\infty u^{-3/2-it} du \right) \right|^2 dt + O(\tilde{h} \log^3 T). \end{aligned}$$

Inserting $\Psi_U(t/T)$ into the integral and extending the range of integration to the whole real line, we have

$$\begin{aligned} 4\pi F(x, T) &= \frac{1}{x^2} I_1(x, T) + x^2 I_2(x, T) - \frac{2}{x^2} I_3(x, T) - 2x^2 I_4(x, T) \\ &\quad + O\left(\frac{T(\log T)^2}{U}\right) + O\left(\frac{x^{1+6\epsilon}}{T}\right) \end{aligned} \tag{12}$$

where

$$\begin{aligned} I_1(x, T) &= \int_{-\infty}^{\infty} \left| A_h\left(-\frac{1}{2} + it\right) - \int_1^x (1+u^{ih})u^{1/2-it} du \right|^2 \Psi_U\left(\frac{t}{T}\right) dt, \\ I_2(x, T) &= \int_{-\infty}^{\infty} \left| A_h^*\left(\frac{3}{2} + it\right) - \int_x^\infty (1+u^{ih})u^{-3/2-it} du \right|^2 \Psi_U\left(\frac{t}{T}\right) dt, \\ I_3(x, T) &= \int_{-\infty}^{\infty} \left| A\left(-\frac{1}{2} + it\right) - \int_1^x u^{1/2-it} du \right|^2 \Psi_U\left(\frac{t}{T}\right) dt, \\ I_4(x, T) &= \int_{-\infty}^{\infty} \left| A^*\left(\frac{3}{2} + it\right) - \int_x^\infty u^{-3/2-it} du \right|^2 \Psi_U\left(\frac{t}{T}\right) dt \end{aligned}$$

by Lemma 1 of [7] with modification $V = -T/U$ and $T - T/U$, and $W = 2T/U$. The contribution from the cross terms are estimated via Theorem 3 of [6]. Note

that by partial summation with the Riemann Hypothesis and **TPC**,

$$\begin{aligned} \sum_{n \leq x} \Lambda(n)(1 + n^{ih}) &= \int_1^x (1 + u^{ih}) du + O(\tilde{h}x^{1/2+\epsilon}), \\ \sum_{n \leq x} \Lambda(n)\Lambda(n+k)(1 + n^{ih})(1 + (n+k)^{-ih}) \\ &= \mathfrak{S}(k) \int_1^x (1 + u^{ih})(1 + (u+k)^{-ih}) du + O(\tilde{h}x^{1/2+\epsilon}). \end{aligned}$$

By Corollary 1 of [6] (see also the calculations at the end of [6] and [7]),

$$\begin{aligned} I_1(x, T) &= \hat{\Psi}_U(0)T \sum_{n \leq x} \Lambda^2(n)n|1 + n^{ih}|^2 \\ &\quad + 4\pi \left(\frac{T}{2\pi}\right)^3 \int_{T/2\pi x}^{\infty} \left[\sum_{k \leq 2\pi xv/T} \mathfrak{S}(k)k^2 \left(1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right) \right. \\ &\quad \times \left. \left(1 + \left(\frac{kT}{2\pi v} + k\right)^{-ih}\right) \right] \operatorname{Re} \hat{\Psi}_U(v) \frac{dv}{v^3} \\ &\quad - 4\pi \left(\frac{T}{2\pi}\right)^3 \int_{T/2\pi \tau x}^{\infty} \left[\int_0^{2\pi xv/T} u^2 \left|1 + \left(\frac{uT}{2\pi v}\right)^{ih}\right|^2 du \right] \operatorname{Re} \hat{\Psi}_U(v) \frac{dv}{v^3} \\ &\quad + O\left(\tilde{h} \frac{x^{3+6\epsilon}}{T}\right) + O(\tilde{h}x^{5/2+7\epsilon}). \end{aligned}$$

Note that

$$\begin{aligned} &\left(1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right) \left(1 + \left(\frac{kT}{2\pi v} + k\right)^{-ih}\right) \\ &= \left|1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right|^2 + \left(1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right) \left(\left(\frac{kT}{2\pi v} + k\right)^{-ih} - \left(\frac{kT}{2\pi v}\right)^{-ih}\right) \\ &= \left|1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right|^2 + \left(1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right) \left(\frac{kT}{2\pi v}\right)^{-ih} \left(\left(1 + \frac{2\pi v}{T}\right)^{-ih} - 1\right) \\ &= \left|1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right|^2 + O\left(\min\left(\frac{hv}{T}, 1\right)\right). \end{aligned} \tag{13}$$

Thus,

$$\begin{aligned} &\int_{T/2\pi x}^{\infty} \left[\sum_{k \leq 2\pi xv/T} \mathfrak{S}(k)k^2 \left(1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right) \left(1 + \left(\frac{kT}{2\pi v} + k\right)^{-ih}\right) \right] \operatorname{Re} \hat{\Psi}_U(v) \frac{dv}{v^3} \\ &= \int_{T/2\pi x}^{\infty} \left[\sum_{k \leq 2\pi xv/T} \mathfrak{S}(k)k^2 \left|1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right|^2 \right] \operatorname{Re} \hat{\Psi}_U(v) \frac{dv}{v^3} \\ &\quad + O\left(\int_{T/2\pi x}^{T^{2-\epsilon/x}} \left(\frac{xv}{T}\right)^3 \frac{hv}{T} \frac{1}{v} \frac{dv}{v^3} + \int_{T^{2-\epsilon/x}}^{\infty} \left(\frac{xv}{T}\right)^3 \frac{1}{\Delta v^2} \frac{dv}{v^3}\right) \\ &= \int_{T/2\pi x}^{\infty} \left[\sum_{k \leq 2\pi xv/T} \mathfrak{S}(k)k^2 \left|1 + \left(\frac{kT}{2\pi v}\right)^{ih}\right|^2 \right] \operatorname{Re} \hat{\Psi}_U(v) \frac{dv}{v^3} + O\left(\frac{hx^2}{T^{2+\epsilon}} + \frac{x^4}{\Delta T^{5-\epsilon}}\right) \end{aligned}$$

as $\sum_{k \leq x} \mathfrak{S}(k) \sim x$ and $Re \hat{\Psi}_U(v) \ll \min\left(\frac{1}{v}, \frac{1}{\Delta v^2}\right)$. Therefore,

$$\begin{aligned}
I_1(x, T) &= T \sum_{n \leq x} \Lambda^2(n) n (2 + 2 \cos(h \log n)) \\
&\quad + 4\pi \left(\frac{T}{2\pi}\right)^3 \int_{T/2\pi x}^{\infty} \left[\sum_{k \leq 2\pi xv/T} \mathfrak{S}(k) k^2 (2 + 2 \cos(h \log \frac{kT}{2\pi v})) \right. \\
&\quad \left. - \int_0^{2\pi xv/T} u^2 (2 + 2 \cos(h \log \frac{uT}{2\pi v})) du \right] Re \hat{\Psi}_U(v) \frac{dv}{v^3} \\
&\quad - 4\pi \left(\frac{T}{2\pi}\right)^3 \int_0^{T/2\pi x} \int_0^{2\pi xv/T} u^2 (2 + 2 \cos(h \log \frac{uT}{2\pi v})) du Re \hat{\Psi}_U(v) \frac{dv}{v^3} \\
&\quad + O\left(\frac{\tilde{h}x^{3+6\epsilon}}{T}\right) + O(\tilde{h}x^{5/2+7\epsilon}) + O(\tilde{h}x^2 T^{1-\epsilon}) + O\left(\frac{x^4}{\Delta T^{2-\epsilon}}\right).
\end{aligned}$$

Similarly, by Corollary 2 of [6],

$$\begin{aligned}
I_2(x, T) &= T \sum_{x < n} \frac{\Lambda^2(n)}{n^3} (2 + 2 \cos(h \log n)) \\
&\quad + \frac{8\pi^2}{T} \int_0^{TH^*/2\pi x} \left[\sum_{2\pi xv/T \leq k \leq H^*} \frac{\mathfrak{S}(k)}{k^2} (2 + 2 \cos(h \log \frac{kT}{2\pi v})) \right. \\
&\quad \left. - \int_{2\pi xv/T}^{H^*} \frac{1}{u^2} (2 + 2 \cos(h \log \frac{uT}{2\pi v})) du \right] Re \hat{\Psi}_U(v) v dv \\
&\quad + O(\tilde{h}T^{-1}x^{-1+6\epsilon}) + O(\tilde{h}x^{-3/2+7\epsilon}) + O(\tilde{h}T^{1-\epsilon/2}x^{-2}) + O\left(\frac{\tilde{h}H^*}{\Delta x^2}\right)
\end{aligned}$$

where the last error term comes from the error term in (13). $I_3(x, T)$ and $I_4(x, T)$ are computed in [3] or one can simply set $h = 0$ in $I_1(x, T)$ and $I_2(x, T)$, and divide by 4. Putting these into (12) with a substitution $y = \frac{2\pi xv}{T}$ and using

Lemma 2.8,

$$\begin{aligned}
& 4\pi F_h(x, T) \\
&= 2T \left[\frac{4 \cos(h \log x)}{4 + h^2} \log x - \frac{8h \sin(h \log x)}{(4 + h^2)^2} \right] \\
&+ 4T \int_1^\infty \left[\sum_{k \leq y} \mathfrak{S}(k) k^2 \cos(h \log \frac{kx}{y}) - \int_0^y u^2 \cos(h \log \frac{ux}{y}) du \right] \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) \frac{dy}{y^3} \\
&- 4T \int_0^1 \int_0^y u^2 \cos(h \log \frac{ux}{y}) du \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) \frac{dy}{y^3} \\
&+ 4T \int_1^{H^*} \left[\sum_{y \leq k \leq H^*} \frac{\mathfrak{S}(k) \cos(h \log \frac{kx}{y})}{k^2} - \int_y^{H^*} \frac{\cos(h \log \frac{ux}{y})}{u^2} du \right] \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) y dy \\
&+ 4T \int_0^1 \left[\sum_{k \leq H^*} \frac{\mathfrak{S}(k) \cos(h \log \frac{kx}{y})}{k^2} - \int_y^{H^*} \frac{\cos(h \log \frac{ux}{y})}{u^2} du \right] \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) y dy \\
&+ O\left(\frac{\tilde{h}x^{1+6\epsilon}}{T}\right) + O(\tilde{h}x^{1/2+7\epsilon}) + O\left(\frac{\tilde{h}x^2}{T^{2-2\epsilon}}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right)
\end{aligned}$$

From Lemma 2.7,

$$\int e^{ax} \cos bx = \frac{a}{a^2 + b^2} e^{ax} \cos bx + \frac{b}{a^2 + b^2} e^{ax} \sin bx. \quad (14)$$

Also,

$$\begin{aligned}
\int_0^1 \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) dy &= \frac{x}{T} \int_0^{T/x} \frac{\sin u}{u} (1 + O(\Delta^2 u^2)) du \\
&= \frac{x}{T} \int_0^{T/x} \frac{\sin u}{u} du + O\left(\frac{\Delta^2 T}{x}\right).
\end{aligned} \quad (15)$$

By appropriate change of variables, (14) and (15),

$$\begin{aligned}
& \int_0^1 \int_0^y u^2 \cos(h \log \frac{ux}{y}) du \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) \frac{dy}{y^3} \\
&= \frac{x}{T} \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{3}{9 + h^2} \cos(h \log x) + \frac{h}{9 + h^2} \sin(h \log x) \right] + O\left(\frac{\Delta^2 T}{x}\right). \\
& \int_0^1 \left[\sum_{k \leq H^*} \frac{\mathfrak{S}(k) \cos(h \log \frac{kx}{y})}{k^2} - \int_y^{H^*} \frac{\cos(h \log \frac{ux}{y})}{u^2} du \right] \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) y dy \\
&= \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \int_0^1 y \cos(h \log \frac{kx}{y}) \frac{\sin \frac{Ty}{x}}{y} dy + O\left(\frac{1}{H^*}\right) + O\left(\frac{\Delta^2 T}{x}\right) \\
&- \frac{x}{T} \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{1}{1 + h^2} \cos(h \log x) - \frac{h}{1 + h^2} \sin(h \log x) \right] + O\left(\frac{\Delta^2 T}{x}\right).
\end{aligned}$$

Therefore, with the notation $S_\alpha^h(y)$ and $T_\alpha^h(y)$,

$$\begin{aligned}
& 4\pi F_h(x, T) \\
&= 2T \left[\frac{4 \cos(h \log x)}{4 + h^2} \log x - \frac{8h \sin(h \log x)}{(4 + h^2)^2} \right] \\
&+ 4T \int_1^\infty S_2^h(y) \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) \frac{dy}{y^3} \\
&+ 4T \int_1^{H^*} (T_2^h(y) - T_2^h(H^*)) \operatorname{Re} \hat{\Psi}_U \left(\frac{Ty}{2\pi x} \right) y dy \\
&- 4x \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{3 \cos(h \log x)}{9 + h^2} + \frac{h \sin(h \log x)}{9 + h^2} \right] \\
&- 4x \int_0^{T/x} \frac{\sin u}{u} du \left[\frac{\cos(h \log x)}{1 + h^2} - \frac{h \sin(h \log x)}{1 + h^2} \right] \\
&+ 4T \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \int_0^1 y \cos(h \log \frac{kx}{y}) \frac{\sin \frac{Ty}{x}}{\frac{Ty}{x}} dy \\
&+ O\left(\frac{\tilde{h}x^{1+6\epsilon}}{T}\right) + O(\tilde{h}x^{1/2+7\epsilon}) + O\left(\frac{\tilde{h}x^2}{T^{2-2\epsilon}}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right).
\end{aligned}$$

By (3) and (5), $T_2^h(H^*) \ll \frac{h(\log H^*)^{2/3}}{(H^*)^2}$. It follows that the contribution from $T_2^h(H^*)$ in the second integral is $O(hT^{-\epsilon})$. Also, one can extend the upper limit of the second integral to ∞ with an error $O(hT^{-\epsilon})$ by (3) and (5) again. Finally, we obtain the theorem by applying Lemma 4.2, Lemma 4.3, Lemma 4.4, (6) and dividing by 4π .

6 Proof of Theorem 1.3 and 1.4

Proof of Theorem 1.3: It follows directly from Theorem 1.2 by observing that all the other main terms besides the first one are $O(x)$ because of (6).

Before proving Theorem 1.4, we need the following lemmas.

Lemma 6.1.

$$\int_1^\infty \frac{\sin ax}{x^{2n}} dx = \frac{a^{2n-1}}{(2n-1)!} \left[\sum_{k=1}^{2n-1} \frac{(2n-k-1)!}{a^{2n-k}} \sin\left(a + (k-1)\frac{\pi}{2}\right) + (-1)^n ci(a) \right]$$

where $ci(x) = -\int_x^\infty \frac{\cos t}{t} dt = C_0 + \log x + \int_0^x \frac{\cos t - 1}{t} dt$ and C_0 is Euler's constant.

Proof: This is formula 3.761(3) on P.430 of [8] which can be proved by integration by parts repeatedly.

Lemma 6.2. *If $F(y) \ll y^{-3/2+\epsilon}$ for $y \geq 1$, then for $T \leq x$,*

$$\int_1^\infty F(y) \frac{\sin \frac{T}{x} y}{\frac{T}{x} y} dy = \int_1^\infty F(y) dy + O\left(\left(\frac{T}{x}\right)^{1/2-\epsilon}\right).$$

Proof: This is Lemma 5.2 in [3].

Lemma 6.3.

$$\begin{aligned} I &= \int_1^\infty \frac{1}{y^3} \int_0^y f(u) \left[(2-h^2) \cos\left(h \log \frac{ux}{y}\right) - 3h \sin\left(h \log \frac{ux}{y}\right) \right] du dy \\ &= \int_0^1 f(u) [\cos(h \log ux) - h \sin(h \log ux)] du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2} du [\cos(h \log x) - h \sin(h \log x)]. \end{aligned}$$

Proof: Because of (6), we can change the order of integration.

$$\begin{aligned} I &= \int_0^1 f(u) \int_1^\infty \frac{1}{y^3} \left[(2-h^2) \cos\left(h \log \frac{ux}{y}\right) - 3h \sin\left(h \log \frac{ux}{y}\right) \right] dy du \\ &\quad + \int_1^\infty f(u) \int_u^\infty \frac{1}{y^3} \left[(2-h^2) \cos\left(h \log \frac{ux}{y}\right) - 3h \sin\left(h \log \frac{ux}{y}\right) \right] dy du \\ &= \int_0^1 f(u) \left\{ (2-h^2) \left[\frac{2}{4+h^2} \cos(h \log ux) + \frac{h}{4+h^2} \sin(h \log ux) \right] \right. \\ &\quad \left. - 3h \left[\frac{2}{4+h^2} \sin(h \log ux) - \frac{h}{4+h^2} \cos(h \log ux) \right] \right\} du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2} \left\{ (2-h^2) \left[\frac{2}{4+h^2} \cos(h \log x) + \frac{h}{4+h^2} \sin(h \log x) \right] \right. \\ &\quad \left. - 3h \left[\frac{2}{4+h^2} \sin(h \log x) - \frac{h}{4+h^2} \cos(h \log x) \right] \right\} du. \end{aligned}$$

by substituting $v = \log \frac{ux}{y}$ and applying Lemma 2.7. The lemma follows after some simple algebra.

Lemma 6.4.

$$\begin{aligned} J &= \int_1^\infty y \int_y^\infty \frac{f(u)}{u^4} \left[(6-h^2) \cos\left(h \log \frac{ux}{y}\right) + 5h \sin\left(h \log \frac{ux}{y}\right) \right] du dy \\ &= - \int_1^\infty \frac{f(u)}{u^4} [3 \cos(h \log ux) + h \sin(h \log ux)] du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2} du [3 \cos(h \log x) + h \sin(h \log x)]. \end{aligned}$$

Proof: Again, because of (6), we can change the order of integration.

$$\begin{aligned}
J &= \int_1^\infty \int_1^u y \left[(6 - h^2) \cos \left(h \log \frac{ux}{y} \right) + 5h \sin \left(h \log \frac{ux}{y} \right) \right] dy du \\
&= \int_1^\infty \frac{f(u)}{u^4} \left\{ (6 - h^2) \left[\frac{-2}{4 + h^2} \cos (h \log ux) + \frac{h}{4 + h^2} \sin (h \log ux) \right] \right. \\
&\quad + 5h \left[\frac{-2}{4 + h^2} \sin (h \log ux) - \frac{h}{4 + h^2} \cos (h \log ux) \right] \\
&\quad - (6 - h^2) \left[\frac{-2}{4 + h^2} \cos (h \log x) + \frac{h}{4 + h^2} \sin (h \log x) \right] \\
&\quad \left. - 5h \left[\frac{-2}{4 + h^2} \sin (h \log x) - \frac{h}{4 + h^2} \cos (h \log x) \right] \right\} du.
\end{aligned}$$

by substituting $v = \log \frac{ux}{y}$ and applying Lemma 2.7. The lemma follows after some simple algebra.

Lemma 6.5.

$$\begin{aligned}
S &= \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \int_0^1 y \cos \left(h \log \frac{kx}{y} \right) dy \\
&= \left[\frac{1}{1 + h^2} \cos (h \log x) - \frac{h}{1 + h^2} \sin (h \log x) \right] \\
&\quad - \left[\frac{4 - h^2}{2(4 + h^2)^2} \cos (h \log x) - \frac{2h}{(4 + h^2)^2} \sin (h \log x) \right] \\
&\quad + \frac{B}{2} \left[\frac{2}{4 + h^2} \cos (h \log x) - \frac{h}{4 + h^2} \sin (h \log x) \right] + \left(1 + \frac{B}{2} \right) \cos (h \log x) \\
&\quad + \int_1^\infty \frac{f(u)}{u^4} \left[3 \cos (h \log ux) + h \sin (h \log ux) \right] du.
\end{aligned}$$

Proof: By substituting $v = \log \frac{kx}{y}$ and Lemma 2.7,

$$S = \frac{2}{4 + h^2} \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \cos (h \log kx) - \frac{h}{4 + h^2} \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2} \sin (h \log kx).$$

Recall the definition of $S_0(u)$ from (3) and use partial summation,

$$\begin{aligned}
S &= \frac{2}{4+h^2} \int_1^\infty \frac{S_0(u)+u}{u^3} [2 \cos(h \log ux) + h \sin(h \log ux)] du \\
&\quad - \frac{h}{4+h^2} \int_1^\infty \frac{S_0(u)+u}{u^3} [-h \cos(h \log ux) + 2 \sin(h \log ux)] du \\
&= \int_1^\infty \frac{S_0(u)+u}{u^3} \cos(h \log ux) du \\
&= \int_1^\infty \frac{u - \frac{1}{2} \log u + \epsilon(u)}{u^3} \cos(h \log ux) du \\
&= \int_1^\infty \frac{1}{u^2} \cos(h \log ux) du - \frac{1}{2} \int_1^\infty \frac{\log u}{u^3} \cos(h \log ux) du \\
&\quad + \frac{B}{2} \int_1^\infty \frac{1}{u^3} \cos(h \log ux) du + \int_1^\infty \frac{\epsilon(u) - \frac{B}{2}}{u^3} \cos(h \log ux) du \\
&= I_1 - \frac{1}{2} I_2 + \frac{B}{2} I_3 + I_4.
\end{aligned}$$

By appropriate substitution and Lemma 2.7,

$$\begin{aligned}
I_1 &= \frac{1}{1+h^2} \cos(h \log x) - \frac{h}{1+h^2} \sin(h \log x), \\
I_2 &= \frac{4-h^2}{(4+h^2)^2} \cos(h \log x) - \frac{2h}{(4+h^2)^2} \sin(h \log x), \\
I_3 &= \frac{2}{4+h^2} \cos(h \log x) - \frac{h}{4+h^2} \sin(h \log x).
\end{aligned}$$

Finally, by integration by parts,

$$\begin{aligned}
I_4 &= \int_1^\infty \frac{\cos(h \log ux)}{u^3} df(u) \\
&= \left(1 + \frac{B}{2}\right) \cos(h \log x) + \int_1^\infty \frac{f(u)}{u^4} [3 \cos(h \log ux) + h \sin(h \log ux)] du
\end{aligned}$$

because $f(1) = -1 - \frac{B}{2}$. Combining the results for I_1 , I_2 , I_3 and I_4 , we have the lemma.

Lemma 6.6.

$$\begin{aligned}
&\int_0^1 f(u) [\cos(h \log ux) - h \sin(h \log ux)] du \\
&= -\frac{1}{2} \left[\frac{4+3h^2}{(4+h^2)^2} \cos(h \log x) + \frac{h^3}{(4+h^2)^2} \sin(h \log x) \right] \\
&\quad - \left(\frac{1}{2} + \frac{B}{2} \right) \left[\frac{2+h^2}{4+h^2} \cos(h \log x) - \frac{h}{4+h^2} \sin(h \log x) \right] \\
&\quad - \frac{1}{2} \left[\frac{3+h^2}{9+h^2} \cos(h \log x) - \frac{2h}{9+h^2} \sin(h \log x) \right].
\end{aligned}$$

Proof: The key is $\epsilon(u) = \frac{1}{2} \log u - u$ when $0 \leq u \leq 1$ (see (3)). So,

$$f(u) = \int_0^u \epsilon(v) - \frac{B}{2} dv = \frac{1}{2} u \log u - \left(\frac{1}{2} + \frac{B}{2}\right)u - \frac{1}{2}u^2.$$

Putting this into the integral and evaluating the integral piece by piece with suitable substitution and Lemma 2.7, one gets the lemma.

Proof of Theorem 1.4: First observe that when $T \leq x \leq T^{2-29\epsilon}$, the error terms in Theorem 1.2 is $O\left(\frac{hT}{\log^{M-2}T}\right)$. Rewrite Theorem 1.2 as

$$F_h(x, T) = T_1 + T_2 + T_3 + T_4 + T_5 + O\left(\frac{\tilde{h}T}{\log^{M-2}T}\right).$$

Since $\frac{\sin u}{u} = 1 + O(u^2)$,

$$\begin{aligned} T_3 &= -\frac{T}{\pi} \left[\frac{3 \cos(h \log x)}{9 + h^2} + \frac{h \sin(h \log x)}{9 + h^2} \right] + O\left(T\left(\frac{T}{x}\right)^2\right), \\ T_4 &= -\frac{T}{\pi} \left[\frac{\cos(h \log x)}{1 + h^2} - \frac{h \sin(h \log x)}{1 + h^2} \right] + O\left(T\left(\frac{T}{x}\right)^2\right). \end{aligned}$$

By Lemma 6.5,

$$\begin{aligned} T_5 &= \frac{T}{\pi} \sum_{k=1}^{\infty} \frac{\mathfrak{S}(k)}{k^2} \int_0^1 y \cos\left(h \log \frac{kx}{y}\right) dy + O\left(T\left(\frac{T}{x}\right)^2\right) \\ &= \frac{T}{\pi} \left[\frac{\cos(h \log x)}{1 + h^2} - \frac{h \sin(h \log x)}{1 + h^2} \right] \\ &\quad - \frac{T}{\pi} \left[\frac{4 - h^2}{2(4 + h^2)^2} \cos(h \log x) - \frac{2h}{(4 + h^2)^2} \sin(h \log x) \right] \\ &\quad + \frac{T}{\pi} \frac{B}{2} \left[\frac{2 \cos(h \log x)}{4 + h^2} - \frac{h \sin(h \log x)}{4 + h^2} \right] + \frac{T}{\pi} \left(1 + \frac{B}{2}\right) \cos(h \log x) \\ &\quad + \frac{T}{\pi} \int_1^{\infty} \frac{f(u)}{u^4} \left[3 \cos(h \log ux) + h \sin(h \log ux) \right] du + O\left(T\left(\frac{T}{x}\right)^2\right). \end{aligned}$$

By Lemma 4.3, (6), Lemma 6.2, Lemma 6.3, Lemma 6.4 and Lemma 6.6,

$$\begin{aligned}
T_2 &= -\frac{2x \cos(h \log x)}{\pi(4+h^2)} \int_1^\infty \frac{\sin \frac{Ty}{x}}{y^2} dy - \frac{4T}{\pi} \cos(h \log x) \int_1^\infty \frac{f(y)}{y^2} dy \\
&\quad + \frac{T}{\pi} \int_1^\infty (G_1(y) + G_2(y)) dy + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^M T}\right) \\
&= -\frac{2x \cos(h \log x)}{\pi(4+h^2)} \int_1^\infty \frac{\sin \frac{Ty}{x}}{y^2} dy + O\left(\frac{\tilde{h}T}{\log^M T}\right) \\
&\quad + \frac{T}{\pi} \int_0^1 f(u) [\cos(h \log ux) - h \sin(h \log ux)] du \\
&\quad - \frac{T}{\pi} \int_1^\infty \frac{f(u)}{u^4} [3 \cos(h \log ux) + h \sin(h \log ux)] du + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) \\
&= -\frac{2x \cos(h \log x)}{\pi(4+h^2)} \int_1^\infty \frac{\sin \frac{Ty}{x}}{y^2} dy + O\left(\frac{\tilde{h}T}{\log^M T}\right) \\
&\quad - \frac{T}{\pi} \frac{1}{2} \left[\frac{4+3h^2}{(4+h^2)^2} \cos(h \log x) + \frac{h^3}{(4+h^2)^2} \sin(h \log x) \right] \\
&\quad - \frac{T}{\pi} \left(\frac{1}{2} + \frac{B}{2} \right) \left[\frac{2+h^2}{4+h^2} \cos(h \log x) - \frac{h}{4+h^2} \sin(h \log x) \right] \\
&\quad - \frac{T}{\pi} \frac{1}{2} \left[\frac{3+h^2}{9+h^2} \cos(h \log x) - \frac{2h}{9+h^2} \sin(h \log x) \right] \\
&\quad - \frac{T}{\pi} \int_1^\infty \frac{f(u)}{u^4} [3 \cos(h \log ux) + h \sin(h \log ux)] du + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right).
\end{aligned}$$

Therefore, with miraculous cancellations,

$$\begin{aligned}
T_2 + T_3 + T_4 + T_5 &= -\frac{2x \cos(h \log x)}{\pi(4+h^2)} \int_1^\infty \frac{\sin \frac{Ty}{x}}{y^2} dy + \frac{T}{\pi} \frac{2B \cos(h \log x)}{4+h^2} \\
&\quad + \frac{T}{\pi} \frac{4h \sin(h \log x)}{(4+h^2)^2} + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^M T}\right).
\end{aligned}$$

By Lemma 6.1 and $B = -C_0 - \log 2\pi$,

$$\begin{aligned}
F_h(x, T) &= \frac{T}{\pi} \left[\frac{2 \cos(h \log x)}{4+h^2} \log x \right] - \frac{2T \cos(h \log x)}{\pi(4+h^2)} \left[\frac{\sin(T/x)}{T/x} - ci\left(\frac{T}{x}\right) \right] \\
&\quad + \frac{T}{\pi} \frac{2B \cos(h \log x)}{4+h^2} + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right) \\
&= \frac{T}{\pi} \left[\frac{2 \cos(h \log x)}{4+h^2} \log x \right] - \frac{2T \cos(h \log x)}{\pi(4+h^2)} \left[1 - C_0 - \log \frac{T}{x} \right. \\
&\quad \left. + C_0 + \log 2\pi \right] + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right) \\
&= \frac{T}{2\pi} \log \frac{T}{2\pi e} \left[\frac{4 \cos(h \log x)}{4+h^2} \right] + O\left(T\left(\frac{T}{x}\right)^{1/2-\epsilon}\right) + O\left(\frac{\tilde{h}T}{\log^{M-2} T}\right).
\end{aligned}$$

7 Sketch for Conjecture 1.2

Fix $\alpha > 0$. Let $r(u)$ be an even function which is almost the characteristic function of the interval $[-\alpha, \alpha]$ with $\hat{r}(\alpha) \ll \frac{1}{\alpha^2}$ (see page 87 of [1] for detail construction). We use Conjecture 1.1 to compute the right hand side of (1).

$$\begin{aligned}
I &= \int_{-\infty}^{\infty} F_h(\alpha) \hat{r}(\alpha) d\alpha = 2 \int_0^{\infty} F_h(\alpha) \hat{r}(\alpha) d\alpha \\
&= 2(1 + o(1)) \log T \int_0^1 T^{-2\alpha} \hat{r}(\alpha) d\alpha + 2 \frac{4}{4 + h^2} \int_0^1 \alpha \cos(h \log T \alpha) \hat{r}(\alpha) d\alpha \\
&\quad + 2 \frac{4}{4 + h^2} \int_1^{\infty} \cos(h \log T \alpha) \hat{r}(\alpha) d\alpha + O\left(\frac{1}{A}\right) + o(1) \\
&= \frac{4}{4 + h^2} \int_{-\infty}^{\infty} \cos(h \log T \alpha) \hat{r}(\alpha) d\alpha - \frac{4}{4 + h^2} \int_{-1}^1 (1 - |\alpha|) \cos(h \log T \alpha) \hat{r}(\alpha) d\alpha \\
&\quad + (1 + o(1)) \log T \int_{-\infty}^{\infty} T^{-2|\alpha|} \hat{r}(\alpha) d\alpha + O\left(\frac{1}{A}\right) + o(1) \\
&= \frac{4}{4 + h^2} \int_{-\infty}^{\infty} \hat{r}_1(\alpha) d\alpha - \frac{4}{4 + h^2} \int_{-1}^1 (1 - |\alpha|) \hat{r}_1(\alpha) d\alpha \\
&\quad + (1 + o(1)) \log T \int_{-\infty}^{\infty} T^{-2|\alpha|} \hat{r}(\alpha) d\alpha + O\left(\frac{1}{A}\right) + o(1) \\
&= \frac{4}{4 + h^2} I_1 - \frac{4}{4 + h^2} I_2 + (1 + o(1)) I_3 + O\left(\frac{1}{A}\right) + o(1)
\end{aligned}$$

where $r_1(u) = r(u + \frac{h \log T}{2\pi})$. As $\int_{-\infty}^{\infty} \hat{r}_1(\alpha) d\alpha = r_1(0)$,

$$I_1 = r_1(0) = r\left(\frac{h \log T}{2\pi}\right).$$

By $\int f \hat{g} = \int \hat{f} g$, the transform pair and the definition of $r(u)$,

$$f(t) = \max(1 - |t|, 0), \quad \hat{f}(u) = \left(\frac{\sin \pi u}{\pi u}\right)^2,$$

$$I_2 = \int_{-\infty}^{\infty} r_1(u) \left(\frac{\sin \pi u}{\pi u}\right)^2 du = \int_{-\alpha + h \log T / (2\pi)}^{\alpha + h \log T / (2\pi)} \left(\frac{\sin \pi u}{\pi u}\right)^2 du + o(1).$$

Similarly, by the transform pair

$$f(t) = e^{-2a|t|}, \quad \hat{f}(u) = \frac{4a}{4a^2 + (2\pi u)^2},$$

$$I_3 = \int_{-\alpha}^{\alpha} \frac{4 \log^2 T}{4 \log^2 T + (2\pi u)^2} du + o(1) = \int_{-\alpha + h \log T / (2\pi)}^{\alpha + h \log T / (2\pi)} 1 du + o(1).$$

Therefore,

$$I = \frac{4}{4 + h^2} r\left(\frac{h \log T}{2\pi}\right) + \int_{-\alpha + h \log T / (2\pi)}^{\alpha + h \log T / (2\pi)} 1 - \frac{4}{4 + h^2} \left(\frac{\sin \pi u}{\pi u}\right)^2 du + O\left(\frac{1}{A}\right) + o(1). \quad (16)$$

Now, the left hand side of (1) is

$$\frac{4}{4+h^2} r\left(\frac{h \log T}{2\pi}\right) + \left(\frac{T}{2\pi} \log T\right)^{-1} \sum_{\substack{0 < \gamma \neq \gamma' \leq T \\ |\gamma - \gamma' - h| \leq 2\pi\alpha / \log T}} (1 + o(1)). \quad (17)$$

Combining (16) and (17), we have Conjecture 1.2 by making A arbitrarily large. The only shaky point in the above argument is the error analysis. All of these become rigorous following page 87 – 90 of [1].

References

- [1] T.H. Chan, *Pair Correlation and Distribution of Prime Numbers*, Thesis, U. of Mich., Ann Arbor, 2002.
- [2] T.H. Chan, *On a conjecture of Liu and Ye*, to appear.
- [3] T.H. Chan, *More precise Pair Correlation Conjecture on the zeros of the Riemann zeta function*, submitted.
- [4] J.B. Friedlander and D.A. Goldston, *Some singular series averages and the distribution of Goldbach numbers in short intervals*, Illinois J. Math. **39** (1995), 158-180.
- [5] D.A. Goldston, *Large Differences between Consecutive Prime Numbers*, Thesis, U. of Calif., Berkeley, 1981.
- [6] D.A. Goldston and S.M. Gonek, *Mean value theorems for long Dirichlet polynomials and tails of Dirichlet series*, Acta Arith. (2) **84** (1998), 155-192.
- [7] D.A. Goldston, S.M. Gonek, A.E. Ozluk and C. Synder, *On the pair correlation of zeros of the Riemann zeta-function*, Proc. London Math. Soc. (3) **80** (2000), 31-49.
- [8] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press, sixth edition, 2000.
- [9] H.L. Montgomery, *The pair correlation of zeros of the zeta function*, Analytic Number Theory (St. Louis Univ., 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc., Providence, 1973, pp. 181-193.
- [10] H.L. Montgomery and R.C. Vaughan, *Hilbert's inequality*, J. London Math. Soc. (2) **8** (1974), 73-82.

Tsz Ho Chan
 Case Western Reserve University
 Mathematics Department, Yost Hall 220
 10900 Euclid Avenue

Cleveland, OH 44106-7058
USA
txc50@po.cwru.edu