

## LOCAL COHOMOLOGY AND GORENSTEIN INJECTIVE DIMENSION OVER LOCAL HOMOMORPHISMS

LEILA KHATAMI, MASSOUD TOUSI, AND SIAMAK YASSEMI

ABSTRACT. Let  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a local homomorphism of commutative noetherian local rings. Suppose that  $M$  is a finitely generated  $S$ -module. A generalization of Grothendieck's non-vanishing theorem is proved for  $M$  (i.e. the Krull dimension of  $M$  over  $R$  is the greatest integer  $i$  for which the  $i$ th local cohomology module of  $M$  with respect to  $\mathfrak{m}$ ,  $H_{\mathfrak{m}}^i(M)$ , is non-zero). It is also proved that the Gorenstein injective dimension of  $M$ , if finite, is bounded below by dimension of  $M$  over  $R$  and is equal to the supremum of  $\text{depth } R_{\mathfrak{p}}$ , where  $\mathfrak{p}$  runs over the support of  $M$  as an  $R$ -module.

### INTRODUCTION

Let  $R$  be a non-trivial commutative noetherian local ring. An  $R$ -module  $M$  is called *finite over a local homomorphism* if there exists a local homomorphism of noetherian local rings  $R \rightarrow S$  such that  $M$  is a finite (that is finitely-generated)  $S$ -module and the  $S$ -action is compatible with the action of  $R$ . Studied by Apassov, Avramov, Christensen, Foxby, Iyengar, Miller, Sather-Wagstaff and others, cf. [1, 3, 5, 10, 13], homological properties of finite modules over (local) homomorphisms are shown to extend the those of finite modules.

It is well-known that over a local ring, injective dimension of a finite module is either infinite or equals the depth of the ring and is not less than the dimension of the module. This is known as the *Bass formula*.

A similar formula is also showed to be true for a module finite over a local homomorphism (cf. [5, 18, 14]).

In this paper, we deal with *Gorenstein injective* dimension of a module which is finite over a local homomorphism. Introduced by Enochs and Jenda [11, 12], Gorenstein injective dimension is the dual notion to the G-dimension, due to Auslander and Bridger [2].

In [14] and [19], a generalized Bass formula is proved for finite modules of finite Gorenstein injective dimension.

**Theorem.** Let  $R$  be a noetherian local ring and  $M$  be a finite module of finite Gorenstein injective dimension. Then

$$\dim_R M \leq \text{Gid}_R(M) = \sup\{\text{depth } R_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\}.$$

Our main goal is to extend this theorem to modules finite over a local homomorphism. To this end, we provide some preliminary results. The most interesting one among them, is a generalization of the Grothendieck's non-vanishing theorem (Theorem 1.3).

2000 *Mathematics Subject Classification.* 13D05, 13D45, 14B15.

*Key words and phrases.* Local cohomology, Gorenstein injective dimension, Bass' formula.

Emails: [lkhatami@ictp.it](mailto:lkhatami@ictp.it); [mtousi@ipm.ir](mailto:mtousi@ipm.ir); [yassemi@ipm.ir](mailto:yassemi@ipm.ir).

**Theorem.** Let  $(R, \mathfrak{m})$  be a noetherian local ring and  $M$  an  $R$ -module finite over a local homomorphism. Then  $n = \dim_R M$  is the greatest integer  $i$  for which the  $i$ -th local cohomology module  $H_{\mathfrak{m}}^i(M)$  is non-zero.

Using the auxiliary results, we prove our main theorem (Theorem 2.3) with similar techniques as in [14] and [19]. As a corollary, we also prove that if a local ring  $R$  admits a module  $M$  finite over a local homomorphism with finite Gorenstein injective dimension and maximal dimension over  $R$ , then  $R$  is Cohen-Macaulay.

**Convention.** Throughout this notes all rings are assumed to be commutative, noetherian and local.

## 1. LOCAL COHOMOLOGY OVER A LOCAL HOMOMORPHISM

Let  $(R, \mathfrak{m})$  be a noetherian local ring and  $M$  a finite  $R$ -module of the Krull dimension  $n$ . Then the Grothendieck's non-vanishing theorem says that the local cohomology module  $H_{\mathfrak{m}}^n(M)$  does not vanish (cf. [6, 6.1.4]). In this section, we prove the same non-vanishing result for a module finite over a local homomorphism. First, we prove a couple of auxiliary lemmas.

The main tool we use in our proofs in this paper, is the so-called *Cohen factorization* of a homomorphism.

A local homomorphism  $\varphi : R \rightarrow S$  is said to have a Cohen factorization if it can be represented as a composition  $R \rightarrow R' \rightarrow S$  of local homomorphisms, where  $R'$  is a complete local ring, the map  $R \rightarrow R'$  is flat with regular closed fiber  $R'/\mathfrak{m}R'$  and  $R' \rightarrow S$  is a surjective homomorphism. For any local homomorphism  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  its semi-completion  $\varphi' : R \rightarrow S \rightarrow \hat{S}$  admits a Cohen factorization (cf. [4]).

**Lemma 1.1.** *Let  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a flat local homomorphism. If  $M$  is an  $S$ -module, then*

$$\begin{aligned} \text{Supp}_R(M) &= \{\varphi^{-1}(\mathfrak{q}) \mid \mathfrak{q} \in \text{Supp}_S(M)\} \text{ and} \\ \text{Supp}_S(M) &= \{\mathfrak{q} \in \text{Spec}(S) \mid \varphi^{-1}(\mathfrak{q}) \in \text{Supp}_R(M)\}. \end{aligned}$$

*Proof.* Suppose that  $\mathfrak{q} \in \text{Spec}(S)$  and set  $\mathfrak{p} = \varphi^{-1}(\mathfrak{q})$ . Therefore  $\varphi$  induces a (faithfully) flat local homomorphism  $R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$  and the following isomorphisms hold.

$$\begin{aligned} M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} S_{\mathfrak{q}} &\cong (M \otimes_R R_{\mathfrak{p}}) \otimes_{R_{\mathfrak{p}}} S_{\mathfrak{q}} \\ &\cong M \otimes_R S_{\mathfrak{q}} \\ &\cong M \otimes_R (S \otimes_S S_{\mathfrak{q}}) \\ &\cong M_{\mathfrak{q}} \otimes_R S \end{aligned}$$

If  $\mathfrak{q} \in \text{Supp}_S(M)$  then,  $\varphi$  being faithfully flat,  $M_{\mathfrak{q}} \otimes_R S$  and thus  $M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} S_{\mathfrak{q}}$  are non-zero. Hence  $\mathfrak{p} = \varphi^{-1}(\mathfrak{q}) \in \text{Supp}_R(M)$ .

On the other hand, if  $\mathfrak{p} \in \text{Supp}_R(M)$ , using faithfully flatness of  $\varphi$  we can find a prime  $\mathfrak{q} \in \text{Spec}(S)$  such that  $\varphi^{-1}(\mathfrak{q}) = \mathfrak{p}$  and the above isomorphisms give  $M_{\mathfrak{q}} \otimes_R S \neq 0$  and hence  $\mathfrak{q} \in \text{Supp}_S(M)$ .  $\square$

**Lemma 1.2.** *Let  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a flat local homomorphism and  $M$  a non-zero  $S$ -module. Then*

$$\dim_S M \geq \dim_R M + \dim_S S/\mathfrak{m}S.$$

*Proof.* First note that since  $\varphi$  is flat, every prime ideal of  $S$  which contains  $\mathfrak{m}S$  minimally, contracts to  $\mathfrak{m}$  via  $\varphi$ . Let  $\mathfrak{q} \in \text{Spec}(S)$  be one such prime. By Lemma 1.1,  $\mathfrak{q} \in \text{Supp}_S(M)$ .

Suppose that  $\dim_R M = n$  and

$$\mathfrak{p}_0 \subset \cdots \subset \mathfrak{p}_n = \mathfrak{m}$$

is a maximal chain of prime ideals in  $\text{Supp}_R(M)$ . Since  $\varphi$  is flat and  $\varphi^{-1}(\mathfrak{q}) = \mathfrak{m}$  we get a strict chain in  $\text{Spec}(S)$ ,

$$\mathfrak{q}_0 \subset \cdots \subset \mathfrak{q}_n = \mathfrak{q}$$

such that  $\varphi^{-1}(\mathfrak{q}_i) = \mathfrak{p}_i$  for  $0 \leq i \leq n$ . Using Lemma 1.1 again, this is a chain in  $\text{Supp}_S(M)$  which gives the inequalities

$$\dim_S M \geq \dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}} + \dim_S S/\mathfrak{q} \geq n + \dim_S S/\mathfrak{q}.$$

In particular, we can choose  $\mathfrak{q}$  such that  $\dim_S S/\mathfrak{m}S = \dim_S S/\mathfrak{q}$  to get the desired inequality.  $\square$

Now we are ready to prove the main theorem of this section.

**Theorem 1.3.** *Let  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a local homomorphism and  $M$  a non-zero finite  $S$ -module. If  $n = \dim_R M$  then*

$$H_{\mathfrak{m}}^n(M) \neq 0.$$

*Proof.* First suppose that  $\varphi$  is a flat homomorphism.

As in the proof of Lemma 1.2, we pick a prime  $\mathfrak{q} \in \text{Supp}_S(M)$  which contains  $\mathfrak{m}S$  minimally. Therefore, the induced homomorphism  $R \rightarrow S_{\mathfrak{q}}$  is faithfully flat and has zero-dimensional fiber  $S_{\mathfrak{q}}/\mathfrak{m}S_{\mathfrak{q}}$ . Furthermore, the inequality  $\dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}} \geq \dim_R M$  holds.

On the other hand, the isomorphism  $H_{\mathfrak{m}}^i(M_{\mathfrak{q}}) \cong H_{\mathfrak{q}S_{\mathfrak{q}}}^i(M_{\mathfrak{q}})$  holds for every  $i \geq 0$ . Since  $M_{\mathfrak{q}}$  is a finite  $S_{\mathfrak{q}}$ -module,  $H_{\mathfrak{q}S_{\mathfrak{q}}}^d(M_{\mathfrak{q}}) \neq 0$  for  $d = \dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}}$  and therefore we get  $\dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}} \leq \dim_R M_{\mathfrak{q}}$ .

Note that  $\text{Supp}_R(M_{\mathfrak{q}}) \subseteq \text{Supp}_R(M)$  and hence

$$\dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}} \leq \dim_R M_{\mathfrak{q}} \leq \dim_R M \leq \dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}}.$$

Thus the equality  $\dim_{S_{\mathfrak{q}}} M_{\mathfrak{q}} = \dim_R M = n$  holds.

Finally, the following isomorphisms give the desired non-vanishing in this case.

$$H_{\mathfrak{m}}^n(M_{\mathfrak{q}}) \cong H_{\mathfrak{m}S_{\mathfrak{q}}}^n(M_{\mathfrak{q}}) \cong H_{\mathfrak{m}}^n(M) \otimes_R S_{\mathfrak{q}}$$

In general,  $R \rightarrow S \rightarrow \hat{S}$  admits a Cohen factorization  $R \rightarrow R' \rightarrow \hat{S}$ . Since  $M$  is finite over  $S$ , so is  $\hat{M} = M \otimes_S \hat{S}$  as an  $\hat{S}$ -module and thus as an  $R'$ -module. It is also easy to see that  $\text{Supp}_R(\hat{M}) = \text{Supp}_R(M)$  and consequently we have  $\dim_R \hat{M} = \dim_R M = n$ . So using the first part of the proof, we get

$$H_{\mathfrak{m}}^n(\hat{M}) \neq 0.$$

Using the isomorphisms

$$H_{\mathfrak{m}}^n(\hat{M}) \cong H_{\mathfrak{m}\hat{S}}^n(\hat{M}) \cong H_{\mathfrak{m}S}^n(M) \otimes_S \hat{S}$$

we conclude that  $H_{\mathfrak{m}S}^n(M) \cong H_{\mathfrak{m}}^n(M)$  is non-zero.  $\square$

## 2. A GENERALIZED BASS FORMULA

This section is dedicated to proving a Bass-type formula for a module which is finite over a local homomorphism and has finite Gorenstein injective dimension.

**Definition 2.1.** *An  $R$ -module  $G$  is said to be Gorenstein injective if and only if there exists an exact complex of injective  $R$ -modules,*

$$I = \cdots \rightarrow I_2 \rightarrow I_1 \rightarrow I_0 \rightarrow I_{-1} \rightarrow I_{-2} \rightarrow \cdots$$

*such that the complex  $\text{Hom}_R(J, I)$  is exact for every injective  $R$ -module  $J$  and  $G$  is the kernel in degree 0 of  $I$ . The Gorenstein injective dimension of an  $R$ -module  $M$ ,  $\text{Gid}_R(M)$ , is defined to be the infimum of integers  $n$  such that there exists an exact sequence*

$$0 \rightarrow M \rightarrow G_0 \rightarrow G_{-1} \rightarrow \cdots \rightarrow G_{-n} \rightarrow 0$$

*with all  $G_i$ 's Gorenstein injective.*

**Proposition 2.2.** *Let  $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a local homomorphism and  $M$  a finite  $S$ -module. If  $\mathfrak{p} \in \text{Supp}_R(M)$  then  $M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}} \neq 0$ .*

*Proof.* First suppose that  $\varphi$  admits a Cohen factorization  $R \rightarrow R' \rightarrow S$ . Since  $R' \rightarrow S$  is surjective,  $M$  is also finite as an  $R'$ -module. Thus we can assume that  $\varphi$  is flat. By Lemma 1.1, there exists a  $\mathfrak{q} \in \text{Supp}_S(M)$  such that  $\varphi^{-1}(\mathfrak{q}) = \mathfrak{p}$ . By NAK we have  $M_{\mathfrak{q}}/\mathfrak{p}M_{\mathfrak{q}} \neq 0$ . Since  $M_{\mathfrak{q}}/\mathfrak{p}M_{\mathfrak{q}} \cong (T^{-1}M/\mathfrak{p}T^{-1}M)_{T^{-1}\mathfrak{q}}$ , where  $T = \varphi(R - \mathfrak{p})$ , we get  $M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}} \cong T^{-1}M/\mathfrak{p}T^{-1}M \neq 0$ .

In general  $R \rightarrow S \rightarrow \hat{S}$  admits a Cohen factorization. If  $\mathfrak{p} \in \text{Supp}_R(M)$  then  $\mathfrak{p} \in \text{Supp}_R(M \otimes_S \hat{S})$  as well and then, setting  $\hat{M} = M \otimes_S \hat{S}$ , we have  $M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}} \otimes_S \hat{S} \cong \hat{M}_{\mathfrak{p}}/\mathfrak{p}\hat{M}_{\mathfrak{p}} \neq 0$  and we are done.  $\square$

**Theorem 2.3.** *Let  $R$  be a noetherian local ring and  $M$  be an  $R$ -module finite over a local homomorphism with  $\text{Gid}_R(M) < \infty$ . Then*

$$\dim_R M \leq \text{Gid}_R(M) = \sup\{\text{depth } R_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\}.$$

*Proof.* To prove the first inequality we use the proof of [15, 3.4] to see that  $H_{\mathfrak{m}}^i(M) = 0$  for  $i > \text{Gid}_R(M)$ . Thus, by Theorem 1.3, we have

$$\dim_R M \leq \text{Gid}_R(M).$$

To prove the formula, we use [9, 2.18], to get an exact sequence

$$0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$$

with  $K$  Gorenstein injective and  $\text{inj.dim}_R(L) = \text{Gid}_R(M)$ . By [7], we have

$$\text{inj.dim}_R(L) = \sup\{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(L)\},$$

where  $\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \inf\{i \mid \text{Tor}_i^{R_{\mathfrak{p}}}(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}, M_{\mathfrak{p}}) \neq 0\}$ .

For any  $\mathfrak{p} \in \text{Supp}_R(M)$ , we get an exact sequence  $L_{\mathfrak{p}}/\mathfrak{p}L_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}} \rightarrow 0$  which gives  $L_{\mathfrak{p}}/\mathfrak{p}L_{\mathfrak{p}} \neq 0$ , by Proposition 2.2. Therefore we have

$$\begin{aligned} \sup\{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(L)\} &\geq \\ \sup\{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\} &= \\ \sup\{\text{depth } R_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\} &\geq 0. \end{aligned}$$

This proves the desired formula for  $M$  a Gorenstein injective  $R$ -module.

For any  $\mathfrak{p} \in \text{Supp}_R(L) - \text{Supp}_R(M)$ , we get  $L_{\mathfrak{p}} \cong K_{\mathfrak{p}}$ . Since  $K$  is Gorenstein injective, there exists an exact sequence

$$\cdots \rightarrow I_1 \rightarrow I_0 \rightarrow L_{\mathfrak{p}} \rightarrow 0,$$

where for each  $\ell \geq 0$ ,  $I_{\ell}$  is an injective  $R_{\mathfrak{p}}$ -module.

Set  $K_{\ell} = \ker(I_{\ell-1} \rightarrow I_{\ell})$ . Then for any  $R_{\mathfrak{p}}$ -module  $T$  with  $\text{proj.dim}_{R_{\mathfrak{p}}}(T) = t < \infty$  we have  $\text{Ext}_{R_{\mathfrak{p}}}^i(T, L_{\mathfrak{p}}) \cong \text{Ext}_{R_{\mathfrak{p}}}^{i+t}(T, K_t) = 0$  for  $i > 0$ . Thus, by [8, 5.3(c)], we get

$$\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \leq 0.$$

Therefore, if  $\text{Gid}_R(M) = \text{inj.dim}_R(L) > 0$  then we have

$$\begin{aligned} \text{Gid}_R(M) &= \text{inj.dim}_R(L) = \\ &= \sup\{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(L)\} = \\ &= \sup\{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} L_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\} = \\ &= \sup\{\text{depth } R_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Supp}_R(M)\}. \end{aligned}$$

□

The following corollary of Theorem 2.3, gives a sufficient condition for Cohen-Macaulayness of a noetherian local ring. This result generalizes [17, 3.5] as well as [19, 1.3].

**Corollary 2.4.** *Let  $R$  be a local ring and  $M$  an  $R$ -module finite over a local homomorphism. If  $M$  has finite Gorenstein dimension and maximal Krull dimension over  $R$  (i.e.  $\text{Gid}_R(M) < \infty$  and  $\dim_R M = \dim R$ ), then  $R$  is Cohen-Macaulay.*

*Proof.* By 2.3, there is a prime ideal  $\mathfrak{p} \in \text{Supp}_R(M)$  such that

$$\dim R = \dim_R M \leq \text{Gid}_R(M) = \text{depth } R_{\mathfrak{p}} \leq \dim R_{\mathfrak{p}} \leq \dim R.$$

Therefore,  $\mathfrak{p}$  must be the maximal ideal of  $R$  and thus  $R$  is Cohen-Macaulay. □

**Corollary 2.5.** *Let  $R$  be an almost Cohen-Macaulay local ring (i.e.  $\dim(R) - \text{depth}(R) \leq 1$ ) and let  $M$  be an  $R$ -module finite over a local homomorphism. If  $M$  has finite Gorenstein injective dimension over  $R$  then  $\text{Gid}_R(M) = \text{depth } R$ .*

*Proof.* It is enough to use Theorem 2.3 and the fact that over an almost Cohen-Macaulay ring  $R$  the inequality  $\text{depth } R_{\mathfrak{p}} \leq \text{depth } R_{\mathfrak{q}}$  holds for  $\mathfrak{p} \subseteq \mathfrak{q}$  in  $\text{Spec}(R)$ . □

Note that Corollary 2.5 can be considered as a generalized Bass formula for Gorenstein injective dimension.

We conclude this paper with a change of ring result for Gorenstein injective dimension.

**Corollary 2.6.** *Let  $(R, \mathfrak{m})$  be a local ring and  $M$  be an  $R$ -module finite over a local homomorphism. If  $x \in \mathfrak{m}$  is an  $R$ - and  $M$ -regular element, then*

$$\text{Gid}_{R/xR}(M/xM) \leq \text{Gid}_R(M) - 1.$$

Furthermore, the equality holds when  $R$  is almost Cohen-Macaulay and  $\text{Gid}_R(M) < \infty$ .

*Proof.* If  $M$  has finite Gorenstein injective dimension over  $R$  then the proof of [16, Lemma 2] shows that  $\text{Gid}_{R/xR}(M/xM) < \infty$ , too. By 2.2, for any  $\mathfrak{p} \in \text{Supp}_R(M)$  with  $x \in \mathfrak{p}$ , the  $R_{\mathfrak{p}}$ -module  $M_{\mathfrak{p}}/xM_{\mathfrak{p}}$  is non-zero. Therefore,

$$\text{Supp}_{R/xR}(M/xM) = \{\mathfrak{p}/xR \mid \mathfrak{p} \in \text{Supp}_R(M) \text{ with } x \in \mathfrak{p}\}.$$

Thus Theorem 2.3 gives the desired inequality. In the case of an almost Cohen-Macaulay base ring the equality is a consequence of 2.5.  $\square$

#### REFERENCES

1. D. Apassov, *Almost finite modules*, Comm. Algebra **27** (1999), 919–931.
2. M. Auslander and M. Bridger, *Stable module theory*, Memoirs of the American Mathematical Society, No. **94**, American Mathematical Society, Providence, R.I. 1969.
3. L. L. Avramov and H-B. Foxby, *Homological dimensions of unbounded complexes*, J. Pure Appl. Algebra **71** (1991), 129–155.
4. L. L. Avramov, H-B. Foxby, and B. Herzog, *Structure of local homomorphisms*, J. Algebra **164** (1994), 124–145.
5. L. L. Avramov, S. Iyengar, and C. Miller, *Homology over local homomorphisms*, Amer. J. Math. **128** (2006), 23–90.
6. Brodmann, M. P. and Sharp, R. Y. *Local cohomology: an algebraic introduction with geometric applications*, Cambridge Studies in Advanced Mathematics, **60**, Cambridge University Press, Cambridge, 1998.
7. L. G. Chouinard, *On finite weak and injective dimension*, Proc. Amer. Math. Soc. **60** (1976), 57–60.
8. L. W. Christensen, H-B. Foxby and A. Frankild, *Restricted homological dimensions and Cohen-Macaulayness*, J. Algebra **251** (2002) (1), 479–502.
9. L. W. Christensen, A. Frankild, and H. Holm, *On Gorenstein projective, injective and flat dimensions-A functorial description with applications*, To appear in J. Algebra.
10. L. W. Christensen and S. Iyengar, *Gorenstein dimension of modules over homomorphisms*, to appear in J. Pure Appl. Algebra.
11. E. E. Enochs and O. M. .G. Jenda, *Resolutions by Gorenstein injective and projective modules and modules of finite injective dimension over Gorenstein rings*, Comm. Algebra **23** (1995), 869–877.
12. E. E. Enochs and O. M. .G. Jenda, *Gorenstein injective and flat dimensions*, Math. Japon. **26**(1998), 3489–3496.
13. S. Iyengar and S. Sather-Wagstaff, *G-dimension over local homomorphisms. Applications to the Frobenius endomorphism*, Illinois J. Math. **48** (2004), 241–272.
14. L. Khatami and S. Yassemi, *A Bass formula for Gorenstein injective dimension*, to appear in Comm. Algebra.
15. R. Sazeeleh, *Gorenstein injective modules and local cohomology*, Proc. Amer. Math. Soc. **132** (2004), no. 10, 2885–2891.
16. S. Salarian, S. Sather-Wagstaff and S. Yassemi, *Characterizing local rings via homological dimensions and regular sequences*, to appear in J. Pure Appl. Algebra.
17. R. Takahashi, *The existence of finitely generated modules of finite Gorenstein injective dimension*, to appear in Proc. Amer. Math. Soc.
18. R. Takahashi, Y. Yoshino, *Characterizing Cohen-Macaulay local rings by Frobenius maps*, Proc. Amer. Math. Soc. **132** (2004), 3177–3187
19. S. Yassemi, *A generalization of a theorem of Bass*, to appear in Comm. Algebra.

LEILA KHATAMI, THE ABDUS SALAM ICTP, STRADA COSTIERA 11, 34100 TRIESTE, ITALY

MASSOUD TOUSI, DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, TEHRAN, IRAN,, AND INSTITUTE FOR THEORETICAL PHYSICS AND MATHEMATICS (IPM).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEHRAN, TEHRAN, IRAN,, AND INSTITUTE FOR THEORETICAL PHYSICS AND MATHEMATICS (IPM).