THE RELATIVE EXTREMAL FUNCTION FOR BOREL SETS IN COMPLEX MANIFOLDS

Armen Edigarian* and Ragnar Sigurdsson

July 12, 2006

Abstract

We study a disc formula for the relative extremal function for Borel sets in complex manifolds.

Keywords: Analytic disc, relative extremal function, pluriregular set, pluripolar set, Josefson manifold, weakly regular domain.

Subject Classification (2000): Primary 32U15. Secondary 32U10, 32U30.

1 Introduction

Let X be a complex manifold, PSH(X) be the class of all plurisubharmonic functions on X, and $PSH^{-}(X)$ be the subclass of all non-positive functions. For any subset A of X we define

$$\omega(\cdot, A, X) = \sup\{u \in \mathrm{PSH}^-(X) ; u | A \le -1\} = \sup\{u \in \mathrm{PSH}(X) ; u \le -\chi_A\},\$$

where χ_A denotes the characteristic function of the set A. The least upper semicontinuous majorant $\omega^*(\cdot, A, X)$ of $\omega(\cdot, A, X)$ is plurisubharmonic and it is called the *relative extremal* function for A in X.

Observe that if A is a Borel set, $u \in \mathrm{PSH}(X)$, $u \leq -\chi_A$, x = f(0), where $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$, i.e., f is an analytic disc which extends to a continuous map from the closure $\overline{\mathbb{D}}$ of the unit disc \mathbb{D} to X, then the subaverage property of u implies

$$u(x) \le \frac{1}{2\pi} \int_0^{2\pi} u(f(e^{i\theta})) d\theta \le \frac{1}{2\pi} \int_0^{2\pi} (-\chi_A)(f(e^{i\theta})) d\theta = -\sigma(f^{-1}(A) \cap \mathbb{T}) = -\sigma_f(A),$$

where σ denotes the normalized arc length measure on the unit circle \mathbb{T} and σ_f the image measure (push-forward) of σ under the map f. By taking supremum over all plurisubharmonic $u \leq -\chi_A$ and infimum over all $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$, we get

$$\omega(x, A, X) \leq \Omega(x, A, X) = \inf\{-\sigma_f(A) : f \in \emptyset(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X), f(0) = x\}$$
$$= -\sup\{\sigma_f(A) : f \in \emptyset(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X), f(0) = x\}.$$

^{*}This work is a part of the Research Grant No. 1 PO3A 005 28, which is supported by public means in the programme for promoting science in Poland in the years 2005-2008. The first author is a fellow of Krzyżanowski Fund at the Jagiellonian University. This work was also supported by the Research Fund at the University of Iceland.

In this paper we are mainly concerned with a possible converse of this inequality. If A is an open subset of X, then $\omega(\cdot, A, X) = \Omega(\cdot, A, X)$. This is a special case of Poletsky's theorem which states that if φ is an upper semicontinuous function on X, then for every x in X

$$\sup\{u(x)\,;\,u\in\mathrm{PSH}(X),u\leq\varphi\}=\inf\{\int_{\mathbb{T}}\varphi\circ f\,d\sigma\,;\,f\in\emptyset(\overline{\mathbb{D}},X),f(0)=x\}.$$

See [10], [11], [13], and [15]. Here $\emptyset(\overline{\mathbb{D}},X) \subset \emptyset(\mathbb{D},X) \cap C(\overline{\mathbb{D}},X)$ denotes the set of all closed analytic discs in X, i.e., analytic discs which extend to holomorphic maps in some neighbourhood of $\overline{\mathbb{D}}$. With $-\chi_A$ in the role of φ we get $\omega(\cdot,A,X) = \Omega(\cdot,A,X)$ for every open set A

We say that the subset A of X is pluriregular at the point $x \in A$ if $\omega^*(x, A, X) = -1$, we say that A is locally pluriregular at the point x in A if $\omega^*(x, A \cap U, U) = -1$ for every neighbourhood U of x, and finally we say that A is (locally) pluriregular if A is (locally) pluriregular at each of its points. Note that if A is locally pluriregular, then A is pluriregular and that if A is pluriregular, then $\omega(\cdot, A, X) = \omega^*(\cdot, A, X)$. Our main result of Section 2 is that if A is a locally pluriregular subset of X, then $\Omega(\cdot, \overline{A}, X) \leq \omega(\cdot, A, X)$. (See Th. 2.1.) This is a generalization of Th. 7.2 in Poletsky [13].

Let E be a subset of a complex manifold X. We say that E is pluripolar or locally pluripolar if for any $a \in E$ there exists a neighbourhood U of a in X and $u \in PSH(U)$, $u \not\equiv -\infty$, such that $E \cap U \subset \{x \in U : u(x) = -\infty\}$. We say that $E \subset X$ is globally pluripolar if there exists $u \in PSH(X), u \not\equiv -\infty$, with $E \subset \{x \in X : u(x) = -\infty\}$. Note that any globally pluripolar set is locally pluripolar. Josefson [8] proved that in \mathbb{C}^n every pluripolar set is globally pluripolar. We say that a complex manifold X is a $Josefson \ manifold$ if any locally pluripolar set is globally pluripolar. Bedford [3] has generalized Josefson's theorem to a certain class of complex spaces including Stein manifolds. He also showed that examples, originally given by Grauert [7], of complex manifolds which possess no non-constant holomorphic functions are Josefson manifolds.

In Section 3 we prove that if X is a relatively compact domain in a Josefson manifold and A is a Borel subset of X, then $\Omega(\cdot, A, X) \leq \omega^*(\cdot, A, X)$. (See Th. 3.1.) The main result of Section 3 is that a Josefson manifold X has the property that every bounded plurisubharmonic function on X is constant if and only if for every $p \in X$, every non-pluripolar Borel subset A of X, and every $\varepsilon > 0$ there exists $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ such that f(0) = p and

$$\sigma_f(A) = \sigma(f^{-1}(A) \cap \mathbb{T}) > 1 - \varepsilon.$$

(See Th. 3.3.) As a consequence we get a characterization of pluripolar sets in terms of analytic discs.

In Section 4 we look at Borel subsets A of the boundary ∂D of a relatively compact domain in a complex manifold X. We define the relative extremal function for an open subset U of the boundary as $\omega(\cdot, U, D) = u_{-\chi_U, D}$, where $u_{f,D}$ is the Perron-Bremermann envelope of the boundary function f, and for any subset A of ∂D we define $\omega(\cdot, A, D)$ as the supremum over all $\omega(\cdot, U, D)$ for U open containing A. We call the domain weakly regular if the upper semicontinuous extension $u_{-\chi_U, D}^*$ of $u_{-\chi_U, D}$ to the closure \overline{D} is less than or equal to $-\chi_U$ on ∂D . For any Borel subset A of ∂D we define $\Omega(x, A, D)$ as the infimum over $-\sigma_f(A)$ for $f \in \emptyset(\mathbb{D}, D) \cap C(\overline{\mathbb{D}}, \overline{D})$ with f(0) = x. We prove (see Th. 4.3) that for a weakly regular domain D and every open subset U of ∂D we have $\omega(\cdot, U, D) = \Omega(\cdot, U, D)$ and (see Th. 4.10) $\omega(\cdot, \overline{A}, D) \leq \Omega(\cdot, \overline{A}, D) \leq \omega^*(\cdot, A, D)$, if A is a Borel subset of the form $A = A_1 \cup E$, where A_1 is locally pluriregular with respect to D and E is such that there exists $u \in \mathrm{PSH}^-(D)$, $u \not\equiv -\infty$, and $u^*|_E \equiv -\infty$. It remains an open question if the last result holds for every Borel set A.

2 Construction of analytic discs

We have already seen that for every manifold X and every Borel subset A of X we have $\omega(\cdot, A, X) \leq \Omega(\cdot, A, X)$ and that Poletsky's theorem implies that equality holds if A is open.

Theorem 2.1 Let X be a complex manifold and A be any locally pluriregular subset of X. Then $\Omega(\cdot, \overline{A}, X) \leq \omega(\cdot, A, X)$, in particular, $\Omega(\cdot, A, X) = \omega(\cdot, A, X)$ if A is also closed.

The main argument of the proof consists of an approximation of analytic discs and it appears a few times in this paper. We therefore state it as a separate result. A similar result for domains in \mathbb{C}^n and, more generally, for domains in Banach spaces is proved by Poletsky in [14]. Our proof uses the existence of Stein neighbourhoods of certain sets which was proved by Rosay [15]. For a simplification of his arguments and further development see [6] and [11].

If X is a complex manifold and $d: X \times X \to [0, +\infty)$ is a continuous function vanishing on the diagonal, i.e., d(x,x) = 0 for all $x \in X$, then for any subset A of X we define the diameter of A with respect to d as $\sup\{d(x,y); x,y \in A\}$. In the proof of Theorem 2.1 we will take d as a complete hermitian metric defining the topology of X.

Theorem 2.2 Let X be a complex manifold, $d: X \times X \to [0, +\infty)$ be a continuous function vanishing on the diagonal, $\delta > 0$, and $\{B_j\}$ be a countable family of open subsets in X of diameter less than δ with respect to d. Assume that U and V are open subsets of X and

$$V \subset \cup_j \{x \in B_j ; \omega(x, U \cap B_j, B_j) < -a\},\$$

where $a \in (0,1)$. Let $h \in \mathcal{O}(\overline{\mathbb{D}},X)$ and assume that $\Delta \subset h^{-1}(V) \cap \mathbb{T}$ is a non-empty open set. Then for every $\varepsilon \in (0,1)$ there exist $g \in \mathcal{O}(\overline{\mathbb{D}},X)$ and an open set $\widetilde{\Delta} \subseteq \Delta$ such that

- (1) g(0) = h(0),
- (2) $d(g,h) = \sup_{z \in \overline{\mathbb{D}}} d(g(z), h(z)) \le \delta + \varepsilon$,
- (3) $\sigma(\tilde{\Delta}) \ge (1 \varepsilon)a\sigma(\Delta)$, and
- (4) $g(\tilde{\Delta}) \subset U$.

Proof: For r > 0 we let D_r be the open disc in $\mathbb C$ with radius r and centre at the origin and we assume that $h \in \mathcal O(D_s, X)$ for some s > 1. Fix $\Delta_0 \subset \Delta$ a union of closed arcs such that $\sigma(\Delta_0) > (1 - \varepsilon)\sigma(\Delta)$.

Take $w_0 \in \Delta$. Then $x_0 = h(w_0) \in V$, so there exists a j_0 such that $x_0 \in B_{j_0}$ and $\omega(x_0, U \cap B_{j_0}, B_{j_0}) < -a$. Since U is open, Poletsky's theorem implies that there exists $f_0 \in \emptyset(\overline{\mathbb{D}}, B_{j_0})$ such that $f_0(0) = x_0$ and $\sigma_{f_0}(U) > a$. Let $I_0 \subset f_0^{-1}(U) \cap \mathbb{T}$ be a union

of finite number of closed arcs such that $\sigma(I_0) > a$. By Lemma 2.3 in [10], there exists an open neighbourhood V_0 of $x_0 = f_0(0)$ in X, r > 1, and $f \in \mathcal{O}(D_r \times V_0, B_{j_0})$ such that $f(z, x_0) = f_0(z)$ for all $z \in D_r$ and f(0, x) = x for all $x \in V_0$. By choosing r > 1 sufficiently small and shrinking the neighbourhood V_0 of x_0 , we may assume that $f(z, x) \in U$ for all $z \in I_0$ and $x \in V_0$. We set $F_0(z, w) = f(z, h(w))$ and note that F_0 is defined on $D_r \times h^{-1}(V_0)$ and that $h^{-1}(V_0)$ is a neighbourhood of w_0 .

We apply a compactness argument on Δ_0 and conclude that we may find:

- Open discs U_1, \ldots, U_m centred on \mathbb{T} with mutually disjoint closures such that $U_j \cap \mathbb{T} \subset \Delta$ and $\sigma((U_1 \cup \cdots \cup U_m) \cap \mathbb{T}) > (1 \varepsilon)\sigma(\Delta)$.
- $r_j > 1$, j = 1, ..., m, and holomorphic maps $F_j : D_{r_j} \times U_j \to B_{k(j)}$ with $F_j(0, w) = h(w)$ for all $w \in U_j$.
- Finite unions I_j , j = 1, ..., m, of closed arcs on \mathbb{T} with $\sigma(I_j) > a$ and $F_j(z, w) \in U$ for all $z \in I_j$ and $w \in U_j$.

Take closed arcs J_1, \ldots, J_m in \mathbb{T} such that $J_j \subset U_j \cap \mathbb{T}$ and $\sigma(J_1 \cup \cdots \cup J_m) > (1 - \varepsilon)\sigma(\Delta)$. Let

$$K_0 = \{(w, 0, 0, 0, h(w)); w \in \overline{\mathbb{D}}\} \subset \mathbb{C}^4 \times X$$

and

$$K_j = \{(w, z, 0, 0, F_j(z, w)); w \in J_j, z \in \overline{\mathbb{D}}\} \subset \mathbb{C}^4 \times X, \qquad j = 1, \dots, m.$$

By the proof of Th. 1.2 in [11], there exists a Stein neighbourhood Z of $K_0 \cup K_1 \cup \cdots \cup K_m$ in $\mathbb{C}^4 \times X$. Let $\tau : Z \to \mathbb{C}^N$ be an embedding, $\varkappa : W \to \tau(Z)$ be a holomorphic retraction from a Stein neighbourhood W of $\tau(Z)$ in \mathbb{C}^N , and $\varphi = \operatorname{pr} \circ \tau^{-1} \circ \varkappa : W \to X$ be the holomorphic submersion, where $\operatorname{pr} : \mathbb{C}^4 \times X \to X$ is the projection.

We let $\rho: \mathbb{T} \to [0,1]$ be a C^{∞} function such that $\rho = 0$ on $\mathbb{T} \setminus (\cup_j J_j)$ and $\rho = 1$ on a subset of $\cup_j J_j$ such that $\sigma(\{w \in \mathbb{T} : \rho(w) = 1\}) > (1 - \varepsilon)\sigma(\Delta)$. We define a C^{∞} map $F: D_s \times \mathbb{T} \to X$ by

$$F(z,w) = \begin{cases} F_j(\rho(w)z, w), & w \in J_j, \ j = 1, \dots, m, \\ h(w), & w \notin \cup_j J_j. \end{cases}$$

Since Z is a neighbourhood of $K_0 \cup \cdots \cup K_m$ we can replace s > 1 by a smaller number and can define $\widetilde{F}: D_s \times \mathbb{T} \to W$ by $\widetilde{F}(z,w) = \tau(w,z,0,0,F(z,w))$ and $\widetilde{h}: D_s \to W$ by $\widetilde{h}(w) = \tau(w,0,0,0,h(w))$. We note that $\widetilde{F}(0,w) = \widetilde{h}(w)$ for all $w \in \mathbb{T}$.

In exactly the same way as in the proof of Lemma 2.6 in [10] we construct a sequence $\widetilde{F}_j \in \mathcal{O}(D_s \times A_j, W), j \geq j_0$, where A_j is an open annulus containing \mathbb{T} , such that

- $\widetilde{F}_j \to \widetilde{F}$ uniformly on $D_s \times \mathbb{T}$ as $j \to \infty$,
- there is an integer $k_j \geq j$ such that for all $k \geq k_j$ the map $\widetilde{F}_j(zw^k, w)$ can be extended to a map $\widetilde{G}_j \in \mathcal{O}(D_{s_j} \times D_{s_j}, W)$, where $s_j \in (1, s)$, and
- $\widetilde{G}_j(0, w) = \widetilde{h}(w)$ for all $w \in D_{s_j}$.

We need to estimate $\sup_{w \in \mathbb{T}} \sup_{z \in \overline{\mathbb{D}}} d(\varphi(\widetilde{F}_j(z, w)), h(w))$. Since $\widetilde{F}_j \to \widetilde{F}$ uniformly on $\overline{\mathbb{D}} \times \mathbb{T}$ as $j \to \infty$ and $\varphi(\widetilde{F}) = F$ we have

$$\sup_{z\in\overline{\mathbb{D}},w\in\mathbb{T}}d(\varphi(\widetilde{F}_{j}(z,w)),h(w))\to \sup_{z\in\overline{\mathbb{D}},w\in\mathbb{T}}d(F(z,w),h(w)), \qquad j\to\infty.$$

We have $\sup_{z\in\overline{\mathbb{D}}} d(F(z,w),h(w)) = 0$ for all $w\in\mathbb{T}\setminus(\cup_j J_j)$ and since F_j takes values in $B_{k(j)}$, we have $\sup_{z\in\overline{\mathbb{D}}} d(F(z,w),h(w)) \leq \delta$ for all $w\in J_j$. Hence

$$\limsup_{j\to\infty} \left[\sup_{z\in\overline{\mathbb{D}},w\in\mathbb{T}} d(\varphi(\widetilde{F}_j(z,w)),h(w)) \right] \leq \delta.$$

Take $j \geq j_0$ so that $\sup_{z \in \overline{\mathbb{D}}, w \in \mathbb{T}} d(\varphi(\widetilde{F}_j(z, w)), h(w)) \leq \delta + \varepsilon/2$. There exists $t \in (0, 1)$ such that $\sup_{z \in \overline{\mathbb{D}}, w \in [t, 1]} d(\varphi(\widetilde{F}_j(z, w)), h(w)) < \delta + \varepsilon$. Note that

$$\sup_{z \in \overline{\mathbb{D}}, |w| \le t} d(\varphi(\widetilde{F}_j(zw^k, w)), h(w)) \to 0$$

as $k \to \infty$, so for sufficiently large k we have

$$\sup_{z,w\in\overline{\mathbb{D}}} d(\varphi(\widetilde{F}_j(zw^k,w)),h(w)) < \delta + \varepsilon.$$

We set $G(z,w) = \varphi(\widetilde{F}_j(zw^k,w))$. Then $G \in \mathcal{O}(\overline{\mathbb{D}}^2,X)$ and G(0,w) = h(w) for all $w \in \mathbb{D}$. Put $C = \bigcup_j (I_j \times \widetilde{J}_j)$, where $\widetilde{J}_j = \{w \in J_j ; \rho(w) = 1\}$. If $\sigma_2 = \sigma \times \sigma$ is the product measure on the torus \mathbb{T}^2 , then

$$\sigma_2(C) = \sum_j \sigma_2(I_j \times \widetilde{J}_j) = \sum_j \sigma(I_j)\sigma(\widetilde{J}_j) > a(1 - \varepsilon)\sigma(\Delta).$$

The map $\mathbb{T}^2 \ni (z,w) \to (zw^k,w) \in \mathbb{T}^2$ is an automorphism with the absolute value of the Jacobian equal to 1. Therefore the measure of the set $\widetilde{C} = \{(z,w) \in \mathbb{T}^2 : (zw^k,w) \in C\}$ is equal to $\sigma_2(C)$. By Fubini's theorem there is a $\theta \in [0,2\pi)$ such that $\sigma(C') \geq \sigma_2(\widetilde{C})$, where $C' = \{w \in \mathbb{T} : (e^{i\theta}w,w) \in \widetilde{C}\}$.

Now we finally define $g(w) = G(e^{i\theta}w, w)$ for $w \in \overline{\mathbb{D}}$ and $\widetilde{\Delta} = g^{-1}(U) \cap \Delta$. Then (1) and (4) are obvious and (2) holds because

$$d(g,h) \leq \sup_{z,w \in \overline{\mathbb{D}}} d(G(z,w),h(w)) < \delta + \varepsilon.$$

For proving (3) we take $w \in C'$ and observe that $(e^{i\theta}w, w) \in \widetilde{C}$ and therefore $(e^{i\theta}w \cdot w^k, w) \in C$. This implies that $e^{i\theta}w^{k+1} \in I_j$, $w \in \widetilde{J}_j$ for some j, and consequently $g(w) \in U$. Hence $C' \subset \widetilde{\Delta}$ and

$$\sigma(\widetilde{\Delta}) \ge \sigma(C') \ge \sigma_2(C) > (1 - \varepsilon)a\sigma(\Delta).$$

Proof of Th. 2.1: Let $x_0 \in X$. It is sufficient to prove that if $a \in (0,1)$ and $\omega(x_0, A, X) < -a$, then $\Omega(x_0, \overline{A}, X) \le -a$. This inequality will in turn follow if we prove that for every $\varepsilon \in (0,1)$ there exists $h \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ such that $h(0) = x_0$ and $\sigma_h(\overline{A}) > (1 - \varepsilon)a$.

We take $\varepsilon_m \searrow 0$ such that $\prod_m (1 - \varepsilon_m) \ge \sqrt{1 - \varepsilon}$. For every m we find a covering $\{B_j^m\}$ of X by countably many balls of diameter less than ε_m and set

$$U_m = \bigcup_j \{ x \in B_j^m ; \, \omega^*(x, A \cap B_j^m, B_j^m) < -1 + \varepsilon_m \}.$$

Since A is locally pluriregular, U_m is a neighbourhood of A and the inequality

$$\omega(\cdot, A \cap B_j^m, B_j^m) \ge \omega(\cdot, U_{m+1} \cap B_j^m, B_j^m)$$

implies

$$U_m \subseteq \bigcup_j \{x \in B_j^m ; \omega(x, U_{m+1} \cap B_j^m, B_j^m) < -1 + \varepsilon_m \}.$$

Since A is locally pluriregular and U_1 is an open neighbourhood of A we have

$$-a \ge \omega^*(x_0, A, X) \ge \omega(x_0, U_1, X) = \Omega(x_0, U_1, X)$$

and there exists $h_1 \in \mathcal{O}(\overline{\mathbb{D}}, X)$ such that $h_1(0) = x_0$ and $\sigma_{h_1}(U_1) > a$. We set $\Delta_1 = h_1^{-1}(U_1) \cap \mathbb{T}$ and observe that by the definition of the measure σ_{h_1} we have $\sigma(\Delta_1) > a$. We apply Th. 2.2 and get inductively a sequence h_m in $\mathcal{O}(\overline{\mathbb{D}}, X)$ and a decreasing sequence Δ_m of open subsets of \mathbb{T} such that $h_m(0) = x_0$, $h_m(\Delta_m) \subset U_m$, $\sigma(\Delta_{m+1}) > (1 - \varepsilon_m)^2 \sigma(\Delta_m)$, and $d(h_{m+1}, h_m) < 2\varepsilon_m$.

The last condition implies that h_m converges uniformly on $\overline{\mathbb{D}}$ to some $h \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$. We set $\Delta = \cap_m \Delta_m$. Since $h_m(\Delta_m) \subset U_m$ and the points of U_m are at a distance less than or equal to ε_m from A, we have $h(\Delta) \subset \overline{A}$ and since $\sigma(\Delta_{m+1}) > (1 - \varepsilon_m)^2 \sigma(\Delta_m)$ we get

$$\sigma_h(\overline{A}) \ge \sigma(\Delta) > \prod_m (1 - \varepsilon_m)^2 \sigma(\Delta_1) > (1 - \varepsilon)a.$$

3 Characterization of pluripolar sets

Let X be a complex manifold. We say that X is a Josefson manifold if any locally pluripolar subset of X is globally pluripolar. Note that any domain in a Josefson manifold is a Josefson manifold. In particular, any domain in \mathbb{C}^n is a Josefson manifold. As a direct consequence of Th. 2.1 we get (cf. Cor. 7.2 in [13])

Theorem 3.1 Let X be a relatively compact domain in a Josefson manifold and A be a Borel subset of X. Then $\Omega(\cdot, A, X) \leq \omega^*(\cdot, A, X)$.

Before we prove the theorem we prove the following auxiliary result.

Lemma 3.2 [See Th. 8.3 in [4] or Th. 7.3 in [13])]. Let μ be a Borel probability measure which is zero on every pluripolar set. Then the set function $c = c_{\mu}$ defined by

$$c(A) = c_{\mu}(A) = -\int_{X} \omega^{*}(\cdot, A, X) d\mu, \qquad A \subset X,$$

is a Choquet capacity, i.e.,

- (1) $c(A_1) \leq c(A_2)$ if $A_1 \subset A_2$;
- (2) $c(K) = \lim_{j \to \infty} c(K_j)$, where $K_1 \supset K_2 \supset \cdots \supset K$ are compact sets and $K = \cap_j K_j$;
- (3) $c(A) = \lim_{i \to \infty} c(A_i)$, where $A_1 \subset A_2 \subset \cdots \subset A$ are arbitrary sets and $A = \bigcup A_i$.

Proof: Since $\omega^*(\cdot, A, X) = \omega(\cdot, A, X)$ on $X \setminus P$ for some pluripolar set P, we have

$$c(A) = c_{\mu}(A) = -\int_{X} \omega(\cdot, A, X) d\mu, \qquad A \subset X.$$

Since $-\omega(\cdot, A_1, X) \leq -\omega(\cdot, A_2, X)$ if $A_1 \subset A_2$, (1) holds. For proving (2) we first observe that (1) implies $c(K) \leq \lim_{j \to \infty} c(K_j)$. If $\{V_k\}_{k \in \mathbb{N}}$ is a decreasing basis of neighbourhoods of K, then $\omega(x, V_k, X)$ increases to $\omega(x, K, X)$ and the monotone convergence theorem implies that $\lim_{k \to \infty} c(V_k) = c(K)$. For every $k \geq 1$ there exists a j_k such that $K_j \subset V_k$ for all $j \geq j_k$, so

$$\lim_{j \to \infty} c(K_j) \le \lim_{k \to \infty} c(V_k) = c(K).$$

Note that (3) is clear for open sets. Fix $\varepsilon > 0$ and put $V_j = \{x \in X : \omega^*(x, A_j, X) < -1 + \varepsilon\}$ and $V = \bigcup_j V_j$. Then $c(V_j) \to c(V)$. Note that $V_j \supset A_j \setminus P_j$, where P_j is a pluripolar set. Hence, $V \supset A \setminus P$, where $P = \bigcup_j P_j$. The set P is pluripolar, so $c(V) \ge c(A)$.

We have $\omega(\cdot, V_j, X) \ge \omega^*(\cdot, A_j, X)/(1-\varepsilon)$ and, therefore, $c(V_j) \le c(A_j)/(1-\varepsilon)$. Hence,

$$c(A) \le \frac{1}{1-\varepsilon} \lim_{j \to \infty} c(A_j).$$

Since $\varepsilon > 0$ is arbitrary, this proves (3).

Proof of Th. 3.1: If A is pluripolar, then there exists $u \in \mathrm{PSH}^-(X)$ such that $u \not\equiv -\infty$ and $A \subset \{u = -\infty\}$. This implies that $\omega^*(\cdot, A, X) = 0$ and the inequality holds.

From now on we assume that A is non-pluripolar. Let us first take A compact. It is sufficient to show that A can be written as $A = A_1 \cup E_1$, where A_1 is locally pluriregular and $E_1 \subset \{u = -\infty\}$ for some $u \in \mathrm{PSH}^-(X)$, $u \not\equiv -\infty$. Indeed, then Theorem 2.1 gives

$$\omega^*(\cdot, A, X) = \omega^*(\cdot, A_1, X) \ge \Omega(\cdot, \overline{A}_1, X) \ge \Omega(\cdot, \overline{A}, X).$$

In order to prove that $A = A_1 \cup E_1$, we choose a countable dense subset $\{a_k\}$ of A and set

$$E_1 = \bigcup_k \bigcup_m \{ x \in A \cap B_d(a_k, 1/m) ; \ \omega^*(x, A \cap B_d(a_k, 1/m), B_d(a_k, 1/m)) > -1 \}$$

where $B_d(a,r)$ denotes the ball with centre a and radius r with respect to a complete hermitian metric d defining the topology of X. Note that E_1 is locally pluripolar and therefore by assumption globally pluripolar. Moreover, since X is relatively compact in a Josefson manifold, we can find $u \in \mathrm{PSH}^-(X)$ so that $E_1 \subset \{u = -\infty\}$. Now we put $A_1 = A \setminus E_1$. Then A_1 is locally pluriregular, for if $x \in A_1$ and U is a neighbourhood of x, then there exists a ball $B_d(x, 1/m) \subset U$ and $a_k \in B_d(x, 1/2m)$ such that $B_d(a_k, 1/2m) \subset B_d(x, 1/m)$ and we get

$$-1 \le \omega^*(x, A \cap U, U) \le \omega^*(x, A \cap B_d(a_k, 1/2m), B_d(a_k, 1/2m)) = -1.$$

Now we let A be any Borel subset of X. We fix $x_0 \in X$ and are going to show that

$$\Omega(x_0, A, X) \le \omega^*(x_0, A, X).$$

It suffices to show that there exists a sequence of compact sets $K_1 \subset K_2 \subset \cdots \subset A$ so that

$$\omega^*(x_0, K_i, X) \to \omega^*(x_0, A, X), \qquad j \to \infty.$$

Let us construct a probability measure on X which is zero on every pluripolar set. Fix a covering $\{U_j\}$ of X so that (U_j, ψ_j) is a holomorphic chart and $\psi_j(U_j) \subset \mathbb{C}^m$ is a bounded domain. (We assume that X is m-dimensional.) For any Borel set A we put

$$\mu(A) = \sum_{j=1}^{\infty} \frac{1}{2^j} \cdot \frac{\lambda_m(\psi_j(A \cap U_j))}{\lambda_m(\psi_j(U_j))},$$

where λ_m is the Lebesgue measure in \mathbb{C}^m . It is easy to see that μ is a probability measure on X. Moreover, for any pluripolar set P we have $\mu(P) = 0$.

By Lemma 3.2 c_{μ} is a Choquet capacity. The Choquet capacitability theorem states that

$$c_{\mu}(A) = \sup\{c_{\mu}(K); K \subset A \text{ is compact}\}\$$

for all Borel subsets A of X. Hence, for a fixed Borel set A there exists a sequence $K_1 \subset K_2 \subset \cdots \subset A$ of compact sets such that $c_{\mu}(K_j) \to c_{\mu}(A)$. It is easy to see that $\omega^*(\cdot, K_j, X) \to \omega^*(\cdot, A, X)$.

The equivalence of (1) and (3) in the following theorem is well known and it indeed holds on every manifold. See Edigarian [5] and Rosay [15]. Using the theorem above we are able to refine this result.

Theorem 3.3 Let X be a Josefson manifold. Then the following conditions are equivalent

- (1) Any bounded plurisubharmonic function on X is constant.
- (2) For every $p \in X$, every nonpluripolar Borel subset A of X, and every $\varepsilon > 0$ there exists $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ such that f(0) = p and

$$\sigma_f(A) = \sigma(f^{-1}(A) \cap \mathbb{T}) > 1 - \varepsilon.$$

(3) For every $p \in X$, every nonempty open subset U of X, and every $\varepsilon > 0$ there exists $f \in \emptyset(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ such that f(0) = p and

$$\sigma_f(U) > 1 - \varepsilon$$
.

Proof: The proof that (2) implies (3) is trivial. In order to prove that (3) implies (1), we let u be a negative plurisubharmonic function on X. Assume that u is non-constant. Then there exist $x_1, x_2 \in X$ such that $u(x_1) < u(x_2)$. Take an $a \in (u(x_1), u(x_2))$. Put $U = \{x \in X : u(x) < a\}$. Then U is an open set and $x_1 \in U$. By (3) we have $\Omega(\cdot, U, X) \equiv -1$. Since $-1 \le \omega(\cdot, U, X) \le \Omega(\cdot, U, X)$, we have $\omega(\cdot, U, X) \equiv -1$. But $\frac{1}{|a|}u(\cdot) \le \omega(\cdot, U, X)$, which implies $u(x_2) \le a$, a contradiction.

In order to prove that (1) implies (2), we take a sequence of subdomains $X_1 \subseteq X_2 \subseteq \cdots \subseteq X$ such that $\bigcup_{n=1}^{\infty} X_n = X$. There exists a compact set $K \subset A$ such that K is nonpluripolar. Without loss of generality we may assume that $K \subset X_1$. For any $n \geq 1$ we have by Theorem 3.1 that

$$\Omega(\cdot, K, X_n) \le u_n = \omega^*(\cdot, K, X_n)$$
 on X_n .

There exists $x_1 \in K$ such that $u_1(x_1) = -1$. Note that the sequence $\{u_n\}$ is decreasing. Put $u = \lim u_n \in \mathrm{PSH}(X)$. Hence u is a constant, $u(x_1) = -1$, so $u \equiv -1$.

Fix a $p \in X$ and $\varepsilon > 0$. Then $\Omega(p, K, X_n) \to -1$ as $n \to \infty$, so there exists $n \in \mathbb{N}$ such that

$$\Omega(p, F, X_n) \le \Omega(p, K, X_n) < -1 + \varepsilon.$$

Observe that if $u \in \mathrm{PSH}(X)$ is such that $E \subseteq \{x \in X : u(x) = -\infty\} = \tilde{E}, x \in X \setminus \tilde{E}$, and $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ with f(0) = x, then

$$-\infty < u(x) \le \int_{\mathbb{T}} u \circ f \, d\sigma,$$

and we conclude that $\sigma_f(\tilde{E}) = \sigma(\{t \in \mathbb{T}; u(f(t)) = -\infty\}) = 0$. Hence we have

Theorem 3.4 Let X be a complex manifold and let E be a globally pluripolar subset. Then there exists a globally pluripolar $\tilde{E} \supseteq E$ such that for every $x \in X \setminus \tilde{E}$ and every $f \in \mathcal{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ with f(0) = x we have $\sigma_f(\tilde{E}) = 0$.

As a direct consequence of Theorems 3.3 and 3.4 we get a characterization of pluripolar sets by analytic discs.

Corollary 3.5 Let X be a Josefson manifold and assume that every bounded plurisubharmonic function on X is constant. Let E be a Borel subset in X. Then E is pluripolar if and only if

$$\{x \in X \; ; \; \exists f \in \emptyset(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X), f(0) = x, \sigma_f(E) > 0\} \neq X.$$

Observe that even in \mathbb{C}^n this corollary gives a new characterization of pluripolar sets.

4 Analytic discs with images in boundaries of domains

Let X be a complex manifold and let $D \subset X$ be a domain. If $u \in PSH(D)$, then we extend u to an upper semicontinuous function u^* on the closure \overline{D} by the formula

$$u^*(x) = \limsup_{D \ni y \to x} u(y), \qquad x \in \partial D.$$

For every bounded function $f: \partial D \to \mathbb{R}$ the function

$$u_{f,D} = \sup\{v \in \mathrm{PSH}(D); v^*|_{\partial D} \le f\}$$

is called the *Perron-Bremermann envelope* of f on D. We say that D is weakly regular if for every relatively open subset U of ∂D we have

$$u_{-\chi_U,D}^* \le -\chi_U$$
 on ∂D ,

where χ_U is the characteristic function of U. We put $\omega(\cdot, U, D) = u_{-\chi_U, D}$. Note that $\omega(\cdot, U, D)$ is a maximal plurisubharmonic function on D and $\omega^*(\cdot, U, D) \leq -\chi_U$ on ∂D , if D is weakly regular.

We say that D is locally weakly regular if for any $x \in \partial D$ there exists a neighbourhood basis $\{V_j\}_{j=1}^{\infty}$ of x in X such that $D \cap V_j$ is weakly regular for all j.

Note that every locally weakly regular domain is weakly regular. Indeed, for any $D_1 \subset D_2$ and any open subset U_i of ∂D_i such that $U_1 \subset U_2$ we have

$$u_{-\chi_{U_1},D_1} \ge u_{-\chi_{U_2},D_2}.$$

Proposition 4.1 (cf. [2]). Any bounded domain D in \mathbb{C}^n which is regular for the Dirichet problem (as a domain in \mathbb{R}^{2n}) for the Laplace operator is locally weakly regular. In particular, any hyperconvex domain is locally weakly regular.

Proof: The intersection of two Dirichlet regular domains is Dirichlet regular, so it is enough to show that D is weakly regular. For a Dirichlet regular domain it is well-known that for any $f \in C(\partial D)$ we have $u_{f,D}^* \leq f$ on ∂D . Since $-\chi_U$ is upper semicontinuous on ∂D it is sufficient to show that $u_{f,D}^* \leq f$ on ∂D for any upper semicontinuous function f. Let f_j be a sequence of continuous functions decreasing to f. Then $u_{f,D}^* \leq u_{f_j,D}^* \leq f_j$ on ∂D . We let $j \to \infty$ and get $u_{f,D}^* \leq f$.

For any subset $A \subset \partial D$ we put

$$\omega(x, A, D) = \sup \{ \omega(x, U, D) ; U \text{ is open and } A \subset U \subset \partial D \}, \qquad x \in D,$$

and

$$\Omega(x, A, D) = -\sup\{\sigma_f(A) \, ; \, f \in \emptyset(\mathbb{D}, D) \cap C(\overline{\mathbb{D}}, \overline{D}), f(0) = x\}, \qquad x \in D.$$

We have a natural inequality between $\omega(\cdot, A, D)$ and $\Omega(\cdot, A, D)$ as in the case when A is in the interior of D.

Lemma 4.2 Let X be a complex manifold, $D \subset X$ be a weakly regular domain and $A \subset \partial D$ be a Borel set. Then $\omega(\cdot, A, D) \leq \Omega(\cdot, A, D)$.

Proof: Let $x \in D$. If U is an open set in ∂D such that $A \subset U \subset \partial D$, and $f \in \emptyset(\mathbb{D}, D) \cap C(\overline{\mathbb{D}}, \overline{D})$ such that f(0) = x, then for $u = \omega(\cdot, U, D)$ we have

$$u(x) \le \int_{\mathbb{T}} u \circ f \, d\sigma \le \int_{f^{-1}(A) \cap \mathbb{T}} u \circ f \, d\sigma \le -\sigma_f(A).$$

If we take supremum over U in the left-hand side and infimum over f in the right-hand side, then the inequality follows.

Now we will give a new proof of an improved version of Lemma 9.1 in Poletsky [13].

Theorem 4.3 Let X be a complex manifold, D be a relative compact weakly regular domain in X, and $U \subset \partial D$ be an open set. Then $\omega(\cdot, U, D) = \Omega(\cdot, U, D)$.

The proof is in several steps each of which we state as a lemma.

Lemma 4.4 Assume that $U_1 \subset U_2 \subset \cdots \subset \partial D$ are open sets. Put $U = \cup_j U_j$. Then

$$\lim_{j \to \infty} \omega(x, U_j, D) = \omega(x, U, D), \qquad x \in D.$$

Proof: Put $u(x) = \lim_{j \to \infty} \omega(x, U_j, D)$ for $x \in D$. Note that the sequence is decreasing, so $u \in \mathrm{PSH}(D)$ and $u \geq \omega(\cdot, U, D)$. On the other hand, $u^* \leq \omega^*(\cdot, U_j, D) \leq -\chi_{U_j}$ on ∂D for all $j \geq 1$, so $u^* \leq -\chi_U$ on ∂D and $u \leq \omega(\cdot, U, D)$.

Lemma 4.5 For every $x_0 \in U$ and $\varepsilon > 0$ there exists r > 0 such that

$$B(x_0, r) \cap D \subset \{x \in D : \omega(x, U, D) < -1 + \varepsilon\}.$$

Proof: Assume that for any $n \in \mathbb{N}$ there exists $x_n \in B(x_0, \frac{1}{n}) \cap D$ such that $\omega(x_n, U, D) \ge -1 + \varepsilon$. Then $x_n \to x_0$ and $\omega^*(x_0, U, D) \ge -1 + \varepsilon$. But $\omega^*(\cdot, U, D) \le -\chi_U$ on ∂D , a contradiction.

Lemma 4.6 Assume that $V \subset D$ is an open set such that for any $x_0 \in U$ there exists an r > 0 with $B(x_0, r) \cap D \subset V$. Then $\omega(\cdot, V, D) \leq \omega(\cdot, U, D)$.

Proof: We have
$$\omega^*(\cdot, V, D) \leq -\chi_U$$
 on ∂D . Hence, $\omega(\cdot, V, D) \leq \omega(\cdot, U, D)$ on D .

Lemma 4.7 We have $\Omega(\cdot, \overline{U}, D) \leq \omega(\cdot, U, D)$.

Proof: Fix $x_0 \in D$ and $\varepsilon > 0$ and take $a > \omega(x_0, U, D)$. Let $\varepsilon_m > 0$ be a sequence such that $\varepsilon_m \searrow 0$. Assume that $\{B_j^m\}_j$ is a countable covering of U with balls of radii $< \varepsilon_m$ and centres in U for any $m \ge 1$. Put

$$U_m = \bigcup_j \{ x \in D \cap B_j^m ; \, \omega(x, U \cap B_j^m, D \cap B_j^m) < -1 + \varepsilon_m \}.$$

Let us show that

$$U_m \subset \cup_j \{x \in D \cap B_j^m : \omega(x, U_{m+1} \cap B_j^m, D \cap B_j^m) < -1 + \varepsilon_m \}.$$

For this, it suffices to show that

$$\omega(\cdot, U_{m+1} \cap B_j^m, D \cap B_j^m) \le \omega(\cdot, U \cap B_j^m, D \cap B_j^m)$$
 on $D \cap B_j^m$,

for any $m \ge 1$. It follows from Lemmas 4.5 and 4.6.

We have $\omega(x_0, U, D) \geq \omega(x_0, U_1, D)$. So, there exists an $f_1 \in \mathcal{O}(\overline{\mathbb{D}}, D)$ such that $f_1(0) = x_0$ and $\sigma_{f_1}(U_1) > |a|$. Put $\Delta_1 = f_1^{-1}(U_1) \cap \mathbb{T}$. Now we construct inductively (f_m, Δ_m) as in the proof of Theorem 2.1 and get the analytic disc f as its limit.

Proof of Theorem 4.3: By Lemma 4.2, we have $\omega(\cdot, U, D) \leq \Omega(\cdot, U, D)$. Take open sets $U_1 \in U_2 \in \cdots \in U$ such that $U = \bigcup_{j=1}^{\infty} U_j$. According to Lemma 4.7 we have $\Omega(\cdot, U, D) \leq \omega(\cdot, U_m, D)$ on D for any $m \geq 1$. Take $m \to \infty$ and get $\Omega(\cdot, U, D) \leq \omega(\cdot, U, D)$ on D.

We let X be a complex manifold, D be a relatively compact weakly regular domain in X, and $A \subset \partial D$ be a Borel set. Note that, in general, it is not true that $\{z \in A : \omega^*(z, K, D) > -1\}$ is pluripolar.

Example 4.8 Let $K \subset \mathbb{T}$ be a non-polar compact set of measure zero (take for example a Cantor-type set on the unit circle). Then $\omega^*(\cdot, K, \mathbb{D})$ is given as a Poisson integral over the set K which is of measure zero and therefore $\omega^*(\cdot, K, \mathbb{D}) \equiv 0$.

Nevertheless we have the following result, which is stated in Sadullaev [16], Theorem 27.3.

Lemma 4.9 Let $A \subset \partial D$ be a compact set. Then $\omega^*(\cdot, A, D) \equiv 0$ if and only if there exists a $u \in \mathrm{PSH}^-(D)$, $u \not\equiv -\infty$, such that $u^*|_A \equiv -\infty$.

Proof: If $\omega^*(\cdot, A, D) = 0$, then there exists $x_0 \in D$ such that $\omega(x_0, A, D) = 0$. By the definition of $\omega(\cdot, A, D)$, there exists a sequence of open sets $U_n \supset A$ on ∂D so that $\omega(x_0, U_n, D) > -2^{-n}$. Now put $u = \sum_n \omega(\cdot, U_n, D)$. Then $u \in \text{PSH}(D)$, $u(x_0) > -1$, and $u^*|_A = -\infty$.

If, on the other hand, there exists a function $u \in PSH(D)$ such that $u^*|_A = -\infty$, then we consider the neighbourhoods $U_n = \{u^* < -n\} \cap \partial D$ of A. Then

$$\frac{u}{n} \le \omega(\cdot, U_n, D) \le \omega(\cdot, A, D).$$

So, $\omega(\cdot, A, D) = 0$ on the set $\{u \neq -\infty\}$.

We say that a set $A \subset \partial D$ is locally pluriregular at $x_0 \in \overline{A}$ with respect to D if there exists a sequence $r_i \searrow 0$ such that $\omega^*(x_0, B(x_0, r_i) \cap A, B(x_0, r_i) \cap D) = -1$ for any $j \geq 1$.

Theorem 4.10 Let $A \subset \partial D$ be a Borel set. Assume that $A = A_1 \cup E$, where A_1 is locally pluriregular with respect to D and E is such that there exists a $u \in \mathrm{PSH}^-(D)$, $u \not\equiv -\infty$, with $u^*|_E \equiv -\infty$. Then

$$\omega(x, \overline{A}, D) \le \Omega(x, \overline{A}, D) \le \omega^*(x, A, D) \quad x \in D.$$

Proof: Note that $\omega^*(\cdot, A, D) = \omega^*(\cdot, A_1, D)$ on D. Fix $x_0 \in D$ and $a > \omega^*(x_0, A, D)$. Let $\varepsilon_m > 0$ be a sequence such that $\varepsilon_m \searrow 0$. Assume that $\{B_j^m\}_j$ is a countable covering of A_1 with balls of radii $< \varepsilon_m$ and centres in A_1 for any $m \ge 1$. Put

$$U_m = \bigcup_j \{ x \in D \cap B_j^m ; \omega^*(x, A_1 \cap B_j^m, D \cap B_j^m) < -1 + \varepsilon_m \}.$$

Let us show that

$$U_m \subset \cup_j \{x \in D \cap B_j^m ; \, \omega(x, U_{m+1} \cap B_j^m, D \cap B_j^m) < -1 + \varepsilon_m \}.$$

For this, it suffices to show that

$$\omega(\cdot, U_{m+1}\cap B_j^m, D\cap B_j^m) \leq \omega^*(\cdot, A_1\cap B_j^m, D\cap B_j^m) \quad \text{ on } D\cap B_j^m,$$

for any $m \geq 1$.

We have $\omega^*(x_0, A_1, D) \geq \omega(x_0, U_1, D)$. So, there exists an $f_1 \in \mathcal{O}(\overline{\mathbb{D}}, D)$ such that $f_1(0) = x_0$ and $\sigma_{f_1}(U_1) > |a|$. Put $\Delta_1 = f_1^{-1}(U_1) \cap \mathbb{T}$. Now we construct inductively (f_m, Δ_m) as in the proof of Theorem 2.1 and get the analytic disc f as its limit.

Question 4.11 Is Theorem 4.10 true for any Borel subset A of ∂D ?

Acknowledgment. This paper was written while the first author was visiting the University of Iceland. He likes to thank the Science Institute for its warm hospitality.

References

- [1] Z. BŁOCKI, The complex Monge-Ampère operator in hyperconvex domains, Ann. Scuola Normale Sup. di Pisa, Serie IV. 23, 721–747 (1996).
- [2] Z. Blocki, The complex Monge-Ampere operator in pluripotential theory, preprint (2002)
- [3] E. BEDFORD, The operator $(dd^c)^n$ on complex spaces, Springer Lecture Notes in Math. 919, 294-323, Springer Verlag 1982.
- [4] E. Bedford and B. A. Taylor, A new capacity for plurisubharmonic functions, *Acta Math.* **149**, 1–40, (1982).
- [5] A. EDIGARIAN, Analytic discs method in complex analysis, Dissertationes Math. (Rozprawy Mat.) 402, 56 pp. (2002).
- [6] A. EDIGARIAN, A note on J. P. Rosay's paper, Ann. Polon. Math. 80, 125–132 (2003).
- [7] H. Grauert, Bemerkenswerte pseudokonvexe Mannigfaltigkeiten, Math. Z. 81, 377-391, (1963).
- [8] B. JOSEFSON, On the equivalence between locally polar and globally polar for plurisubharmonic functions on \mathbb{C}^n , Arkiv för mat. 16, 109-115 (1976).
- [9] M. Klimek, Pluripotential theory. Oxford University Press, London, 1991.
- [10] F. LÁRUSSON AND R. SIGURDSSON, Plurisubharmonic functions and analytic discs on manifolds, J. reine angew. Math. 501, 1-39, (1998).
- [11] F. LÁRUSSON AND R. SIGURDSSON, Plurisubharmonicty of envelopes of disc functionals on manifolds, *J. reine angew. Math.* **555**, 27-38, (2003).
- [12] N. LEVENBERG AND E.A. POLETSKY Pluripolar hulls, Michigan Math. J. 46, 151-162 (1999).
- [13] E.A. Poletsky, Holomorphic currents, Indiana Univ. Math. J. 42, no. 1, 85-144, (1993).
- [14] E.A. Poletsky, Disk envelopes of functions II, J. Functional Anal. 163, 111-132, (1999).
- [15] J.-P. Rosay, Poletsky theory of disks on holomorphic manifolds. *Indiana Univ. Math. J.* **52**, no. 1, 157–169, (2003).
- [16] A. SADULLAEV, Plurisubharmonic measures and capacities on complex manifolds, Uspekhi Mat. Nauk, 36, no. 4, 53-105 (1981). English translation in Russian Math. Surveys, 36, no. 4, 61-119 (1981).

[17] N. Sibony, Une classe de domaines pseudoconvexes, Duke Math. J., 55, no. 2, 299-319 (1987).

Armen Edigarian

Institute of Mathematics, Jagiellonian University, Reymonta 4, 30-059 Kraków, Poland

E-mail: Armen.Edigarian@im.uj.edu.pl

Ragnar Sigurdsson

Science Institute, University of Iceland, Dunhaga 3, IS-107 Reykjavik, Iceland

E-mail: ragnar@hi.is