ON A COMBINATORIAL IDENTITY

JINTAI DING AND NAIHUAN JING

ABSTRACT. Recently the second named author discovered a combinatorial identity in the context of vertex representations of quantum Kac-Moody algebras. We give a direct and elementary proof of this identity. Our method is to show a related identity of distributions.

1. INTRODUCTION

In [4] one of us proved the following combinatorial identity in the context of quantum Kac-Moody algebras. For any integer $m \ge 0$ we have

(1.1)
$$\sum_{\sigma \in S_{m+1}} \sum_{r=0}^{m+1} \sigma \cdot {m+1 \choose r} (w - q^m z_1) \cdots (w - q^m z_r) \\ \cdot (z_{r+1} - q^m w) \cdots (z_m - q^m w) \prod_{i < j} \frac{z_i - q^2 z_j}{z_i - z_j} = 0,$$

where the symmetric group S_{m+1} acts on the variables z_i by permuting the indices while fixing q and w. Here the q-Gaussian number $\begin{bmatrix} m+1\\r \end{bmatrix}$ is defined as follows.

$$\begin{bmatrix} m+1\\r \end{bmatrix} = \frac{[m+1]!}{[r]![m+1-r]!}, \qquad [n]! = \prod_{1 \le i \le n} [i], \qquad [i] = \frac{q^i - q^{-i}}{q - q^{-1}},$$

and [0] = 1. Observe that this identity is valid in the ring of polynomials in z_1, \dots, z_{m+1} over $\mathbb{Z}[q, q^{-1}]$.

From combinatorial viewpoint the identity (1.1) is equivalent to m+2 identities of linear relations among Hall-Littlewood polynomials [6] associated to certain tuples (not necessary partitions). The first and the last of these linear relations are actually equivalent to the well-known q-binomial identity:

$$\sum_{r=0}^{m+1} (-1)^r \begin{bmatrix} m+1\\ r \end{bmatrix} = 0.$$

This identity was proved in [4] by interpreting it as the Serre relation with the help of vertex representation of the quantum Kac-Moody algebra. The special cases of m = 1, 2, 3 were known earlier and used in the level one vertex representations of quantum affine algebras [2, 3, 4].

¹⁹⁹¹ Mathematics Subject Classification. Primary: 05E; Secondary: 17B.

 $Key\ words\ and\ phrases.$ q-analysis, q-conformal field theory, combinatorics, symmetric functions.

Jing gratefully acknowledges support from NSA grant MDA904-97-1-0062 and Mathematical Sciences Research Institute through NSF grant DMS-9701755.

Later Tarasov [7] generalized the combinatorial identity in the context of the elliptic quantum algebra and he proved a more general identity using a different method. In [1] Enriquez proposed a distribution "generalization" of the identity using the quantum shuffle algebra and he verified the distribution identity for m = 1, 2. These cover all the quantum shuttle algebra associated the quantum affine algebras except $U_q(G_2^{(2)})$, which corresponds to the distribution identity of m = 3.

The purpose of this paper is to give an elementary (combinatorial) proof of the identity (1.1) for arbitrary m. The key of our proof is that the distribution identity is equivalent to the combinatorial identity (1.1).

To state our result we let the formal delta-function $\delta(z, w)$ be the following infinite series:

$$\delta(z,w) = \sum_{n \in \mathbb{Z}} z^{i-1} w^i$$

We also make the convention that the rational function $\frac{1}{z-w}$ represents the formal series in the direction |z| > |w|: $\sum_{n=0}^{\infty} z^{-1} (\frac{w}{z})^n$. Then we have

(1.2)
$$\delta(z, w) = \frac{1}{z - w} + \frac{1}{w - z}$$

Using (1.1), we will prove the following theorem.

Theorem 1.1. For any non-negative integer number m we have:

(1.3)
$$\sum_{\sigma \in S_{m+1}} \sigma \cdot \sum_{k=0}^{m+1} {m+1 \brack k} \frac{1}{(q^{-m}z_1 - w) \cdots (q^{-m}z_k - w)} \\ \cdot \frac{1}{(q^{-m}w - z_{k+1}) \cdots (q^{-m}w - z_{m+1})} \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} \\ = q^{m-1} \sum_{\sigma \in S_{m+1}} \sigma \cdot \delta(w, q^{-m}z_1) \delta(z_1, q^2 z_2) \cdots \delta(z_m, q^2 z_{m+1})$$

where the symmetric group S_{m+1} acts on the indices of the z_i .

The above identity is of great importance in representation theory, because it provides an analytical method to describe the Serre relations for quantum affine algebras [1]. As an example the distribution identity for m = 3 completes Eneriquez's argument for the exceptional case of $U_q(G_2^{(1)})$.

We will see that each step in our proof of theorem 1.1 is reversible, which means that we also obtain a new and elementary proof of (1.1).

2. Delta-function identity

The delta function plays an important role in the theory of vertex operator algebras and is characterized by the following important property. For any formal distribution $f(z) \in \mathbb{C}[[z, z^{-1}]]$ one has that

$$f(z)\delta(z,w) = f(w)\delta(z,w)$$

To simplify the presentation we will use $\operatorname{Sym}_{S_{m+1}}$ to the denote the symmetrization operator $\sum_{\sigma \in S_{m+1}} \sigma$ on the ring of formal series in z_1, z_2, \dots, z_{m+1} . Let ω_i $(i \geq 2)$ be the transposition (1*i*), and S_m be the symmetric group on letters $2, 3, \dots, m+1$. Then we have the coset decomposition of S_{m+1} : $S_{m+1} = \bigcup_{i=1}^{m+1} \omega_i S_m$, where $\omega_1 = 1$.

On the other hand, we need to do some preparation to interpret the formal power series in a slightly different way. Recall that the function $\frac{1}{z-w}$ represents a formal power series in the region |z| > |w|, which we say for simplicity that $\frac{1}{z-w}$ lives in the region |z| > |w|. We now proceed the argument in several steps.

Let us now fix the following condition on q:

$$(2.1)$$
 $|q| > 1$

Let 0 < i < j. We know that the function $\frac{1}{q^2 z_i - z_j}$ lives in the region $|q^2 z_i| > |z_j|$, which is equivalent to the region $|z_i| > |z_j||q^{-2}|$. This region contains the subset of $|z_i| > |z_j|$. Let us denote the region $|z_i| > |z_j||q^{-2}|$ by R_{ij} . Note that the function $\frac{1}{q^2 z_j - z_i}$ lives in the region $|q^2 z_j| > |z_i|$, which is equivalent

Note that the function $\frac{1}{q^2 z_j - z_i}$ lives in the region $|q^2 z_j| > |z_i|$, which is equivalent to the region $|z_j| > |z_i| |q^{-2}|$. We denote this region by R_{ji} . Clearly the subset $|z_j| > |z_i|$ is contained in R_{ji} .

On the other hand $R_{ij} \bigcap R_{ji}$ is not empty:

(2.2)
$$R_{ij} \bigcap R_{ji} = \{z_i, z_j, |q^2| |z_j| > |z_i| > |z_j| |q^{-2}| \}.$$

This says that there is a common region $R_{ij} \bigcap R_{ji}$, in which both $\frac{1}{q^2 z_j - z_i}$ and $\frac{1}{q^2 z_i - z_j}$ live.

On the other hand, the function $\frac{1}{q^{-m}z_i-w}$ lives in the region $|q^{-m}z_i| > |w|$, which is equivalent to the region $|z_i| > |w||q^m|$. We denote this region by R_{iw} . The function $\frac{1}{q^{-m}w-z_i}$ lives in the region $|q^{-m}w| > |z_i|$, which is equivalent to the region $|w| > |z_i||q^m|$. We denote this region by R_{wi} .

Clearly we have a situation different from what we had above:

(2.3)
$$R_{wi} \bigcap R_{iw} = \emptyset$$

Step 1

Let's first change the expansion direction for the series $\frac{1}{(q^{-m}z_1-w)\cdots(q^{-m}z_k-w)}$ by using

(2.4)
$$\frac{1}{q^{-m}z_i - w} = -\frac{1}{w - q^{-m}z_i} + \delta(w, q^{-m}z_i).$$

Then we have

$$\begin{split} \text{LHS} &= \text{LHS of Eqn (1.3)} = \\ &\sum_{\sigma \in S_{m+1}} \sigma \cdot \sum_{k=0}^{m+1} {m+1 \brack k} (-1)^k \frac{1}{(w-q^{-m}z_1)\cdots(w-q^{-m}z_k)} \\ &\cdot \frac{1}{(q^{-m}w-z_{k+1})\cdots(q^{-m}w-z_{m+1})} \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} + \\ & \underset{S_{m+1}}{\text{Sym}} \sum_{k=0}^{m+1} {m+1 \brack k} \sum_{l=1}^k \frac{\delta(w,q^{-m}z_l)}{(q^{-m}z_1 - w)\cdots(w-q^{-m}z_{l-1})} \cdot \\ &\cdot \frac{(-1)^{k-1}}{(w-q^{-m}z_{l+1})(q^{-m}z_k - w)(q^{-m}w - z_{k+1})\cdots(q^{-m}w - z_{m+1})} \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} \\ &= X_1 + \sum_{n=1}^{m+1} (1,n) \underset{S_m}{\text{Sym}} \sum_{k=0}^{m+1} {m+1 \brack k} \sum_{l=1}^k \frac{\delta(w,q^{-m}z_l)}{(q^{-m}z_1 - w)\cdots(w-q^{-m}z_{l-1})} \cdots \\ &= X_1 + \sum_{n=1}^{m+1} \sum_{k=0}^{m+1} {m+1 \brack k} \delta(q^{-m}z_n,w) \sum_{l=1}^k (1,n) \underset{S_m}{\text{Sym}} (-1)^{k-1} (1,l) \cdot \\ &\frac{1}{(q^{-m}z_1 - w)\cdots(w-q^{-m}z_{l-1})} \cdots \end{split}$$

where we have decomposed the action of S_{m+1} into the summation of the cosets of S_m . Here S_m is the symmetric group of the letters $2, \ldots, m$; and by X_1 we mean the first summation where there is no delta functions.

From the preparation of expansion region, we see that the term X_1 lives in the region:

(2.5)
$$Y_1 = R_{w1} \bigcap R_{w2} \bigcap ... R_{w(m+1)} \bigcap \{z_1, ... z_{m+1} | \bigcap_{0 < i < j < m+1} (R_{i,j} \bigcap R_{ji}) \}.$$

It follows from (1.1) that on Y_1 we have

(2.6)
$$X_1 = 0.$$

Then we have

LHS =

$$\sum_{n=1}^{m+1} \sum_{k=0}^{m+1} {m+1 \brack k} \delta(q^{-m} z_n, w) \sum_{l=1}^{k} (1, n) \operatorname{Sym}_{S_m} (-1)^{k-1} (1, l) \cdot \frac{1}{(q^{-m} z_1 - w) \cdots (w - q^{-m} z_{l-1})} \cdots$$

Now let us take a look at the first term of the summation. We have

4

$$\begin{split} \sum_{k=0}^{m+1} \begin{bmatrix} m+1\\ k \end{bmatrix} \delta(q^{-m}z_1, w) & \sum_{l=1}^{k} \operatorname{Sym} \frac{(-1)^{k-1}}{(w-q^{-m}z_2)\cdots(w-q^{-m}z_k)} \cdot \\ & \cdot \frac{1}{(q^{-m}w-z_{k+1})\cdots(q^{-m}w-z_{m+1})} (1, l) \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} \\ &= \delta(q^{-m}z_1, w) \sum_{k=0}^{m+1} \begin{bmatrix} m+1\\ k \end{bmatrix} \sum_{l=1}^{k} \operatorname{Sym} \frac{(-1)^{k-1}q^{m(k-1)}}{(z_1 - z_2)\cdots(z_1 - z_k)} \cdot \\ & \cdot \frac{1}{(q^{-2m}z_1 - z_{k+1})\cdots(q^{-2m}z_1 - z_{m+1})} (1, l) \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} \\ &= \delta(q^{-m}z_1, w) \sum_{k=0}^{m+1} \begin{bmatrix} m+1\\ k \end{bmatrix} \sum_{l=1}^{k} \operatorname{Sym} \frac{(-1)^{k-1}q^{m(k-1)}}{(z_1 - z_2)\cdots(z_1 - z_k)} \cdot \\ & \cdot \frac{1}{(q^{-2m}z_1 - z_{k+1})\cdots(q^{-2m}z_1 - z_{m+1})} (1, 2, \dots, l) \prod_{i < j} \frac{z_i - z_j}{q^2 z_i - z_j} \end{split}$$

since (1, 2, ..., l)(1, 2) is an element in $S_m = S_m(z_2, ..., z_{m+1})$. Then the above is simplified to the following expression:

$$\delta(q^{-m}z_1,w)\sum_{k=0}^{m+1} {m+1 \brack k} \sum_{l=1}^k \operatorname{Sym}_{S_m} \frac{(-1)^{k-1}q^{m(k-1)}(z_1-z_{k+1})\cdots(z_1-z_{m+1})}{(q^{-2m}z_1-z_{k+1})\cdots(q^{-2m}z_1-z_{m+1})} \cdot \frac{1}{(q^2z_2-z_1)\cdots(q^2z_l-z_1)(q^2z_1-z_{l+1})\cdots(q^2z_1-z_{m+1})} \prod_{2\leq i< j\leq m+1} \frac{z_i-z_j}{q^2z_i-z_j},$$

which we will denote by $\delta(q^{-m}z_1, w)X_{1,w}$.

It is clear as an analytic function

(2.7)
$$X_{1,w} = \frac{1}{2\pi\sqrt{-1}} \int_{W} \operatorname{around}_{q^{-m}z_1} - LHS \ d_w.$$

Therefore based upon (1.1) we have obtained the following result.

Proposition 2.1. As an analytic function,

(2.8)
$$X_{1,w} = 0$$

Step 2 Now let's change the expansion direction of z_1 in $\frac{1}{q^2 z_i - z_1}$ for i = 2, ..., l. Following the same argument as in **Step 1**, and we see that the above expression

becomes:

$$\begin{split} \delta(q^{-m}z_1,w) \sum_{k=0}^{m+1} \begin{bmatrix} m+1\\ k \end{bmatrix} \sum_{l=1}^k \operatorname{Sym} \frac{(-1)^{k-1}q^{m(k-1)}(z_1-z_{k+1})\cdots(z_1-z_{m+1})}{(q^{-2m}z_1-z_{k+1})\cdots(q^{-2m}z_1-z_{m+1})} \cdot \\ \cdot \sum_{l_1=1}^l \delta(z_1,q^2z_{l_1}) \frac{(-1)^{l-1}}{(z_1-q^2z_2)\cdots(z_1-q^2z_l)(q^2z_1-z_{l+1})\cdots(q^2z_1-z_{m+1})} \\ \prod_{2\leq i< j\leq m+1} \frac{z_i-z_j}{q^2z_i-z_j} \\ = \delta(q^{-m}z_1,w) \sum_{k=0}^{m+1} \begin{bmatrix} m+1\\ k \end{bmatrix} \sum_{l=1}^k \operatorname{Sym} \frac{(-1)^{k-1}q^{m(k-1)}(z_1-z_{k+1})\cdots(z_1-z_{m+1})}{(q^{-2m}z_1-z_{k+1})\cdots(q^{-2m}z_1-z_{m+1})} \\ \cdot \sum_{l_1=1}^l \delta(z_1,q^2z_{l_2}) \frac{(-1)^{l-1}}{(z_1-q^2z_2)\cdots(z_1-q^2z_l)(q^2z_1-z_{l+1})\cdots(q^2z_1-z_{m+1})} (2,l_1) \\ \prod_{2\leq i< j\leq m+1} \frac{z_i-z_j}{q^2z_i-z_j} \end{split}$$

Step 3 After this, we can use the same argument to change the expansion direction of z_2 , and then z_3 and so on. The final term will be $q^{m-1}\delta(z_m, q^2 z_{m+1})$, which gives us a proof of (1.3).

This gives us the proof under the condition that q > 1. Using the argument of analytic continuation, it is clear that (1.3) also holds for the case that q < 1. This gives us the proof for our Theorem.

If we observe carefully the proof above, we can see that each step step above is reversible. This means that we can start with the right hand side of the delta identity and follow the proof steps backward to the beginning. After finishing Step 1 in the end, replace the delta function by 0, which actually gives us a new proof of identity (1.1). This is a direct and elementary proof of identity (1.1).

Remark. By replacing the delta function by 0, we also obtain an elementary proof of the combinatorial identity (6.1) of [4].

References

- B. Enriquez, On correlation functions of Drinfeld currents and shuffle algebras, math.QA/9809036.
- [2] I. B. Frenkel and N. Jing, Vertex representations of quantum affine algebras, Proc. Nat'l. Acad. Sci. USA 85 (1988), 9373–9377.
- [3] N. Jing, Twisted vertex representations of quantum affine algebras, Invent. Math. 102 (1990), 663–690.
- [4] N. Jing, Quantum Kac-Moody algebras and vertex representations, Lett. Math. Phys. 44 (1998), 261–271.
- [5] N. Jing, Level one representations of $U_q(G_2^{(1)})$, Proc. Amer. Math. Soc. 127 (1999), 21–27.
- [6] I.G. Macdonald, Symmetric functions and Hall polynomials, 2nd edition, Clarendon Press, Oxford, 1995.
- [7] V. Tarasov, Combinatorial identities related to representations of $U_q(\hat{\mathfrak{gl}}_2)$, math.QA/9811050.

 $\mathbf{6}$

Ding: Department of Mathematical Sciences, University of Cincinnati, Cincinnati, OH 45221

 $E\text{-}mail \ address: \ \texttt{dingQmath.uc.edu}$

Jing: Mathematical Sciences Research Institute, 1000 Centennial Drive, Berkeley, CA 94720-5070

Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205

 $E\text{-}mail\ address: \ \texttt{jing@math.ncsu.edu}$