

# Three-centre cluster structure in $^{11}\text{C}$ and $^{11}\text{B}$

N Soić†‡, M Freer‡, L Donadille‡, N M Clarke‡, P J Leask‡, W N Catford§, K L Jones§, D Mahboub§, B R Fulton||, B J Greenhalgh||, D L Watson|| and D C Weisser¶

†Ruđer Bošković Institute, Bijenička 54, HR-10000 Zagreb, Croatia

‡School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

§School of Electronics and Physical Sciences, University of Surrey, Guildford GU2 7XH, United Kingdom

|| Department of Physics, University of York, Heslington, York, YO10 5DD, United Kingdom


¶Department of Nuclear Physics, The Australian National University, Canberra ACT 0200, Australia

## Abstract.

Studies of the  $^{16}\text{O}(^9\text{Be},\alpha^7\text{Be})^{14}\text{C}$ ,  $^7\text{Li}(^9\text{Be},\alpha^7\text{Li})^5\text{He}$  and  $^7\text{Li}(^9\text{Be},\alpha t)^5\text{He}$  reactions at  $E_{beam}=70$  and 55 MeV have been performed using resonant particle spectroscopy techniques. The  $^{11}\text{C}$  excited states decaying into  $\alpha+^7\text{Be}(\text{gs})$  are observed between 8.5 and 13.5 MeV. The  $\alpha+^7\text{Li}(\text{gs})$ ,  $\alpha+^7\text{Li}^*(4.652\text{ MeV})$  and  $t+^8\text{Be}(\text{gs})$  decays of  $^{11}\text{B}$  excited states between 9 and 19 MeV are observed. The decay processes are used to indicate the possible three-centre  $2\alpha+^3\text{He}$  ( $2\alpha+^3\text{H}$ ) cluster structure of observed states. This cluster structure is more prominent in the positive-parity states, where two rotational bands with large deformations are suggested. Excitations of some of the observed  $T=1/2$  resonances coincide with the energies of previously measured  $T=3/2$  isobaric analogs of the  $^{11}\text{Be}$  states, indicating that these states may have mixed isospin.

Many light nuclei possess well developed cluster structure made of  $\alpha$ -particles as main building block. The best known examples are  $2\alpha$  and  $3\alpha$  cluster structures of  $^8\text{Be}$  and  $^{12}\text{C}$ . In recent years attention has been focused on neutron-rich Be and C nuclei, where evidence for developed cluster structures was found (for example [1, 2, 3, 4] and references therein). It is interesting to investigate influence of  $\alpha$ -clustering on structure of the neutron deficient  $^{11}\text{C}$  nucleus and its mirror nucleus  $^{11}\text{B}$  for which the experimental evidence for cluster structures is scarce. Here we present results of the experimental studies which probe cluster structure of  $^{11}\text{C}$  and  $^{11}\text{B}$  via the  $\alpha$ -decay of their excited states and for  $^{11}\text{B}$  also via  $2\alpha+t$  decay.


The measurements were performed at the Australian National University's 14UD tandem accelerator facility. 70 MeV and 55 MeV  $^9\text{Be}$  beam were incident on a thin  $\text{Li}_2\text{O}_3$  foil. Reaction products were detected in an array of four charged-particle telescopes which provided charge and mass resolution up to Be. Telescopes 1 and 2 were located with their centres at  $\sim 17.5^\circ$  from the beam axis and telescopes 3 and 4 at  $\sim 29^\circ$ . The measurement of the energies and angles of detected particles permitted the kinematics



**Figure 1.**  $^{11}\text{B}$  excitation energy spectrum from the  $^7\text{Li}(^9\text{Be}, \alpha ^7\text{Li})^5\text{He}$  reaction (a) and  $^{11}\text{C}$  excitation energy spectrum from the  $^{16}\text{O}(^9\text{Be}, \alpha ^7\text{Be})^{14}\text{C}$  reaction (b) at  $E_{\text{beam}}=70$  MeV. Spectra are shifted so that the  $7/2^+, 5/2^+$  doublets are aligned. Lines mark positions of the states with the same value of the spin and parity populated in both nuclei. The positions of the  $K=5/2^+$  band members are marked with full circles and triangles,  $K=3/2^+$  band members with full squares and inverted triangles, possible  $9/2^+$  states in  $K=3/2^+$  bands with open circles. The excitations of the isobaric analogues of the lowest three  $^{11}\text{Be}$  states are marked with stars.

of the many-body reactions to be fully reconstructed. Further experimental details can be found in [4, 5].

The  $\alpha$ -decay of excited states in  $^{11}\text{B}$  has been studied using the  $^7\text{Li}(^9\text{Be}, ^{11}\text{B}^* \rightarrow \alpha + ^7\text{Li})^5\text{He}$  reaction ( $Q=-2.4$  MeV). The  $^{11}\text{B}$  excitation energy spectra show resonances at 9.2, 10.3, 10.55, 11.2, (11.4), 11.8, 12.5, (13.0), 13.1, (14.0), 14.35, (17.4) and (18.6) MeV. A study of the  $\alpha$ -decay of  $^{11}\text{C}$  excited states has been performed using the  $^{16}\text{O}(^9\text{Be}, ^{11}\text{C}^* \rightarrow \alpha + ^7\text{Be})^{14}\text{C}$  reaction ( $Q=-14.602$  MeV). The  $^{11}\text{C}$  excitation energy spectra provide evidence for peaks at 8.65, 9.85, 10.7 and 12.1 MeV and indications for peaks at 12.6 and 13.4 MeV. Detailed description and discussion of the results are



**Figure 2.**  $^{11}\text{B}$  excitation energy spectra from the  ${}^7\text{Li}({}^9\text{Be}, \alpha t){}^5\text{He}$  reaction at  $E_{\text{beam}}=70$  and 55 MeV. Numbers at the top denote detectors combination in which events were detected: (ijk) i,j are for  $\alpha$ -particles and k for triton.

presented in [5]. Analysis of these data shows that main reaction mechanism for the population of the  $\alpha$ -decaying states in both nuclei was two-nucleon transfer onto the cluster nucleus  ${}^9\text{Be}$ . Figure 1 presents comparison of the  $^{11}\text{B}$  and  $^{11}\text{C}$  excitation energy spectra from these measurements. We observe the same series of excited states at the lower excitations in both nuclei: unresolved doublet of  $7/2^+$ ,  $5/2^+$  states (marked 'a'), a  $3/2^-$ ,  $5/2^-$ ,  $7/2^+$  triplet ('b') which is  $\sim 1.2$  MeV above the doublet,  $9/2^+$  state ('c') at an excitation of 2 MeV higher than the doublet, then the state which is believed to be the isobaric analogue state of the  $^{11}\text{Be}$  ground state ('d') which is  $\sim 3.4$  MeV above the doublet, and then weak states, which are probably  $7/2^+$  ('e') and  $11/2^+$  ('f'), and which are 3.9 and 4.8 MeV above the doublet's excitation. An interesting feature of these results is the observation of the  $\alpha$ -decay, which means isospin  $T=1/2$ , of the excited states proposed to be the isobaric analogue states of the  $^{11}\text{Be}$  states, which have  $T=3/2$ , in both nuclei. If the observed states are indeed those identified with  $T=3/2$  character, then our results show that the lowest three  $T=3/2$  levels in  $^{11}\text{B}$  and the first  $T=3/2$  level in  $^{11}\text{C}$  probably have large isospin mixing. Alternatively, the peaks may have a genuine  $T=1/2$  character and may be linked to rotational bands. The present results show that six observed states in  $^{11}\text{B}$  and six in  $^{11}\text{C}$  can be grouped in two rotational bands

[5]. Rotational bands with  $K=5/2^+$  have been proposed beginning at 7.286 and 6.905 MeV in  $^{11}\text{B}$  and  $^{11}\text{C}$  respectively, with rotational members at 9.185 and 8.655 ( $7/2^+$ ), 11.265 and 10.679 ( $9/2^+$ ) and 14.04 and 13.4 MeV ( $11/2^+$ ). The moment of inertia  $I$  of these bands is very large, with a rotational parameter  $\hbar^2/2I$  of 0.25 MeV which would correspond to very deformed structure. Other possible positive-parity bands in both  $^{11}\text{B}$  and  $^{11}\text{C}$  have been proposed with  $K=3/2^+$ , beginning at excitation energies 7.97784 and 7.4997 MeV, respectively. Rotational members are at 9.274 and 8.699 MeV ( $5/2^+$ ) and 10.597 and 10.083 MeV ( $7/2^+$ ) and the  $9/2^+$  members of these bands would be at 12.6 MeV in  $^{11}\text{B}$  and 12.1 MeV in  $^{11}\text{C}$ , very close to the excitations of the proposed  $1/2^+$   $T=3/2$  states and exactly at excitations where resonances are in the present spectra. The rotational parameter  $\hbar^2/2I$  of these two bands would be 0.215 MeV.

Triple coincidence data provide clear evidence for the  $^7\text{Li}(^9\text{Be},\alpha\alpha t)^5\text{He}$  reaction ( $Q=-4.9$  MeV) at both  $E_{beam}=70$  and 55 MeV [6]. Reconstructed relative energy spectra for all possible combinations of 2 and 3 particles in the exit channel provide evidence for decays of  $^{11}\text{B}$  excited states into  $2\alpha+t$ , the  $^8\text{Be}$  ground state and the first excited state into  $2\alpha$  and for  $\alpha+t$  decay of  $^7\text{Li}$  states at 4.652 and 6.6 MeV. The  $^{11}\text{B}$  excitation energy spectra reconstructed from the energies and momenta of all three detected decay products presented in Figure 2 show peaks at 13.1, 14.4 and 17.5 MeV. All three states are also observed in the  $\alpha+^7\text{Li}$  data. An analysis gives evidence for the  $t+^8\text{Be}(gs)$  decay of the  $^{11}\text{B}$  states at 13.1 and 14.4 MeV and decays via  $^8\text{Be}(3.03$  MeV) and  $^7\text{Li}(4.652$  MeV) states of the 14.4 and 17.5 MeV states [6]. These results are the first direct evidence for decay of  $^{11}\text{B}$  excited states into these three-body channels.

Given the nature of the reaction processes, two-nucleon transfer onto the  $2\alpha+n$  nucleus  $^9\text{Be}$ , the observed strong  $\alpha$ -decay of these mirror states at excitations where various decay channels are open and known  $\alpha+t(^3\text{He})$  structure of  $^7\text{Li}(^7\text{Be})$ , as well as observed decays of  $^{11}\text{B}$  states into  $2\alpha+t$ , it is possible that these states possess the three-centre  $2\alpha+t(^3\text{He})$  cluster structure. These two nuclei may be an example of the three-centre structure in which holes, rather than nucleons, are being exchanged between  $\alpha$ -cores. The present measurements did not provide information on the spin and parity of the observed states, which are unknown for some of the states, and also on partial widths, which are crucial for the understanding of their structure. Additional measurements capable of determining this information will be performed in the near future.

## References

- [1] Soić N et al 1996 *Europhys. Lett.* **34** 7 ; Milin M et al 1999 *Europhys. Lett.* **48** 616 ; Milin M et al 2004 *Nucl. Phys. A* **730** 285 ; Curtis N et al 2004 *Phys. Rev. C* **70** 014305
- [2] Freer M et al 1999 *Phys. Rev. Lett.* **82** 1383 ; Freer M et al 2001 *Phys. Rev. C* **63** 034301
- [3] Milin M and von Oertzen W 2002 *Eur. Phys. J. A* **14** 295
- [4] Soić N et al 2003 *Nucl. Phys. A* **728** 12 ; Soić N et al 2003 *Phys. Rev. C* **68** 014321
- [5] Soić N et al 2004 *Nucl. Phys. A* **742** 271
- [6] Soić N et al, to be published