## Equation of state extracted from the root-mean-square-radius-isospin

Zbigniew Sosin\*

Jagellonian University, M. Smoluchowski Institute of Physics, Reymonta 4, PL-30059 Kraków, Poland

## Abstract

Experimental correlation between the rms nucleus charge radius and the neutron-proton asymmetry is discussed. Simple attempt of explanation of this correlation using a semi-empirical liquid drop model which takes into account nuclear compressibility, nuclear deformation and neutron skin effects is presented. It seems that such procedure could be used as a tool for determination of the nuclear equation of state.

PACS numbers: 21.10.Ft 21.60. -n 21.65.+f

<sup>\*</sup>Electronic address: ufsosin@cyf-kr.edu.pl

The nuclear equation of state (EOS) and the nuclear reaction kinetics determine stellar structure and evolution. In particular, the EOS allows to estimate the critical parameters at which the collapse process is halted and the shock wave in a core collapse supernova becomes formed [1, 2]. But the equation of state is not only relevant in astrophysical aspects. Knowledge of EOS is a way to better understand the nuclear heavy-ion reactions [3, 4] and some fundamental properties of heavy nuclei, like for instance binding energy or stability of neutron-rich nuclei [5].

The equation of state for infinite, asymmetric nuclear matter usually represents the relation between the energy density per nucleon (baryon) e and densities reached by neutrons and protons in the nuclear matter. In this relation the Coulomb energy is not taken into account. Many theoretical approaches [6] have shown that the properties of infinite asymmetric nuclear matter, at relatively low temperature, can be described by an approximate form of EOS:

$$e(\rho,\delta) = e(\rho,\delta=0) + \delta^2 e_{sym}(\rho) \tag{1}$$

where the baryon density  $\rho$  is a sum of neutron  $\rho_n$  and proton  $\rho_p$  density. Isospin asymmetry  $\delta$ , is defined as

$$\delta = \frac{\rho_n - \rho_p}{\rho_n + \rho_p} = \frac{\rho_n - \rho_p}{\rho}.$$
(2)

First term in the eq. (1) represents energy density per baryon associated to symmetric nuclear matter ( $\delta = 0$ ). This energy can be expanded around normal nuclear matter density  $\rho_0$  as

$$e(\rho, 0) = e(\rho_0, 0) + \frac{K_{\delta=0}}{18} \left(\frac{\rho - \rho_0}{\rho_0}\right)^2 + \dots$$
(3)

In the formula above  $K_{\delta=0}$  is the nuclear compressibility for symmetric matter, defined as

$$K_{\delta=0} = 9\rho_0^2 \frac{\partial^2 e(\rho, 0)}{\partial \rho^2} |_{\rho=\rho_0} .$$
(4)

Second term in eq. (1) stands for the symmetry energy and can be also expanded around  $\rho_0$  as

$$e_{sym}(\rho) = e_{sym}(\rho_0) + \frac{L}{3} \left(\frac{\rho - \rho_0}{\rho_0}\right) + \frac{K_{sym}}{18} \left(\frac{\rho - \rho_0}{\rho_0}\right)^2 + \dots$$
(5)

where variables L (slope) and  $K_{sym}$  (curvature) characterizing the density dependence of the nuclear symmetry energy are given by

$$L = 3\rho_0 \frac{\partial e_{sym}(\rho)}{\partial \rho} \mid_{\rho = \rho_0}$$
(6)

and

$$K_{sym} = 9\rho_0^2 \frac{\partial^2 e_{sym}(\rho)}{\partial \rho^2} \mid_{\rho=\rho_0}$$
(7)

The coefficients  $\rho_0$ ,  $e(\rho_0, 0)$ ,  $e_{sym}(\rho_0)$ ,  $K_{\delta=0}$ , L,  $K_{sym}$  allow us to determine the EOS around the normal nuclear density. Unfortunately, values of the slope L and the curvature  $K_{sym}$  are neither well determined experimentally nor accurately estimated theoretically [6]. Data from the giant monopole resonances suggest that  $K_{sym}$  varies from -566 ±1350 MeV to 34 ±159 MeV [7]. So far, experimental predictions for the L value are rather scarce. In the recent paper [8] the isospin transport calculations compared with the MSU diffusion data suggest  $L \approx 60$ . In theoretical models L varies from -50 up to 200 MeV [9]. Moreover, theoretical estimates for  $K_{sym}$  are model-dependent and vary from about -700 to +466 MeV [10].

In the present paper we will show a way to put some constraints for possible values of these parameters. In order to do it, we compare the experimentally measured correlation between root mean square radii of nuclei and isospin asymmetry, to the similar correlation resulting from a modified liquid-drop model (LDM).

One of the most elementary properties of a nucleus is the spatial nucleon density distribution. Unfortunately, this distribution cannot be obtained experimentally in a straightforward way. In the case of protons, knowledge of the Coulomb interaction allows to measure with a high precision the momenta of the charge distribution. From experimental data one usually fixes the root-mean-square (rms) radius  $\langle r_p^2 \rangle^{1/2}$ . In this paper we will use a part of the set of 799 ground state nuclear charge radii presented in [11]. This experimental data were obtained from the elastic electron scattering, muonic atom X-rays,  $K_{\alpha}$  isotope shift, and optical isotope shift. Errors associated with the data are usually much lower than 1%. For neutrons, the differences between neutron and proton density distributions at large nuclear radii in stable nuclei may be determined e.g. from antyprotonic atoms [16]

In the first column of Fig. 1 we plot a correlation between the experimentaly measured

 $\langle r_p^2 \rangle^{1/2} / \sqrt[3]{A}$  and isospin asymmetry I = (N - Z)/(N + Z), where N and Z denote neutron and proton numbers, respectively. In Fig. 1a we plot these correlations for all available nuclei with masses A = N + Z > 100. We can notice here some branches which are roughly mutually parallel. These branches are much better visible on the next drawings in this column (Fig. 1c, e, g) showing the correlations for nuclei with selected mass intervals.

Let us try to explain the nature of the observed correlations. We consider here the influence of three effects: nuclear deformation, dependence of nucleon density on isospin, and neutron skin.

At the beginning we will study the influence of nucleus deformation on the proton rms radius. In our analysis we neglect the diffuseness of the surface and we assume only groundstate quadrupole deformation described by " $\beta$  parametrization" (a spherical harmonic expansion) [12]. Such an aproach should be sufficient for rather small ground state deformations. Now we can write the proton mean-square radius in the form

$$\left\langle r_{p}^{2} \right\rangle = \frac{3}{5} R_{0}^{2} \left( 1 + \frac{5}{8\pi} \beta^{2} \left( A, I \right) \right)$$
 (8)

where  $R_0$  is the average radius. Parameters  $\beta(A, I)$  for considered set of nuclei are taken from model calculations used in [12]. Obtained correlations between deformation  $\beta^2$  and isospin I are presented in the right column in Fig. 1. As we can see from the upper graph (Fig. 1b) for almost all masses  $\beta^2$  is less than 0.1. Formula (8) suggests that for a given  $R_0$ , nuclear deformation does not modify the  $\langle r_p^2 \rangle^{1/2}$  value more than 1%, however (see Figs 1d,f,h) it can enhance or suppress the observed correlations depending on the selected mass range. Even more importantly, it can also change the correlation sign (see e.g. Fig 1e and Fig 1f)).

One can suspect, that such a strong correlation observed in the first column could be associated with the form of the nuclear equation of state. To examine this possibility we assume that the considered correlation can be a result of differences in the density of nuclear matter in the central region of nuclei due to different values of the surface tension and Coulomb interaction.

Saturation of the nuclear matter density in the central region of nuclei provides a base for description of the nuclear binding energy in the liquid-drop model. This model was formulated by Weizsäcker (1935), Bethe (1936), and Bohr (1936). Up to now various modifications were introduced in this model and some of them consider also compressibility of nuclear matter [13].

The liquid-drop model in its simplest version describes the nuclear binding energy B as a sum of the volume, surface and Coulomb terms.

$$B = B_v + B_s + B_C,\tag{9}$$

where  $B_v$  is the dominant volume ingredient usually written as,

$$B_v = a_v (1 - k_v I^2) A, (10)$$

and  $a_v$  and  $k_v$  are coefficients. The term  $k_v I^2$  takes into account the lowering of the binding energy caused by different number of protons and neutrons in the nucleus.

Next term is (9) takes into account lower binding energy for nucleons located on the nuclear surface, and can be expressed as

$$B_s = a_s (1 - k_s I^2) A^{2/3} f = \left[ a_s (1 - k_s I^2) A^{-1/3} \right] A f,$$
(11)

where  $a_s$  and  $k_s$  are coefficients. Factor  $f = (1 + \frac{2}{5}\beta^2 - \frac{4}{105}\beta^3)$  takes into account the nuclear deformation (see [14]). We can notice that the surface correction per one nucleon (square bracket in (11)) vanishes with the increasing A.

Coulomb interaction between protons appears in the next correction, which is proportional to  $Z^2$  and inversely proportional to the radius of the nucleus. This term can be written as

$$B_c = a_c \frac{Z^2}{A^{1/3}} g = \left[ a_c \frac{Z^2}{A^{4/3}} \right] g A \tag{12}$$

where  $a_c$  is a parameter, and  $g = (1 - \frac{1}{5}\beta^2 - \frac{4}{105}\beta^3)$  similarly like f in (11) describes possible influence of the nuclear deformation. Comparison of (9) to the experimental data (e.g. [15] allows to determine coefficients presented in formulas (10)- (12) as:

$$a_v = -15.677 \text{ MeV}, a_s = 18.560 \text{ MeV}, a_c = 0.717 \text{ MeV}$$
  
 $k_v = k_s = 1.79 \text{ MeV}.$  (13)

It should be stressed, that LDM is dedicated to a finite drop of nuclear matter with a constant density. From the analysis of experimental data we can deduce that apart from nuclear surface the nucleus density is roughly constant and for mass A > 40 it has a Fermi-like shape distribution. The central region density itself somewhat varies for different nuclei.

The form of the volume part of the binding energy (10) is in fact related to the EOS at the ground state density. For  $a_v = e(\rho_0, 0)$ ,  $a_v k_v = e_{sym}(\rho_0)$ ,  $\rho = \rho_0$ , and  $\delta^2 = I^2$  the energy obtained from EOS (1) is equivalent to the volume energy per nucleon in the LDM (10). For our consideration we reformulate the liquid-drop model in a way that allows to determine the central region density for a given nucleus. We want to obtain the central region density as a function of coefficients  $\rho_0$ ,  $e(\rho_0, 0)$ ,  $e_{sym}(\rho_0)$ ,  $K_{\delta=0}$ , L,  $K_{sym}$ , and  $\delta = I$ , determining the nuclear EOS. In principle neutron and proton density distributions may have different radii,  $R_n$ ,  $R_p$ , what gives the so called skin effect. Lets assume for the beginning  $R_n = R_p$ .

Modifying the LDM we assume that density in the central region is a function of global values N and Z (i.e. there is no local dependence). According to the idea of the LDM, we can say that the central region density  $\rho$  results mainly from nuclear interaction present in infinite nuclear matter, further being modified by the surface tension and Coulomb repulsion. Therefore, reformulating the liquid-drop model, we write the binding energy per nucleon  $b(\rho, I, A)$  in a form similar to (9):

$$b(\rho, I, A) = b_v(\rho, I) + b_s(\rho, I, A) + b_c(\rho, I, A).$$
(14)

In the above formula the volume part of the binding energy is defined by EOS (1).

$$b_v(\rho, I) = e(\rho, \delta). \tag{15}$$

Let  $b_{v0} = e(\rho_0, \delta)$ . Now, the correction for the surface effect we can rewrite in a new, density dependent form

$$b_{s}(\rho, I, A) = a'_{s}(1 - k'_{s}I^{2}) \frac{b_{v}(\rho, I)}{b_{v0}} \left(\frac{\rho_{0}}{\rho}\right)^{2/3} A^{-1/3} f$$
$$= \frac{a'_{s}}{a'_{v}} b_{v}(\rho, I) \left(\frac{\rho_{0}}{\rho}\right)^{2/3} A^{-1/3} f.$$
(16)

Factor  $\left(\frac{\rho_0}{\rho}\right)^{2/3}$  in (16) is responsible for the surface change due to the change in the core

density. The corresponding coefficients are marked by prime to distinguish from parameters of the standard LDM. In principle their values can slightly differ from the original ones.

The part describing the Coulomb interaction is modified only by factor  $\left(\frac{\rho_0}{\rho}\right)^{1/3}$  according to the radius change induced by core density changes (Z = Z(I, A)).

$$b_c(\rho, I, A) = a'_c \frac{Z^2}{A^{4/3}} \left(\frac{\rho_0}{\rho}\right)^{1/3} g.$$
 (17)

Here the binding energy is determined by the density  $\rho$  and a new set of coefficients  $a'_{v}$ ,  $a'_{s}, a'_{c}, \rho_{0}, e(\rho_{0}, 0), e_{sym}(\rho_{0}), K_{\delta=0}, L, K_{sym}$ . The density  $\rho$  for a given nucleus is found from a condition  $\partial b(\rho, I, A)/\partial \rho |_{I,A=const} = 0$ . In this way  $\rho$  and binding energy values are uniquely determined by a selected set of coefficients. Here, as a first approximation, we are taking the value of coefficients with the sign prime equal to corresponding coefficients without prime which are given by (13). Coefficients  $\rho_{0}, e(\rho_{0}, 0), e_{sym}(\rho_{0})$ , are rather well known. In Fig. 2 we present the binding energy  $b(\rho, I, A)$ , as a function of density, for nuclei with A = 150and three different Z values equal to 50, 60 and 75 (I = 1/3, I = 1/5 and I = 0). For these test calculations we take  $K_{\delta=0} = 240$  MeV and we check energy density behavior for all possible combinations of  $L = \pm 100$  MeV and  $K_{sym} = \pm 500$  MeV what determine the symmetry energy (5).

The saturation of nuclear density in the central region is characteristic for heavier nuclei. For light nuclei we can say that almost all nucleons are located on their surface. In order to omit the problem of high contribution coming from the surface we shall test formula (14) only for heavier nuclei, A>100.

Assuming  $R_0 \sim \sqrt[3]{A/\rho}$  and using (8) we can rewrite relation between  $r_{rms}$  and  $\rho$  in the form

$$\frac{\left\langle r_p^2 \right\rangle^{1/2}}{\sqrt[3]{A}} = \alpha \frac{\sqrt{\left(1 + \frac{5}{8\pi}\beta^2\right)}}{\sqrt[3]{\rho}} \tag{18}$$

where  $\alpha$  is a normalization constant. Formula (18) neglects a neutron skin, which because of different radii of neutron and proton density distributions, surrounds the larger A nuclei. Lets define thickness of the neutron skin as

$$d = R_n - R_p. \tag{19}$$

where  $R_p = \sqrt{\frac{5}{3}} \langle r_p^2 \rangle^{1/2}$ , and  $R_n = \sqrt{\frac{5}{3}} \langle r_n^2 \rangle^{1/2}$  is the effective proton and neutron radius, respectively. The central region nucleon density is now

$$\rho = \rho_n + \rho_p = \frac{N}{\frac{4}{3}\pi R_n^3} + \frac{Z}{\frac{4}{3}\pi R_p^3}$$
(20)

and instead of I=(N-Z)/(N+Z) one gets for the central region isospin  $\widetilde{I}$ 

$$\widetilde{I} = \frac{N - Z(1 + d/R_p)^3}{N + Z(1 + d/R_p)^3} \simeq I\left(1 - \frac{3Zd}{r_0 \left(\frac{\rho_0}{\rho}\right)^{1/3} A^{4/3}}\right)$$
(21)

For numerical calculations we use  $r_0 = 1.2049$  (see [14]).

As d seems to be much smaller than  $R_p$ , one gets from (19) and (20)

$$R_{p} = \sqrt[3]{\frac{3A}{4\pi\rho}} \left( 1 - \frac{Nd}{r_{0} \left(\frac{\rho_{0}}{\rho}\right)^{1/3} A^{4/3}} \right)$$
(22)

The Coulomb energy is inversely proportional to the radius and consequently  $b_c$  in (17) has to be written as:

$$b_c(\rho, I, A) = a'_c \frac{Z^2}{A^{4/3}} \left(\frac{\rho_0}{\rho}\right)^{1/3} \left(1 + \frac{Nd}{r_0 \left(\frac{\rho_0}{\rho}\right)^{1/3} A^{4/3}}\right) g.$$
 (23)

As an alternative for eq. (18) one should use now:

$$\frac{\left\langle r_p^2 \right\rangle^{1/2}}{\sqrt[3]{A}} = \alpha \frac{\sqrt{\left(1 + \frac{5}{8\pi}\beta^2\right)}}{\sqrt[3]{\rho}} \left(1 - \frac{Nd}{r_0 \left(\frac{\rho_0}{\rho}\right)^{1/3} A^{4/3}}\right)$$
(24)

Lets try to discuss the origin and slope of the observed correlation's as a function of different factors.

In Figs. 3 and 4 in columns marked "model 1", "model 2", and "model 3", we plot correlation's given by formula (24), for different sets of the EOS parameters. For calculations named "model 1" we neglect the neutron skin and deformations of nuclei ( $\beta = d = 0$  in formula (24)) whereas in "model 2" deformations are taken into account ( $\beta \neq 0, d = 0$ ). In "model 3" we are trying to estimate the influence of the neutron skin ( $\beta \neq 0, d \neq 0$ ). Calculations were performed for the soft ( $K_{\delta=0} = 120$  MeV, red points) and for the hard ( $K_{\delta=0} = 500$  MeV, blue points) EOS. In Fig. 3 L = -150 MeV and in Fig. 4 L = 60 MeV. As for nuclei  $\rho \approx \rho_0$  and in formula (5) the third term is a rather small correction, in both cases we take  $K_{sym} = 500$  MeV.

For "model 1" the  $\langle r_p^2 \rangle^{1/2} / \sqrt[3]{A}$  versus *I* correlation already contains a number of separated chains because of discrete values of the *A* and *Z* numbers. Nuclear deformations modify

slightly slopes of the observed chains and may result in additional splitting (see "model 2", 180 < A < 210). As seen from the "model 3" calculations taking into account the neutron skin introduces quite important changes.

## Conclusions

An experimental correlation between the nucleus rms charge radius and the isospin of heavier nuclei has been observed. It seems to be related to interactions characteristic for the nuclear matter equation of state modified by confinement inside of an atomic nucleus. These interactions are responsible for nuclear deformations and creation of the neutron skin. We propose explanation of the observed correlation on the basis of a modified nuclear liquid drop model. Inspection of studied correlation's (Figs 3 and 4) suggests a possibility of evaluating the EOS parameters. A precise determination of these values requires a more precise search and a more complex analysis which will be subject of a future work. Such investigation probably requires an extension of the LDM by additional terms taking into account such corrections as for example pairing, shell effects and diffuseness of the nucleus surface. Although  $b_s$  in formula (16) decreases for large A values, it could be interesting to introduce here an additional correction related to the local isospin of the surface.

## Acknowledgments

The author is indebted to P. Pawłowski for solving some computing problems and to J. Błocki, K. Grotowski, M. Misiaszek, and P. Pawłowski for careful reading of manuscript and helpful discussions. This work was supported by the Polish-French collaboration IN2P3, grant no. CSI8.

Figure 1: Correlation between the neutron-proton asymmetry and (first column) the experimentally measured nuclear rms radius, or (second column) the calculated parameter of deformation.

Figure 2: Binding energy per nucleon vs density  $\rho$  for different equations of state (see text).

Figure 3: Comparison between the measured and calculated nuclear rms radius vs I, for L = -150MeV and  $K_{sym} = 500$ MeV (see text for details).

Figure 4: Comparison between the measured and calculated nuclear rms radius vs I, for L = 60 MeVand  $K_{sym} = 500 \text{MeV}$  (see text for details).

- [1] G. McLaughlin, Nuclear Physics News, Vol. 15, No 3, 2005, Nuclear Physics News.
- [2] J. M. Lattimer and M. Prakash, Astrophys. J. 550, 426 (2001)
- [3] B.-A. Li, Phys. Rev. Lett. 88, 192701 (2002), and references therein
- [4] P. Danielewicz, R. Lacey, and W. G. Lynch, Science 298, 1592 (2002)
- [5] B. Todd and J. Piekarewicz, Phys. Rev. C 67, 044317 (2003)
- [6] L.-W. Chen, Phys. Rev. C 68, 014605 (2003), and references therein
- [7] S. Shlomo and D.H. Youngblood, Phys. Rev. C 47, 529 (1993)
- [8] B.-A. Li and L.-W. Chen, Phys. Rev. C 72, 064611 (2005).
- [9] R.J. Furnstahl, Nucl. Phys. A706, 85 (2002)
- [10] I. Bombaci and U. Lombardo, Phys. Rev. C 44, 1892 (1991)
- [11] I. Angeli, Atomic Data Nucl. Data Tbl, 87, 185 (2004)
- [12] P. Möller, J.R. Nix W.D. Myers and W.J. Świątecki, Atomic Data Nucl. Data Tbl. 59, 185 (1995)
- [13] G.Baym, H.A. Bethe and C.J. Pethick, Nucl. Phys. A175 225(1971)
- [14] W.D. Myers and W.J. Świątecki, UGRL 11980/1965
- [15] W.D. Myers and W.J. Świątecki, Nucl. Phys. 81, 1 (1966)
- [16] A. Trzcińska et al., Phys. Rev. Lett. 87, 082501 (2001)