Astronomy & Astrophysics manuscript no. 42263
November 6, 2021
©ESO 2021

Ultra-deep 31.0-50.3 GHz spectral survey of IRC+10216 ⋆

J. R. Pardo[1], J. Cernicharo[1], B. Tercero[2], C. Cabezas[1], C. Bermúdez[1], M. Agúndez[1] , J. D. Gallego[2], F. Tercero[2], M. Gómez-Garrido[2], P. de Vicente[2], J. A. López-Pérez[2]

[1] Consejo Superior de Investigaciones Cientifícas, Instituto de Física Fundamental, Serrano 123, 28006 Madrid, Spain
e-mail: jr.pardo@csic.es
[2] Instituto Geográfico Nacional, Centro de Desarrollos Tecnológicos, Observatorio de Yebes, Apartado 148, 19080 Yebes, Spain

Received September 20, 2021; accepted October 29, 2021

ABSTRACT

Context. IRC+10216, the carbon-rich envelope of the asymptotic giant branch (AGB) star CW Leo, is one of the richest molecular sources in the sky. Available spectral surveys below 51 GHz are more than 25 years old and new work is needed.
Aims. Characterizing the rich molecular content of this source, specially for heavy species, requires to carry out very sensitive spectral surveys at low frequencies. In particular in this work we have achieved an rms in the range 0.2-0.6 mK per MHz.
Methods. long Q-band (31.0-50.3 GHz) single dish integrations were carried out with the Yebes 40m telescope using specifically built receivers. State of the art line catalogs are used for line identification.
Results. A total of 652 spectral features corresponding to 713 transitions from 81 species (we count as different the isomers, isotopologues and ortho/para species) are present in the data. Only 57 unidentified lines remain with signal to noise ratios ⩾3. Some new species and/or vibrational modes have been discovered for the first time with this survey.
Conclusions. This IRC+10216 spectral survey is, by far, the most sensitive one carried out to this date in the Q-band. It therefore provides the most complete view of IRC+10216 from 31.0 to 50.3 GHz, giving unique information on its molecular content, specially for heavy species. Rotational diagrams built from the data provide valuable information on the physical conditions and chemical content of this circumstellar envelope.

Key words. line:identification — radio lines: stars — stars: AGB and post-AGB — stars: carbon — stars: individual (IRC+10216) — surveys

arXiv:2111.02902v1 [astro-ph.GA] 3 Nov 2021

1. Introduction

The combination of proximity (123 ± 14 pc from the Sun; Groenewegen et al. 2012) and high mass-loss rate ($\dot{M} = (2\text{-}4) \times 10^{-5}$ $M_\odot$ yr^{-1}; Cernicharo et al. 2015; Guélin et al. 2018) makes IRC+10216 the main stellar target for molecule hunting in space (Cernicharo et al. 2000). In this context, the largest work at millimeter wavelengths (4.3-1.0 mm) has been carried out with the IRAM-30m telescope. Accumulated data achieved during more than three decades are now in their final phase of analysis for publication (Cernicharo et al. 2021, in prep.). In order to complement those data at the longer end of the millimeter range (6.0-9.7 mm, or 31.0 to 50.3 GHz in frequency), which is specially important for the detection of heavy molecules, we decided to undergo an extremely deep integration in the so-called Q-band towards IRC+10216 and several other stellar sources known for their molecular richness. For this work it was important to have access to a large antenna in order to keep angular resolutions similar to those provided by the IRAM-30m (but here at wavelengths 1.4 to 10 times larger), state of the art receivers with very broad instantaneous band to reduce telescope time for a given rms goal, and large spectral resolutions to allow for degradation if desired.

The importance of sensitive surveys for understanding the rich molecular complexity found in the Universe, eventually leading to the appearance of life, is unquestionable and has made a large progress in recent years due to two key factors. On one hand the rapid and large increase in the instantaneous bands of receivers, allowing to accumulate large amounts of integration times over large frequency bands because short-spaced frequency tunings are no longer necessary. On the other hand, large interferometers now in operation such as ALMA or NOEMA, allow to explore the sources with rich molecular content at narrower angular scales thus providing a new molecular view of them since some molecular species, specially for stellar sources, appear only in the innermost regions of their envelopes.

The spectral region below the atmospheric 60 GHz O_2 spin-rotation band generally lacks coverage by the most important millimeterwave single dish telescopes or interferometers. In addition, due to wavelength, large antennas are needed to keep a reasonably coupling to stellar sources. As a consequence, available spectral surveys of IRC+10216 in the Q-band are more than 25 years old (Kawaguchi et al. 1995). This is the reason that has motivated our work, in which we present a big improvement with respect to the past. The step forward is not only due to technical reasons related to the observational capabilities themselves, but also to a continuous effort on extending the spectroscopic databases with the latest laboratory and theoretical results. The observations are described in section 2. The systematic procedure for data reduction, line identification, assignment and fitting is described in section 3. We present the results in section 4, although an extended series of figures and tables, with the full results, has been left for appendix A. The procedure for line identification and handling this large amount of information is described in section 5. A summary stressing the legacy im-

⋆ Based on observations carried out with the Yebes 40m radiotelescope at Yebes Observatory, operated by the Spanish Geographic Institute (IGN, Ministerio de Transportes, Movilidad y Agenda Urbana).



Fig. 1. Noise level in the spectra presented in this work at their final resolution (see section 3) and typical atmospheric brightness temperature and average receiver temperature across the band.

portance of the results obtained in this work, in the framework of our knowledge of the physical conditions and chemistry of IRC+10216, is finally given in section 6.

2. Observations

This IRC+10216 31.0-50.3 GHz spectral survey was carried out in several runs from May 2019 to February 2020, and in April 2021, with the 40 meter antenna of Yebes Observatory (IGN, Spain), hereafter Yebes-40m, after several improvements in its equipment funded by the Nanocosmos project[1]. This large antenna provides now a main beam efficiency from 0.6 at 31 GHz to 0.43 at 50 GHz, and a beam size in the range 36-56" for those frequencies. Although the experimental setup is described in detail in a separate technical paper (Tercero et al. 2021), the most relevant information for this work is the use of a receiver consisting of two HEMT cold amplifiers covering the 31.0-50.3 GHz band with horizontal and vertical polarizations. The receiver temperature ranges from 22 K at 31 GHz to 42 K at 50 GHz. The backends are 16×2.5 GHz fast Fourier transform spectrometers (FFTS), with a primary spectral resolution of 38.1 kHz (later smoothed in the data reduction for this work), providing the whole coverage of the observed band.

The observing mode was position switching with an off position at 300" in azimuth. The reference position used for IRC+10216 has been J2000 RA 09:47:57.36 Dec +13:16:44.4. The HPBW of the antenna at 39.2 GHz is 45". Pointing corrections were obtained by observing the SiO masers of R Leo, and errors were always within 2-3". The intensity scale of the first-calibrated is antenna temperature (T_A^*) corrected for atmospheric absorption using the ATM package (Cernicharo 1985; Pardo et al. 2001). A conversion to main beam temperature (T_{MB}) is done in order to generate the rotational diagrams of Figure 4. Calibration uncertainties are estimated to be within 10 %. More details on the observational procedure can be found in Pardo et al. (2020)

[1] https://nanocosmos.iff.csic.es/

3. Data reduction and analysis

The GILDAS package[2] was used to add up the raw spectra provided by the telescope calibration software, to fit individual baselines in the different FFTS sections, and to stitch all these sections into a final spectrum using the appropriate weights according to system temperatures and integration times of each individual scan. If necessary, additional baselines have been fitted to narrow sections of the data, but these interventions have been kept to the strict minimum. The resulting spectrum was smoothed to a resolution of 0.229 MHz, for which the sensitivity ranges from ~ 0.4 to 1.2 mK in the surveyed frequency range (see Figure 1). The average data noise every 500 MHz shown in the latter figure is the basis for a first automatic procedure that checks which observed spectral features are above the 3σ level. Obviously, a second check is done by hand to spot features that were missed in the first step. Once the detected spectral features are well established, the next step has been to assign them to molecular lines. For this task we rely upon the latest version of the MADEX line catalog (Cernicharo 2012) with up to 6150 molecular species an 23.5 million transitions. We proceed in several steps. First, as part of the automatic procedures, there is a cross-check between the data and the catalog to assign as many lines as possible. Second, all the assignments are checked by hand in order to eliminate errors and to disentangle blended lines from different species, lines spread due to hyperfine structure, etc. A third step consists of assigning lines not present in the catalog to new species, isotopologues or vibrational levels. This step has resulted in the discovery for the first time in space of the ν_{19} vibrational mode of HC_9N (Pardo et al. 2020) and the Mg-containing carbon chains MgC_5N and MgC_6H (Pardo et al. 2021). This careful work has allowed to identify 93 % of all lines present in the data. In addition, features at the 2-3 σ level with a linewidth compatible with the ±14.5 kms^{-1} typical of the source have also been individually checked by hand, and this has resulted on extra assignments, mainly to ^{13}C substitutions of HC_7N.

The overall summary of the identification work described in this section is the following: 652 spectral features belonging

[2] http://www.iram.fr/IRAMFR/GILDAS



Fig. 2. Overall view of the data presented in this paper with a zoom around 45.6 GHz. Several weak spectral features (∼1-3 mK), revealed in this work, are shown in different colors in one of the panels.

to 713 identified lines and 57 unidentified lines (U-lines). An overview is given in Table 1; the details on the results of individual fits to all lines, using the SHELL method implemented in the GILDAS software, can be found in Table A.1; and the remaining U-lines are listed in Table A.2. The final spectra presented in this paper (Figures 2, and A.1 to A.39) improve by a factor of ≈10 the sensitivity of those presented in the previous work conducted with the Nobeyama 45m telescope (Kawaguchi et al. 1995) in a similar frequency range. In total they label 132 lines and report 33 U-lines. Among the latter, 22 of them could be labeled with the spectroscopic data available nowadays (they correspond to C_8H, C_6H^- C_5N^-, C_6H, HC_9N and HC_7N v_{15}), 6 are spurious features as they are not present in our data, and 5 are still U-lines for us in 2021. The strongest observed feature (at 14σ in our data) that still remains unidentified stands at 41712 MHz.

4. Results

The detected lines can be divided into different families of carriers. We present the results for each one of these separately.

4.1. Cyanopolyynes

This molecular family, $HC_{2n+1}N$, dominates the millimeter wave spectrum of this source, as it happens with other C-rich evolved stars such as CRL 618. In IRC+10216, up to 304 lines (43% of all detected lines in the Q-band) belong to this family, which comprises HCN, HC_3N, HC_5N, HC_7N, and HC_9N in their main species, isotopic substitutions, vibrationally excited states, together with a few isomers. The upper panel in Figures 3 and 4 provide a quick view of all lines from this family in this work. The difference with respect to Kawaguchi et al. (1995) is quite large since they cannot detect isotopic substitutions of HC_5N nor the vibrational states of HC_5N, HC_7N and HC_9N. Sometimes, v_{11} lines of HC_5N are present in their data but labeled as U probably due to lack of information in line catalogs available in 1995.

The alternative single and triple C-bond is, in fact, the dominant chemical feature in this source since most radicals, anions, and metal-bearing species, are carbon chains of this type.

Concerning HCN, although there are no lines from its main rotational ladder in the Q-band, three *l*-doubling transitions from the v_2=1 vibrational state are present in the data (J=12-12,13-13,14-14). The upper energy levels of these transitions lie between 1300 and 1500 K and, therefore, the signal detected arises from much closer to the photosphere of CW Leo than most lines in this work. They are narrower too. Although this is the first Q-band detection in IRC+10216 of HCN *l*-doubling lines, higher-J ones have been previously reported (Cernicharo et al. 2011).

4.2. Carbon chain Radicals

The list of radicals detected that are related with the cyanopolyynes is also long: C_3H, C_4H, C_5H, C_6H, C_7H, C_8H, C_3N ($^{13}CCCN$, $C^{13}CCN$, $CC^{13}CN$) and C_5N. They are responsible for a total of 167 lines or 23 % of all detected and labeled spectral lines in this work. Their open structure makes that several configurations are close in energy, and hyperfine structure is also common for them. C_3H presents 6 hyperfine components (two of them blended) in its ground vibrational state, from 32617 to 32668 MHz. Another set of hyperfine components from 44857 to 44939 GHz, corresponding to the v_4 vibrational mode (at 42.4 K) does not show up in the data. C_4H presents a total of 9 doublets corresponding to its ground, v_7 and $2v_7$ vibrational states. C_5H presents 4 doublets for each one of the lowest states: $^2\Pi_{3/2}$ (not resolved) and $^2\Pi_{1/2}$ (resolved). C_6H is detected as resolved doublets from its $^2\Pi_{3/2}$ and $^2\Pi_{1/2}$ states, and other lines from the vibronic states $^2\Sigma^-$, $^2\Delta_{3/2}$, and $^2\Delta_{5/2}$ belonging to the v_{11} vibrational state are also detected. The total number of lines from C_6H is 56 , the largest number in this family. Finally, C_7H and C_8H also display doublets in the electronic states $^2\Pi_{3/2}$ and $^2\Pi_{1/2}$, with only the last ones resolved in both cases, although the lowest energy state is $^2\Pi_{1/2}$ for C_7H and $^2\Pi_{3/2}$ for C_8H.

Table 1. Molecules and total number of spectral features (N) and lines (L) detected in this work for each one of them. μ is the molecule's electric dipole moment in Debyes (1 Debye = $3.33564 \cdot 10^{-30}$ Coulomb·meter).

Molecule	μ	N/L	References for spectroscopy	Molecule	μ	N/L	References for spectroscopy
HCN	2.985	3/ 3	Zelinger et al. (2003), Ebenstein and Muenter (1984)	HC_3N	3.732	8/10	Mbosei et al. (2000), Thorwirth et al. (2000)
$H^{13}CCCN$	3.732	2/ 2	Creswell et al. (1977), Mallinson & de Zafra (1978), Thorwirth et al. (2001)	$HC^{13}CCN$	3.732	2/ 2	Same as $H^{13}CCCN$.
$HCC^{13}CN$	3.732	2/ 2	Same as $H^{13}CCCN$.	HCCNC	2.957	2/ 2	Guarnieri et al. (1992), Gripp et al. (2000)
$HCCN^{13}C$	2.957	2/ 2	Same as HCCNC	HNCCC	5.665	2/ 2	Kawaguchi et al. (1992), Hirahara et al. (1993), Botschwina et al. (1992)
HC_5N	4.330	20/22	Bizzocchi et al. (2004), Alexander et al. (1976)	$H^{13}CCCCCN$	4.330	8/ 8	Same as HC_5N.
$HC^{13}CCCCN$	4.330	8/ 8	Same as HC_5N.	$HCC^{13}CCCN$	4.330	7/ 7	Same as HC_5N.
$HCCC^{13}CCN$	4.330	7/ 7	Same as HC_5N.	$HCCCC^{13}CN$	4.330	8/ 8	Same as HC_5N.
HC_7N	4.820	72/74	McCarthy et al. (2000), Degli Esposti et al. (2004), Botschwina et al. (1997)	$H^{13}CCCCCCCN$	4.820	10/11	McCarthy et al. (2000), Botschwina et al. (1997)
$HC^{13}CCCCCCN$	4.820	8/ 8	Same as $H^{13}CCCCCCCN$	$HCC^{13}CCCCCN$	4.820	10/10	Same as $H^{13}CCCCCCCN$
$HCCC^{13}CCCCN$	4.820	10/10	Same as $H^{13}CCCCCCCN$	$HCCCC^{13}CCCN$	4.820	6/ 6	Same as $H^{13}CCCCCCCN$
$HCCCCC^{13}CCN$	4.820	11/11	Same as $H^{13}CCCCCCCN$	$HCCCCCC^{13}CN$	4.820	9/ 9	Same as $H^{13}CCCCCCCN$
HC_9N	5.200	80/80	McCarthy et al. (2000), Botschwina et al. (1997), Pardo et al. (2020)	HCCN	3.600	1/ 1	Allen et al. (2001), Aoki et al. (1993)
l-HC_4N	4.330	6/ 6	Tang et al. (1999), Ikuta et al. (2000)	o-H_2C_4	4.457	6/ 6	Killian et al. (1990), Oswald and Botschwina (1995)
p-H_2C_4	4.457	4/ 5	Killian et al. (1990), Oswald and Botschwina (1995)	o-H_2C_3	4.100	2/ 2	Vrtilek et al. (1990)
p-H_2C_3	4.100	1/ 1	Vrtilek et al. (1990)	C_3H	3.550	6/ 6	Yamamoto et al. (1990a), Gotlieb et al. (1985), Kaifu et al. (2004), Woon (1995)
C_4H	2.100	18/19	Guélin et al. (1982), Gottlieb al. (1983), Chen al. (1995), Oyama et al. (2020)	C_5H	4.881	12/16	McCarthy et al. (1999), Woon (1995)
C_6H	5.540	56/56	Linnartz et al. (1999), Woon (1995)	C_7H	5.945	23/33	McCarthy et al. (1997), Woon (1995)
C_8H	6.500	31/46	McCarthy et al. (1996), McCarthy et al. (1999)	C_3N	2.850	4/ 4	McCarthy et al. (1995), Guélin et al. (1982), Gottlieb al. (1983)
$CC^{13}CN$	2.850	4/ 6	McCarthy et al. (1995), McCarthy et al. (2003)	$C^{13}CCN$	2.850	4/ 4	Same as $CC^{13}CN$
$^{13}CCCN$	2.850	5/ 6	Same as $CC^{13}CN$	C_5N	3.385	4/ 4	Kasai et al. (1997), Botschwina (1996)
C_5N^-	5.200	6/ 6	Cernicharo et al. (2008)	C_6H^-	8.200	7/ 8	McCarthy et al. (2007), Blanksby et al. (2001)
C_8H^-	10.400	13/13	Gupta et al. (2007), Blanksby et al. (2001)	c-C_3H_2 (ortho)	3.430	2/ 2	Thaddeus et al. (1985), Bogey et al. (1986), Vrtilek et al. (1987), Spezzano et al. (2012), Kanata et al. (1987)
c-C_3H_2 (para)	3.430	2/ 2	Same as c-C_3H_2 (ortho)	SiS	1.735	1/ 1	Müller et al. (2007), Tiemann et al. (1972), Sanz et al. (2003)
^{29}SiS	1.735	1/ 1	Same as $^{(28)}SiS$.	^{30}SiS	1.735	1/ 1	Same as $^{(28)}SiS$.
$Si^{33}S$	1.735	1/ 1	Same as $^{(28)}SiS$.	$Si^{34}S$	1.735	1/ 1	Same as $^{(28)}SiS$.
SiO	3.098	1/ 1	Manson et al. (1977), Mollaaghababa et al. (1991), Sanz et al. (2003), Cho and Saito (1998), Raymonda et al. (1970)	^{29}SiO	3.098	1/ 1	Same as $^{(28)}SiO$.
^{30}SiO	3.098	1/ 1	Same as $^{(28)}SiO$.	$Si^{17}O$	3.098	1/ 1	Same as $^{(28)}SiO$.
SiC_2	2.393	2/ 2	Gotlieb et al. (1989), Suenram et al. (1989)	$Si^{13}CC$	2.393	2/ 2	Cernicharo et al. (1991)
$^{29}SiC_2$	2.393	1/ 1	Suenram et al. (1989), Cernicharo et al. (1991), Kokkin et al. (2011)	$^{30}SiC_2$	2.393	2/ 2	Same as $^{29}SiC_2$
SiC_4	6.420	6/ 6	Ohishi et al. (1989), Gordon et al. (2000)	$^{29}SiC_4$	6.420	5/ 5	Same as $^{29}SiC_4$
SiC_6	8.220	7/ 7	Gordon et al. (2000)	CS	1.958	1/ 1	Gottlieb et al. (2003), Winnewisser & Cook (1968)
^{13}CS	1.958	1/ 1	Same as CS.	$C^{33}S$	1.958	1/ 1	Same as CS.
$C^{34}S$	1.958	1/ 1	Same as CS.	$C^{36}S$	1.958	1/ 1	Same as CS.
CCS	2.855	4/ 4	Saito et al. (1987), Yamamoto et al. (1990b), Lovas et al. (1992), Lee (1997)	C_3S	3.704	3/ 3	Yamamoto et al. (1987), Tang et al. (1995), Lovas et al. (1992), Ohshima & Endo (1992), Suenram & Lovas (1994)
NaCN	8.850	7/ 8	Van Vaals (1984), Halfen & Ziurys (2011)	MgCN	5.150	2/ 2	Anderson et al. (1994), Hirano et al. (2002)
MgNC	5.308	4/ 4	Kawaguchi et al. (1993), Steimle and Bousquet (2001)	MgC_3N	6.250	14/14	Cernicharo et al. (2019)
MgC_5N	7.300	20/30	Pardo et al. (2021)	MgC_4H	2.100	10/10	Cernicharo et al. (2019)
MgC_6H	2.500	23/30	Pardo et al. (2021)	CH_3CN	3.925	2/ 3	Pavone et al. (1990), Cazzoli and Puzzarini (2006), Simeckova et al. (2004), Müller et al. (2009), Anttila et al. (1993)
CH_3C_3N	4.750	5/ 5	Moises et al. (1982), Bester et al. (1983), Bester et al. (1984)	KCl	10.270	2/ 2	Clouser and Gordy (1964), Caris et al. (2004)
NaCl	9.002	1/ 1	Clouser and Gordy (1964), Caris et al. (2002), Timp et al. (2012)	$Na^{37}Cl$	9.002	1/ 1	Same as NaCl.
AlCl	1.500	1/ 1	Wyse & Gordy (1972), Lide (1965), Hensel et al. (1993)	U Lines		57	

The total number of spectral features of C_7H and C_8H detected is 23 and 31, respectively, with a counterpart of 33 and 46, respectively, rotational transitions.

Other radicals detected in this family are C_3N (two doublets at 39571/39590 and 49466/49485 MHz), the ^{13}C isotopic substitutions of it (marginal), and C_5N through two extremely weak doublets at 39280/39291 and 42086/42097 MHz. For a graphic summary of all lines from carbon chain radicals, see the second panel of Figures 3 and 4.

4.3. Anions

Up to date, IRC+10216 is the only circumstellar environnment where anions have been detected. The survey presented in this paper provided a good opportunity for detecting heavy linear anions and the results have confirmed those expectations: C_5N^-, C_6H^-, and C_8H^- are detected through 6, 8 and 13 lines, respectively. The lighter anion C_4H^- could be present but its two transitions at 37239.41 and 46549.16 MHz are very close in frequency with lines of $HCC^{13}CCCN$ and $HCCC^{13}CCN$, for the first one, and HC_9N ν_{19}, for the second one, to the point of making impossible to confirm its detection. The same happens to the C_3N^- line at 48515.87 (blended with a strong C_6H line). The other line of C_3N^- at 38812.79 MHz is not seen, a result that is compatible with its low abundance revealed from its discovery in this source at higher frequencies (Thaddeus et al. 2008). C_5N^- and C_6H^- are expected (and confirmed) to display the strongest lines among anions in this survey. A graphic summary of all lines from anions can be found on the third panel of Figures 3 and 4.

4.4. Si-bearing species

The only oxygen-bearing species detected in our line survey is SiO. The J=1-0 transition is detected for the main isotopologue as well as for ^{29}SiO and ^{30}SiO. On the other hand $Si^{17}O$ at 41794.658 MHz and $Si^{18}O$ at 40352.771 could be there but their detection is still marginal at the actual noise level.

Another Si bearing molecule with only one rotational transition in this range (J=1-0) is SiS. It has been detected for its main isotopologue as well as for ^{29}SiS, ^{30}SiS, $Si^{33}S$ and $Si^{34}S$.

The cyclic C_{2v} molecule SiC_2 has two strong enough lines for detection in the surveyed frequency range. Except $^{29}SiC_2$ (1 line detected), all other substitutions (SiC_2, $^{30}SiC_2$ and $^{13}CSiC$) exhibit two detected lines here.

The linear molecule SiC_4 has 6 rotational transitions in the surveyed frequency range (from J=11-10 to J=16-15). The multiple possible isotopic substitutions make it a candidate for a large number of lines in this source but only $^{29}SiC_4$ can be confirmed as detected here with five of its individual six rotational transitions present in the Q-band range just marginally seen.

Finally, the detection of SiC_6 is marginal, although possible, through several transitions from J=26→25 to J=38→37 at the 1-2 mK level (sometimes blended with lines from other species). If confirmed, this would be the first time that this Si-bearing carbon chain is seen in space.

4.5. S-bearing species

Apart from SiS (see section 4.4) other S-bearing molecules detected are CS, CCS and C_3S. The diatomic molecule CS is detected in 5 isotopic substitutions, the main one and ^{13}CS, $C^{33}S$, $C^{34}S$ and $C^{36}S$. The linear species C_3S has 3 transitions in the surveyed frequency range (J=6-5,7-6,8-7), all three detected, and the $^3\Sigma$ species CCS displays also four detected lines. The isotopic substitutions of CCS and C_3S are not seen.

4.6. Metal chlorides and cyanides

These interesting molecules appear in most cases with just one or two lines (MgCN, KCl, NaCl, $Na^{37}Cl$ and AlCl) given their molecular weight and the fact that they are mostly diatomic or linear. NaCN, due to its bent structure, has more rotational lines, 7 of them detected in this work. Two doublets have been detected corresponding to the isomer MgNC with the first one (35793/35809 MHz) within a blended complex of lines from other molecules.

Related to the cyanides, we should mention the detection of CH_3CN (1 line), CH_3C_3N (5 lines), and the radicals MgC_3N and MgC_5N. The last two species, together with MgC_4H and MgC_6H, have been discovered in space for the first time in IRC+10216 (Cernicharo et al. 2019; Pardo et al. 2021).

4.7. Cyclic molecules

Besides SiC_2 (section4.4), the only cyclic molecule we can report as detected here is c-C_3H_2 with just 2 lines for both the ortho and para configuration of H spins. Only one line (the one involving the lowest energy levels within the Q-band) for each species is strong and nicely detected. The other line, at 42231.25 MHz and 42139.19 GHz for the para and ortho species respectively, is almost 10 times weaker. In addition, the 42231.25 MHz is blended with another line from an unidentified species. For these reasons, a proper analysis using rotational diagrams needs other lines observed, for example, with the IRAM-30m telescope.

It is interesting to mention that IRC+10216 remains poor in cyclic molecules. A similar Q-band survey with the same telescope on TMC-1 has discovered several cyclic molecules not seen in the envelope of CW Leo, such as indene and benzyne (Cernicharo et al. 2021a,b).

4.8. Latest findings

As the line identification goes on and the number of unidentified features reduces, it becomes easier to find harmonical relations in other subsets of U-lines. The great progress made in line catalogs in recent years, based on both laboratory work and ab initio calculations is obvious. Nevertheless, spectral surveys of space objects such as this one are also part of this progress as some species are directly discovered in them thanks to these subsets of harmonically related U-lines. This is the case, for example, of the ν_{19} vibrational level of HC_9N, for which a series of 26 doublets, harmonically related with integer quantum numbers ranging from J_{up} = 54 to 80, were found among the U-lines in an early stage of analysis (Pardo et al. 2020). A fit of the observed central frequencies and line intensities allowed us to derive the rotational constants and, therefore, incorporate HC_9N ν_{19} to MADEX and the identifications to this paper.

Similarly to HC_9N ν_{19}, another harmonically related series of doublets was discovered later resulting on the first space detection of MgC_5N. This discovery follows the previous one by Cernicharo et al. (2019) of MgC_3N and and MgC_4H, which motivated the search for MgC_6H in in IRC+10126. However, given the line intensities of MgC_4H, the expected ones for MgC_6H would be quite in the limit for detection. Extra observations were carried out in 2021 in order to confirm the detection of MgC_6H, and finally both species were presented in Pardo et al. (2021).

Table 2. Results from rotational diagram fits in this work. T_{rot}: Rotational temperature, N_{col}: Column density, Z_{rot}: Partition function. The resulting fits are plotted in Figure 4, on top of the data points used for those fits (extracted from the individual line fits as listed in Table A.1). Species with unreliable fits, due to a very low number of data points or extremely high data disperison in them, are not shown. These species, and others with a very low number of detected lines here, will be revisited in future publications of similar 4.3 to 1.0 mm surveys carried out with the IRAM 30m telescope.

Molecule	# lines	T_{rot}(K)	N_{col}(cm^{-2})	Z_{rot}	Comments
HC_5N	7	10.1 ± 0.6	$(4.2 \pm 0.4) \times 10^{14}$	158.80	Reliable fit
HC_5N $v_{11} = 1$	12	6.2 ± 1.8	$(7 \pm 5) \times 10^{12}$	195.20	Large data dispersion
$H^{13}CCCCCN$	8	10.0 ± 1.7	$(9.7 \pm 2.7) \times 10^{12}$	161.52	Reliable fit
$HC^{13}CCCCN$	8	14.4 ± 2.6	$(8.2 \pm 1.7) \times 10^{12}$	228.04	Reliable fit
$HCC^{13}CCCN$	7	10.4 ± 1.3	$(1.00 \pm 0.19) \times 10^{13}$	162.42	Reliable fit
$HCCC^{13}CCN$	7	10.6 ± 1.7	$(1.00 \pm 0.25) \times 10^{13}$	166.20	Reliable fit
$HCCCC^{13}CN$	8	9.2 ± 0.6	$(1.09 \pm 0.13) \times 10^{13}$	145.42	Reliable fit
HC_7N	17	15.8 ± 0.4	$(1.95 \pm 0.11) \times 10^{14}$	583.44	Reliable fit
HC_7N $v_{15} = 1$	31	21.3 ± 1.7	$(2.0 \pm 0.3) \times 10^{13}$	1572.97	Reliable fit (3 points removed)
HC_7N $v_{15} = 2$	26	12.6 ± 2.5	$(1.0 \pm 0.5) \times 10^{13}$	1358.51	Large data dispersion
$H^{13}CCCCCCCN$	11	23 ± 4	$(3.0 \pm 0.8) \times 10^{12}$	879.75	Reliable fit (1 point removed)
$HC^{13}CCCCCCN$	8	15 ± 7	$(3.31 \pm 3.41) \times 10^{12}$	559.13	Large data dispersion
$HCC^{13}CCCCCN$	10	19 ± 6	$(3.5 \pm 2.1) \times 10^{12}$	713.79	Large data dispersion
$HCCC^{13}CCCCN$	10	23 ± 4	$(2.7 \pm 0.8) \times 10^{12}$	836.22	Reliable fit (2 points removed)
$HCCCC^{13}CCCN$	6	16 ± 3	$(3.9 \pm 1.7) \times 10^{12}$	573.14	Reliable fit
$HCCCCC^{13}CCN$	11	22 ± 6	$(2.7 \pm 1.3) \times 10^{12}$	824.82	Large data dispersion
$HCCCCCC^{13}CN$	9	31 ± 10	$(3.2 \pm 1.1) \times 10^{12}$	1156.63	Reliable fit (1 point removed)
HC_9N	30	22.9 ± 1.1	$(4.5 \pm 0.6) \times 10^{13}$	1644.03	Reliable fit (1 point removed)
HC_9N $v_{19} = 1$	50	28.0 ± 2.2	$(2.4 \pm 0.4) \times 10^{13}$	4009.46	Reliable fit (1 point removed)
l-HC_4N	6	5.1 ± 2.5	$(3.9 \pm 3.7) \times 10^{12}$	126.85	Large data dispersion
C_4H	4	4.9 ± 0.6	$(1.84 \pm 0.25) \times 10^{14}$	86.42	Reliable fit
C_4H $v_7 = 1$	10	10 ± 5	$(6.6 \pm 2.0) \times 10^{13}$	314.65	Large data dispersion
C_4H $v_7 = 2$	4	6.4 ± 1.3	$(2.2 \pm 0.4) \times 10^{13}$	113.77	Reliable fit
C_5H $^2\Pi_{1/2}$	8	5.0 ± 0.4	$(2.7 \pm 0.5) \times 10^{13}$	172.74	Reliable fit
C_5H $^2\Pi_{3/2}$	8	5.9 ± 0.8	$(2.52 \pm 2.48) \times 10^{14}$	10.67	Moderate data dispersion
C_6H $^2\Pi_{3/2}$	14	11.0 ± 0.8	$(1.51 \pm 0.14) \times 10^{13}$	334.91	Reliable fit
C_6H $^2\Pi_{1/2}$	14	12.4 ± 0.6	$(1.23 \pm 0.18) \times 10^{13}$	66.83	Reliable fit
C_6H $v_{11} = 1$	14	10.4 ± 0.5	$(9.3 \pm 0.7) \times 10^{12}$	310.61	Reliable fit (2 points removed)
C_6H $^2\Delta_{5/2}$ $v_{11} = 1$	6	13 ± 3	$(2.0 \pm 0.5) \times 10^{12}$	201.51	Moderate data dispersion
C_6H $^2\Delta_{3/2}$ $v_{11} = 1$	7	21 ± 13	$(1.4 \pm 0.7) \times 10^{12}$	311.75	Large data dispersion
C_7H $^2\Pi_{1/2}$	22	13.2 ± 1.4	$(5.1 \pm 1.0) \times 10^{12}$	1260.39	Reliable fit (1 point removed)
C_7H $^2\Pi_{3/2}$	11	14.7 ± 2.2	$(3.2 \pm 2.0) \times 10^{12}$	110.99	Moderate data dispersion
C_8H $^2\Pi_{3/2}$	30	14.8 ± 0.6	$(8.9 \pm 0.9) \times 10^{12}$	2111.08	Reliable fit (1 point removed)
C_8H $^2\Pi_{1/2}$	16	8.0 ± 1.4	$(1.15 \pm 1.38) \times 10^{13}$	35.16	Large data dispersion
C_5N^-	6	8.8 ± 1.4	$(4.9 \pm 1.6) \times 10^{12}$	132.62	Reliable fit
C_6H^-	7	11.4 ± 1.0	$(4.3 \pm 0.6) \times 10^{12}$	173.51	Reliable fit
C_8H^-	13	23 ± 5	$(1.1 \pm 0.4) \times 10^{12}$	810.24	Reasonable fit. Large data dispersion
SiC_4	6	12.9 ± 1.8	$(6.2 \pm 1.0) \times 10^{12}$	175.36	Reliable fit
$^{29}SiC_4$	5	10 ± 6	$(5.73 \pm 5.06) \times 10^{11}$	134.88	Large data dispersion
SiC_6	7	17.3 ± 2.5	$(1.3 \pm 0.4) \times 10^{12}$	588.57	Reliable fit (1 point removed)
CCS	4	7.9 ± 1.1	$(4.0 \pm 0.7) \times 10^{13}$	44.48	Reliable fit
C_3S	3	23.60 ± 96.85	$(2.58 \pm 3.60) \times 10^{13}$	170.50	Very high data dispersion
NaCN	7	24 ± 13	$(8.5 \pm 1.7) \times 10^{12}$	425.76	Large data dispersion (1 point removed)
MgC_3N	14	11.1 ± 0.9	$(5.2 \pm 0.6) \times 10^{12}$	336.00	Reliable fit (1 point removed)
MgC_5N	30	15.4 ± 1.8	$(4.7 \pm 1.3) \times 10^{12}$	1109.27	Reliable fit (1 point removed)
MgC_4H	10	4.7 ± 1.2	$(1.5 \pm 1.3) \times 10^{13}$	140.43	Large data dispersion
MgC_6H	30	24.8 ± 8.9	$(2.0 \pm 0.9) \times 10^{13}$	966.24	Large data dispersion



Fig. 3. Graphic summary of all lines detected in this work ($\int T_A^* dv$ versus frequency). Dotted line marks the 1 σ limit in the data according to Fig. 1. The remaining U-lines appear in the bottom panel.



Fig. 4. A more physical view of most lines detected in this work, in the form of rotational diagrams. See fit results in Table 2. Note that the changing telescope beam across the Q-band is taken into account here through W=$\int T_{MB} dv$ in the y-axis, instead of $\int T_A^* dv$ as in Figure 3. The values of W for each individual line can be found in Table A.1. The fits shown correspond to molecules listed in Table 2.

4.9. Most significant U-lines

A careful, long and painstaking work, has allowed us to reduce to a minimum the number of unidentified features in this work. They are summarized in Table A.2.

Among all the remaining U-Lines, the strongest one, at 14σ appears at a frequency of 41711.90 MHz and was already present and not identified by Kawaguchi et al. (1995). MADEX does not provide any candidate to be the carrier, and the same applies for other catalogs.

After this strong U-line we have to go down to 6σ for the next unidentified line, standing at a central frequency of 42340 MHz. This line appears as a nice and isolated spectral feature also present at the same position on all receiver tunings (reference central frequencies: 42300.0, 42280.0 and 41370.0 MHz). A look at MADEX catalog does not provide any plausible carrier as the only candidate, $Ag^{37}Cl$, does not have any emission at other frequencies in this survey, around 35284 and 49396 MHz, and $Ag^{35}Cl$ is not found either.

At the 5σ level there are just 2 unidentified lines at the following frequencies in round numbers: 35849 and 37663 MHz. These two lines, again, appear isolated and well detected on all three different receiver tunings at the same position. Again, no answers are found in MADEX about possible carriers. It is worth noting, in any case, that we are talking about features with peak intensities below 3 mK in the T_A^* scale.

At the 4σ level we have only one unidentified line, at 43457.12 MHz. Again, it appears at the same position in all three receiver tunings but no carrier has been found for it yet.

4.10. Unidentified lines at 3σ level

Apart from the most significant unidentified lines discussed in section 4.9, 52 more features remain unidentified just at the 3σ level and with line widths within ±20% from that typical of the source. Most of these features are in fact at ∼1 mK level so that some of them could just be artifacts that would disappear on deeper integrations (that we plan to do) or might be baseline effects although we do not have evidence to remove any of the remaining ones, as we have done with several other in our careful data reduction process. Table A.2 should therefore be kept as a reference for follow-up in future works.

5. Procedure for line identification

The first thing to say in this section is that not all the identifications are 100% sure, although we are quite confident about most of them. The main doubts arise when (i) the frequency coincidence is not within ∼1 MHz typically, (ii) the line width is too narrow or too broad with respect to that typical of the source, (iii) there is a close blending with a stronger line, and (iv) only one line for a given species is found and we cannot confirm it from other lines detected at higher frequencies with other telescopes. These circumstances are pointed out in our line identification Table A.1, when necessary.

For our data analysis and line identification process, we have defined a reference file containing the data smoothed to a resolution of 229 kHz (6 channel smoothing from the original output), from which we have removed all artifacts (spikes, baseline undulations) as much as we could. Then we fitted all the spectral features in the data using the method SHELL implemented in GILDAS package, which is adequate for the CW Leo expanding envelope. We left the expansion velocity as a free parameter in this fit as an extra check for the line identification process (see below). The fitted line profiles were stored as a third column in the reference file so that we can incorporate them to the figures for the convenience of the reader. We created a separate file containing the MADEX catalog entries that we identified as corresponding to the observed spectral features, and another file storing the SHELL method fitting parameters. The first file (irc10216_qband.dat) is used to create figures . The other two files (irc10216_qband.lines and irc10216_qband.parameters) are used as inputs for a fortran script, specially designed for this paper, that automatically creates Tables 1 and A.1, calculating in particular $W=\int T_{MB} dv$ where the changing telescope beam across the Q-band is taken into account, and also updates the numbers and percentages of identified lines given in the abstract and in the main text. This allows us to introduce criteria in our analysis, and new fits or baseline corrections, that are immediately checked throughout all the data in a consistent consistent way, with the results updated in our manuscript without having to check "by hand" again and again. This also provides a systematic way of re-checking the whole work if, for example, more observations are added and, therefore, the noise level is improved and/or, more frequencies are added.

6. Legacy and summary

The main purpose of this work has been to provide a reference in the Q-band for further investigations on the molecular content of the circumstellar envelope of CW Leo using other bands. For that reason we provide the full data set and line fits as on-line material available for the scientific community.

Nevertheless, we can say a few things about the results published here. First of all, the large amount of lines and molecules detected provides an opportunity to give a very good diagnostics for the rotational temperature. To do that we use those molecules in Table 2 with reliable fits. We provide three results, the first two are weighted averages using the number of lines detected and the derived column density. The third value is just an arithmetic average. The values are: 16.9, 11.4, and 14.1 K respectively.

Concerning the isotopic ratios, the only one we can give in this work is $^{12}C/^{13}C$ using HC_5N due to the weakness of the lines from the HC_7N isotopologues and the small number of lines (2) from HC_3N and its isotopologues. The result ($^{12}C/^{13}C$ = 43.0± 4.7 is in good agreement with previous works, 44 ± 3 from Kahane et al. (1992) and 45 ± 3 from Cernicharo et al. (2000).

The other important contribution of this work is that it provides a large amount of data on molecular column densities to be used for comparisons with results from chemical models of this source, together with the full 4.3-1.0 mm data from the IRAM-30m telescope that should also be published soon. Detected molecules with very few lines in the Q-band are left for analysis in that future publication, combining the data from them published here (Table A.1 and Figures A.1 to A.39) with those obtained with the IRAM-30m telescope.

In this paper we have presented the full results of Q-band (31.0-50.3 GHz) deep integration towards IRC+10216, one of the richest and more exhaustively studied molecular stellar envelopes in the sky. This work represents a big improvement over previously available surveys in the same frequency range, in terms of both noise level and line identification due to the use of state of the art line catalogs. Prior to this publication, a series of papers has presented the identification from these data of new molecular species and/or vibrational states never before detected in space of known molecular species. The list of U-lines is almost entirely reduced to features at 3-4 σ level with the exception of three lines that still defy identification.

In order to provide a useful legacy from this project, we have proceeded to fit all the spectroscopic features in the data to provide detailed figures of data versus fit; tables with line identifications with theoretical versus experimental line parameters; and, finally, rotational diagrams to explore the physical conditions of IRC+10216 through the molecular species present in this work.

Acknowledgements. We thank Ministerio de Ciencia e Innovación of Spain for funding support through projects PID2019-106110GB-I00, PID2019-107115GB-C21 / AEI / 10.13039/501100011033, PID2019-106235GB-I00, and grant FJCI-2016-27983 for CB. We also thank ERC for funding through grant ERC-2013-Syg- 610256-NANOCOSMOS. M.A. thanks MICIU for grant RyC-2014-16277.

References

Alexander, A. J., Kroto, H. W., & Walton, D. R. M., 1976, J. Mol. Spectrosc., 62, 175.
Allen, M.D., Evenson, K.M., Brown, J.M., et al., 2001, J. Mol. Spectrosc., 209, 143
Anderson, M. A., Steimle, T. C., & Ziurys, L. M., 1994, ApJ, 429, L41
Anttila, R., Horneman, V.M., Koivusaari, M., et al., 1993, J. Mol. Spectrosc., 157, 198
Aoki, K., and Ikuta, S., 1993, J. Chem. Phys., 99, 3809.
Bester, M., Tanimoto, M., Vowinkel, B., et al. 1983, Z. Naturforschung., 38a, 64
Bester, M., Yamada, K., Winnewisser, G., et al. 1984, A&A, 137, L20
Bizzocchi, L., Degli Esposti, C., Botschwina, P., 2004, J. Mol. Spectrosc., 225, 145
Blanksby, S. J., McAnoy, A. M., Dua, S., et al., 2001, MNRAS, 328, 89
Bogey, M., Demuynck, C., & Destombes, J. L., 1986, Chem. Phys. Letters, 125, 383
Botschwina, P., Horn, M., Seeger, S., et al., 1992, Chem. Phys. Letters, 195, 427.
Botschwina, P., 1996, Chem. Phys. Lett., 259, 627
Botschwina, P., Horn, M., Markey, K., et al., 1997, Mol. Phys. 92, 381.
Caris, M., Lewen, F., & Winnewisser, G., 2002, Z. Naturforschung, 57a, 663
Caris, M., Lewen, F., Muller, H.S.P., Winnewisser, G., 2004, J. Mol. Struct., 2004, 695, 243
Cazzoli, G., & Puzzarini, C., 2006, J. Mol. Spectrosc., 240, 153
Cernicharo, J. 1985, Internal IRAM report (Granada: IRAM)
Cernicharo, J., Guélin, M., Kahane, K., et al. 1991, A&A, 246, 213
Cernicharo, J., Guélin, M., Kahane, C., 2000, A&AS, 142, 181
Cernicharo, J., Guélin, M., Agúndez, M., et al. 2008, ApJ, 688, L83
Cernicharo, J., Agúndez, M., Kahane, C., Guélin, M., Goicoechea, J.R., Marcelino, N., de Beck, E.,& Decin, L., 2011, A&A, 529, L3
Cernicharo, J., 2012, in ECLA 2011: Proc. of the European Conference on Laboratory Astrophysics, EAS Publications Series, 2012, Ed.: C. Stehl, C. Joblin, & L. d'Hendecourt (Cambridge: Cambridge Univ. Press), 251; https://nanocosmos.iff.csic.es/?page_id=1619
Cernicharo, J., Marcelino, N., Agúndez, M., et al. 2015, A&A, 575, A91
Cernicharo, J., Cabezas, C., Pardo, J. R., et al. 2019 A&A, 63, L2
Cernicharo, J., Agúndez, Cabezas, C., Tercero, B., Marcelino, M., Pardo, J.R., & de Vicente, P., 2021, A&A, 649, L15
Cernicharo, J., Agúndez, Kaiser, R. I., Cabezas, C., Tercero, B., Marcelino, M., Pardo, J.R., & de Vicente, P., 2021, A&A, 652, L9
Chen, W., Novick, S. E., McCarthy, M. C., Gottlieb, C. A., & Thaddeus, P., 1995, J. Chem. Phys. 103, 7828
Cho, S.-H., & Saito, S., 1998, ApJ, 496, L51.
Clouser, P.L., & Gordy, W., Phys. Rev. 1964, 134A, 863
Creswell, R. A., Winnewisser, G., & Gerry, M. C. L., 1977, J. Mol. Spectrosc. 65, 420
Degli Esposti, C., and Bizzocchi, L., 2004, ApJ., 614, 518
Ebenstein, W.L., & Muenter, J.S., 1984, J. Chem. Phys., 80, 3989
Gripp, J., Guarnieri, A., Stahl, W., et al., 2000, J. Mol. Struct., 526, 81
Gordon, V.D., Nathan, E.S., Apponi, A.J., et al., 2000, J. Chem. Phys., 113, 5311
Gottlieb, C. A., Gottlieb, E. W., Thaddeus, P., & Kawamura, H. 1983, ApJ 275, 916
Gottlieb, C. A., Vrtilek, J. M., Gottlieb, E. W., Thaddeus, P., & Hjalmarson, A., 1985, ApJ, 294, L55
Gottlieb, C.A., Vrtilek, J. M., & Thaddeus, P., 1989, ApJ, 343, L29
Gottlieb, C., Myers, P.C., & Thaddeus, P., 2003, ApJ, 588, 655
Groenewegen, M. A. T., Barlow, M. J., Blommaert, J. A. D. L., et al. 2012, A&A, 543, L8
Guarnieri, A., Hinze, R., Krüger, M., et al., 1992, J. Mol. Spectrosc., 156, 39
Guélin, M., P. Friberg, and A. Mezaoui, 1982, A&A, 109, 23
Guélin, M., Patel, N. A., Bremer, M., et al., 2018, A&A, 610, A4
Gupta, H., Brünken, S., Tamassia, F., et al., 2007, ApJ, 655, L57
Halfen, D.T., & Ziurys, L.M., 2011, ApJ, 730, 107
Hensel, K.D., Styger, C., Jäger, W., et al., 1993, J. Chem. Phys., 99, 3320
Hirahara, Y., Ohshima, Y., & Endo, Y., 1993, ApJ, 403, L83
Hirano, T., Ishii, K., Odaka, T.E., et al., 2002, J. Mol. Spectrosc., 215, 42
Ikuta, S. Tsuboi, T., & Aoki, K., 2000, J. Mol. Struct., 528, 297
Kahane, C., Cernicharo, J., Gómez-González, J., & Guélin, M. 1992, A&A, 256, 235
Kaifu, N., Ohishi, M., Kawaguchi, K., Saito, S., Yamamoto, S., Miyaji, T., Miyazawa, K., Ishikawa, S. I., Noumara, C., Harasawa, S., Okuda, M., & Suzuki, H., 2004, PASJ, 56, 69
Kanata, H., Yamamoto, S., & Saito, S., 1987, Chem. Phys. Letters, 140, 221
Kasai, Y., Sumiyoshi, Y., Endo, Y., 1997, ApJ, 477, L65
Kawaguchi, K., Takano, S., Ohishi, M., et al., 1992, ApJ, 396, L49
Kawaguchi, K., Kagi, E., Hirano, T., et al., 1993, ApJ, 406, L39
Kawaguchi, K., Kasai, Y., Ishikawa, S., & Kaifu, N., 1995, PASJ, 47, 853
Killian, T. C., Vrtilek, T. C., Gottlieb, J. M., 1990 , ApJ, 365, L89
Kokkin, D. L., Brünken, S., Young, K. H., et al., 2011, ApJS, 196, 17
Lee, S., 1997, Chem. Phys. Letters, 268, 69.
Lide, D. R., 1965, J. Chem. Phys., 42, 1013.
Linnartz, H., Motylewski, T., Vaizert, O., Maier, J. P., Apponi, A. J., McCarthy, M. C., Gottlieb, C. A., & Thaddeus, P. 1999, J. Mol. Spectrosc. 197, 1
Lovas, F. J., Suenram, R. D., Ogata, T., & Yamamoto, S., 1992, ApJ, 399, 325
Mallinson, P. D., & de Zafra, R. L., 1978, Mol. Phys. 36, 827.
Manson Jr, E. L., Clark, W. W., DeLucia F. C., & Gordy, W., 1977, Phys. Rev. A, 15, 223.
Mbosei, L., Fayt, A., Dréan, P., & Cosléou, J., 2000 , J. Mol. Struc., 517, 271.
McCarthy, M. C., Gottlieb, C. A., Thaddeus, P., Horn, M., & Botschwina, P., 1995, J. Chem. Phys., 103, 7820
McCarthy, M. C., Travers, M. J., Kovács, A., Gottlieb, C. A., & Thaddeus, P., 1996, A&A, 309, L31
McCarthy, M. C., Travers, M. J., Kovács, A., Gottlieb, C. A., & Thaddeus, P., 1997, ApJ, 113, 105
McCarthy, M. C. Chen, W., Apponi, A. J., Gottlieb, C. A., & Thaddeus, P., 1999, ApJ, 520, 158
McCarthy, M. C., Levine, E. S., Apponi, A. J., & Thaddeus, P., 2000, J. Mol. Spectrosc. 203, 75
McCarthy, M. C., Fuchs, G. W., Kucera, J., Winnewisser, G., & Thaddeus, P., 2003, J. Chem. Phys., 118, 3549
McCarthy, M. C., Gottlieb, C. A., Gupta, H., & Thaddeus, P., 2006, ApJ, 652, L141.
Moises, A., Boucher, D., Burie, J., et al., 1982, J. Mol. Spectrosc., 92, 497
Mollaaghababa, R., Gottlieb C. A., Vrtilek J. M., & Thaddeus, P., 1991, ApJ, 368, L19
Müller H. S. P., McCarthy M. C., Bizzocchi L., et al., 2007, Phys. Chem. Chem. Phys., 9, 1579
Müller, H. S. P., Drouin, B. J., & Pearson, J. C., 2009, A&A, 506, 1487
Ohishi, M., Kaifu, N., Kawaguchi, K., et al. 1989, ApJ, 345, L83
Ohshima, Y, & Endo, Y., 1992, J. Mol. Spectrosc., 153, 627
Oswald, M., & Botschwina, P., 1995, J. Mol. Spectrosc. 169, 181
Oyama et al., 2020, ApJ, 890, A39
Pardo, J. R., Cernicharo, J., Serabyn, E. 2001, IEEE Trans. Antennas and Propagation, 49, 12
Pardo, J. R., Bermúdez, C., Cabezas, C., Agúndez, M., Gallego, J. D., Fonfría, J. P., Velilla-Prieto, L., Quintana-Lacaci, G., Tercero, B., Guélin, M., & Cernicharo, J., 2020, A&A, 640, L13
Pardo, J. R., Cabezas, C., Fonfría, J. P., Agúndez, M., Tercero, B., de Vicente, P., Guélin, M., & Cernicharo, J., 2021, A&A, 652, L13
Pavone, F. S., Zink, L. R., Prevedelli, M., et al., 1990, J. Mol. Spectrosc., 144, 45
Raymonda, J. W., Muenter, J. S., & Kemplerer W. A., 1970, J. Chem. Phys., 52, 3458
Saito, S., Kawaguchi, K., Yamamoto, S., et al., 1987, ApJ, 317, L115
Sanz, M. E., McCarthy, M. C., Thaddeus, P., 2003, J. Chem. Phys., 119, 11715
Simeckova, M., Urban, S., Fuchs, U., et al., 2004, J. Mol. Spectrosc., 226, 123
Spezzano, S., Tamassia, F., Thorwirth, S., et al. 2012, ApJS, 200, 1
Steimle, T. C., & Bousquet, R. R., 2001, J. Chem. Phys., 115, 5203
Suenram, R. D., Lovas, F. J., & Matsumura, K., 1989, ApJ, 342, L103
Suenram, R. D., & Lovas, F. J., 1994, ApJ, 429, L89
Tang, J. A., Saito, S., 1995, J. Mol. Spectrosc., 169, 92
Tang, J. A., Sumiyoshi, Y., & Endo, Y., 1999 , Chem. Phys. Letters, 315, 69
Tercero, F., López-Pérez, J. A., Gallego, J. D., et al., 2021, A&A, 645, A37
Thaddeus, P., Vrtilek, J. M., & Gottlieb, C. A., 1985, ApJ, 299, L63
Thaddeus, P., Gottlieb, C. A., Gupta, H., Brünken, S., McCarthy, M. C., Agúndez, M., Guèlin, M., & Cernicharo, J., 2008, 677, 1139
Thorwirth, S., Müller, H. S. P., & Winnewisser, G., 2000, J. Mol. Spectrosc. 204, 133
Thorwirth, S., Müller, H. S. P., & Winnewisser, G., 2001, Phys. Chem. Chem. Phys., 3, 1236
Tiemann, E., Renwanz, E., Hoeft, J., et al., 1972, Z. Naturforschung, 27a, 1566
Timp, B. A., Doran, J. L., Iyer, S., et al., 2012, J. Mol. Spectrosc., 271, 20
Van Vaals, J. J., Meerts, W. L., & Dymanus, A., 1984, Chem. Phys., 86, 147
Vrtilek, J. M., Gottlieb, C. A., & Thaddeus, P., 1987, ApJ, 314, 716
Vrtilek, J. M., Gottlieb, C. A., Gottlieb, E. W., Killian, T. C. & Thaddeus, P., 1990, ApJ, 364, L53
Winnewisser, G., & Cook, R. L., 1968, J. Mol. Spectrosc. 28, 266.
Woon, D. E., 1995, Chem. Phys. Lett. 244, 45
Wyse, F. C., & Gordy, W., 1972, J. Chem. Phys., 56, 2130.
Yamamoto, S., Saito, S., Kawaguchi, K., et al., 1987, ApJ, 317, L119
Yamamoto, S. Saito, S., Suzuki, H., Deguchi, S., Kaifu, N., Ishikawa, S. I., & Ohishi, M., 1990, ApJ, 348, 363
Yamamoto, S., Saito, S., & Kawaguchi, K., 1990, ApJ, 361, 318
Zelinger, Z., Amano, T., Ahrens, V., et al., 2003, J. Mol. Spectrosc., 220, 233

Appendix A: Tables and Figures

This appendix is devoted to provide the exhaustive information of our IRC+10216 Q-band survey with the Yebes 40m telescope, in two forms. First, the most significant line parameters (theoretical + observed) of all spectral lines that can be assigned to an identified molecular species are given in Table A.1. The accumulated number of distinguishable lines for each species is provided in the last column of Table A.1. This table is the basis to build Figures 3 and 4. The observed values in this table, including errors in parenthesis affecting the last digits of the observed frequency and the expansion velocity, come from the line fits performed over the data with the Gildas SHELL method as shown in Figures A.1 to A.39, that can be found in this appendix after the tables. An error of 0 or 0.0 means the given observed frequency (ν_{obs}) and/or expansion velocity (Vexp) has been forced in the fit. The figures themselves have labels to identify the positions of all the lines that appear in Table A.1, and they also label the U-lines listed in Table A.2. The frequency scale in all tables and figures assumes a velocity of the source with respect to the local standard of rest of -26.5 kms^{-1}. The changing telescope beam across the Q-band is taken into account in the calculation of W=$\int T_{MB} dv$.

Table A.1. Line identification and associated parameters from our 31.0-50.3 GHz observations presented in this work.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	W=$\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ a	31093.03	31093.03(0)	20.4	52.80	509	20.926	14.5(0.0)	A.1a	1
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ b	31093.42	31093.42(0)	20.4	52.80	339	20.521	14.5(0.0)	A.1a	1
$H^{13}CCCCCN$	J=12→11	31120.03	31119.94(23)	9.7	12.00	761	23.622	13.5(2.2)	A.1a	1
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ a	31151.11	31150.84(23)	48.3	53.00	169	19.816	14.0(2.2)	A.1a	2
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ b	31160.98	31161.29(19)	48.4	53.00	337	20.508	16.3(2.4)	A.1a	3
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{13}{2}\rightarrow\frac{11}{2}$ a	31241.77	31241.77(0)	40.1	11.30	1421	24.063	14.5(0.0)	A.1b	1
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{13}{2}\rightarrow\frac{11}{2}$ b	31242.33	31242.33(0)	40.1	13.20	1254	23.782	14.5(0.0)	A.1b	1
NaCN	$J_{Ka,Kc}=2_{0,2}\rightarrow1_{0,1}$	31262.33	31262.36(23)	2.3	2.00	751	23.966	14.5(2.2)	A.1b	1
MgC_6H	$N_J=27_{\frac{53}{2}}\rightarrow26_{\frac{51}{2}}$	31305.40	31305.41(13)	20.9	26.50	332	23.094	14.5(0.0)	A.1b	1
MgC_6H	$N_J=27_{\frac{55}{2}}\rightarrow26_{\frac{53}{2}}$	31307.36	31307.36(10)	20.9	27.50	332	23.056	14.5(0.0)	A.1b	2
HC_9N	J=54→53	31375.44	31375.45(23)	41.4	54.00	1568	22.466	13.9(2.2)	A.1b	1
HC_9N	ν^e_{19} J=54→53	31407.07	31406.86(23)	113.5	54.00	411	21.127	13.2(2.2)	A.1b	2
HC_9N	ν^f_{19} J=54→53	31422.87	31422.82(11)	113.6	54.00	246	20.614	13.9(1.1)	A.1b	3
HC_7N	J=28→27	31583.71	31583.69(0)	22.0	28.00	10352	25.156	13.9(0.0)	A.2a	1
HC_7N	ν^e_{15} J=28→27	31623.21	31623.31(23)	111.2	28.00	483	22.090	12.7(2.2)	A.2a	2
$HC^{13}CCCCN$	J=12→11	31624.34	31624.36(23)	9.9	12.00	644	23.440	14.6(2.2)	A.2a	1
$HCCCC^{13}CN$	J=12→11	31636.13	31636.04(0)	9.9	12.00	966	23.844	12.8(0.3)	A.2a	1
HC_7N	ν^f_{15} J=28→27	31643.34	31643.18(3)	111.2	28.00	241	21.395	13.6(1.3)	A.2a	3
SiC_6	J=26→25	31784.66	31782.93(10)	20.6	26.00	238	20.383	14.4(0.0)	A.2b	1
C_4H	ν_7 $N_J=3_{\frac{7}{2}}\rightarrow2_{\frac{5}{2}}$ a	31786.25	31785.94(3)	247.1	5.60	873	25.947	14.3(0.4)	A.2b	1
C_4H	ν_7 $N_J=3_{\frac{7}{2}}\rightarrow2_{\frac{5}{2}}$ b	31830.35	31830.32(3)	247.1	5.60	790	25.846	15.5(0.4)	A.2b	2
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$ a	31881.86	31881.81(1)	9.3	22.60	2752	23.757	13.3(0.1)	A.2b	1
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$ b	31885.54	31885.53(2)	9.3	22.60	2672	23.728	13.8(0.1)	A.2b	2
$HCC^{13}CCCN$	J=12→11	31918.69	31918.64(2)	10.0	12.00	1019	23.888	13.4(0.3)	A.2b	1
$HCCC^{13}CCN$	J=12→11	31922.57	31922.55(1)	10.0	12.00	783	23.625	14.3(0.3)	A.2b	1
HC_5N	J=12→11	31951.78	31951.75(0)	10.0	12.00	43975	27.652	13.8(0.0)	A.3a	1
HC_9N	J=55→54	31956.44	31956.45(1)	42.9	55.00	1249	22.202	14.2(0.2)	A.3a	4
C_6H	ν_{11} $^2\Delta_{\frac{5}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$	31967.06	31967.45(22)	75.9	11.00	312	22.297	17.3(0.9)	A.3a	3
HC_9N	ν^e_{19} J=55→54	31988.66	31988.63(23)	115.1	55.00	234	20.524	13.8(2.1)	A.3a	5
HC_5N	ν^e_{11} J=12→11	32004.59	32004.59(0)	163.5	11.90	233	22.419	14.5(0.0)	A.3a	2
HC_9N	ν^f_{19} J=55→54	32004.75	32004.75(0)	115.1	55.00	233	20.522	14.5(0.0)	A.3a	6
HC_5N	ν^f_{11} J=12→11	32032.49	32032.71(7)	163.5	11.90	388	22.927	15.1(0.9)	A.3a	3
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$ a	32095.24	32095.24(1)	31.7	23.00	1850	23.336	13.9(0.1)	A.3a	4
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$ b	32125.56	32125.56(0)	31.8	23.00	1998	23.412	14.2(0.1)	A.3a	5
l-HC_4N	$N_J=7_7\rightarrow6_6$	32233.42	32233.42(0)	7.1	6.86	228	22.942	14.5(0.0)	A.3b	1
C_6H	ν_{11} $^2\Delta_{\frac{3}{2}}$ J=$\frac{23}{2}\rightarrow\frac{21}{2}$	32234.82	32234.82(0)	93.4	11.30	152	21.544	14.2(0.0)	A.3b	6
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{55}{2}\rightarrow\frac{53}{2}$ a	32266.33	32266.33(0)	22.0	54.90	455	20.739	14.5(0.0)	A.3b	4
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{55}{2}\rightarrow\frac{53}{2}$ b	32266.74	32266.74(0)	22.0	54.80	455	20.740	14.5(0.0)	A.3b	4
MgC_5N	$N_J=28_{\frac{55}{2}}\rightarrow27_{\frac{53}{2}}$	32278.59	32278.52(7)	22.5	27.50	152	20.098	14.5(0.0)	A.3b	1
MgC_5N	$N_J=28_{\frac{57}{2}}\rightarrow27_{\frac{55}{2}}$	32280.40	32280.53(7)	22.5	28.50	379	20.978	14.5(0.0)	A.3b	2
$HC^{13}CCCCCCN$	J=29→28	32324.86	32324.86(0)	23.3	29.00	302	21.562	14.5(0.0)	A.3b	1
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{55}{2}\rightarrow\frac{53}{2}$ a	32326.79	32326.79(0)	49.9	55.00	226	20.036	14.5(0.0)	A.3b	5
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{55}{2}\rightarrow\frac{53}{2}$ b	32336.63	32336.79(5)	49.9	55.00	452	20.728	14.1(0.4)	A.3b	6
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{37}{2}\rightarrow\frac{35}{2}$ a	32354.93	32354.93(0)	15.1	36.90	376	21.121	14.5(0.0)	A.3b	1
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{37}{2}\rightarrow\frac{35}{2}$ b	32356.56	32356.56(0)	15.1	36.90	301	20.898	14.5(0.0)	A.3b	2
NaCN	$J_{Ka,Kc}=2_{1,1}\rightarrow1_{1,0}$	32373.67	32373.79(9)	4.7	1.50	300	23.303	14.6(1.3)	A.3b	2
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{37}{2}\rightarrow\frac{35}{2}$	32429.51	32429.49(4)	52.7	73.50	448	20.605	13.8(0.4)	A.3b	3
MgC_6H	$N_J=28_{\frac{55}{2}}\rightarrow27_{\frac{53}{2}}$	32464.88	32464.93(3)	22.5	27.50	223	22.623	14.5(0.0)	A.4a	3
MgC_6H	$N_J=28_{\frac{57}{2}}\rightarrow27_{\frac{55}{2}}$	32466.84	32466.84(0)	22.5	28.50	74	21.489	14.5(0.0)	A.4a	4
HC_9N	J=56→55	32537.45	32537.44(1)	44.5	56.00	962	21.905	13.9(0.1)	A.4a	7
HC_9N	ν^e_{19} J=56→55	32570.25	32570.32(9)	116.6	56.00	443	21.128	14.7(0.4)	A.4a	8
$HCC^{13}CCCCCN$	J=29→28	32577.82	32577.88(32)	23.5	29.00	295	21.531	14.2(0.3)	A.4a	1

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
HCCCCC^{13}CCN	J=29→28	32585.58	32585.55(26)	23.5	26.00	147	20.946	14.5(0.0)	A.4a	1
HC_9N	v_{19}^f J=56→55	32586.63	32586.76(5)	116.7	54.00	368	20.980	14.5(0.0)	A.4a	9
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=1→-1 F=1→1	32617.02	32617.01(23)	1.6	0.33	588	27.298	14.1(2.1)	A.4a	1
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=1→-1 F=2→1	32627.30	32627.30(0)	1.6	1.66	3082	27.349	14.1(0.1)	A.4a	2
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=1→-1 F=1→0	32634.39	32634.34(1)	1.6	0.67	1173	27.298	13.6(0.1)	A.4a	3
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=-1→1 F=2→1	32660.65	32660.68(23)	1.6	1.66	3439	27.457	14.4(2.1)	A.4a	4
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=-1→1 F=1→0	32663.36	32663.35(23)	1.6	0.67	1244	27.355	15.2(2.1)	A.4a	5
C_3H	J=$\frac{3}{2}$ → $\frac{1}{2}$ p=-1→1 F=1→1	32667.67	32667.48(23)	1.6	0.33	1024	27.855	15.7(2.1)	A.4a	6
HCCC^{13}CCCCN	J=29→28	32693.21	32693.05(23)	23.5	29.00	292	21.517	14.5(0.0)	A.4a	1
HCCCC^{13}CCCN	J=29→28	32696.08	32695.92(23)	23.5	29.00	219	21.229	14.5(0.0)	A.4a	1
HC_7N	J=29→28	32711.67	32711.65(0)	23.5	29.00	9759	25.027	13.8(0.0)	A.4b	4
HC_7N	v_{15}^e J=29→28	32752.58	32752.58(23)	112.7	29.00	580	22.203	13.7(2.1)	A.4b	5
HC_7N	v_{15}^f J=29→28	32773.43	32773.42(23)	112.7	29.00	507	22.067	13.9(2.1)	A.4b	6
HC_7N	v_{15}=2 J=29→28 (l=0)	32812.96	32812.96(0)	202.2	29.00	72	20.116	14.5(0.0)	A.4b	7
HC_7N	v_{15}=2 J=29→28 (l=-2)	32814.39	32814.39(0)	202.6	28.90	72	20.120	14.5(0.0)	A.4b	7
HC_7N	v_{15}=2 J=29→28 (l=+2)	32815.98	32815.98(0)	202.6	28.90	72	20.120	14.5(0.0)	A.4b	8
C_6H^-	J=12→11	33044.49	33044.45(0)	10.3	12.00	1767	23.127	13.7(0.1)	A.5a	1
CH_3C_3N	N_J=8_3→7_3	33048.75	33049.13(23)	74.5	6.88	71	21.556	15.4(2.1)	A.5a	1
CH_3C_3N	N_J=8_0→7_0	33051.62	33051.60(23)	7.1	8.00	71	21.405	12.9(2.1)	A.5a	2
H^{13}CCCCCCCN	J=30→29	33098.20	33098.00(2)	24.6	30.00	141	20.742	14.6(0.7)	A.5a	1
HC_9N	J=57→56	33118.45	33118.44(23)	46.1	57.00	1053	21.960	13.7(2.1)	A.5a	10
MgC_3N	N_J=$12_{\frac{23}{2}}$→$11_{\frac{21}{2}}$	33138.61	33138.59(23)	10.3	11.50	561	22.562	13.6(2.1)	A.5a	1
MgC_3N	N_J=$12_{\frac{25}{2}}$→$11_{\frac{23}{2}}$	33142.93	33142.98(23)	10.3	12.50	631	22.596	14.2(2.1)	A.5a	2
HC_9N	v_{19}^e J=57→56	33151.84	33151.84(0)	118.2	57.00	350	20.858	14.5(0.0)	A.5a	11
MgC_4H	N_J=$12_{\frac{23}{2}}$→$11_{\frac{21}{2}}$	33153.43	33153.43(0)	10.3	11.50	210	23.761	14.5(0.0)	A.5a	1
MgC_4H	N_J=$12_{\frac{25}{2}}$→$11_{\frac{23}{2}}$	33158.12	33158.12(0)	10.3	12.50	210	23.677	14.5(0.0)	A.5a	2
HC_9N	v_{19}^f J=57→56	33168.52	33168.59(6)	118.3	57.00	350	20.856	13.4(0.5)	A.5a	12
$^{29}SiC_4$	J=11→10	33224.79	33224.75(7)	9.6	11.00	139	21.157	13.0(0.4)	A.5b	1
MgC_5N	N_J=$29_{\frac{57}{2}}$→$28_{\frac{55}{2}}$	33431.38	33431.36(5)	24.1	28.50	273	20.616	14.5(0.0)	A.5b	3
MgC_5N	N_J=$29_{\frac{59}{2}}$→$28_{\frac{57}{2}}$	33433.19	33433.20(6)	24.1	29.50	273	20.582	14.5(0.0)	A.5b	4
HC^{13}CCCCCCN	J=30→29	33439.49	33439.49(0)	24.9	30.00	137	20.701	14.5(0.0)	A.5b	2
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ a	33439.61	33439.61(0)	23.6	56.80	478	20.718	14.5(0.0)	A.5b	7
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ b	33440.05	33440.05(0)	23.6	56.80	478	20.718	14.5(0.0)	A.5b	7
HCCCCCC^{13}CN	J=30→29	33451.94	33451.90(3)	24.9	30.00	273	21.393	13.6(0.4)	A.6a	1
C_6H	v_{11} N_J=$12_{\frac{25}{2}}$→$11_{\frac{23}{2}}$	33462.22	33462.24(1)	26.3	12.50	886	23.167	14.2(0.2)	A.6a	7
C_6H	v_{11} N_J=$12_{\frac{23}{2}}$→$11_{\frac{21}{2}}$	33479.82	33479.79(1)	26.3	11.50	748	23.082	13.9(0.2)	A.6a	8
$^{30}SiC_2$	$J_{Ka,Kc}$=$9_{2,7}$→$9_{2,8}$	33499.56	33497.91(5)	58.7	0.73	543	27.190	17.5(0.5)	A.6a	1
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ a	33502.47	33502.28(6)	51.5	57.00	272	20.147	13.8(0.5)	A.6a	8
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ b	33512.28	33512.05(5)	51.5	57.00	271	20.146	15.2(0.5)	A.6a	9
MgC_6H	N_J=$29_{\frac{57}{2}}$→$28_{\frac{55}{2}}$	33624.36	33624.36(0)	24.1	28.50	67	21.351	14.5(0.0)	A.6a	5
MgC_6H	N_J=$29_{\frac{59}{2}}$→$28_{\frac{57}{2}}$	33626.32	33626.31(5)	24.1	29.50	67	21.316	14.5(0.0)	A.6a	6
HC_9N	J=58→57	33699.45	33699.43(1)	47.7	58.00	934	21.806	13.9(0.1)	A.6a	13
HCC^{13}CCCCCN	J=30→29	33701.16	33701.09(7)	25.1	30.00	200	21.076	13.6(0.6)	A.6b	2
HCCCCC^{13}CCN	J=30→29	33709.19	33709.14(39)	25.1	30.00	133	20.670	13.8(0.6)	A.6b	2
H^{13}CCCCCN	J=13→12	33713.33	33713.30(2)	11.3	13.00	867	23.592	13.5(0.2)	A.6b	2
HC_9N	v_{19}^e J=58→57	33733.43	33733.51(4)	119.8	58.00	466	21.109	15.8(0.3)	A.6b	14
SiC_4	J=11→10	33742.68	33742.66(1)	9.7	11.00	1463	23.494	14.0(0.1)	A.6b	1
HC_9N	v_{19}^f J=58→57	33750.40	33750.30(0)	119.9	58.00	532	21.240	14.5(0.0)	A.6b	15
CCS	N_J=2_3→1_2	33751.37	33751.35(23)	3.2	2.98	3322	27.240	14.5(0.0)	A.6b	1
HCCC^{13}CCCCN	J=30→29	33820.53	33820.53(0)	25.2	30.00	132	20.657	14.5(0.0)	A.6b	2
C_8H^-	J=29→28	33833.31	33833.21(4)	24.4	29.00	330	20.068	13.2(0.5)	A.6b	1
HC_7N	J=30→29	33839.63	33839.61(0)	25.2	30.00	9496	24.931	13.7(0.0)	A.6b	9
HC_7N	v_{15}^e J=30→29	33881.95	33881.85(7)	114.4	30.00	657	22.259	14.5(0.2)	A.6b	10
HC_7N	v_{15}^f J=30→29	33903.52	33903.52(5)	114.4	30.00	525	22.034	14.1(0.2)	A.6b	11
HC_7N	v_{15}=2 J=30→29 (l=0)	33944.29	33944.29(0)	203.8	30.00	131	20.643	14.5(0.0)	A.6b	12
HC_7N	v_{15}=2 J=30→29 (l=-2)	33945.89	33945.89(0)	204.2	29.90	392	21.744	14.5(0.0)	A.6b	12
HC_7N	v_{15}=2 J=30→29 (l=+2)	33947.65	33947.65(0)	204.2	29.90	131	20.646	14.5(0.0)	A.6b	13
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{39}{2}$ → $\frac{37}{2}$ a	34103.86	34103.75(1)	16.7	39.00	258	20.636	12.8(0.3)	A.7a	4
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{39}{2}$ → $\frac{37}{2}$ b	34105.50	34105.37(1)	16.7	39.00	387	21.041	15.4(0.3)	A.7a	4
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{39}{2}$ → $\frac{37}{2}$	34182.42	34182.42(0)	54.4	77.50	384	20.345	14.5(0.0)	A.7a	5
H^{13}CCCCCCCN	J=31→30	34201.44	34201.50(11)	26.3	31.00	192	20.986	14.9(0.9)	A.7b	2
HC^{13}CCCCN	J=13→12	34259.67	34259.62(1)	11.5	13.00	955	23.672	13.6(0.1)	A.7b	2
HCCCC^{13}CN	J=13→12	34272.44	34272.43(22)	11.5	13.00	1081	23.796	14.3(0.1)	A.7b	2
HC_9N	J=59→58	34280.46	34280.46(3)	49.4	59.00	1080	21.916	14.5(0.1)	A.7b	16
HC_9N	v_{19}^e J=59→58	34315.01	34315.14(3)	121.5	59.00	317	20.689	13.6(0.3)	A.7b	17
HC_9N	v_{19}^f J=59→58	34332.28	34332.21(5)	121.5	59.00	379	20.869	13.0(0.3)	A.7b	18
HC^{13}CCCCCCN	J=31→30	34554.10	34554.10(7)	26.5	31.00	62	19.848	12.1(0.9)	A.8a	3
HCCCCCC^{13}CN	J=31→30	34566.97	34567.09(1)	26.5	31.00	124	20.539	13.4(0.1)	A.8a	2
HCC^{13}CCCN	J=13→12	34578.54	34578.56(2)	11.6	13.00	867	23.567	14.0(0.1)	A.8a	2
HCCC^{13}CCN	J=13→12	34582.75	34582.84(0)	11.6	13.00	1177	23.872	15.0(0.0)	A.8a	2
MgC_5N	N_J=$30_{\frac{59}{2}}$→$29_{\frac{57}{2}}$	34584.17	34584.20(23)	25.7	29.50	310	20.673	14.5(0.0)	A.8a	5
MgC_5N	N_J=$30_{\frac{61}{2}}$→$29_{\frac{59}{2}}$	34585.98	34586.10(23)	25.7	30.50	433	20.976	14.5(0.0)	A.8a	6
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{59}{2}$ → $\frac{57}{2}$ a	34612.90	34612.90(0)	25.2	58.80	433	20.549	13.9(0.0)	A.8a	10
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{59}{2}$ → $\frac{57}{2}$ b	34613.40	34613.40(0)	25.2	58.80	432	20.549	13.9(0.0)	A.8a	10

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
HC_5N	J=13→12	34614.39	34614.34(0)	11.6	13.00	38612	27.362	13.8(0.0)	A.8a	4
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$ a	34654.04	34654.01(0)	11.0	24.60	2586	23.527	14.2(0.0)	A.8a	9
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$ b	34658.38	34658.36(0)	11.0	24.60	2462	23.478	14.2(0.0)	A.8a	10
HC_5N	v_{11}^e J=13→12	34671.60	34672.97(3)	165.2	12.90	553	23.123	14.5(0.0)	A.8a	5
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{59}{2}\rightarrow\frac{57}{2}$ a	34678.14	34678.02(1)	53.2	59.00	307	20.202	14.4(0.3)	A.8a	11
C_3S	J=6→5	34684.37	34684.37(0)	5.8	6.00	2518	25.716	14.5(0.0)	A.8a	1
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{59}{2}\rightarrow\frac{57}{2}$ b	34687.93	34687.93(0)	53.2	59.00	307	20.201	14.5(0.0)	A.8a	12
HC_5N	v_{11}^f J=13→12	34701.83	34701.76(4)	165.2	12.90	245	22.309	13.9(0.5)	A.8b	6
C_6H	$v_{11}\ ^2\Delta_{\frac{5}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$	34746.86	34746.94(19)	77.6	12.00	428	22.443	14.2(0.4)	A.8b	11
MgC_6H	N_J=30$_{\frac{59}{2}}\rightarrow$29$_{\frac{57}{2}}$	34783.84	34784.02(7)	25.7	29.50	244	22.571	14.5(0.0)	A.8b	7
MgC_6H	N_J=30$_{\frac{61}{2}}\rightarrow$29$_{\frac{59}{2}}$	34785.79	34785.76(5)	25.7	30.50	122	21.844	14.5(0.0)	A.8b	8
C_4H	v_7 $N_{J,F}$=4$_{\frac{7}{2},3}\rightarrow$3$_{\frac{5}{2},2}$	34824.89	34824.89(0)	252.2	2.84	546	26.067	14.5(0.0)	A.8b	3
C_4H	v_7 $N_{J,F}$=4$_{\frac{7}{2},4}\rightarrow$3$_{\frac{5}{2},3}$	34825.45	34825.45(0)	252.2	3.83	425	25.516	14.5(0.0)	A.8b	3
HC_9N	J=60→59	34861.46	34861.47(1)	51.0	60.00	908	21.709	13.8(0.1)	A.8b	19
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$ a	34887.11	34887.10(1)	33.4	25.00	2054	23.274	13.8(0.1)	A.8b	12
HC_9N	v_{19}^e J=60→59	34896.59	34896.60(1)	123.2	60.00	302	20.607	14.0(0.0)	A.8b	20
HC_9N	v_{19}^f J=60→59	34914.15	34914.10(0)	123.2	60.00	362	20.787	14.5(0.0)	A.8b	21
C_4H	v_7 N_J=4$_{\frac{7}{2}}\rightarrow$3$_{\frac{5}{2}}$ b	34917.40	34917.16(0)	252.2	6.68	1024	25.837	14.5(0.0)	A.8b	4
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$ b	34917.89	34917.89(0)	33.4	25.00	1928	23.210	14.5(0.0)	A.8b	13
$HCCC^{13}CCCCN$	J=31→30	34947.85	34947.85(0)	26.8	31.00	60	19.804	14.5(0.0)	A.8b	3
HCN	v_2=1 J=12→12 l=1→-1	34953.77	34953.73(7)	1355.6	0.08	60	26.681	5.3(0.7)	A.9a	1
HC_7N	J=31→30	34967.59	34967.59(0)	26.9	31.00	8822	24.792	14.1(0.0)	A.9a	14
C_8H^-	J=30→29	34999.94	34999.94(6)	26.0	30.00	180	19.392	11.7(0.6)	A.9a	2
HC_7N	v_{15}^e J=31→30	35011.32	35011.24(4)	116.0	31.00	598	22.100	13.1(0.4)	A.9a	15
^{30}SiS	J=2→1	35021.16	35021.20(0)	2.5	2.00	657	26.978	14.2(0.0)	A.9a	1
HC_7N	v_{15}^f J=31→30	35033.60	35033.60(0)	116.1	31.00	537	21.992	14.5(0.0)	A.9a	16
C_6H	$v_{11}\ ^2\Delta_{\frac{3}{2}}$ J=$\frac{25}{2}\rightarrow\frac{23}{2}$	35037.68	35037.68(0)	95.1	12.30	358	22.232	14.5(0.0)	A.9a	14
$H^{13}CCCN$	J=4→3	35267.40	35267.35(0)	4.2	4.00	4276	26.619	13.7(0.0)	A.9b	1
$H^{13}CCCCCCCN$	J=32→31	35304.68	35304.68(0)	28.0	32.00	175	20.832	14.5(0.0)	A.9b	3
$Si^{34}S$	J=2→1	35315.28	35315.28(4)	2.5	2.00	934	27.321	14.0(0.1)	A.9b	1
HC_9N	J=61→60	35442.45	35442.43(1)	52.7	61.00	982	21.755	13.8(0.1)	A.9b	22
SiC_6	J=29→28	35452.02	35451.91(4)	25.5	29.00	231	20.134	12.9(0.4)	A.10a	2
HC_9N	v_{19}^e J=61→60	35478.18	35478.12(4)	126.6	62.00	288	20.511	13.4(0.3)	A.10a	23
HC_9N	v_{19}^f J=61→60	35496.03	35496.01(3)	126.6	62.00	345	20.691	13.2(0.2)	A.10a	24
o-H_2C_4	$J_{Ka,Kc}$=4$_{1,4}\rightarrow$3$_{1,3}$	35577.01	35577.05(3)	3.8	3.75	1086	24.949	14.5(0.3)	A.10a	1
^{29}SiS	J=2→1	35642.47	35642.45(1)	2.6	2.00	967	27.346	13.8(0.1)	A.10a	1
$HC^{13}CCCCCCN$	J=32→31	35668.72	35668.38(27)	28.2	32.00	113	20.387	17.0(1.7)	A.10a	4
o-H_2C_4	$J_{Ka,Kc}$=4$_{3,1}\rightarrow$3$_{3,0}$+4$_{3,2}\rightarrow$3$_{3,1}$	35716.54	35716.41(1)	112.0	1.75	396	24.698	13.2(0.2)	A.10b	2
p-H_2C_4	$J_{Ka,Kc}$=4$_{2,3}\rightarrow$3$_{2,2}$	35722.91	35722.91(0)	58.4	3.00	169	23.311	14.5(0.0)	A.10b	1
p-H_2C_4	$J_{Ka,Kc}$=4$_{2,2}\rightarrow$3$_{2,1}$	35723.06	35723.06(0)	58.4	3.00	169	23.311	14.5(0.0)	A.10b	1
p-H_2C_4	$J_{Ka,Kc}$=4$_{0,4}\rightarrow$3$_{0,3}$	35727.38	35727.38(2)	4.3	4.00	282	23.533	13.5(0.2)	A.10b	2
MgC_5N	N_J=31$_{\frac{61}{2}}\rightarrow$30$_{\frac{59}{2}}$	35736.95	35736.93(23)	27.4	30.50	282	20.514	14.5(0.0)	A.10b	7
MgC_5N	N_J=31$_{\frac{63}{2}}\rightarrow$30$_{\frac{61}{2}}$	35738.76	35738.74(23)	27.4	31.50	169	19.971	14.5(0.0)	A.10b	8
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{61}{2}\rightarrow\frac{59}{2}$ a	35786.18	35786.17(0)	26.9	60.80	337	20.233	14.5(0.0)	A.10b	13
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{61}{2}\rightarrow\frac{59}{2}$ b	35786.68	35786.68(0)	26.9	60.80	394	20.387	14.5(0.0)	A.10b	13
MgNC	N_J=3$_{\frac{5}{2}}\rightarrow$2$_{\frac{3}{2}}$	35793.32	35794.09(0)	3.4	2.50	1292	25.173	14.5(0.0)	A.10b	1
$Si^{33}S$	J=2→1	35796.67	35796.70(0)	2.6	2.00	618	26.894	14.5(0.0)	A.10b	1
C_6H^-	J=13→12	35798.15	35798.20(0)	12.0	13.00	1460	22.776	14.5(0.0)	A.10b	2
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{15}{2}\rightarrow\frac{13}{2}$ a	35802.89	35802.82(11)	7.2	14.90	2526	24.226	14.2(0.2)	A.10b	2
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{15}{2}\rightarrow\frac{13}{2}$ b	35806.94	35806.88(0)	7.2	14.90	3311	24.496	14.4(0.0)	A.10b	3
MgNC	N_J=3$_{\frac{7}{2}}\rightarrow$2$_{\frac{5}{2}}$	35808.53	35808.63(2)	3.4	3.50	1235	24.790	15.7(0.2)	A.10b	2
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{41}{2}\rightarrow\frac{39}{2}$ a	35852.78	35852.78(0)	18.5	40.90	280	20.619	15.7(0.0)	A.10b	6
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{61}{2}\rightarrow\frac{59}{2}$ a	35853.82	35853.82(0)	54.9	61.00	168	19.530	15.7(0.0)	A.10b	14
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{41}{2}\rightarrow\frac{39}{2}$ b	35854.42	35854.22(0)	18.5	40.90	280	20.619	15.7(0.0)	A.10b	6
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{61}{2}\rightarrow\frac{59}{2}$ b	35863.57	35863.67(0)	54.9	61.00	224	19.816	14.5(0.0)	A.10b	15
o-H_2C_4	$J_{Ka,Kc}$=4$_{1,3}\rightarrow$3$_{1,2}$	35875.78	35875.73(3)	3.9	3.75	782	24.612	13.6(0.3)	A.10b	3
MgC_3N	N_J=13$_{\frac{25}{2}}\rightarrow$12$_{\frac{23}{2}}$	35900.24	35900.21(1)	12.1	12.50	501	22.287	13.9(0.1)	A.10b	3
MgC_3N	N_J=13$_{\frac{27}{2}}\rightarrow$12$_{\frac{25}{2}}$	35904.56	35904.56(3)	12.1	13.50	557	22.315	13.7(0.3)	A.10b	4
MgC_4H	N_J=13$_{\frac{25}{2}}\rightarrow$12$_{\frac{23}{2}}$	35916.32	35916.32(0)	12.1	12.50	56	22.269	14.5(0.0)	A.10b	3
MgC_4H	N_J=13$_{\frac{27}{2}}\rightarrow$12$_{\frac{25}{2}}$	35921.00	35921.00(0)	12.1	13.50	56	22.192	14.5(0.0)	A.10b	4
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{41}{2}\rightarrow\frac{39}{2}$	35935.32	35935.27(2)	56.1	81.50	445	20.391	13.2(0.2)	A.10b	7
MgC_6H	N_J=31$_{\frac{61}{2}}\rightarrow$30$_{\frac{59}{2}}$	35943.31	35943.32(31)	27.4	30.50	111	21.719	14.5(0.0)	A.10b	9
MgC_6H	N_J=31$_{\frac{63}{2}}\rightarrow$30$_{\frac{61}{2}}$	35945.27	35945.15(9)	27.4	31.50	167	22.092	14.5(0.0)	A.10b	10
$HCCCCC^{13}CCN$	J=32→31	35956.41	35956.49(8)	28.5	32.00	222	21.050	13.9(0.8)	A.11a	3
HC_9N	J=62→61	36023.45	36023.44(1)	54.5	62.00	938	21.676	14.1(0.1)	A.11a	25
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{15}{2}\rightarrow\frac{13}{2}$ a	36048.22	36048.21(0)	41.9	13.40	1487	23.795	14.5(0.0)	A.11a	4
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{15}{2}\rightarrow\frac{13}{2}$ b	36048.67	36048.67(0)	41.9	15.30	1432	23.625	14.5(0.0)	A.11a	4
HC_9N	v_{19}^e J=62→61	36059.76	36059.57(4)	124.9	61.00	330	20.647	13.4(0.4)	A.11a	26
$HCCC^{13}CCCN$	J=32→31	36075.17	36074.37(10)	28.6	32.00	165	20.750	11.1(1.2)	A.11a	4

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
HC_9N	ν_{19}^{f} J=62→61	36077.91	36078.01(4)	124.9	61.00	495	21.051	15.2(0.4)	A.11a	27
HC_7N	J=32→31	36095.54	36095.49(0)	28.6	32.00	8342	24.673	13.5(0.1)	A.11a	17
C_5N^-	J=13→12	36110.12	36110.18(2)	12.1	13.00	603	22.794	13.9(0.2)	A.11a	1
HC_7N	ν_{15}^{e} J=32→31	36140.68	36140.68(1)	117.8	32.00	711	22.209	14.0(0.2)	A.11a	18
HC_7N	ν_{15}^{f} J=32→31	36163.68	36163.72(4)	117.8	32.00	710	22.207	14.1(0.2)	A.11a	19
C_8H^-	J=31→30	36166.58	36166.82(2)	27.8	31.00	327	19.927	11.3(0.4)	A.11a	3
HC_7N	ν_{15}=2 J=32→31 (l=0)	36206.92	36206.92(0)	207.3	32.00	163	20.736	14.5(0.0)	A.11b	20
HC_7N	ν_{15}=2 J=32→31 (l=-2)	36208.88	36208.88(0)	207.6	31.90	163	20.739	14.5(0.0)	A.11b	21
HC_7N	ν_{15}=2 J=32→31 (l=+2)	36211.01	36211.01(0)	207.6	31.90	163	20.739	14.5(0.0)	A.11b	22
$HC^{13}CCN$	J=4→3	36237.95	36237.95(0)	4.3	4.00	4071	26.543	14.5(0.0)	A.11b	1
$HCC^{13}CN$	J=4→3	36241.44	36241.44(0)	4.3	4.00	4504	26.644	14.5(0.0)	A.11b	1
$^{29}SiC_4$	J=12→11	36245.16	36245.16(0)	11.3	12.00	217	21.427	14.5(0.0)	A.11b	2
C_6H	ν_{11} N_J=13 $_{\frac{27}{2}}$→12 $_{\frac{25}{2}}$	36251.52	36251.49(1)	28.0	13.50	868	22.990	13.9(0.1)	A.11b	15
C_6H	ν_{11} N_J=13 $_{\frac{25}{2}}$→12 $_{\frac{23}{2}}$	36268.91	36268.87(0)	28.0	12.50	758	22.931	13.8(0.1)	A.11b	16
$H^{13}CCCCCN$	J=14→13	36306.62	36306.95(0)	13.1	14.00	1404	23.926	16.9(0.1)	A.11b	3
SiS	J=2→1	36309.63	36309.62(0)	2.6	2.00	32067	30.829	14.1(0.0)	A.11b	1
HC_3N	J_F=4$_4$→3$_4$	36390.89	36390.89(0)	4.4	0.25	3273	29.093	14.5(0.0)	A.11b	1
HC_3N	J=4→3	36392.33	36392.33(0)	4.4	4.00	129501	29.998	14.5(0.0)	A.11b	1
HC_3N	J_F=4$_3$→3$_3$	36394.18	36394.18(0)	4.4	0.25	2843	28.952	14.5(0.0)	A.11b	2
$H^{13}CCCCCCCN$	J=33→32	36407.92	36407.88(7)	29.7	33.00	161	20.684	11.0(0.6)	A.11b	4
HC_3N	ν_7^e J=4→3	36481.82	36482.06(8)	324.4	3.75	107	22.957	11.5(0.7)	A.12a	3
HC_3N	ν_7^f J=4→3	36534.12	36533.92(23)	324.4	3.75	265	23.868	14.2(1.9)	A.12a	4
HC_9N	J=63→62	36604.45	36604.43(23)	56.2	63.00	845	21.539	14.0(1.9)	A.12a	28
HC_9N	ν_{19}^{e} J=63→62	36641.35	36641.46(11)	128.4	63.00	526	21.066	21.5(1.5)	A.12a	29
HC_9N	ν_{19}^{f} J=63→62	36659.78	36659.89(8)	128.4	63.00	263	20.371	14.5(0.6)	A.12a	30
CH_3CN	$J_{K,F}$=2$_{0,2}$ →1$_{0,2}$+2$_{0,1}$ →1$_{0,0}$	36794.33	36794.33(0)	2.6	1.17	937	26.186	14.5(0.0)	A.12b	1
CH_3CN	$J_{K,F}$=2$_{0,2}$ →1$_{0,1}$+2$_{0,3}$ →1$_{0,2}$	36795.54	36795.54(0)	2.6	4.30	1613	25.428	14.5(0.0)	A.12b	1
CH_3CN	$J_{K,F}$=2$_{0,1}$ →1$_{0,1}$	36797.58	36797.58(0)	2.6	0.50	208	25.532	14.5(0.0)	A.12b	2
SiC_4	J=12→11	36810.14	36810.12(1)	11.5	12.00	1559	23.384	14.0(0.1)	A.12b	2
l-HC_4N	N_J=8$_8$→7$_7$	36838.13	36838.07(7)	8.9	7.87	311	22.981	15.9(0.7)	A.12b	2
MgC_5N	N_J=32 $_{\frac{63}{2}}$→31 $_{\frac{61}{2}}$	36889.72	36889.68(6)	29.2	31.50	155	19.851	14.5(0.0)	A.12b	9
MgC_5N	N_J=32 $_{\frac{65}{2}}$→31 $_{\frac{63}{2}}$	36891.53	36891.22(0)	29.2	32.50	155	19.820	14.5(0.0)	A.12b	10
$HC^{13}CCCCN$	J=14→13	36894.98	36894.94(2)	13.3	14.00	826	23.380	13.6(0.2)	A.12b	3
$HCCCC^{13}CN$	J=14→13	36908.73	36908.67(1)	13.3	14.00	1032	23.602	13.6(0.1)	A.12b	3
l-HC_4N	N_J=8$_9$→7$_8$	36943.21	36943.08(6)	8.3	8.96	154	22.147	14.0(0.7)	A.12b	3
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{63}{2}$ → $\frac{61}{2}$ a	36959.45	36959.45(0)	28.7	62.80	360	20.233	14.5(0.0)	A.13a	16
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{63}{2}$ → $\frac{61}{2}$ b	36959.99	36959.99(0)	28.7	62.80	360	20.233	14.5(0.0)	A.13a	16
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{63}{2}$ → $\frac{61}{2}$ a	37029.49	37029.49(0)	56.7	63.00	102	18.970	14.5(0.0)	A.13a	17
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{63}{2}$ → $\frac{61}{2}$ b	37039.20	37039.20(0)	56.7	63.00	256	19.886	14.5(0.0)	A.13a	18
MgC_6H	N_J=32 $_{\frac{63}{2}}$→31 $_{\frac{61}{2}}$	37102.79	37102.79(0)	29.2	31.50	102	21.567	14.5(0.0)	A.13a	11
MgC_6H	N_J=32 $_{\frac{65}{2}}$→31 $_{\frac{63}{2}}$	37104.74	37104.61(4)	29.2	32.50	305	22.634	14.5(0.0)	A.13a	12
SiC_2	$J_{Ka,Kc}$=9$_{2,7}$→9$_{2,8}$	37108.78	37108.68(1)	60.2	0.73	813	27.502	14.5(0.0)	A.13a	1
CH_3C_3N	N_J=9$_0$→8$_0$	37183.02	37182.86(23)	8.9	9.00	152	21.934	12.8(1.8)	A.13a	3
HC_9N	J=64→63	37185.44	37185.47(23)	58.0	64.00	708	21.331	13.7(1.8)	A.13a	31
HC_7N	J=33→32	37223.49	37223.49(0)	30.4	33.00	8873	24.673	13.9(0.0)	A.13b	23
$HCC^{13}CCN$	J=14→13	37238.38	37238.50(1)	13.4	14.00	1209	23.751	15.0(0.1)	A.13b	3
$HCCC^{13}CN$	J=14→13	37242.91	37242.79(1)	13.4	14.00	1208	23.750	14.1(0.1)	A.13b	3
HC_7N	ν_{15}^{e} J=33→32	37270.04	37269.80(20)	119.6	33.00	402	21.577	11.8(0.2)	A.13b	24
HC_5N	J=14→13	37276.99	37276.98(0)	13.4	14.00	41028	27.275	14.1(0.0)	A.13b	7
HC_7N	ν_{15}^{f} J=33→32	37293.76	37293.69(1)	119.6	33.00	401	21.575	13.0(0.2)	A.13b	25
C_8H^-	J=32→31	37333.21	37333.17(23)	29.6	32.00	400	20.064	13.8(1.8)	A.13b	4
HC_5N	ν_{11}^{e} J=14→13	37338.60	37338.45(5)	167.0	13.90	250	22.179	14.5(0.0)	A.13b	8
HNCCC	J=4(3)→3(3)	37346.08	37346.48(6)	4.5	0.25	150	25.149	14.5(0.0)	A.13b	1
HC_5N	ν_{11}^{f} J=14→13	37371.15	37371.29(10)	167.0	13.90	199	21.953	13.4(0.4)	A.13b	9
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$ a	37426.19	37426.15(1)	12.8	26.70	2484	23.328	13.7(0.0)	A.13b	17
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$ b	37431.25	37431.23(5)	12.8	26.70	2483	23.327	13.4(0.0)	A.13b	18
$H^{13}CCCCCCCN$	J=34→33	37511.15	37511.33(43)	31.5	34.00	49	19.444	11.1(1.2)	A.14a	5
C_6H	ν_{11} $^2\Delta_{\frac{5}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$	37526.68	37526.64(23)	79.4	13.00	296	21.917	13.9(1.8)	A.14a	19
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{43}{2}$ → $\frac{41}{2}$ a	37601.70	37601.70(0)	20.3	43.00	245	20.390	14.5(0.0)	A.14a	8
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{43}{2}$ → $\frac{41}{2}$ b	37603.35	37603.35(0)	20.3	43.00	441	20.978	14.5(0.0)	A.14a	8
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$ a	37678.94	37678.92(0)	35.2	27.00	2146	23.164	14.1(0.0)	A.14a	20
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{43}{2}$ → $\frac{41}{2}$	37688.22	37688.22(0)	57.9	85.50	341	20.031	14.5(0.0)	A.14a	9
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$ b	37710.20	37710.16(0)	35.2	27.00	1947	23.065	13.7(0.0)	A.14b	21
HC_9N	J=65→64	37766.44	37766.43(0)	59.8	65.00	873	21.509	13.9(0.1)	A.14b	32
HC_9N	ν_{19}^{e} J=65→64	37804.50	37804.48(6)	132.0	65.00	242	20.225	14.2(0.5)	A.14b	33
HC_9N	ν_{19}^{f} J=65→64	37823.52	37823.47(4)	132.0	65.00	290	20.405	13.4(0.2)	A.14b	34
C_6H	ν_{11} $^2\Delta_{\frac{3}{2}}$ J=$\frac{27}{2}$ → $\frac{25}{2}$	37840.50	37840.47(3)	96.9	13.30	193	21.458	12.9(0.4)	A.14b	22
SiC_6	J=31→30	37896.90	37896.98(13)	29.1	31.00	96	19.124	14.5(0.0)	A.14b	3
$HC^{13}CCCCCCN$	J=34→33	37897.94	37897.90(7)	31.8	34.00	192	20.792	14.5(0.0)	A.14b	5
$HCCCCCC^{13}CN$	J=34→33	37912.06	37912.03(4)	31.8	34.00	192	20.791	15.4(0.4)	A.14b	3
MgC_5N	N_J=33 $_{\frac{65}{2}}$→32 $_{\frac{63}{2}}$	38042.49	38042.49(5)	31.0	32.50	143	19.705	14.5(0.0)	A.15a	11
MgC_5N	N_J=33 $_{\frac{67}{2}}$→32 $_{\frac{65}{2}}$	38044.30	38044.36(2)	31.0	33.50	285	20.368	14.5(0.0)	A.15a	11
C_4H	$N_{J,F}$=4$_{9/2,4}$-3$_{7/2,3}$+4$_{9/2,5}$-3$_{7/2,4}$	38049.65	38049.61(23)	4.6	8.77	7221	27.432	13.7(1.8)	A.15a	5
C_4H	$N_{J,F}$=4$_{7/2,4}$-3$_{5/2,3}$+4$_{7/2,3}$-3$_{5/2,2}$	38088.46	38088.42(0)	4.6	6.73	5638	27.448	14.1(0.0)	A.15a	6

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{65}{2}\rightarrow\frac{63}{2}$ a	38132.73	38132.73(0)	30.6	64.80	283	19.932	14.5(0.0)	A.15a	19
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{65}{2}\rightarrow\frac{63}{2}$ b	38133.30	38133.30(0)	30.6	64.80	331	20.086	14.5(0.0)	A.15a	19
^{13}CCCN	N=4→3 J=$\frac{9}{2}\rightarrow\frac{7}{2}$ F=5→4	38159.86	38160.04(8)	4.6	3.80	94	23.316	10.7(0.6)	A.15a	1
^{13}CCCN	N=4→3 J=$\frac{9}{2}\rightarrow\frac{7}{2}$ F=4→3	38168.92	38168.92(0)	4.6	3.75	47	22.635	14.5(0.0)	A.15a	2
^{13}CCCN	N=4→3 J=$\frac{7}{2}\rightarrow\frac{5}{2}$ F=4→3	38170.25	38170.25(0)	4.6	3.55	47	22.690	14.5(0.0)	A.15a	2
^{13}CCCN	N=4→3 J=$\frac{7}{2}\rightarrow\frac{5}{2}$ F=3→2	38180.56	38180.56(0)	4.6	3.55	47	22.689	14.5(0.0)	A.15a	3
HCC^{13}CCCCCN	J=34→33	38194.50	38194.64(7)	31.8	34.00	141	20.476	13.5(0.6)	A.15a	3
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{65}{2}\rightarrow\frac{63}{2}$ a	38205.15	38205.15(0)	58.5	65.00	94	18.823	14.5(0.0)	A.15b	20
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{65}{2}\rightarrow\frac{63}{2}$ b	38214.83	38214.83(0)	58.5	65.00	94	18.822	14.5(0.0)	A.15b	21
C_4H	ν_7=2 l=0 N_J=4$_{\frac{9}{2}}\rightarrow$3$_{\frac{7}{2}}$	38228.69	38228.69(1)	540.8	8.77	891	25.335	14.0(0.1)	A.15b	7
Na^{37}Cl	J=3→2	38241.48	38241.48(0)	3.7	3.00	94	21.244	14.5(0.0)	A.15b	1
MgC_6H	N_J=33$_{\frac{65}{2}}\rightarrow$32$_{\frac{63}{2}}$	38262.26	38262.21(9)	31.0	32.50	94	21.422	14.5(0.0)	A.15b	13
MgC_6H	N_J=33$_{\frac{67}{2}}\rightarrow$32$_{\frac{65}{2}}$	38264.21	38264.21(0)	31.0	33.50	140	21.797	14.5(0.0)	A.15b	14
C_4H	ν_7=2 l=0 N_J=4$_{\frac{7}{2}}\rightarrow$3$_{\frac{5}{2}}$	38285.57	38285.51(2)	540.9	6.72	607	25.216	14.0(0.1)	A.15b	8
HCCC^{13}CCCCN	J=34→33	38329.79	38329.79(0)	32.2	34.00	140	20.463	14.5(0.0)	A.15b	5
HC_9N	J=66→65	38347.43	38347.46(3)	61.7	66.00	791	21.381	14.2(0.2)	A.15b	35
HC_7N	J=34→33	38351.44	38351.42(0)	32.2	34.00	7814	24.486	14.1(0.0)	A.15b	26
HC_9N	v_{19}^e J=66→65	38386.08	38385.83(4)	133.8	66.00	186	19.930	15.1(0.4)	A.15b	36
HC_7N	v_{15}^e J=34→33	38399.40	38399.49(4)	121.4	34.00	464	21.660	14.4(0.3)	A.15b	27
HC_9N	v_{19}^f J=66→65	38404.39	38404.55(9)	133.8	66.00	324	20.488	15.1(0.8)	A.15b	37
HC_7N	v_{15}^f J=34→33	38423.83	38423.77(2)	121.4	34.00	602	21.920	14.4(0.2)	A.15b	28
HCCN^{13}C	J=4→3	38438.82	38438.56(17)	6.9	5.00	92	22.941	14.4(1.0)	A.15b	1
KCl	J=5→4	38443.80	38444.02(21)	5.5	5.00	92	20.451	12.3(1.4)	A.15b	1
HC_7N	v_{15}=2 J=34→33 (l=0)	38469.48	38469.48(0)	210.9	34.00	92	20.044	14.5(0.0)	A.16a	29
HC_7N	v_{15}=2 J=34→33 (l=-2)	38471.85	38471.85(0)	211.3	33.90	92	20.047	14.5(0.0)	A.16a	30
HC_7N	v_{15}=2 J=34→33 (l=+2)	38474.41	38474.41(0)	211.3	33.90	92	20.047	14.5(0.0)	A.16a	31
C_8H^-	J=33→32	38499.83	38499.87(23)	31.4	33.00	276	19.632	14.5(1.8)	A.16a	5
C_6H^-	J=14→13	38551.81	38551.77(0)	13.9	14.00	1468	22.633	13.8(0.1)	A.16a	3
H^{13}CCCCCCCN	J=35→34	38614.39	38614.04(12)	32.2	34.00	137	20.436	16.2(0.4)	A.16a	6
MgC_3N	N_J=14$_{\frac{27}{2}}\rightarrow$13$_{\frac{25}{2}}$	38661.85	38661.77(1)	13.9	13.50	592	22.302	13.9(0.1)	A.16a	5
MgC_3N	N_J=14$_{\frac{29}{2}}\rightarrow$13$_{\frac{27}{2}}$	38666.17	38666.19(2)	13.9	14.50	592	22.230	13.6(0.1)	A.16a	6
MgC_4H	N_J=14$_{\frac{27}{2}}\rightarrow$13$_{\frac{25}{2}}$	38679.18	38679.18(0)	13.9	13.50	91	22.610	14.5(0.0)	A.16a	5
MgC_4H	N_J=14$_{\frac{29}{2}}\rightarrow$13$_{\frac{27}{2}}$	38683.86	38683.86(0)	13.9	14.50	45	21.844	14.5(0.0)	A.16a	6
CCS	N_J=3$_3\rightarrow$2$_2$	38866.42	38866.40(1)	12.4	2.67	988	25.996	13.9(0.1)	A.16b	2
C_5N^-	J=14→13	38887.77	38887.69(0)	14.0	14.00	852	22.991	14.2(0.1)	A.16b	2
H^{13}CCCCCN	J=15→14	38899.90	38899.86(0)	14.9	15.00	985	23.434	13.9(0.1)	A.16b	4
HC_9N	J=67→66	38928.42	38928.42(2)	63.5	67.00	760	21.311	13.8(0.1)	A.16b	38
HC_9N	v_{19}^e J=67→66	38967.66	38967.61(1)	135.7	67.00	312	20.420	14.4(0.1)	A.17a	39
HC_9N	v_{19}^f J=67→66	38987.26	38987.22(5)	135.7	67.00	223	20.081	14.2(0.4)	A.17a	40
HC^{13}CCCCCCN	J=35→34	39012.54	39013.10(3)	32.2	34.00	178	20.686	10.4(0.3)	A.17a	6
HCCCCCC^{13}CN	J=35→34	39027.08	39026.79(1)	32.2	34.00	178	20.685	15.3(0.1)	A.17a	4
C_6H	ν_{11} N_J=14$_{\frac{29}{2}}\rightarrow$13$_{\frac{27}{2}}$	39040.82	39040.79(1)	29.9	14.50	843	22.815	13.8(0.1)	A.17a	23
C_6H	ν_{11} N_J=14$_{\frac{27}{2}}\rightarrow$13$_{\frac{25}{2}}$	39057.97	39057.93(0)	29.9	13.50	709	22.713	13.9(0.1)	A.17a	24
NaCl	J=3→2	39077.29	39077.29(0)	3.8	3.00	443	22.775	14.5(0.0)	A.17a	1
MgC_5N	N_J=34$_{\frac{67}{2}}\rightarrow$33$_{\frac{65}{2}}$	39195.26	39195.18(5)	32.9	33.50	220	20.077	14.5(0.0)	A.17a	12
MgC_5N	N_J=34$_{\frac{69}{2}}\rightarrow$33$_{\frac{67}{2}}$	39197.06	39197.00(3)	32.9	34.50	307	20.384	14.5(0.0)	A.17a	12
$^{29}SiC_4$	J=13→12	39265.53	39266.31(3)	13.2	13.00	87	20.358	11.8(0.6)	A.17b	3
C_5N	N_J=14$_{\frac{29}{2}}$-13$_{\frac{27}{2}}$	39280.30	39280.40(0)	14.1	14.50	87	21.527	14.5(0.0)	A.17b	1
C_5N	N_J=14$_{\frac{27}{2}}$-13$_{\frac{25}{2}}$	39291.13	39291.13(0)	14.1	13.50	87	21.598	14.5(0.0)	A.17b	2
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{67}{2}\rightarrow\frac{65}{2}$ a	39306.00	39306.00(0)	32.4	66.90	262	19.789	14.5(0.0)	A.17b	22
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{67}{2}\rightarrow\frac{65}{2}$ b	39306.60	39306.60(0)	32.4	66.90	262	19.789	14.5(0.0)	A.17b	22
HCC^{13}CCCCCN	J=35→34	39317.83	39317.81(0)	32.2	34.00	174	20.658	14.3(0.0)	A.17b	4
HCCCCC^{13}CCN	J=35→34	39327.21	39327.02(9)	32.2	34.00	131	20.369	15.7(1.0)	A.17b	4
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{45}{2}\rightarrow\frac{43}{2}$ a	39350.61	39350.63(3)	22.2	45.00	304	20.515	14.5(0.0)	A.17b	10
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{45}{2}\rightarrow\frac{43}{2}$ b	39352.26	39352.26(0)	22.2	45.00	261	20.360	14.5(0.0)	A.17b	10
C^{13}CCN	N=4→3 J=$\frac{9}{2}\rightarrow\frac{7}{2}$ F=4→3	39362.32	39362.39(8)	4.7	2.85	87	23.491	11.3(0.4)	A.17b	1
C^{13}CCN	N=4→3 J=$\frac{7}{2}\rightarrow\frac{5}{2}$ F=3→2	39375.57	39374.43(6)	4.7	2.53	217	24.525	12.9(0.5)	A.17b	2
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{67}{2}\rightarrow\frac{65}{2}$ a	39380.82	39380.82(0)	60.4	67.00	130	19.088	14.5(0.0)	A.17b	23
C_8H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{67}{2}\rightarrow\frac{65}{2}$ b	39390.46	39390.46(0)	60.4	67.00	130	19.087	14.5(0.0)	A.17b	24
MgC_6H	N_J=34$_{\frac{67}{2}}\rightarrow$33$_{\frac{65}{2}}$	39421.72	39421.72(0)	32.9	33.50	173	21.976	14.5(0.0)	A.17b	15
MgC_6H	N_J=34$_{\frac{69}{2}}\rightarrow$33$_{\frac{67}{2}}$	39423.68	39423.39(23)	32.9	34.50	346	22.640	14.5(0.0)	A.17b	16
MgC_6H	N_J=34$_{\frac{69}{2}}\rightarrow$33$_{\frac{67}{2}}$	39423.68	39423.43(0)	32.9	34.50	130	21.659	14.5(0.0)	A.17b	16
CC^{13}CN	N=4→3 J=$\frac{9}{2}\rightarrow\frac{7}{2}$ F=4→3	39423.69	39423.69(0)	4.7	3.68	130	23.635	14.5(0.0)	A.17b	1
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{45}{2}\rightarrow\frac{43}{2}$	39441.11	39441.02(2)	59.8	89.60	475	20.270	14.5(0.0)	A.17b	11
CC^{13}CN	N=4→3 J=$\frac{7}{2}\rightarrow\frac{5}{2}$ F=3→2	39441.72	39441.93(5)	4.7	1.71	302	25.247	14.5(0.0)	A.17b	2
HCCC^{13}CCCCN	J=35→34	39457.10	39457.10(0)	32.2	34.00	173	20.645	14.5(0.0)	A.18a	6
HCCCC^{13}CCCN	J=35→34	39460.57	39460.37(0)	32.2	34.00	173	20.645	14.5(0.0)	A.18a	2
HC_7N	J=35→34	39479.39	39479.37(0)	34.1	35.00	7885	24.437	13.9(0.0)	A.18a	32
HC_9N	J=68→67	39509.41	39509.40(1)	65.4	68.00	774	21.299	13.9(0.1)	A.18a	41
HC_7N	v_{15}^e J=35→34	39528.75	39528.75(23)	123.3	35.00	258	21.015	12.7(1.7)	A.18a	33
HC^{13}CCCCN	J=15→14	39530.29	39530.22(23)	15.2	15.00	816	23.229	13.7(1.7)	A.18a	4

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
HCCCC^{13}CN	J=15→14	39545.02	39545.01(1)	15.2	15.00	1030	23.462	14.0(0.1)	A.18a	4
HC_9N	v_{19}^e J=68→67	39549.23	39549.18(5)	137.6	68.00	300	20.351	14.1(0.4)	A.18a	42
HC_7N	v_{15}^f J=35→34	39553.90	39553.77(3)	123.3	35.00	429	21.524	13.9(0.2)	A.18a	34
HC_9N	v_{19}^f J=68→67	39569.13	39568.22(8)	137.6	68.00	86	19.097	7.1(0.6)	A.18a	43
C_3N	N=4→3 a	39571.35	39571.33(0)	4.7	12.90	12806	26.969	14.0(0.0)	A.18a	1
C_3N	N=4→3 b	39590.18	39590.15(0)	4.8	9.72	10181	27.022	13.9(0.0)	A.18a	2
HC_7N	v_{15}=2 J=35→34 (l=-2)	39603.33	39603.33(0)	213.2	34.90	85	19.912	14.5(0.0)	A.18a	35
HC_7N	v_{15}=2 J=35→34 (l=+2)	39606.12	39606.12(0)	213.2	34.90	85	19.912	14.5(0.0)	A.18a	36
C_8H^-	J=34→33	39666.46	39666.49(23)	33.3	34.00	383	19.899	14.0(1.7)	A.18a	6
HCCNC	J=4→3	39742.55	39742.54(3)	4.8	4.00	424	24.653	14.2(0.2)	A.18b	1
SiC_4	J=13→12	39877.57	39877.53(1)	13.4	13.00	1637	23.272	14.1(0.1)	A.18b	3
HCC^{13}CCCN	J=15→14	39898.21	39898.20(1)	15.3	15.00	922	23.343	13.8(0.1)	A.18b	4
HCCC^{13}CCN	J=15→14	39903.07	39903.02(1)	15.3	15.00	1006	23.429	13.8(0.1)	A.18b	4
HC_5N	J=15→14	39939.57	39939.55(0)	15.3	15.00	41604	27.151	14.1(0.0)	A.18b	10
HC_5N	v_{11}^e J=15→14	40005.59	40005.60(9)	168.9	14.90	83	20.942	13.1(0.7)	A.19a	11
HC_5N	v_{11}^f J=15→14	40040.46	40040.28(6)	168.9	14.90	125	21.344	12.5(0.5)	A.19a	12
HC_9N	J=69→68	40090.40	40090.40(0)	67.3	69.00	621	21.050	13.8(0.1)	A.19a	44
HC_5N	v_{11}=2 (l=0) J=15→14	40105.99	40105.96(0)	322.7	15.00	41	20.233	14.5(0.0)	A.19a	13
HC_5N	v_{11}=2 (l=-2) J=15→14	40106.52	40106.52(0)	323.7	14.70	41	20.253	14.5(0.0)	A.19a	13
HC_5N	v_{11}=2 (l=+2) J=15→14	40107.44	40107.44(0)	323.7	14.70	41	20.253	14.5(0.0)	A.19a	13
HC_9N	v_{19}^e J=69→68	40130.81	40130.45(10)	139.5	69.00	248	20.130	16.1(0.4)	A.19a	45
HCCCCCC^{13}CN	J=36→35	40142.10	40142.41(9)	15.3	15.00	124	21.114	14.8(0.2)	A.19a	5
HC_9N	v_{19}^f J=69→68	40151.00	40150.95(3)	139.5	69.00	206	19.946	13.7(0.2)	A.19a	46
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$ a	40198.32	40198.29(0)	14.7	28.70	2714	23.273	13.8(0.0)	A.19a	25
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$ b	40204.16	40204.15(0)	14.7	28.70	2672	23.257	14.1(0.0)	A.19b	26
C_6H	v_{11} $^2\Delta_{\frac{5}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$	40306.51	40306.52(3)	81.3	14.10	286	21.730	13.3(0.2)	A.19b	27
MgC_5N	N_J=35 $\frac{69}{2}$ →34 $\frac{67}{2}$	40348.01	40347.94(4)	34.9	34.50	285	20.280	14.5(0.0)	A.19b	13
MgC_5N	N_J=35 $\frac{71}{2}$ →34 $\frac{69}{2}$	40349.82	40349.69(4)	34.9	35.50	285	20.251	14.5(0.0)	A.19b	13
HCC^{13}CCCCCN	J=36→35	40441.15	40441.15(5)	35.9	36.00	162	20.500	13.9(0.4)	A.19b	5
HCCCCC^{13}CCN	J=36→35	40450.80	40450.77(4)	35.9	36.00	162	20.499	13.3(0.3)	A.20a	5
C_3S	J=7→6	40465.01	40465.03(0)	7.8	7.00	1658	24.989	14.2(0.1)	A.20a	2
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$ a	40470.71	40470.68(0)	37.2	29.00	2102	23.000	13.7(0.0)	A.20a	28
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{69}{2}$ → $\frac{67}{2}$ a	40479.26	40479.25(0)	34.4	68.90	283	19.809	14.5(0.0)	A.20a	25
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{69}{2}$ → $\frac{67}{2}$ b	40479.90	40479.90(0)	34.4	68.90	323	19.942	14.5(0.0)	A.20a	25
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$ b	40502.50	40502.46(4)	37.2	29.00	2098	22.997	13.9(0.0)	A.20a	29
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{17}{2}$ → $\frac{15}{2}$ a	40576.81	40576.81(0)	9.2	16.90	2690	24.037	14.5(0.0)	A.20a	5
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{17}{2}$ → $\frac{15}{2}$ b	40580.95	40580.95(0)	9.2	16.90	2730	24.052	14.5(0.0)	A.20a	6
MgC_6H	N_J=35 $\frac{69}{2}$ →34 $\frac{67}{2}$	40581.19	40581.19(0)	34.9	34.50	40	20.457	14.5(0.0)	A.20a	17
MgC_6H	N_J=35 $\frac{71}{2}$ →34 $\frac{69}{2}$	40583.15	40583.15(0)	34.9	35.50	241	22.220	14.5(0.0)	A.20a	17
HCCC^{13}CCCCN	J=36→35	40584.40	40584.40(0)	36.0	36.00	120	20.199	14.5(0.0)	A.20a	7
HCCCC^{13}CCCN	J=36→35	40587.97	40587.97(0)	36.0	36.00	120	20.199	14.5(0.0)	A.20a	3
HC_7N	J=36→35	40607.33	40607.31(0)	36.1	36.00	7173	24.286	13.8(0.0)	A.20a	37
C_6H	v_{11} $^2\Delta_{\frac{3}{2}}$ J=$\frac{29}{2}$ → $\frac{27}{2}$	40643.28	40643.30(5)	98.9	14.30	160	21.127	13.5(0.4)	A.20a	30
HC_7N	v_{15}^e J=36→35	40658.11	40658.13(2)	125.3	36.00	479	21.579	13.5(0.2)	A.20a	38
HC_9N	J=70→69	40671.39	40671.39(1)	69.3	70.00	639	21.049	14.5(0.1)	A.20a	47
HC_7N	v_{15}^f J=36→35	40683.97	40683.94(4)	125.3	36.00	479	21.577	14.0(0.2)	A.20a	39
HC_9N	v_{19}^e J=70→69	40712.38	40712.28(3)	141.5	70.00	279	20.219	14.5(0.0)	A.20b	48
HC_7N	v_{15}=2 J=36→35 (l=0)	40731.97	40731.90(0)	214.7	36.00	80	19.781	14.5(0.0)	A.20b	40
HC_9N	v_{19}^f J=70→69	40732.86	40732.80(0)	141.5	70.00	199	19.881	14.5(0.0)	A.20b	49
HC_7N	v_{15}=2 J=36→35 (l=-2)	40734.81	40734.80(0)	215.1	35.90	80	19.784	14.5(0.0)	A.20b	41
HC_7N	v_{15}=2 J=36→35 (l=+2)	40737.84	40737.80(0)	215.1	35.90	79	19.783	14.5(0.0)	A.20b	42
MgCN	N_J=4 $\frac{7}{2}$ →3 $\frac{5}{2}$	40750.24	40750.24(0)	4.9	3.50	79	21.978	14.5(0.0)	A.20b	1
MgCN	N_J=4 $\frac{9}{2}$ →3 $\frac{7}{2}$	40765.25	40765.25(0)	4.9	3.50	119	22.382	14.5(0.0)	A.20b	2
HCN	v_2=1 J=13→13 l=1→-1	40766.88	40766.88(0)	1411.2	0.08	159	27.578	12.4(0.6)	A.20b	2
H^{13}CCCCCCCN	J=37→36	40820.83	40820.93(1)	37.2	37.00	119	20.152	13.9(0.3)	A.20b	7
C_8H^-	J=35→34	40833.07	40833.05(4)	35.3	35.00	277	19.515	13.9(0.3)	A.20b	7
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{17}{2}$ → $\frac{15}{2}$ a	40854.57	40854.76(1)	43.8	16.40	1539	23.502	15.0(0.0)	A.20b	7
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{17}{2}$ → $\frac{15}{2}$ b	40855.01	40854.83(1)	43.8	16.40	1539	23.502	15.0(0.0)	A.20b	7
C_4H	v_7 N_J=4 $\frac{9}{2}$ → 3 $\frac{7}{2}$ a	40924.21	40923.96(2)	249.0	7.92	1022	25.505	14.6(0.2)	A.20b	9
C_4H	v_7 N_J=4 $\frac{9}{2}$ → 3 $\frac{7}{2}$ b	40994.13	40993.77(1)	249.0	7.92	1213	25.675	14.5(0.1)	A.21a	10
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{47}{2}$ → $\frac{45}{2}$ a	41099.52	41099.52(0)	24.1	47.00	233	20.162	14.5(0.0)	A.21a	12
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{47}{2}$ → $\frac{45}{2}$ b	41101.17	41101.17(0)	24.1	47.00	311	20.449	14.5(0.0)	A.21a	12
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{47}{2}$ → $\frac{45}{2}$	41193.99	41193.95(3)	61.8	93.60	271	19.619	13.5(0.2)	A.21a	13
o-H_2C_3	$J_{Ka,Kc}$=$2_{1,2}$→$1_{1,1}$	41198.33	41198.33(0)	2.0	1.50	155	23.936	14.5(0.0)	A.21a	1
HC^{13}CCCCCCN	J=37→36	41241.74	41241.68(4)	37.6	37.00	116	20.115	13.0(0.3)	A.21b	7
HC_9N	J=71→70	41252.37	41252.40(2)	71.3	71.00	578	20.920	14.0(0.1)	A.21b	50
HC_9N	v_{19}^e J=71→70	41294.00	41293.69(6)	143.4	71.00	269	20.155	17.7(0.5)	A.21b	51
C_6H^-	J=15→14	41305.45	41305.43(1)	15.9	15.00	1612	22.589	14.3(0.1)	A.21b	4
HC_9N	v_{19}^f J=71→70	41314.72	41314.46(6)	143.5	71.00	345	20.404	17.5(0.5)	A.21b	52
MgC_3N	N_J=15 $\frac{29}{2}$ →14 $\frac{27}{2}$	41423.43	41423.40(2)	15.9	14.50	572	22.127	14.0(0.1)	A.21b	7
MgC_3N	N_J=15 $\frac{31}{2}$ →14 $\frac{29}{2}$	41427.75	41427.78(2)	15.9	15.50	610	22.124	14.0(0.2)	A.21b	8
MgC_4H	N_J=15 $\frac{29}{2}$ →14 $\frac{27}{2}$	41442.02	41442.02(0)	15.9	14.50	76	22.291	14.5(0.0)	A.21b	7

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	W=$\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
l-HC_4N	N_J=9$_9$→8$_8$	41442.83	41442.83(0)	10.8	8.89	152	22.026	14.5(0.0)	A.21b	4
MgC_4H	N_J=15 $\frac{31}{2}$ →14 $\frac{29}{2}$	41446.70	41446.70(0)	15.9	15.50	76	22.224	14.5(0.0)	A.21b	8
H^{13}CCCCCN	J=16→15	41493.17	41493.12(1)	16.9	16.00	797	23.093	13.4(0.1)	A.22a	5
MgC_5N	N_J=36 $\frac{71}{2}$ →35 $\frac{69}{2}$	41500.77	41500.78(23)	36.8	35.50	266	20.152	14.5(0.0)	A.22a	14
MgC_5N	N_J=36 $\frac{73}{2}$ →35 $\frac{71}{2}$	41502.57	41502.43(23)	36.9	36.50	152	19.564	14.5(0.0)	A.22a	14
l-HC_4N	N_J=9$_{10}$→8$_9$	41527.91	41527.99(5)	10.2	9.96	114	21.618	13.0(0.7)	A.22a	5
SiC_6	J=34→33	41564.20	41564.20(0)	34.9	34.00	132	19.259	14.5(0.0)	A.22a	4
HCC^{13}CCCCCN	J=37→36	41564.47	41564.46(0)	37.9	37.00	132	20.242	14.5(0.0)	A.22a	6
HCCCCC^{13}CCN	J=37→36	41574.39	41574.39(0)	37.9	37.00	76	19.682	14.5(0.0)	A.22a	6
p-H_2C_3	$J_{Ka,Kc}$=2$_{0,2}$→1$_{0,1}$	41584.68	41584.60(12)	3.0	2.00	75	22.922	12.9(0.8)	A.23a	1
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{71}{2}$ → $\frac{69}{2}$ a	41652.52	41652.51(3)	36.4	70.90	263	19.680	15.2(0.2)	A.22a	26
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{71}{2}$ → $\frac{69}{2}$ a	41652.52	41652.53(3)	36.4	70.90	263	19.680	15.2(0.2)	A.22a	26
C_5N^-	J=15→14	41665.41	41665.51(2)	16.0	15.00	563	22.440	14.1(0.1)	A.22a	3
HCCCC^{13}CCCN	J=37→36	41715.37	41715.67(5)	38.0	37.00	112	20.075	11.3(0.8)	A.22b	4
HC_7N	J=37→36	41735.26	41735.27(23)	38.1	37.00	7257	24.243	14.5(0.0)	A.22b	43
MgC_6H	N_J=36 $\frac{71}{2}$ →35 $\frac{69}{2}$	41740.66	41740.10(23)	36.8	35.50	150	21.715	14.5(0.0)	A.22b	18
MgC_6H	N_J=36 $\frac{73}{2}$ →35 $\frac{71}{2}$	41742.61	41742.61(0)	36.9	36.50	112	21.400	14.5(0.0)	A.22b	18
HC_7N	ν_{15}^e J=37→36	41787.45	41787.41(3)	127.3	37.00	410	21.369	13.9(0.3)	A.22b	44
Si^{17}O	J=1→0	41794.66	41794.66(0)	2.0	1.00	112	24.564	14.5(0.0)	A.22b	1
HC_7N	ν_{15}^f J=37→36	41814.03	41814.03(3)	127.3	37.00	559	21.677	14.7(0.2)	A.22b	45
C_6H	ν_{11} N_J=15 $\frac{31}{2}$ →14 $\frac{29}{2}$	41830.10	41829.80(2)	31.9	15.50	707	22.503	12.4(0.2)	A.22b	31
HC_9N	J=72→71	41833.35	41833.41(0)	73.3	72.00	669	21.040	13.5(0.0)	A.22b	53
C_6H	ν_{11} N_J=15 $\frac{29}{2}$ →14 $\frac{27}{2}$	41847.01	41846.98(2)	31.9	14.50	706	22.569	13.7(0.1)	A.22b	32
HC_7N	ν_{15}=2 J=37→36 (l=0)	41863.19	41863.19(0)	216.8	37.00	74	19.658	14.5(0.0)	A.22b	46
HC_7N	ν_{15}=2 J=37→36 (l=-2)	41866.28	41866.28(0)	217.1	36.90	74	19.660	14.5(0.0)	A.22b	47
HC_7N	ν_{15}=2 J=37→36 (l=+2)	41869.57	41869.57(0)	217.1	36.90	74	19.660	14.5(0.0)	A.22b	48
HC_9N	ν_{19}^e J=72→71	41875.52	41875.56(5)	145.4	72.00	148	19.532	13.9(0.4)	A.22b	54
HC_9N	ν_{19}^f J=72→71	41896.58	41896.53(4)	145.5	72.00	148	19.531	13.8(0.3)	A.22b	55
H^{13}CCCCCCCN	J=38→37	41924.05	41924.05(6)	39.2	38.00	185	20.542	13.9(0.4)	A.22b	8
o-H_2C_3	$J_{Ka,Kc}$=2$_{1,1}$→1$_{1,0}$	41967.67	41967.71(2)	2.0	1.50	221	24.277	14.7(0.6)	A.23a	2
C_8H^-	J=36→35	41999.69	41999.59(3)	37.3	36.00	184	19.052	13.6(0.2)	A.23a	8
C_5N	N_J=15 $\frac{31}{2}$ -14 $\frac{29}{2}$	42086.41	42086.04(14)	16.2	15.50	73	21.216	11.1(1.3)	A.23a	3
C_5N	N_J=15 $\frac{29}{2}$ -14 $\frac{27}{2}$	42097.16	42097.10(6)	16.2	14.50	146	21.975	13.1(0.5)	A.23a	4
c-C_3H_2 (ortho)	$J_{Ka,Kc}$=5$_{4,1}$→5$_{3,2}$	42139.19	42139.06(10)	44.4	4.37	219	23.550	13.4(0.7)	A.23a	1
HC^{13}CCCN	J=16→15	42165.58	42165.52(1)	17.2	16.00	839	23.128	13.9(0.1)	A.23a	5
HCCCC^{13}CN	J=16→15	42181.30	42181.26(1)	17.2	16.00	875	23.169	13.8(0.1)	A.23a	5
c-C_3H_2 (para)	$J_{Ka,Kc}$=4$_{3,1}$→4$_{2,2}$	42231.25	42231.20(5)	30.9	3.25	218	23.839	13.8(0.4)	A.23b	1
$^{29}SiC_4$	J=14→13	42285.88	42285.65(1)	15.2	14.00	109	20.427	13.0(0.0)	A.23b	4
HC^{13}CCCCCN	J=38→37	42356.34	42356.38(15)	39.6	38.00	72	19.591	14.4(1.2)	A.23b	8
HCCCCCC^{13}CN	J=38→37	42372.13	42372.15(0)	39.7	38.00	144	20.282	14.5(0.0)	A.23b	6
^{30}SiO	J=1→0	42373.42	42373.44(1)	2.0	1.00	1622	27.224	14.5(0.0)	A.23b	1
HC_9N	J=73→72	42414.34	42414.35(2)	75.3	73.00	503	20.727	13.8(0.1)	A.23b	56
HC_9N	ν_{19}^e J=73→72	42457.08	42457.10(6)	147.5	73.00	215	19.876	14.1(0.2)	A.24a	57
HC_9N	ν_{19}^f J=73→72	42478.44	42478.51(5)	147.5	73.00	143	19.469	13.3(0.4)	A.24a	58
HCC^{13}CCCN	J=16→15	42558.03	42558.00(1)	17.4	16.00	927	23.219	13.7(0.1)	A.24a	5
HCCC^{13}CCN	J=16→15	42563.21	42563.19(2)	17.4	16.00	856	23.138	14.1(0.1)	A.24a	5
HC_5N	J=16→15	42602.15	42602.15(0)	17.4	16.00	37916	26.929	13.9(0.0)	A.24a	14
MgC_5N	N_J=37 $\frac{73}{2}$ →36 $\frac{71}{2}$	42653.51	42653.55(4)	38.9	36.50	177	19.693	14.5(0.0)	A.24a	15
MgC_5N	N_J=37 $\frac{75}{2}$ →36 $\frac{73}{2}$	42655.32	42655.25(3)	38.9	37.50	177	19.666	14.5(0.0)	A.24a	15
HC_5N	ν_{11}^e J=16→15	42672.57	42672.84(2)	170.9	15.90	602	22.791	13.0(0.2)	A.24a	15
HCC^{13}CCCCCN	J=38→37	42687.79	42687.61(2)	40.0	38.00	71	19.564	13.0(0.1)	A.24a	7
HCCCCC^{13}CCN	J=38→37	42697.98	42698.05(20)	40.0	38.00	71	19.563	13.2(2.6)	A.24a	7
HC_5N	ν_{11}^f J=16→15	42709.76	42709.81(10)	171.0	15.90	141	21.341	14.2(0.8)	A.24b	16
SiC_6	J=35→34	42786.62	42786.79(8)	37.0	35.00	141	19.264	12.4(0.6)	A.24b	5
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{73}{2}$ → $\frac{71}{2}$ a	42825.78	42825.78(0)	38.4	72.90	211	19.402	14.5(0.0)	A.24b	27
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{73}{2}$ → $\frac{71}{2}$ b	42826.50	42826.50(0)	38.4	72.90	211	19.402	14.5(0.0)	A.24b	27
HCCC^{13}CCCCN	J=38→37	42839.00	42839.45(23)	40.0	38.00	140	20.245	13.0(1.6)	A.24b	8
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{49}{2}$ → $\frac{47}{2}$ a	42848.42	42848.39(23)	26.2	49.00	210	19.976	13.9(1.6)	A.24b	14
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{49}{2}$ → $\frac{47}{2}$ b	42850.08	42850.56(23)	26.2	49.00	386	20.582	17.7(1.6)	A.24b	14
HC_7N	J=38→37	42863.20	42863.16(0)	40.1	38.00	6098	24.016	13.8(0.0)	A.24b	49
^{29}SiO	J=1→0	42879.95	42879.96(1)	2.1	1.00	2241	27.536	14.1(0.1)	A.24b	1
MgC_6H	N_J=37 $\frac{73}{2}$ →36 $\frac{71}{2}$	42900.12	42900.43(9)	38.9	36.50	210	21.999	14.5(0.0)	A.24b	19
MgC_6H	N_J=37 $\frac{75}{2}$ →36 $\frac{73}{2}$	42902.07	42901.47(24)	38.9	37.50	70	20.873	14.5(0.0)	A.24b	19
HC_7N	ν_{15}^e J=38→37	42916.79	42916.86(3)	129.3	38.00	559	21.625	13.6(0.2)	A.24b	50
HC_7N	ν_{15}^f J=38→37	42944.09	42944.24(5)	129.3	38.00	454	21.415	14.5(0.0)	A.24b	51
SiC_4	J=14→13	42944.99	42944.98(1)	15.5	14.00	1500	23.037	14.5(0.0)	A.24b	4
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{49}{2}$ → $\frac{47}{2}$	42946.87	42946.94(6)	63.8	97.60	279	19.567	14.5(0.0)	A.24b	15
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$ a	42970.43	42970.44(2)	16.8	30.70	2334	22.988	13.9(0.0)	A.25a	33
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$ b	42977.09	42977.04(1)	16.8	30.70	2472	23.045	14.1(0.0)	A.25a	34
HC_7N	ν_{15}=2 J=38→37 (l=0)	42994.40	42994.00(0)	218.8	38.00	209	20.638	14.5(0.0)	A.25a	52
HC_9N	J=74→73	42995.32	42995.40(0)	77.4	74.00	452	20.593	14.5(0.0)	A.25a	59
HC_7N	ν_{15}=2 J=38→37 (l=-2)	42997.75	42997.54(0)	219.2	37.90	139	20.235	14.5(0.0)	A.25a	53
HC_7N	ν_{15}=2 J=38→37 (l=+2)	43001.31	43001.10(0)	219.2	37.90	35	18.848	14.5(0.0)	A.25a	54
H^{13}CCCCCCCN	J=39→38	43027.27	43027.73(0)	41.3	39.00	139	20.204	14.7(0.0)	A.25a	9

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
HC_9N	v_{19}^e J=74→73	43038.65	43038.65(5)	149.5	74.00	139	19.411	13.2(0.3)	A.25a	60
HC_9N	v_{19}^f J=74→73	43060.30	43060.30(0)	149.6	74.00	139	19.409	14.5(0.0)	A.25a	61
C_6H	ν_{11} $^2\Delta_{\frac{5}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$	43086.35	43085.94(10)	83.4	15.10	381	21.881	16.6(0.8)	A.25a	35
C_8H^-	J=37→36	43166.30	43166.15(4)	39.4	37.00	207	19.113	13.3(0.3)	A.25a	9
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$ a	43262.44	43262.43(1)	39.2	31.00	1851	22.739	14.0(0.0)	A.25b	36
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$ b	43294.79	43294.76(1)	39.3	31.00	1813	22.718	14.0(0.0)	A.25b	37
$Si^{13}CC$	$J_{Ka,Kb}=2_{1,2}\rightarrow1_{1,1}$	43413.98	43414.29(5)	5.0	1.50	136	24.832	11.5(0.4)	A.25b	1
SiO	J=1→0	43423.85	43423.84(0)	2.1	1.00	35364	30.282	14.0(0.0)	A.25b	1
C_6H	ν_{11} $^2\Delta_{\frac{3}{2}}$ J=$\frac{31}{2}$ → $\frac{29}{2}$	43446.00	43446.17(6)	101.0	15.40	271	21.515	14.9(0.3)	A.25b	38
$HCCCCCC^{13}CN$	J=39→38	43487.14	43487.10(7)	41.3	39.00	203	20.573	14.6(0.4)	A.26a	7
HC_9N	J=75→74	43576.30	43576.37(5)	79.5	75.00	472	20.608	13.8(0.2)	A.26a	62
HC_9N	v_{19}^e J=75→74	43620.21	43620.21(0)	151.6	75.00	235	19.912	14.5(0.0)	A.26a	63
HCCN	$N_J=2_3\rightarrow1_2$	43624.20	43624.20(0)	3.1	2.84	504	24.682	14.5(0.0)	A.26a	1
HC_9N	v_{19}^f J=75→74	43642.16	43642.30(6)	151.7	75.00	134	19.350	13.3(0.4)	A.26a	64
AlCl	J=3→2	43731.80	43731.78(7)	4.2	0.93	234	26.777	14.9(0.5)	A.26b	1
MgC_5N	$N_J=38_{\frac{75}{2}}\rightarrow37_{\frac{73}{2}}$	43806.25	43806.45(3)	41.0	37.50	100	19.065	14.5(0.0)	A.26b	16
MgC_5N	$N_J=38_{\frac{77}{2}}\rightarrow37_{\frac{75}{2}}$	43808.06	43807.98(5)	41.0	38.50	200	19.731	14.5(0.0)	A.26b	16
$HCC^{13}CCCCCN$	J=39→38	43811.10	43810.74(6)	41.3	39.00	166	20.366	14.5(0.0)	A.26b	8
$HCCCCC^{13}CCN$	J=39→38	43821.56	43821.56(0)	41.3	39.00	100	19.854	14.1(0.0)	A.26b	8
$HCCC^{13}CCCCN$	J=39→38	43966.29	43966.29(0)	41.3	39.00	99	19.843	14.5(0.0)	A.27a	9
CCS	$N_J=4_3\rightarrow3_2$	43981.02	43981.06(0)	12.9	2.67	989	25.874	14.5(0.1)	A.27a	3
HC_7N	J=39→38	43991.13	43991.15(0)	42.2	39.00	5535	23.867	14.0(0.0)	A.27a	55
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{75}{2}$ → $\frac{73}{2}$ a	43999.03	43999.03(0)	40.5	74.90	231	19.438	14.5(0.0)	A.27a	28
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{75}{2}$ → $\frac{73}{2}$ b	43999.79	43999.79(0)	40.5	74.90	296	19.689	14.5(0.0)	A.27a	28
HC_7N	v_{15}^e J=39→38	44046.13	44046.17(4)	131.4	39.00	361	21.137	13.8(0.3)	A.27a	56
C_6H^-	J=16→15	44059.08	44059.09(0)	18.0	16.00	1576	22.437	14.5(0.0)	A.27a	5
C_6H^-	J=16→15	44059.08	44059.14(23)	18.0	16.00	98	19.665	14.5(0.0)	A.27a	5
MgC_6H	$N_J=38_{\frac{77}{2}}\rightarrow37_{\frac{75}{2}}$	44061.53	44061.53(0)	41.0	38.50	66	20.757	14.5(0.0)	A.27a	20
HC_7N	v_{15}^f J=39→38	44074.15	44074.21(3)	131.5	39.00	492	21.445	14.6(0.2)	A.27a	57
$H^{13}CCCN$	J=5→4	44084.16	44084.16(0)	6.3	5.00	4557	26.236	14.5(0.0)	A.27a	2
$H^{13}CCCCCN$	J=17→16	44086.43	44086.43(0)	19.0	17.00	656	22.776	14.5(0.0)	A.27a	6
c-C_3H_2 (ortho)	$J_{Ka,Kc}=3_{2,1}\rightarrow3_{1,2}$	44104.78	44104.76(1)	15.8	2.09	1998	26.452	13.9(0.0)	A.27a	2
HC_7N	v_{15}=2 J=39→38 (l=0)	44125.58	44125.50(0)	220.9	39.00	33	18.733	14.5(0.0)	A.27a	58
HC_7N	v_{15}=2 J=39→38 (l=-2)	44129.21	44129.20(0)	221.3	38.90	33	18.735	14.5(0.0)	A.27a	59
$H^{13}CCCCCCCN$	J=40→39	44130.48	44130.50(0)	41.3	39.00	131	20.119	14.5(0.0)	A.27a	10
HC_7N	v_{15}=2 J=39→38 (l=+2)	44133.06	44133.20(0)	221.3	38.90	33	18.735	14.5(0.0)	A.27a	60
$H^{13}CCCCCCCN$	J=40→39	44130.48	44130.53(4)	41.3	39.00	131	20.119	14.2(0.4)	A.27a	10
HC_9N	J=76→75	44157.28	44157.26(3)	81.6	76.00	327	20.214	13.4(0.2)	A.27a	65
MgC_3N	$N_J=16_{\frac{31}{2}}\rightarrow15_{\frac{29}{2}}$	44184.99	44185.02(3)	18.0	15.50	424	21.696	14.0(0.2)	A.27a	9
MgC_3N	$N_J=16_{\frac{33}{2}}\rightarrow15_{\frac{31}{2}}$	44189.30	44189.27(3)	18.0	16.50	456	21.708	13.9(0.2)	A.27a	10
HC_9N	v_{19}^e J=76→75	44201.78	44201.78(0)	153.8	76.00	163	19.517	14.5(0.0)	A.27b	66
MgC_4H	$N_J=16_{\frac{31}{2}}\rightarrow15_{\frac{29}{2}}$	44204.83	44204.83(0)	18.0	15.50	33	21.311	14.5(0.0)	A.27b	9
MgC_4H	$N_J=16_{\frac{33}{2}}\rightarrow15_{\frac{31}{2}}$	44209.51	44209.51(0)	18.0	16.50	65	21.941	14.5(0.0)	A.27b	10
HC_9N	v_{19}^f J=76→75	44224.01	44224.54(7)	153.8	76.00	260	19.986	16.0(0.5)	A.27b	67
C_8H^-	J=38→37	44332.91	44333.02(2)	41.5	38.00	291	19.402	15.2(0.2)	A.27b	10
C_5N^-	J=16→15	44443.04	44443.10(23)	18.1	16.00	611	22.392	13.9(1.5)	A.27b	4
o-H_2C_4	$J_{Ka,Kc}=5_{1,5}\rightarrow4_{1,4}$	44471.14	44471.13(23)	6.0	4.80	996	24.392	14.3(1.5)	A.28a	4
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{51}{2}$ → $\frac{49}{2}$ a	44597.31	44597.31(0)	28.3	51.00	191	19.801	14.5(0.0)	A.28a	16
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{51}{2}$ → $\frac{49}{2}$ b	44598.98	44598.98(0)	28.3	51.00	191	19.801	14.5(0.0)	A.28a	16
C_6H	ν_{11} $N_J=16_{\frac{33}{2}}\rightarrow15_{\frac{31}{2}}$	44619.38	44619.36(1)	34.0	16.50	638	22.273	13.8(0.1)	A.28a	39
C_6H	ν_{11} $N_J=16_{\frac{31}{2}}\rightarrow15_{\frac{29}{2}}$	44636.02	44635.98(1)	34.0	15.50	701	22.430	13.8(0.1)	A.28a	40
o-H_2C_4	$J_{Ka,Kc}=5_{3,3}\rightarrow4_{3,2}+5_{3,2}\rightarrow4_{3,1}$	44645.59	44645.47(2)	114.1	3.20	700	24.442	13.6(0.1)	A.28a	5
p-H_2C_4	$J_{Ka,Kc}=5_{2,4}\rightarrow4_{2,3}$	44653.53	44653.62(3)	60.5	4.20	382	23.563	14.1(0.2)	A.28a	3
p-H_2C_4	$J_{Ka,Kc}=5_{0,5}\rightarrow4_{0,4}$	44659.01	44659.00(3)	6.4	5.00	286	23.101	14.0(0.2)	A.28a	4
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{51}{2}$ → $\frac{49}{2}$	44699.74	44699.68(4)	66.0	102.00	254	19.388	13.6(0.2)	A.28a	17
C_4H	v_7 $N_J=5_{\frac{9}{2}}\rightarrow4_{\frac{7}{2}}$ a	44712.64	44712.49(2)	254.3	8.73	888	25.179	15.4(0.1)	A.28b	11
HC_9N	J=77→76	44738.25	44738.28(3)	83.7	77.00	253	19.934	13.7(0.2)	A.28b	68
$HC^{13}CCCCN$	J=17→16	44800.86	44800.82(1)	19.4	17.00	758	22.905	13.7(0.1)	A.28b	6
$HCCCC^{13}CN$	J=17→16	44817.56	44817.53(1)	19.4	17.00	883	23.058	14.0(0.1)	A.28b	6
C_4H	v_7 $N_J=5_{\frac{9}{2}}\rightarrow4_{\frac{7}{2}}$ b	44830.60	44830.67(2)	254.3	8.73	757	25.017	13.0(0.1)	A.28b	12
o-H_2C_4	$J_{Ka,Kc}=5_{1,4}\rightarrow4_{1,3}$	44844.58	44844.54(1)	6.0	4.80	945	24.332	13.7(0.1)	A.28b	6
$HCC^{13}CCCCCN$	J=40→39	44934.41	44934.41(0)	41.3	39.00	94	19.771	14.5(0.0)	A.28b	9
MgC_5N	$N_J=39_{\frac{77}{2}}\rightarrow38_{\frac{75}{2}}$	44958.98	44958.88(18)	43.2	38.50	94	18.952	14.5(0.0)	A.29a	17
MgC_5N	$N_J=39_{\frac{79}{2}}\rightarrow38_{\frac{77}{2}}$	44960.79	44960.46(7)	43.2	39.50	157	19.437	14.5(0.0)	A.29a	17
$HCCCC^{13}CCCN$	J=40→39	45097.55	45098.02(14)	41.3	39.00	93	19.759	11.5(1.0)	A.29a	5
HC_7N	J=40→39	45119.06	45119.07(0)	44.4	40.00	4630	23.638	14.1(0.0)	A.29a	61
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{77}{2}$ → $\frac{75}{2}$ a	45172.28	45172.28(0)	42.7	76.90	186	19.170	14.5(0.0)	A.29a	29
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{77}{2}$ → $\frac{75}{2}$ b	45173.08	45173.08(0)	42.7	76.90	186	19.170	14.5(0.0)	A.29a	29
HC_7N	v_{15}^f J=40→39	45204.20	45204.11(4)	133.6	40.00	371	21.112	13.8(0.2)	A.29b	62
$HCC^{13}CCCN$	J=17→16	45217.84	45217.83(1)	19.5	17.00	742	22.875	14.0(0.1)	A.29b	6
$HCCC^{13}CCN$	J=17→16	45223.35	45223.33(1)	19.5	17.00	773	22.915	13.8(0.1)	A.29b	6
SiC_6	J=37→36	45231.46	45231.90(3)	41.2	37.00	154	19.245	12.3(0.6)	A.29b	6
NaCN	$J_{Ka,Kc}=3_{1,3}\rightarrow2_{1,2}$	45259.08	45259.07(0)	6.7	2.67	370	22.600	14.5(0.0)	A.29b	3

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	W=$\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
NaCN	$J_{Ka,Kc}=3_{1,3}\rightarrow2_{1,2}$	45259.08	45259.08(0)	6.7	2.67	370	22.600	14.5(0.0)	A.29b	3
HC_5N	J=17→16	45264.72	45264.71(0)	19.6	17.00	31463	26.621	13.9(0.0)	A.29b	17
$HC^{13}CCN$	J=5→4	45297.33	45297.29(0)	6.5	5.00	4188	26.125	13.7(0.0)	A.29b	2
$HCC^{13}CN$	J=5→4	45301.71	45301.69(0)	6.5	5.00	4865	26.274	14.2(0.0)	A.29b	2
$^{29}SiC_4$	J=15→14	45306.21	45306.21(0)	17.4	15.00	154	20.637	14.5(0.0)	A.29b	5
HC_9N	J=78→77	45319.22	45319.33(4)	85.9	78.00	461	20.508	15.8(0.2)	A.29b	69
HC_5N	v_{11}^e J=17→16	45339.53	45339.31(10)	173.1	16.90	154	21.303	11.9(0.7)	A.29b	18
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{19}{2}\rightarrow\frac{17}{2}$ a	45350.72	45350.92(1)	11.3	18.90	2027	23.531	15.8(0.1)	A.29b	8
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{19}{2}\rightarrow\frac{17}{2}$ b	45354.96	45354.88(2)	11.3	18.90	1658	23.330	14.4(0.1)	A.29b	9
HC_9N	v_{19}^e J=78→77	45364.90	45365.16(7)	158.1	78.00	123	19.183	12.9(0.5)	A.29b	70
CCS	$N_J=3_4\rightarrow2_3$	45379.03	45378.99(0)	5.4	3.97	3374	26.673	13.8(0.0)	A.29b	4
HC_9N	v_{19}^f J=78→77	45387.72	45387.72(0)	158.1	78.00	92	18.893	14.5(0.0)	A.29b	71
CH_3C_3N	$N_J=11_0\rightarrow10_0$	45445.75	45445.50(9)	13.1	11.00	153	21.540	15.7(0.6)	A.29b	4
HC_3N	$J_F=5_5\rightarrow4_5$	45488.84	45488.84(0)	6.5	0.20	2623	28.871	14.5(0.0)	A.30a	5
HC_3N	J=5→4	45490.31	45490.24(0)	6.5	5.00	109408	29.383	13.6(0.0)	A.30a	5
HC_3N	$J_F=5_4\rightarrow4_4$	45492.11	45492.11(0)	6.5	0.20	2257	28.721	14.5(0.0)	A.30a	6
HC_3N	v_7^e J=5→4	45602.17	45602.06(4)	326.6	4.80	303	23.534	13.8(0.3)	A.30a	7
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{19}{2}\rightarrow\frac{17}{2}$ a	45660.82	45660.98(2)	46.0	18.50	1059	22.896	14.4(0.1)	A.30a	10
C_5H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{19}{2}\rightarrow\frac{17}{2}$ b	45661.37	45661.21(2)	46.0	18.50	1059	22.896	14.4(0.1)	A.30a	10
HC_3N	v_7^f J=5→4	45667.55	45667.42(2)	326.6	4.80	242	23.306	13.0(0.4)	A.30a	8
$HCCCCCC^{13}CN$	J=41→40	45717.14	45717.28(10)	46.1	41.00	151	20.176	13.6(0.7)	A.30b	8
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$ a	45742.52	45742.48(23)	19.0	32.70	1898	22.655	13.7(1.5)	A.30b	41
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$ b	45750.05	45750.01(23)	19.0	32.70	1867	22.639	13.7(1.5)	A.30b	42
$^{30}SiC_2$	$J_{Ka,Kc}=2_{0,2}\rightarrow1_{0,1}$	45754.87	45755.22(23)	3.3	2.00	301	25.287	13.6(1.5)	A.30b	2
C_6H	$v_{11}\ ^2\Delta_{\frac{5}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$	45866.22	45866.07(6)	85.6	16.10	359	21.697	13.4(0.2)	A.30b	43
HC_9N	J=79→78	45900.20	45900.08(5)	88.1	79.00	448	20.454	14.2(0.2)	A.30b	72
HC_9N	v_{19}^e J=79→78	45946.45	45946.45(0)	160.3	79.00	119	19.129	14.5(0.0)	A.30b	73
HC_9N	v_{19}^f J=79→78	45969.57	45969.57(0)	160.3	79.00	179	19.532	14.5(0.0)	A.31a	74
$Si^{13}CC$	$J_{Ka,Kb}=2_{0,2}\rightarrow1_{0,1}$	45970.88	45970.88(0)	3.3	2.00	387	25.534	14.5(0.0)	A.31a	2
SiC_4	J=15→14	46012.39	46012.39(1)	17.7	15.00	1754	23.055	14.2(0.1)	A.31a	5
l-HC_4N	$N_J=10_{10}\rightarrow9_9$	46047.50	46047.58(7)	13.1	9.90	148	21.788	13.6(0.5)	A.31a	6
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$ a	46054.11	46054.03(1)	41.5	33.00	1751	22.559	13.5(0.1)	A.31a	44
$HCCCCC^{13}CCN$	J=41→40	46068.71	46068.62(4)	46.4	41.00	178	20.333	12.7(0.4)	A.31a	9
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$ b	46087.06	46087.04(1)	41.5	33.00	2015	22.698	14.0(0.1)	A.31a	45
MgC_5N	$N_J=40\ \frac{79}{2}\rightarrow39\ \frac{77}{2}$	46111.71	46111.60(1)	45.4	39.50	118	19.132	14.5(0.0)	A.31a	18
MgC_5N	$N_J=40\ \frac{81}{2}\rightarrow39\ \frac{79}{2}$	46113.51	46113.74(7)	45.4	40.50	178	19.512	14.5(0.0)	A.31a	18
KCl	J=6→5	46131.70	46131.52(1)	7.7	6.00	266	21.143	12.3(0.0)	A.31a	2
$HCCC^{13}CCCCN$	J=41→40	46220.87	46220.87(0)	46.6	41.00	118	19.917	14.5(0.0)	A.31b	10
$HCCCC^{13}CCCN$	J=41→40	46224.93	46224.93(0)	46.6	41.00	118	19.917	14.5(0.0)	A.31b	6
C_3S	J=8→7	46245.62	46245.57(23)	10.0	8.00	3675	25.518	14.5(0.0)	A.31b	3
HC_7N	J=41→40	46246.98	46246.96(23)	46.6	41.00	4087	23.463	14.5(0.0)	A.31b	63
^{13}CS	J=1→0	46247.56	46247.41(23)	2.2	1.00	3410	28.798	14.5(0.0)	A.31b	1
C_6H	$v_{11}\ ^2\Delta_{\frac{3}{2}}$ J=$\frac{33}{2}\rightarrow\frac{31}{2}$	46248.68	46248.68(0)	103.2	16.40	294	21.469	14.5(0.0)	A.31b	46
HC_7N	v_{15}^e J=41→40	46304.80	46304.64(7)	135.8	41.00	440	21.233	15.3(0.4)	A.31b	64
HC_7N	v_{15}^f J=41→40	46334.25	46334.16(5)	135.9	41.00	264	20.720	13.1(0.3)	A.31b	65
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ a	46346.20	46346.20(0)	30.6	53.00	234	19.925	14.5(0.0)	A.31b	18
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$ b	46347.88	46347.88(0)	30.6	53.00	234	19.925	14.5(0.0)	A.31b	18
MgC_6H	$N_J=40\ \frac{79}{2}\rightarrow39\ \frac{77}{2}$	46378.49	46378.49(0)	45.4	39.50	146	21.480	14.5(0.0)	A.31b	21
MgC_6H	$N_J=40\ \frac{81}{2}\rightarrow39\ \frac{79}{2}$	46380.45	46380.45(0)	45.4	40.50	29	19.845	14.5(0.0)	A.31b	21
$^{29}SiC_2$	$J_{Ka,Kc}=2_{0,2}\rightarrow1_{0,1}$	46387.49	46387.47(2)	3.3	2.00	672	26.076	14.4(0.1)	A.31b	1
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{53}{2}\rightarrow\frac{51}{2}$	46452.60	46452.60(0)	68.2	106.00	291	19.448	14.5(0.0)	A.32a	19
SiC_6	J=38→37	46453.86	46453.86(0)	43.5	38.00	146	19.133	14.5(0.0)	A.32a	7
HC_9N	J=80→79	46481.17	46481.30(3)	90.4	80.00	494	20.526	14.6(0.3)	A.32a	75
HC_9N	v_{19}^e J=80→79	46528.01	46527.87(8)	162.5	80.00	232	19.769	15.4(0.5)	A.32a	76
HC_9N	v_{19}^f J=80→79	46551.42	46549.29(4)	162.6	80.00	261	19.885	14.6(0.4)	A.32a	77
C_8H^-	J=40→39	46666.11	46666.06(5)	45.9	40.00	317	19.385	14.1(0.3)	A.32a	11
$H^{13}CCCCCN$	J=18→17	46679.68	46679.66(3)	21.3	18.00	720	22.756	14.0(0.2)	A.32a	7
HNCCC	J=5→4	46683.09	46683.02(3)	6.7	5.00	317	22.678	14.7(0.4)	A.32a	2
c-C_3H_2 (para)	$J_{Ka,Kc}=2_{1,1}\rightarrow2_{0,2}$	46755.61	46755.61(1)	8.7	1.05	1521	26.810	14.1(0.0)	A.32b	2
C_6H^-	J=17→16	46812.71	46812.67(0)	20.2	17.00	1431	22.220	13.8(0.1)	A.32b	6
$HCCCCCC^{13}CN$	J=42→41	46832.14	46832.14(0)	48.3	42.00	57	19.158	14.2(0.0)	A.32b	9
NaCN	$J_{Ka,Kc}=3_{0,3}\rightarrow2_{0,2}$	46847.73	46847.68(0)	4.5	3.00	1000	23.443	14.0(0.1)	A.32b	4
NaCN	$J_{Ka,Kc}=3_{2,2}\rightarrow2_{2,1}$	46900.65	46900.72(6)	14.1	1.67	371	23.035	17.9(0.5)	A.32b	5
MgC_3N	$N_J=17\ \frac{33}{2}\rightarrow16\ \frac{31}{2}$	46946.51	46946.36(4)	20.3	16.50	455	21.644	13.1(0.2)	A.32b	11
MgC_3N	$N_J=17\ \frac{35}{2}\rightarrow16\ \frac{33}{2}$	46950.83	46950.71(2)	20.3	17.50	569	21.808	14.6(0.1)	A.33a	12
NaCN	$J_{Ka,Kc}=3_{2,1}\rightarrow2_{2,0}$	46972.35	46972.41(5)	14.1	1.67	512	23.355	14.1(0.4)	A.33a	6
HCN	v_2=1 J=14→14 l=1→-1	47023.15	47023.15(0)	1471.0	0.07	85	26.886	14.5(0.0)	A.33a	3
SiC_2	$J_{Ka,Kc}=2_{0,2}\rightarrow1_{0,1}$	47064.82	47064.82(0)	3.4	2.00	13215	29.040	13.9(0.0)	A.33a	2
$HCCCCC^{13}CCN$	J=42→41	47192.29	47192.15(9)	48.7	42.00	141	20.050	13.7(0.5)	A.33a	10
C_5N^-	J=17→16	47220.65	47220.79(3)	20.4	17.00	450	21.964	14.9(0.2)	A.33b	5
MgC_5N	$N_J=41\ \frac{81}{2}\rightarrow40\ \frac{79}{2}$	47264.42	47264.57(23)	47.6	40.50	168	19.434	14.5(0.0)	A.33b	19
MgC_5N	$N_J=41\ \frac{83}{2}\rightarrow40\ \frac{81}{2}$	47266.23	47266.21(23)	47.6	41.50	84	18.716	14.5(0.0)	A.33b	19
HC_7N	J=42→41	47374.90	47374.84(0)	48.9	42.00	4130	23.426	13.8(0.0)	A.33b	66
C_6H	$v_{11}\ N_J=17\ \frac{35}{2}\rightarrow16\ \frac{33}{2}$	47408.64	47408.63(2)	36.3	17.50	669	22.201	13.8(0.1)	A.33b	47

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
C_6H	ν_{11} N_J=17 $_{\frac{33}{2}}$→16 $_{\frac{31}{2}}$	47425.01	47424.92(1)	36.3	16.50	612	22.172	13.8(0.2)	A.33b	48
$HC^{13}CCCCN$	J=18→17	47436.13	47436.14(2)	21.6	18.00	974	23.042	14.8(0.1)	A.33b	7
$HCCCC^{13}CN$	J=18→17	47453.81	47453.85(0)	21.6	18.00	667	22.663	14.2(0.1)	A.34a	7
HC_7N	ν_{15}^f J=42→41	47464.29	47464.24(1)	138.1	42.00	389	21.062	13.7(0.2)	A.34a	67
$C^{36}S$	J=1→0	47508.78	47507.65(8)	2.3	1.00	166	25.751	14.5(0.0)	A.34a	1
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{81}{2}$ → $\frac{79}{2}$ a	47518.77	47518.77(0)	47.2	80.90	194	19.111	14.5(0.0)	A.34a	30
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{81}{2}$ → $\frac{79}{2}$ b	47519.65	47519.65(0)	47.2	80.90	166	18.957	14.5(0.0)	A.34a	30
MgC_6H	N_J=41 $_{\frac{81}{2}}$→40 $_{\frac{79}{2}}$	47537.94	47537.76(33)	47.6	40.50	55	20.460	14.5(0.0)	A.34a	22
MgC_6H	N_J=41 $_{\frac{83}{2}}$→40 $_{\frac{81}{2}}$	47539.90	47539.43(66)	47.6	41.50	28	19.743	14.5(0.0)	A.34a	22
C_4H	$N_{J,F}$=$5_{11/2,5}$-$4_{9/2,4}$+$5_{11/2,6}$-$4_{9/2,5}$	47566.85	47566.77(0)	6.8	10.80	6750	26.933	13.9(0.0)	A.34a	13
C_4H	$N_{J,F}$=$5_{9/2,5}$-$4_{7/2,4}$+$5_{9/2,4}$-$4_{7/2,3}$	47605.41	47605.47(0)	6.9	8.78	5965	27.016	13.9(0.0)	A.34a	14
HC_9N	J=82→81	47643.11	47643.13(5)	94.9	82.00	248	19.787	13.0(0.3)	A.34a	78
$^{13}CCCN$	N=5→4 J=$\frac{11}{2}$ → $\frac{9}{2}$ F=5→4	47711.04	47711.04(0)	6.9	4.80	55	22.320	14.5(0.0)	A.34a	4
$^{13}CCCN$	N=5→4 J=$\frac{9}{2}$ → $\frac{7}{2}$ F=4→3	47722.07	47722.07(0)	6.9	3.64	55	22.596	14.5(0.0)	A.34a	5
MgNC	N_J=4 $_{\frac{7}{2}}$→3 $_{\frac{5}{2}}$	47726.48	47727.18(23)	5.7	3.50	1730	24.841	15.8(1.4)	A.34b	3
MgNC	N_J=4 $_{\frac{9}{2}}$→3 $_{\frac{7}{2}}$	47741.70	47741.37(23)	5.7	4.50	2224	24.840	15.9(1.4)	A.34b	4
C_4H	ν_7=2 l=0 N_J=5 $_{\frac{11}{2}}$ →4 $_{\frac{9}{2}}$	47792.85	47792.83(1)	543.1	10.80	959	24.976	14.1(0.1)	A.34b	15
C_8H^-	J=41→40	47832.71	47832.57(3)	48.2	41.00	273	19.187	13.4(0.3)	A.34b	12
C_4H	ν_7=2 l=0 N_J=5 $_{\frac{9}{2}}$ →4 $_{\frac{7}{2}}$	47849.61	47849.54(2)	543.1	8.78	710	24.882	13.7(0.1)	A.34b	16
$HCC^{13}CCCN$	J=18→17	47877.64	47877.59(2)	21.8	18.00	737	22.753	14.2(0.1)	A.34b	7
$HCCC^{13}CCN$	J=18→17	47883.47	47883.44(2)	21.8	18.00	737	22.753	14.0(0.1)	A.34b	7
HC_5N	J=18→17	47927.27	47927.24(0)	21.9	18.00	29133	26.430	14.0(0.0)	A.34b	19
HC_5N	ν_{11}^f J=18→17	48048.32	48048.32(0)	175.4	17.90	81	20.551	14.5(0.0)	A.35a	20
$HCCN^{13}C$	J=5→4	48048.41	48048.41(0)	6.9	5.00	108	22.876	14.5(0.0)	A.35a	2
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{55}{2}$ → $\frac{53}{2}$ a	48095.09	48095.09(0)	32.9	55.00	135	19.302	14.5(0.0)	A.35a	20
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{55}{2}$ → $\frac{53}{2}$ b	48096.76	48096.76(0)	32.9	55.00	162	19.484	14.5(0.0)	A.35a	20
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{55}{2}$ → $\frac{53}{2}$	48205.45	48205.45(0)	70.5	110.00	404	19.700	14.5(0.0)	A.35a	21
$C^{34}S$	J=1→0	48206.94	48206.94(0)	2.3	1.00	5003	29.140	14.5(0.0)	A.35b	1
HC_9N	J=83→82	48224.07	48224.14(5)	97.2	83.00	188	19.486	13.7(0.4)	A.35b	79
$HCC^{13}CCCCCN$	J=43→42	48304.31	48305.10(1)	51.0	43.00	188	20.291	10.7(0.0)	A.35b	10
$HCCCCC^{13}CCN$	J=43→42	48315.85	48316.60(4)	51.0	43.00	80	19.443	10.1(0.5)	A.35b	11
MgC_5N	N_J=42 $_{\frac{83}{2}}$→41 $_{\frac{81}{2}}$	48417.14	48417.28(31)	50.0	41.50	53	18.236	14.5(0.0)	A.35b	20
MgC_5N	N_J=42 $_{\frac{85}{2}}$→41 $_{\frac{83}{2}}$	48418.94	48419.07(16)	50.0	42.50	80	18.618	14.5(0.0)	A.35b	20
HC_7N	J=43→42	48502.81	48502.78(1)	51.2	43.00	4250	23.407	13.7(0.1)	A.36a	68
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$ a	48514.58	48514.57(0)	21.3	34.70	2283	22.722	14.0(0.1)	A.36a	49
C_6H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$ b	48523.04	48523.01(0)	21.3	34.70	2256	22.710	13.9(0.0)	A.36a	50
NaCN	$J_{Ka,Kc}$=$3_{1,2}$→$2_{1,1}$	48548.22	48548.22(4)	7.1	2.67	769	23.260	14.2(0.2)	A.36a	7
HC_7N	ν_{15}^e J=43→42	48563.45	48563.49(2)	140.4	43.00	397	21.036	14.1(0.3)	A.36a	69
$C^{33}S$	J=1→0	48586.50	48586.11(23)	2.3	1.00	715	27.186	15.9(1.4)	A.36a	1
HC_7N	ν_{15}^f J=43→42	48594.33	48594.33(0)	140.5	43.00	291	20.724	14.5(0.0)	A.36a	70
C_6H	ν_{11} $^2\Delta_{\frac{5}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$	48646.10	48645.82(6)	87.9	17.10	317	21.452	14.8(0.4)	A.36a	51
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{83}{2}$ → $\frac{81}{2}$ a	48692.00	48692.00(0)	49.6	82.90	184	19.012	14.5(0.0)	A.36a	31
C_8H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{83}{2}$ → $\frac{81}{2}$ b	48692.93	48692.93(0)	49.6	82.90	105	18.452	14.5(0.0)	A.36a	31
MgC_6H	N_J=42 $_{\frac{83}{2}}$→41 $_{\frac{81}{2}}$	48697.40	48697.07(16)	50.0	41.50	158	21.460	14.5(0.0)	A.36a	23
MgC_6H	N_J=42 $_{\frac{85}{2}}$→41 $_{\frac{83}{2}}$	48699.35	48698.96(12)	50.0	42.50	79	20.743	14.5(0.0)	A.36a	23
HC_9N	J=84→83	48805.04	48806.79(16)	99.6	84.00	79	18.590	10.8(2.7)	A.36b	80
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$ a	48845.73	48845.70(1)	43.8	35.00	1597	22.349	13.9(0.1)	A.36b	52
C_6H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$ b	48879.32	48879.33(1)	43.8	35.00	1778	22.456	14.1(0.1)	A.36b	53
CS	J=1→0	48990.96	48990.91(0)	2.4	1.00	88117	31.992	13.9(0.0)	A.37a	1
C_6H	ν_{11} $^2\Delta_{\frac{3}{2}}$ J=$\frac{35}{2}$ → $\frac{33}{2}$	49051.30	49051.19(7)	105.6	17.40	234	21.122	13.2(0.5)	A.37a	54
SiC_4	J=16→15	49079.76	49079.72(1)	20.0	16.00	1296	22.624	13.9(0.1)	A.37a	6
$C^{13}CCN$	N=5→4 J=$\frac{11}{2}$ → $\frac{9}{2}$ F=6→5	49197.23	49197.67(0)	7.1	4.83	77	22.625	11.1(0.1)	A.37a	3
$C^{13}CCN$	N=5→4 J=$\frac{9}{2}$ → $\frac{7}{2}$ F=5→4	49210.54	49210.54(0)	7.1	4.60	103	22.961	14.5(0.0)	A.37a	4
$H^{13}CCCCCN$	J=19→18	49272.91	49272.87(2)	23.6	19.00	566	22.407	13.9(0.2)	A.37b	8
$CC^{13}CN$	N=5→4 J=$\frac{11}{2}$ → $\frac{9}{2}$ F=6→5	49280.90	49280.90(0)	7.1	4.83	103	22.908	14.5(0.0)	A.37b	3
$CC^{13}CN$	N=5→4 J=$\frac{11}{2}$ → $\frac{9}{2}$ F=5→4	49281.41	49281.41(0)	7.1	4.74	103	22.926	14.5(0.0)	A.37b	3
$CC^{13}CN$	N=5→4 J=$\frac{9}{2}$ → $\frac{7}{2}$ F=5→4	49299.11	49299.11(0)	7.1	3.78	51	22.458	14.5(0.0)	A.37b	4
$CC^{13}CN$	N=5→4 J=$\frac{9}{2}$ → $\frac{7}{2}$ F=4→3	49299.64	49299.64(0)	7.1	2.78	51	22.766	14.5(0.0)	A.37b	4
C_3N	N=5→4 a	49466.42	49466.39(0)	7.1	16.00	9904	26.273	14.2(0.0)	A.38a	3
C_3N	N=5→4 b	49485.22	49485.19(0)	7.1	12.90	8085	26.285	14.2(0.0)	A.38a	4
C_6H^-	J=18→17	49566.31	49566.27(0)	22.6	18.00	1169	21.903	13.9(0.1)	A.38a	7
CH_3C_3N	N_J=12_3→11_3	49572.77	49572.77(0)	82.9	11.30	254	21.935	14.5(0.0)	A.38a	5
HC_7N	J=44→43	49630.72	49630.68(0)	53.6	44.00	3347	23.123	13.9(0.0)	A.38a	71
HCCNC	J=5→4	49678.07	49678.04(17)	7.2	5.00	278	23.787	13.5(0.9)	A.38a	2
HC_7N	ν_{15}^e J=44→43	49692.77	49692.86(23)	142.8	44.00	152	20.028	13.6(0.0)	A.38a	72
MgC_3N	N_J=18 $_{\frac{35}{2}}$→17 $_{\frac{33}{2}}$	49708.00	49708.13(7)	22.7	17.50	253	20.940	14.2(0.3)	A.38b	13
MgC_3N	N_J=18 $_{\frac{37}{2}}$→17 $_{\frac{35}{2}}$	49712.32	49712.02(1)	22.7	18.50	430	21.415	16.3(0.1)	A.38b	14
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ a	49843.96	49843.96(0)	35.3	57.00	226	19.746	14.5(0.0)	A.38b	22
C_7H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$ b	49845.64	49845.64(0)	35.3	57.00	75	18.647	14.5(0.0)	A.38b	22
C_7H	$^2\Pi_{\frac{3}{2}}$ J=$\frac{57}{2}$ → $\frac{55}{2}$	49958.30	49958.24(8)	72.9	114.00	225	19.046	13.4(0.5)	A.39a	23
C_5N^-	J=18→17	49998.25	49998.76(9)	22.8	18.00	400	21.733	14.2(0.3)	A.39a	6

Continues on next page.

Molecular species	Quantum numbers of the transition	ν_{rest} (MHz)	ν_{obs} (MHz)	E_{up} (K)	S_{ul}	$W=\int T_{MB} dv$ (mK·kms^{-1})	$ln\frac{3\,k_B\,W}{8\,\pi^3\,\nu_{rest}\,S_{ul}\,\mu^2}$ (Rot. Diagram)	Vexp (kms^{-1})	Fig.	Line count
$HC^{13}CCCCN$	J=19→18	50071.39	50071.38(23)	24.0	19.00	598	22.446	14.1(1.4)	A.39a	8
$HCCCC^{13}CN$	J=19→18	50090.05	50090.09(4)	24.0	19.00	523	22.311	13.5(0.3)	A.39a	8
C_4H	ν_7 N_J=5 $\frac{11}{2}$ → 4 $\frac{9}{2}$ a	50097.69	50097.21(3)	251.4	10.10	896	24.929	14.4(0.2)	A.39a	17
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{21}{2}$ → $\frac{19}{2}$ a	50124.61	50124.63(1)	13.7	20.90	1592	23.089	14.5(0.1)	A.39a	11
C_5H	$^2\Pi_{\frac{1}{2}}$ J=$\frac{21}{2}$ → $\frac{19}{2}$ b	50129.05	50128.90(1)	13.7	20.90	1567	23.073	14.7(0.1)	A.39a	12
C_8H^-	J=43→42	50165.89	50165.63(17)	53.0	43.00	124	18.302	16.9(1.0)	A.39a	13
C_4H	ν_7 N_J=5 $\frac{11}{2}$ → 4 $\frac{9}{2}$ b	50196.75	50196.48(1)	251.5	10.10	794	24.805	14.5(0.0)	A.39a	18
C_6H	ν_{11} N_J=18 $\frac{37}{2}$ →17 $\frac{35}{2}$	50197.90	50197.92(5)	38.7	18.50	422	21.627	14.5(0.0)	A.39b	55
C_6H	ν_{11} N_J=18 $\frac{35}{2}$ →17 $\frac{33}{2}$	50213.97	50213.90(23)	38.7	17.50	347	21.488	13.8(1.4)	A.39b	56

Table A.2. Observational parameters of the remaining unidentified lines in this work.

Label	ν_{obs} (MHz)	$\int T_A^* dv$ (mK·kms^{-1})	Vexp (kms^{-1})	Label	ν_{obs} (MHz)	$\int T_A^* dv$ (mK·kms^{-1})	Vexp (kms^{-1})	Label	ν_{obs} (MHz)	$\int T_A^* dv$ (mK·kms^{-1})	Vexp (kms^{-1})
U31869	31868.98(6)	50	14.6(0.6)	U32036	32035.92(8)	49	13.7(0.9)	U32040	32039.77(11)	49	20.1(0.6)
U32079	32078.51(4)	49	12.1(0.4)	U32119	32118.86(4)	25	12.2(0.4)	U32201	32200.71(0)	24	11.2(0.0)
U32322	32322.47(8)	49	14.5(0.0)	U32971	32970.52(10)	35	13.0(0.9)	U33174	33174.03(19)	35	15.6(1.4)
U33199	33198.51(10)	35	16.4(1.0)	U33202	33201.75(3)	35	12.9(0.2)	U33795	33795.37(6)	33	14.1(0.7)
U34083	34083.01(5)	44	14.0(0.4)	U34312	34312.00(7)	43	14.7(0.5)	U34766	34766.03(6)	52	14.4(0.5)
U34934	34934.28(4)	52	15.5(0.5)	U35849	35848.61(1)	49	13.6(0.4)	U36126	36126.03(5)	39	14.4(0.5)
U36380	36380.43(14)	19	10.5(2.4)	U36745	36744.91(6)	28	10.5(0.6)	U36750	36749.78(11)	28	13.8(0.9)
U36783	36783.37(1)	28	14.9(0.9)	U36952	36951.59(10)	37	11.8(0.7)	U37329	37328.60(23)	36	14.8(1.8)
U37443	37442.57(12)	27	14.1(0.4)	U37663	37663.08(4)	80	14.3(0.2)	U37977	37977.19(5)	44	15.0(0.4)
U38457	38456.75(23)	51	14.9(1.8)	U39021	39020.73(5)	50	15.2(0.4)	U39356	39355.87(18)	25	14.5(0.0)
U39575	39574.86(16)	32	12.4(0.4)	U40245	40245.20(6)	47	14.3(0.4)	U40560	40559.65(13)	39	16.7(0.9)
U40598	40598.39(4)	54	15.6(0.4)	U40714	40714.29(5)	46	14.5(0.0)	U40787	40787.22(7)	38	14.0(0.5)
U41712	41711.90(1)	233	14.6(0.1)	U42229	42229.33(5)	43	13.5(0.4)	U42340	42340.02(2)	92	14.1(0.2)
U42399	42398.99(4)	35	14.1(0.3)	U42443	42443.45(5)	35	13.9(0.3)	U42461	42461.14(7)	42	14.1(0.4)
U42514	42513.56(6)	35	14.1(0.3)	U42781	42780.72(5)	35	14.3(0.4)	U43118	43118.20(5)	48	16.0(0.4)
U43134	43133.53(8)	34	12.7(0.5)	U43457	43457.12(4)	60	13.6(0.3)	U43692	43691.87(3)	47	13.7(0.2)
U44574	44574.19(5)	51	12.4(0.5)	U44857	44857.46(0)	63	14.5(0.0)	U45176	45175.57(23)	75	14.5(0.0)
U46135	46135.23(1)	30	11.4(0.6)	U46613	46613.17(5)	64	14.6(0.4)	U46640	46640.00(3)	41	14.4(0.3)
U46662	46661.51(6)	58	14.5(0.6)	U47584	47584.28(14)	50	11.5(0.4)	U48365	48364.93(21)	33	14.5(0.0)



Fig. A.1. IRC+10216 YEBES 40m data, line fits and labels from 30950 to 31450 GHz.



Fig. A.2. IRC+10216 YEBES 40m data, line fits and labels from 31450 to 31950 GHz.



Fig. A.3. IRC+10216 YEBES 40m data, line fits and labels from 31950 to 32450 GHz.



Fig. A.4. IRC+10216 YEBES 40m data, line fits and labels from 32450 to 32950 GHz.



Fig. A.5. IRC+10216 YEBES 40m data, line fits and labels from 32950 to 33450 GHz.



Fig. A.6. IRC+10216 YEBES 40m data, line fits and labels from 33450 to 33950 GHz.



Fig. A.7. IRC+10216 YEBES 40m data, line fits and labels from 33950 to 34450 GHz.



Fig. A.8. IRC+10216 YEBES 40m data, line fits and labels from 34450 to 34950 GHz.



Fig. A.9. IRC+10216 YEBES 40m data, line fits and labels from 34950 to 35450 GHz.



Fig. A.10. IRC+10216 YEBES 40m data, line fits and labels from 35450 to 35950 GHz.



Fig. A.11. IRC+10216 YEBES 40m data, line fits and labels from 35950 to 36450 GHz.



Fig. A.12. IRC+10216 YEBES 40m data, line fits and labels from 36450 to 36950 GHz.



Fig. A.13. IRC+10216 YEBES 40m data, line fits and labels from 36950 to 37450 GHz.



Fig. A.14. IRC+10216 YEBES 40m data, line fits and labels from 37450 to 37950 GHz.



Fig. A.15. IRC+10216 YEBES 40m data, line fits and labels from 37950 to 38450 GHz.



Fig. A.16. IRC+10216 YEBES 40m data, line fits and labels from 38450 to 38950 GHz.



Fig. A.17. IRC+10216 YEBES 40m data, line fits and labels from 38950 to 39450 GHz.



Fig. A.18. IRC+10216 YEBES 40m data, line fits and labels from 39450 to 39950 GHz.



Fig. A.19. IRC+10216 YEBES 40m data, line fits and labels from 39950 to 40450 GHz.



Fig. A.20. IRC+10216 YEBES 40m data, line fits and labels from 40450 to 40950 GHz.



Fig. A.21. IRC+10216 YEBES 40m data, line fits and labels from 40950 to 41450 GHz.



Fig. A.22. IRC+10216 YEBES 40m data, line fits and labels from 41450 to 41950 GHz.



Fig. A.23. IRC+10216 YEBES 40m data, line fits and labels from 41950 to 42450 GHz.



Fig. A.24. IRC+10216 YEBES 40m data, line fits and labels from 42450 to 42950 GHz.



Fig. A.25. IRC+10216 YEBES 40m data, line fits and labels from 42950 to 43450 GHz.



Fig. A.26. IRC+10216 YEBES 40m data, line fits and labels from 43450 to 43950 GHz.



Fig. A.27. IRC+10216 YEBES 40m data, line fits and labels from 43950 to 44450 GHz.



Fig. A.28. IRC+10216 YEBES 40m data, line fits and labels from 44450 to 44950 GHz.



Fig. A.29. IRC+10216 YEBES 40m data, line fits and labels from 44950 to 45450 GHz.



Fig. A.30. IRC+10216 YEBES 40m data, line fits and labels from 45450 to 45950 GHz.



Fig. A.31. IRC+10216 YEBES 40m data, line fits and labels from 45950 to 46450 GHz.



Fig. A.32. IRC+10216 YEBES 40m data, line fits and labels from 46450 to 46950 GHz.



Fig. A.33. IRC+10216 YEBES 40m data, line fits and labels from 46950 to 47450 GHz.



Fig. A.34. IRC+10216 YEBES 40m data, line fits and labels from 47450 to 47950 GHz.



Fig. A.35. IRC+10216 YEBES 40m data, line fits and labels from 47950 to 48450 GHz.



Fig. A.36. IRC+10216 YEBES 40m data, line fits and labels from 48450 to 48950 GHz.



Fig. A.37. IRC+10216 YEBES 40m data, line fits and labels from 48950 to 49450 GHz.



Fig. A.38. IRC+10216 YEBES 40m data, line fits and labels from 49450 to 49950 GHz.



Fig. A.39. IRC+10216 YEBES 40m data, line fits and labels from 49950 to 50450 GHz.