


**Fig. S1.** Upper panel: The  $O - C$  curve of V1251 Cyg during BJD 2454763–2454776. We used an ephemeris of BJD  $2454765.6790 + 0.0759607E$  for drawing this figure. Middle panel: The amplitude of superhumps. Lower panel: The light curve. The horizontal axis in units of BJD and cycle number is common to all of these panels.

## Supplementary Discussion

V1251 Cyg has shown 5 superoutbursts in 1963, 1991, 1994, 1997 and 2008 (Weber 1966; Wenzel 1991; Kato 1995; Kato et al. 2009). The supercycle seems to be 3–28 yr. Rebrightenings were observed in the 1997 and 2008 superoutbursts and in the 2008 superoutburst, double-wave variations similar to early superhumps were observed (Kato et al. 2009). Kato et al. (2009) analyzed the 2008 superoutburst of V1251 Cyg and found the period of early superhumps to be 0.07433(6).


We reanalyzed the 2008 superoutburst of V1251 Cyg by using VSNET and AAVSO data. The times of maxima of ordinary superhumps are listed in table S3. Figure S1 shows the  $O - C$  curve, the amplitude of superhumps (middle panel) and the light curve (lower panel) of V1251 Cyg during BJD 2454763–2454776. We regarded BJD 2454763.2–2454764.9 ( $0 \leq E \leq 17$ ) as stage A from the variation of the superhump period and the amplitude of superhumps. Figure S2 shows the result of PDM analysis of stage A superhumps (upper panel) and



**Fig. S2.** Upper panel:  $\theta$ -diagram of our PDM analysis of stage A superhumps of V1251 Cyg (BJD 2454763.2–2454764.9). Lower panel: Phase-averaged profile of stage A superhumps.

the mean profile (lower panel). We found the stage A superhump period to be 0.07616(3) d. We also attempt to estimate the period of early superhumps but failed because of a shortage of observational data and contamination by inaccurate data. Although double-wave variations of magnitude similar to early superhumps can be seen in its light curve before the appearance of ordinary superhumps, it is so contaminated that we could not obtain a reliable period of early superhumps.


RZ Leo has shown many superoutbursts. The brightenings of RZ Leo are recorded in 1918, 1935, 1952, 1976, 1984, 1987, 1988, 1989, 1990, 2000, 2006 and 2016 (Wolf 1919; Richter 1985; Ishioka et al. 2001; Kato 2015; Kato et al. 2016). In these recorded brightenings, the 1989 brightening seems to be a normal outburst (Narumi et al. 1989; O’Donoghue et al. 1991). In addition, Ishioka et al. (2001) indicated that the 1987, 1988, 1989 and 1990 brightenings might be short and probably were mistaken as brightenings of another nearby object. Therefore, the superoutbursts seem to be in 1918, 1935, 1952, 1976, 1984, 2000, 2006 and 2016, and thus the supercycle seems to be 6–24 yr. Although RZ Leo had shown many superoutbursts, variations of magnitude similar to early superhumps are recorded only in the 2000 superoutbursts (Ishioka et al. 2001; Kato et al. 2016). The spectroscopic orbital period is 0.07651(26) d (Mennickent, Tappert 2001) and the photometric period is 0.0756(12) d (Mennickent et al. 1999). Ishioka




**Fig. S3.** Upper panel: The  $O-C$  curve of RZ Leo during BJD 2451901–2451915. We used an ephemeris of BJD 2451902.2763+0.0785326 $E$  for drawing this figure. Middle panel: The amplitude of superhumps. Lower panel: The light curve. The horizontal axis in units of BJD and cycle number is common to all of these panels.

et al. (2001) analyzed the 2000 superoutburst of RZ Leo and found the orbital period to be 0.07616(21) d, and estimated the mass ratio to be 0.14 from the period of early superhumps and the superhump excess.


We reanalyzed the 2000 superoutburst of RZ Leo by using VSNET and AAVSO data and estimated the mass ratio by using a new method developed by Kato, Osaki (2013). The times of maxima of ordinary superhumps are listed in table S4. Figure S3 shows the  $O-C$  curve, the amplitude of superhumps (middle panel) and the light curve (lower panel) of RZ Leo during BJD 2451901–2451915. We regarded BJD 2451901.1–2451902.7 ( $0 \leq E \leq 18$ ) as stage A from the variation of the superhump period and the amplitude of superhumps. Figures S4 and S5 show the results of PDM analysis of early superhumps and stage A superhumps (upper panel) and the mean profiles (lower panel), respectively. We found the period of early superhumps to be 0.07626(7) d and the stage A superhump period to be 0.08072(5) d, and then we estimate the mass ratio at 0.165(6)



**Fig. S4.** Upper panel:  $\theta$ -diagram of our PDM analysis of early superhumps of RZ Leo (BJD 2451899.1–2451901.0). Lower panel: Phase-averaged profile of early superhumps.




**Fig. S5.** Upper panel:  $\theta$ -diagram of our PDM analysis of stage A superhumps of RZ Leo (BJD 2451901.1–2451902.7). Lower panel: Phase-averaged profile of stage A superhumps.




**Fig. S6.** Upper panel: The  $O - C$  curve of BC UMa during BJD 2452674–2452685. We used an ephemeris of BJD 2452674.2372+0.0645728 $E$  for drawing this figure. Middle panel: The amplitude of superhumps. Lower panel: The light curve. The horizontal axis in units of BJD and cycle number is common to all of these panels.

by using these periods. The period of early superhumps is consistent with that obtained by Ishioka et al. (2001) within the errors. The mass ratio we obtained is slightly larger than  $q = 0.14$  that Ishioka et al. (2001) calculated from the superhump excess. It should be noted that this estimated mass ratio involves slightly larger uncertainties because of the short baseline of the data of stage A superhumps.


BC UMa has shown many superoutbursts and early superhumps were observed. The superoutbursts of BC UMa were recorded in 1960, 1962, 1982, 1990, 1992, 1994, 2000, 2003 and 2009 (Romano 1964; Kunjaya et al. 1998; Boyd 2003; Maehara et al. 2007; Kato et al. 2009; Kato et al. 2010; Kato 2015). The supercycle of BC UMa is thus about 2–20 yr. Patterson et al. (2003) found the orbital period to be 0.062605(11) d and the period of early superhumps to be 0.06256(8) d from the 2000 superoutburst. Maehara et al. (2007) analyzed the 2003 superoutburst and found the period of early superhumps to be 0.06258(13) d, and estimated the mass



**Fig. S7.** Upper panel:  $\theta$ -diagram of our PDM analysis of early superhumps of BC UMa (BJD 2452672.1–2452673.4). Lower panel: Phase-averaged profile of early superhumps.




**Fig. S8.** Upper panel:  $\theta$ -diagram of our PDM analysis of stage A superhumps of BC UMa (BJD 2452674.0–2452674.6). Lower panel: Phase-averaged profile of stage A superhumps.



**Fig. S9.** Upper panel: The  $O - C$  curve of MASTER J004527 during BJD 2456554–2456565. We used an ephemeris of BJD  $2456556.3815 + 0.0801953E$  for drawing this figure. Middle panel: The amplitude of superhumps. Lower panel: The light curve. The horizontal axis in units of BJD and cycle number is common to all of these panels.

ratio to be  $q = 0.13$  from the superhump excess. They also suggested that BC UMa may be an intermediate class between the WZ Sge type DNe and the SU UMa-type ones from the presence of normal outbursts and two types of superoutbursts. One has long duration and large amplitude comparable to WZ Sge-type DNe and the other has short duration and small amplitude in comparison with WZ Sge-type ones, which are supposed to reflect the stored mass in the disk before each superoutburst.

We reanalyzed the 2003 superoutburst of BC UMa for the purpose of the estimation of its mass ratio by using the new method proposed in Kato, Osaki (2013). The times of maxima of ordinary superhumps are listed in table S5. Figure S6 shows the  $O - C$  curve, the amplitude of superhumps (middle panel) and the light curve (lower panel) of BC UMa during BJD 2452674–2452685. We regarded BJD 2452674.0–2452674.6 ( $0 \leq E \leq 6$ ) as stage A from the variation of the superhump period and the amplitude of superhumps. Figures S7 and S8 show the results of PDM analysis of early superhumps and stage A super-



**Fig. S10.** Upper panel:  $\theta$ -diagram of our PDM analysis of stage A superhumps of MASTER J004527 (BJD 2456554.3–56555.1). Lower panel: Phase-averaged profile of stage A superhumps.

humps (upper panel) and the superhump profiles (lower panel), respectively. We found the period of early superhumps to be  $0.06251(5)$  d, and naturally, this value is consistent with the period in Maehara et al. (2007) within the errors. We obtained the stage A superhump period to be  $0.06476(7)$  d. We then estimated the mass ratio to be  $0.096(6)$ , which is smaller than that in Maehara et al. (2007) and larger than the upper limit of mass ratios for occurrence of the 2:1 resonance.

MASTER OT J004527.52+503213.8 (hereafter MASTER J004527) is regarded as WZ Sge-type DNe based on the large outburst amplitude, and only one superoutburst in 2013 is recorded (Kato et al. 2014). Amplitudes of superhumps is much larger than those of period bouncers and thus this object may not be a period bouncer.

We reanalyzed the 2013 superoutburst of MASTER J004527. The times of maxima of ordinary superhumps are listed in table S6. Figure S9 shows the  $O - C$  curve, the amplitude of superhumps (middle panel) and the light curve (lower panel) of MASTER J004527 during BJD 2456554–2456565. We regarded BJD 2456554.3–2456555.1 ( $0 \leq E \leq 7$ ) as stage A from the variation of the superhump period and the amplitude of superhumps. Figure S10 shows the result of PDM analysis of stage A superhumps (upper panel) and the superhump profile (lower panel). We found the stage A superhump period to be  $0.08136(7)$  d.

QY Per is regarded as a WZ Sge-type DN from the large outburst amplitude and long recurrence time (Kato et al. 2009; Kato et al. 2016). This object showed three superoutbursts in 1999, 2005 and 2015 and thus the supercycle is 6–10 years. The 2015 superoutburst seems to have shown no early superhumps. It appears that this object showed a different type of superoutburst (Kato et al. 2016).

We attempted to obtain the stage A superhump period from the 1999 superoutburst but failed because of the shortage of observational data.

## References

- Boyd, D. 2003, *J. Br. Astron. Assoc.*, 113, 208  
 Ishioka, R., et al. 2001, *PASJ*, 53, 905  
 Kato, T. 1995, *IBVS*, 4152  
 Kato, T. 2015, *PASJ*, 67, 108  
 Kato, T., et al. 2014, *PASJ*, 66, 90  
 Kato, T., et al. 2016, *PASJ*, in press (arXiv/1605.06221)  
 Kato, T., et al. 2009, *PASJ*, 61, S395  
 Kato, T., et al. 2010, *PASJ*, 62, 1525  
 Kato, T., & Osaki, Y. 2013, *PASJ*, 65, 115  
 Kunjaya, C., Kato, T., & Hirata, R. 1998, *Highlights of Astronomy*, 11, 388  
 Maehara, H., Hachisu, I., & Nakajima, K. 2007, *PASJ*, 59, 227  
 Mennickent, R. E., Sterken, C., Gieren, W., & Unda, E. 1999, *A&A*, 352, 239  
 Mennickent, R. E., & Tappert, C. 2001, *A&A*, 372, 563  
 Narumi, H., Korth, S., Dyck, G., Iida, M., & Hurst, G. 1989, *IAU Circ.*, 4757  
 O’Donoghue, D., Chen, A., Marang, F., Mittaz, J. P. D., Winkler, H., & Warner, B. 1991, *MNRAS*, 250, 363  
 Patterson, J., et al. 2003, *PASP*, 115, 1308  
 Richter, G. A. 1985, *IBVS*, 2714  
 Romano, G. 1964, *Mem. Soc. Astron. Ital.*, 35, 101  
 Weber, R. 1966, *IBVS*, 123  
 Wenzel, W. 1991, *IBVS*, 3689  
 Wolf, M. 1919, *Astron. Nachr.*, 209, 85

**Table S1.** Log of observations of ASASSN-16eg.

Start*	End*	mag <sup>†</sup>	error <sup>‡</sup>	$N^{\S}$	obs <sup>  </sup>	sys <sup>#</sup>
89.4093	89.5309	-0.320	0.002	204	Van	C
90.2630	90.3258	-0.762	0.003	156	KU1	C
90.3915	90.3955	13.009	0.005	6	BPO	CV
90.4401	90.4906	12.859	0.003	132	Trt	CV
90.4854	90.6306	-0.153	0.001	330	Van	C
90.6317	90.8780	12.853	0.002	253	UJH	CV
90.6897	90.9264	13.157	0.001	458	BJA	CV
91.3734	91.5569	13.240	0.002	243	Han	CV
91.4236	91.4825	13.018	0.002	157	Trt	CV
91.6305	91.8529	13.028	0.005	133	UJH	CV
92.3479	92.6688	13.448	0.001	383	BPO	CV
92.3834	92.5556	13.452	0.001	253	Han	CV
92.4047	92.5911	0.276	0.001	424	Van	C
92.6635	92.8786	13.232	0.002	242	UJH	CV
92.6854	92.8961	13.315	0.001	250	CMJ	V
92.6897	92.9280	13.531	0.001	443	BJA	CV
92.8935	92.9646	13.203	0.011	42	SGE	CV
93.0910	93.3325	0.384	0.002	340	Aka	C
93.3844	93.5587	13.599	0.001	270	Han	CV
93.4689	93.6118	13.355	0.001	386	Trt	CV
93.5587	93.9208	13.873	0.002	398	SDB	CV
93.6313	93.8754	13.368	0.002	267	UJH	CV
93.6644	93.9031	13.446	0.001	353	CMJ	V
93.7261	93.8841	13.799	0.001	295	MZK	CV
93.7271	93.7868	13.428	0.015	10	Neu	V
93.8262	93.9963	13.343	0.003	152	SGE	CV
94.1120	94.2159	0.519	0.004	74	Aka	C
94.2207	94.3124	13.651	0.003	195	Ioh	C
94.4401	94.6122	13.425	0.002	468	Trt	CV
94.6297	94.8736	13.445	0.003	268	UJH	CV
94.6366	94.9024	13.527	0.003	382	CMJ	V
94.7388	95.0308	13.730	0.003	568	JJI	CV
94.7470	94.8895	13.883	0.004	270	MZK	CV
94.7613	94.9640	13.523	0.012	32	Neu	V
94.8866	95.0007	13.429	0.005	175	SGE	CV
95.0935	95.1498	13.728	0.009	124	Kis	C
95.2966	95.4441	0.115	0.007	200	CRI	C
95.3693	95.5969	0.584	0.006	290	DPV	C
95.3862	95.4936	13.834	0.010	64	Han	CV
95.4556	95.6518	13.598	0.005	359	IMi	CV
95.5360	95.6827	13.604	0.007	202	RPc	V
95.5558	95.8947	14.061	0.005	400	SDB	CV
95.5780	95.9027	13.621	0.005	480	CMJ	V
95.6314	95.8746	13.544	0.006	269	UJH	CV

\*BJD–2457400.

<sup>†</sup>Mean magnitude.

<sup>‡</sup> $1\sigma$  of the mean magnitude.

<sup>§</sup>Number of observations.

<sup>||</sup>Observer’s code: Neu (V. Neustroev team), Van (T. Vanmunster), KU1 (Kyoto U., campus obs.), Trt (T. Tordai), Han (Hankasalmi Obs., by A. Oksanen), Aka (H. Akazawa, OUS), Ioh (H. Itoh), Kis (S. Kiyota), CRI (Crimean Astrophys. Obs.), DPV (P. Dubovsky), IMi (I. Miller), RPc (R. Pickard) and AAVSO collaboration.

<sup>#</sup>Filter. “C” means no filter (clear).

**Table S1.** Log of observations of ASASSN-16eg (continued).

Start*	End*	mag <sup>†</sup>	error <sup>‡</sup>	N <sup>§</sup>	obs <sup>  </sup>	sys <sup>#</sup>
95.6683	95.8868	13.465	0.005	439	RIT	CV
95.7350	95.9283	13.593	0.018	32	Neu	V
95.7366	96.0347	13.796	0.005	384	JJI	CV
95.7462	95.8886	13.992	0.006	270	MZK	CV
95.8664	96.0175	13.484	0.007	225	SGE	CV
95.9414	95.9742	13.635	0.023	30	FJQ	V
95.9439	95.9766	13.451	0.023	30	FJQ	B
95.9464	95.9791	13.606	0.024	30	FJQ	I
96.0548	96.1916	0.056	0.007	235	KU1	C
96.0578	96.2629	13.787	0.008	404	Kis	C
96.1035	96.3189	0.598	0.008	183	Aka	C
96.1277	96.2813	13.746	0.007	274	Ioh	C
96.2862	96.4033	0.156	0.007	159	CRI	C
96.3865	96.6420	0.632	0.005	403	Van	C
96.4372	96.5458	13.579	0.006	257	Trt	CV
96.4803	96.6364	13.616	0.005	300	IMi	CV
96.5763	96.9059	13.663	0.005	456	CMJ	V
96.5828	96.9283	14.104	0.005	415	SDB	CV
96.6250	96.8747	13.580	0.006	276	UJH	CV
96.6661	96.8412	13.701	0.006	291	DKS	CV
96.7195	96.9013	13.513	0.005	394	RIT	CV
96.7291	97.0213	13.857	0.004	551	JJI	CV
96.7344	96.7916	13.604	0.032	10	Neu	V
96.7624	97.0165	13.564	0.005	375	SGE	CV
96.7732	96.9646	13.705	0.007	230	GFB	V
96.8806	96.9795	13.720	0.011	71	FJQ	V
96.8831	96.9820	13.604	0.012	75	FJQ	B
96.8856	96.9770	13.696	0.011	72	FJQ	I
97.2118	97.3228	0.756	0.009	78	Aka	C
97.3548	97.6611	14.019	0.004	372	BPO	CV
97.3786	97.4280	0.802	0.009	82	Van	C
97.7311	98.0273	14.050	0.003	575	JJI	CV
97.7597	98.0115	13.748	0.004	375	SGE	CV
97.8657	97.9663	13.908	0.014	39	FJQ	V
97.8682	97.9689	13.809	0.017	39	FJQ	B
97.8735	97.9638	13.878	0.015	36	FJQ	I
98.1123	98.3056	14.106	0.004	413	Ioh	C
98.1250	98.3075	0.363	0.004	508	KU1	C
98.3869	98.6416	0.940	0.004	404	Van	C
98.4471	98.6381	13.917	0.004	370	IMi	CV
98.5048	98.6048	13.913	0.004	270	Trt	CV
98.5857	98.7635	13.994	0.005	239	CMJ	V
98.6752	98.8918	13.814	0.003	471	RIT	CV
98.7009	98.8318	13.972	0.016	22	Neu	V

\*BJD–2457400.

†Mean magnitude.

‡1 $\sigma$  of the mean magnitude.

§Number of observations.

||Observer's code: RIT (M. Richmond), Neu (V. Neustroev team), KU1 (Kyoto U., campus obs.), Kis (S. Kiyota), Aka (H. Akazawa, OUS), Ioh (H. Itoh), CRI (Crimean Astrophys. Obs.), Van (T. Vanmunster), Trt (T. Tordai) IMi (I. Miller) and AAVSO collaboration.

#Filter. "C" means no filter (clear).

**Table S1.** Log of observations of ASASSN-16eg (continued).

Start*	End*	mag <sup>†</sup>	error <sup>‡</sup>	N <sup>§</sup>	obs <sup>  </sup>	sys <sup>#</sup>
98.7310	98.7638	14.242	0.006	65	JJI	CV
98.7569	99.0205	13.845	0.004	400	SGE	CV
99.4120	99.6418	1.014	0.004	363	Van	C
100.3629	100.4249	14.388	0.008	57	Han	CV
100.5045	100.5554	14.225	0.005	71	Trt	CV
100.7198	100.9118	14.189	0.015	32	Neu	V
101.0525	101.3187	0.663	0.003	711	KU1	C
102.3653	102.4669	14.565	0.008	89	Han	CV
102.6945	102.8869	14.444	0.012	32	Neu	V
103.4847	103.6973	14.688	0.004	258	deM	C
103.7050	103.8968	14.497	0.013	32	Neu	V
104.1938	104.3125	0.936	0.004	329	KU1	C
104.5506	104.6985	14.775	0.005	169	deM	C
104.7136	104.7944	14.595	0.016	14	Neu	V
105.6888	105.8832	14.748	0.009	32	Neu	V
106.5020	106.6276	1.821	0.003	153	Van	C
107.8014	107.9083	15.069	0.010	18	Neu	V
108.1021	108.3142	2.293	0.008	102	Aka	C
108.2283	108.3125	1.787	0.003	366	KU1	C
108.7084	108.9025	15.971	0.022	32	Neu	V
109.1621	109.3116	3.597	0.016	72	Aka	C
109.1985	109.2805	2.908	0.005	229	KU1	C
110.1797	110.1929	3.622	0.033	32	KU1	C
111.6673	111.6673	17.791	0.05	6	Neu	V
112.4593	112.6131	4.024	0.012	128	Van	C
112.7011	112.8398	16.264	0.072	15	Neu	V
113.1365	113.1419	1.901	0.008	13	KU1	C
113.3565	113.6137	2.634	0.003	318	Van	C
113.6668	113.8069	15.934	0.019	15	Neu	V
114.6896	114.6896	17.466	0.04	6	Neu	V
115.8809	115.8809	18.401	0.04	6	Neu	V
119.8117	119.8117	18.710	0.18	6	Neu	V
120.1786	120.1939	18.500	0.000	2	KU1	C
122.7220	122.7220	18.840	0.15	6	Neu	V
152.8352	152.8352	19.100	0.04	6	Neu	V
154.7665	154.7665	19.162	0.04	6	Neu	V
155.6973	155.6973	19.194	0.04	6	Neu	V
157.6497	157.6497	19.198	0.04	6	Neu	V
160.6502	160.6502	19.220	0.04	6	Neu	V
204.6500	204.6500	19.470	0.04	6	Neu	V
236.6107	236.6107	19.472	0.04	6	Neu	V
242.6929	242.6929	19.484	0.01	6	Neu	V
267.5930	267.5930	19.635	0.08	6	Neu	V
301.3372	301.3372	19.560	0.01	6	Neu	V

\*BJD–2457400.

†Mean magnitude.

‡1 $\sigma$  of the mean magnitude.

§Number of observations.

||Observer's code: Neu (V. Neustroev team), Van (T. Vanmunster), Han (Hankasalmi Obs., by A. Oksanen), Trt (T. Tordai), KU1 (Kyoto U., campus obs.), deM (E. de Miguel), Aka (H. Akazawa, OUS) and AAVSO collaboration.

#Filter. "C" means no filter (clear).

**Table S2.** Times of superhump maxima in ASASSN-16eg.

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
0	57494.1637	0.0010	-0.0183	44
1	57494.2401	0.0005	-0.0198	111
4	57494.4828	0.0003	-0.0109	170
5	57494.5578	0.0002	-0.0138	170
6	57494.6424	0.0006	-0.0071	106
7	57494.7194	0.0004	-0.0080	182
8	57494.7982	0.0003	-0.0071	410
9	57494.8814	0.0003	-0.0018	326
10	57494.9628	0.0002	0.0017	217
15	57495.3530	0.0002	0.0023	102
16	57495.4307	0.0001	0.0021	180
17	57495.5085	0.0001	0.0020	197
18	57495.5878	0.0002	0.0033	391
19	57495.6645	0.0002	0.0022	392
20	57495.7434	0.0001	0.0032	460
21	57495.8200	0.0001	0.0018	484
22	57495.8982	0.0001	0.0021	419
23	57495.9750	0.0002	0.0010	293
25	57496.1284	0.0005	-0.0015	330
26	57496.2084	0.0005	0.0007	353
27	57496.2856	0.0002	-0.0001	134
28	57496.3642	0.0001	0.0006	95
29	57496.4418	0.0001	0.0003	187
30	57496.5188	0.0001	-0.0006	348
31	57496.5979	0.0001	0.0005	348
32	57496.6748	0.0001	-0.0004	299
33	57496.7526	0.0001	-0.0005	624
34	57496.8293	0.0001	-0.0018	704
35	57496.9067	0.0001	-0.0022	584
36	57496.9859	0.0002	-0.0010	286
40	57497.2971	0.0002	-0.0014	41
41	57497.3787	0.0006	0.0023	113
42	57497.4525	0.0004	-0.0019	86
43	57497.5309	0.0003	-0.0014	78
44	57497.6082	0.0003	-0.0019	76
46	57497.7658	0.0002	-0.0002	180
47	57497.8429	0.0002	-0.0011	218
48	57497.9192	0.0002	-0.0026	282
49	57497.9967	0.0002	-0.0031	191
51	57498.1561	0.0002	0.0005	302
52	57498.2309	0.0001	-0.0026	306
55	57498.4655	0.0002	-0.0018	189
56	57498.5424	0.0001	-0.0028	392
57	57498.6200	0.0002	-0.0030	320
58	57498.6971	0.0002	-0.0039	192
59	57498.7752	0.0002	-0.0037	289
60	57498.8524	0.0002	-0.0044	239

\*BJD-2400000.

 $^\dagger C = 2457496.4415 + 0.077913E$ . $^\ddagger$ Number of points used to determine the maximum.**Table S2.** Times of superhump maxima in ASASSN-16eg (continued).

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
61	57498.9302	0.0002	-0.0045	96
62	57499.0080	0.0003	-0.0047	65
68	57499.4765	0.0003	-0.0036	98
69	57499.5553	0.0002	-0.0027	101
70	57499.6331	0.0003	-0.0028	63
80	57500.4154	0.0004	0.0003	38
89	57501.1178	0.0004	0.0015	163
90	57501.1989	0.0005	0.0047	172
91	57501.2760	0.0002	0.0039	167
106	57502.4492	0.0005	0.0084	42
120	57503.5348	0.0004	0.0032	76
121	57503.6121	0.0004	0.0025	75
122	57503.6896	0.0005	0.0021	39
129	57504.2322	0.0002	-0.0006	172
130	57504.3097	0.0004	-0.0010	101
134	57504.6195	0.0004	-0.0028	70
135	57504.7011	0.0008	0.0008	42
159	57506.5620	0.0004	-0.0082	78
180	57508.1908	0.0013	-0.0156	30
181	57508.2681	0.0006	-0.0162	294

\*BJD-2400000.

 $^\dagger C = 2457496.4415 + 0.077913E$ . $^\ddagger$ Number of points used to determine the maximum.

**Table S3.** Times of superhump maxima in V1251 Cyg.

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
0	54763.3947	0.0002	-0.0055	48
1	54763.4681	0.0004	-0.0080	80
3	54763.6234	0.0014	-0.0047	39
4	54763.7010	0.0004	-0.0030	93
5	54763.7784	0.0003	-0.0016	89
12	54764.3114	0.0002	-0.0003	273
13	54764.3862	0.0004	-0.0014	296
14	54764.4629	0.0002	-0.0007	326
16	54764.6151	0.0002	-0.0005	28
17	54764.6918	0.0001	0.0003	220
18	54764.7670	0.0001	-0.0005	201
26	54765.3739	0.0004	-0.0012	108
27	54765.4514	0.0003	0.0003	131
28	54765.5271	0.0004	-0.0000	141
29	54765.6038	0.0003	0.0008	183
30	54765.6786	0.0002	-0.0004	174
31	54765.7539	0.0002	-0.0010	118
32	54765.8296	0.0003	-0.0013	40
38	54766.2831	0.0003	-0.0036	80
39	54766.3603	0.0003	-0.0024	143
40	54766.4363	0.0004	-0.0023	86
42	54766.5860	0.0006	-0.0045	20
43	54766.6621	0.0007	-0.0044	28
44	54766.7391	0.0011	-0.0033	29
45	54766.8149	0.0005	-0.0035	26
47	54766.9679	0.0007	-0.0025	127
49	54767.1186	0.0004	-0.0036	178
54	54767.5010	0.0003	-0.0011	99
56	54767.6526	0.0003	-0.0014	104
57	54767.7286	0.0003	-0.0014	106
60	54767.9571	0.0008	-0.0007	419
63	54768.1858	0.0004	0.0001	74
64	54768.2610	0.0003	-0.0006	122
65	54768.3374	0.0003	-0.0002	126
66	54768.4136	0.0003	0.0001	126
67	54768.4878	0.0006	-0.0018	79
69	54768.6412	0.0004	-0.0003	79
70	54768.7177	0.0004	0.0003	79
73	54768.9473	0.0011	0.0020	87
76	54769.1724	0.0004	-0.0008	46
77	54769.2510	0.0004	0.0018	86
78	54769.3265	0.0004	0.0014	87
79	54769.4022	0.0004	0.0011	80
80	54769.4758	0.0005	-0.0012	83
86	54769.9294	0.0007	-0.0034	268
87	54770.0064	0.0005	-0.0024	373
90	54770.2353	0.0004	-0.0014	81

\*BJD-2400000.

 $^\dagger C = 2454765.6790 + 0.0759607E$ . $^\ddagger$ Number of points used to determine the maximum.**Table S3.** Times of superhump maxima in V1251 Cyg (continued).

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
91	54770.3104	0.0004	-0.0022	76
92	54770.3875	0.0004	-0.0011	59
93	54770.4626	0.0005	-0.0019	58
113	54771.9752	0.0013	-0.0085	278
114	54772.0501	0.0003	-0.0096	157
117	54772.2791	0.0005	-0.0085	76
122	54772.6556	0.0005	-0.0118	79
123	54772.7308	0.0005	-0.0125	71
144	54774.3232	0.0006	-0.0154	136
145	54774.3968	0.0004	-0.0177	135
146	54774.4716	0.0006	-0.0189	46

\*BJD-2400000.

 $^\dagger C = 2454765.6790 + 0.0759607E$ . $^\ddagger$ Number of points used to determine the maximum.



**Table S4.** Times of superhump maxima in RZ Leo.

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
0	51901.1568	0.0009	-0.0201	110
1	51901.2307	0.0014	-0.0246	145
2	51901.3041	0.0007	-0.0298	148
5	51901.5444	0.0008	-0.0252	171
6	51901.6245	0.0006	-0.0236	185
13	51902.1980	0.0003	0.0003	208
14	51902.2755	0.0003	-0.0008	203
15	51902.3542	0.0003	-0.0006	171
18	51902.5902	0.0006	-0.0003	134
23	51902.9853	0.0005	0.0022	41
24	51903.0665	0.0003	0.0049	41
25	51903.1455	0.0006	0.0053	117
26	51903.2191	0.0003	0.0005	169
27	51903.2972	0.0004	-0.0000	205
28	51903.3772	0.0006	0.0014	108
39	51904.2464	0.0005	0.0068	83
40	51904.3248	0.0003	0.0066	114
51	51905.1840	0.0003	0.0019	183
52	51905.2611	0.0003	0.0006	176
53	51905.3400	0.0004	0.0009	146
64	51906.2083	0.0005	0.0054	150
65	51906.2856	0.0003	0.0041	215
66	51906.3637	0.0009	0.0037	122
76	51907.1514	0.0004	0.0061	146
77	51907.2294	0.0007	0.0055	108
84	51907.7799	0.0005	0.0064	35
85	51907.8599	0.0008	0.0078	17
86	51907.9387	0.0006	0.0080	17
89	51908.1751	0.0005	0.0089	149
90	51908.2559	0.0004	0.0111	285
91	51908.3329	0.0003	0.0096	257
99	51908.9644	0.0004	0.0128	40
100	51909.0404	0.0004	0.0103	40
111	51909.9016	0.0006	0.0076	33
112	51909.9750	0.0007	0.0025	40
113	51910.0559	0.0003	0.0049	42
128	51911.2277	0.0011	-0.0013	221
129	51911.3055	0.0014	-0.0021	219
153	51913.1815	0.0009	-0.0108	200
154	51913.2620	0.0007	-0.0089	219
155	51913.3423	0.0007	-0.0070	225
167	51914.2836	0.0022	-0.0082	64
168	51914.3615	0.0021	-0.0088	88

\*BJD-2400000.

 $^\dagger C = 2451901.175544 + 0.078533E$ . $^\ddagger$ Number of points used to determine the maximum.**Table S5.** Times of superhump maxima in BC UMa.

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
0	52674.1092	0.0005	0.0011	55
1	52674.1745	0.0002	0.0018	70
2	52674.2384	0.0002	0.0012	118
3	52674.3020	0.0003	0.0003	74
4	52674.3679	0.0002	0.0015	156
5	52674.4330	0.0003	0.0021	66
6	52674.4979	0.0002	0.0024	67
31	52676.1071	0.0005	-0.0027	57
32	52676.1718	0.0006	-0.0025	68
33	52676.2372	0.0006	-0.0017	150
34	52676.2992	0.0007	-0.0043	51
35	52676.3646	0.0003	-0.0035	117
36	52676.4291	0.0004	-0.0036	67
37	52676.4945	0.0005	-0.0027	61
47	52677.1405	0.0005	-0.0025	173
48	52677.2052	0.0005	-0.0024	181
49	52677.2683	0.0004	-0.0038	199
50	52677.3353	0.0006	-0.0014	135
52	52677.4652	0.0013	-0.0006	42
63	52678.1748	0.0008	-0.0014	107
65	52678.3023	0.0013	-0.0030	92
93	52680.1127	0.0006	-0.0006	178
95	52680.2403	0.0010	-0.0021	148
96	52680.3078	0.0014	0.0008	66
97	52680.3712	0.0008	-0.0004	81
98	52680.4368	0.0006	0.0006	133
99	52680.5002	0.0011	-0.0005	80
108	52681.0751	0.0017	-0.0069	80
109	52681.1407	0.0010	-0.0058	76
110	52681.2066	0.0002	-0.0045	316
113	52681.3981	0.0007	-0.0067	67
124	52682.1098	0.0020	-0.0053	102
127	52682.2962	0.0014	-0.0126	63
128	52682.3596	0.0014	-0.0138	85
139	52683.0692	0.0015	-0.0145	115
140	52683.1330	0.0022	-0.0153	159
141	52683.1961	0.0011	-0.0167	333
142	52683.2625	0.0005	-0.0149	262
143	52683.3230	0.0009	-0.0189	142
144	52683.3923	0.0016	-0.0143	87
145	52683.4547	0.0010	-0.0164	54
160	52684.4141	0.0014	-0.0256	44
161	52684.4808	0.0026	-0.0235	44

\*BJD-2400000.

 $^\dagger C = 2452674.2372 + 0.0645728E$ . $^\ddagger$ Number of points used to determine the maximum.

**Table S6.** Times of superhump maxima in MASTER J004527.

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
0	56554.3615	0.0004	-0.0151	95
4	56554.6863	0.0002	-0.0111	234
5	56554.7697	0.0003	-0.0079	147
6	56554.8514	0.0003	-0.0064	129
7	56554.9367	0.0008	-0.0013	74
12	56555.3362	0.0001	-0.0027	204
13	56555.4176	0.0001	-0.0015	287
14	56555.4968	0.0001	-0.0026	381
15	56555.5763	0.0001	-0.0033	312
16	56555.6567	0.0002	-0.0030	319
19	56555.8984	0.0003	-0.0019	62
20	56555.9783	0.0002	-0.0022	64
25	56556.3799	0.0001	-0.0016	280
26	56556.4608	0.0001	-0.0009	279
27	56556.5419	0.0002	0.0000	172
28	56556.6222	0.0001	0.0001	200
29	56556.7031	0.0002	0.0009	187
30	56556.7821	0.0003	-0.0004	188
31	56556.8632	0.0002	0.0005	203
32	56556.9438	0.0003	0.0009	63
36	56557.2658	0.0002	0.0021	312
40	56557.5857	0.0004	0.0012	62
41	56557.6658	0.0003	0.0011	69
42	56557.7471	0.0003	0.0023	61
43	56557.8253	0.0003	0.0003	71
44	56557.9070	0.0004	0.0018	71
47	56558.1487	0.0002	0.0029	120
48	56558.2291	0.0002	0.0031	212
49	56558.3103	0.0002	0.0041	237
50	56558.3881	0.0001	0.0018	187
51	56558.4681	0.0001	0.0015	196
52	56558.5481	0.0002	0.0013	223
53	56558.6279	0.0002	0.0009	255
54	56558.7087	0.0002	0.0015	139
55	56558.7886	0.0003	0.0013	62
60	56559.1900	0.0001	0.0016	210
61	56559.2689	0.0002	0.0004	263
62	56559.3492	0.0002	0.0005	163
63	56559.4307	0.0002	0.0018	128
64	56559.5097	0.0002	0.0005	170
65	56559.5889	0.0004	-0.0004	67
66	56559.6693	0.0002	-0.0002	206
67	56559.7488	0.0002	-0.0009	189
68	56559.8289	0.0003	-0.0010	74
69	56559.9088	0.0005	-0.0013	71
72	56560.1490	0.0002	-0.0017	110
73	56560.2292	0.0002	-0.0016	137

\*BJD-2400000.

 $^\dagger C = 2456556.3815 + 0.0801953E$ . $^\ddagger$ Number of points used to determine the maximum.**Table S6.** Times of superhump maxima in MASTER J004527 (continued).

$E$	max*	error	$O - C^\dagger$	$N^\ddagger$
77	56560.5501	0.0005	-0.0016	57
80	56560.7899	0.0005	-0.0023	61
81	56560.8685	0.0005	-0.0039	69
87	56561.3516	0.0007	-0.0020	104
88	56561.4289	0.0002	-0.0049	108
90	56561.5884	0.0007	-0.0058	69
91	56561.6684	0.0004	-0.0060	69
92	56561.7487	0.0007	-0.0059	53
93	56561.8290	0.0005	-0.0058	72
94	56561.9088	0.0006	-0.0061	71
98	56562.2279	0.0003	-0.0078	154
99	56562.3124	0.0004	-0.0035	126
102	56562.5482	0.0007	-0.0083	57
103	56562.6290	0.0006	-0.0077	64
104	56562.7083	0.0008	-0.0086	52
105	56562.7885	0.0007	-0.0086	64
106	56562.8679	0.0005	-0.0094	70
110	56563.1897	0.0004	-0.0084	74
111	56563.2715	0.0006	-0.0068	84
112	56563.3480	0.0005	-0.0105	139
113	56563.4279	0.0004	-0.0108	104
115	56563.5883	0.0010	-0.0108	63
116	56563.6714	0.0008	-0.0079	121
117	56563.7501	0.0007	-0.0094	88
118	56563.8278	0.0010	-0.0118	72
119	56563.9077	0.0012	-0.0122	71
122	56564.1450	0.0003	-0.0154	65
123	56564.2246	0.0009	-0.0160	84

\*BJD-2400000.

 $^\dagger C = 2456556.3815 + 0.0801953E$ . $^\ddagger$ Number of points used to determine the maximum.