

Supplemental Material for “Inversion symmetry breaking induced triply degenerate points in orderly arranged PtSeTe family materials”

R. C. Xiao,^{1,2,*} C. H. Cheung,^{3,*} P. L. Gong,⁴ W. J. Lu,^{1,†} J. G. Si,^{1,2} and Y. P. Sun^{1,5,6,‡}

¹Key Laboratory of Materials Physics, Institute of Solid State Physics, Chinese Academy of Sciences, Hefei 230031, China

²University of Science and Technology of China, Hefei 230026, China

³Graduate Institute of Applied Physics, National Taiwan University, Taipei 10617, Taiwan

⁴Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China

⁵High Magnetic Field Laboratory, Chinese Academy of Sciences, Hefei 230031, China

⁶Collaborative Innovation Center of Microstructures, Nanjing University, Nanjing 210093, China

I. CHARACTER TABLE OF C_{3v}

Character table for point group C_{3v} is shown in Table S1, and Mulliken symbols (A_1 , A_2 , E) for some irreducible representations are also listed.

TABLE S1: Character table for double point group C_{3v} .

$C_{3v}/3m$	\mathbb{I}	\mathbb{I}	$2C_3$	$2\bar{C}_3$	$3\sigma_v$	$3\bar{\sigma}_v$	Bases
Γ_1/A_1	1	1	1	1	1	1	z
Γ_2/A_2	1	1	1	1	-1	-1	S_z
Γ_3/E	2	2	-1	-1	0	0	(x, y)
Γ_4	2	-2	1	-1	0	0	$\{ 1/2, 1/2\rangle, 1/2, -1/2\rangle\}$
Γ_5	1	-1	-1	1	i	-i	$(3/2, -3/2\rangle - i 3/2, 3/2\rangle)$
Γ_6	1	-1	-1	1	-i	i	$(i 3/2, -3/2\rangle - 3/2, 3/2\rangle)$

Note: data are from Ref. 1. \mathbb{I} means identity, S_z stands for the z competent of an axial vector ($\mathbf{S} = \mathbf{r} \times \mathbf{p}$).

According to the above table:

$$\begin{cases} A_1 \otimes D^{1/2} = \Gamma_4, \\ A_2 \otimes D^{1/2} = \Gamma_4, \\ E \otimes D^{1/2} = \Gamma_4 \oplus \Gamma_5 \oplus \Gamma_6, \end{cases} \quad (\text{S1})$$

where $D^{1/2}$ is the representation matrix of spin. From Eq. S1, we can see that A_1/A_2 state transfers into Γ_4 , while E state splits into Γ_4 , Γ_5 , and Γ_6 states when spin orbit coupling (SOC) effects are considered. Γ_5 and Γ_6 transfer into each other under the time reversal operator T :

$$\begin{cases} T\Gamma_5 = -\Gamma_6, \\ T\Gamma_6 = \Gamma_5, \end{cases} \quad (\text{S2})$$

and the matrix of time reversal operator for the bases of (Γ_5, Γ_6) is

$$D(T) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} K, \quad (\text{S3})$$

where K is complex conjugate operator. Therefore, these two states are Kramers pair, and they are degenerate at time reversal symmetry point. Along the $\Gamma - A$ (Δ) path, we prefer Δ_4 , Δ_5 , and Δ_6 instead of Γ_4 , Γ_5 , and Γ_6 .

II. $k \cdot p$ DERIVATION

The little group of A point is $C_{3v} \otimes 1'$ ($1' = \{\mathbb{I}, T\}$). There are two generators of C_{3v} including a threefold rotational symmetry $C_{3z}(-x/2 - \sqrt{3}y/2, \sqrt{3}x/2 - y/2, z)$ and mirror symmetry $\sigma_{yz}(-x, y, z)$. Consequently, the related symmetry representation matrices under the bases of $\{\Delta_4, \Delta_5, \Delta_6\}$ are given by

$$D(C_{3z}) = \begin{pmatrix} e^{i\pi/3} & 0 \\ 0 & e^{-i\pi/3} \end{pmatrix} \oplus \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (\text{S4})$$

$$D(\sigma_{yz}) = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \oplus \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad (\text{S5})$$

$$D(T) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} K \oplus \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} K. \quad (\text{S6})$$

According to invariant theory:

$$D(R)H(\mathbf{k})D(R)^{-1} = H(R\mathbf{k}), \quad (\text{S7})$$

where R is a symmetry operator, the derived effective Hamiltonian around A point that considers first-order k terms for off-diagonal matrix elements and second-order k terms for diagonal matrix elements is:

$$H_{eff}(\mathbf{q}) = \varepsilon_0(\mathbf{q}) + \begin{bmatrix} M(\mathbf{q}) & iCq_+ & Aq_- & -A^*q_- \\ -iCq_- & M(\mathbf{q}) & -Aq_+ & -A^*q_+ \\ A^*q_+ & -A^*q_- & -M(\mathbf{q}) + Bq_z & 0 \\ -Aq_+ & -Aq_- & 0 & -M(\mathbf{q}) - Bq_z \end{bmatrix}, \quad (\text{S8})$$

where $\varepsilon_0(\mathbf{q}) = C_0 + C_1q_z^2 + C_2(q_x^2 + q_y^2)$, $M(\mathbf{q}) = M_0 - M_1q_z^2 - M_2(q_x^2 + q_y^2)$, $\mathbf{q} = \mathbf{k} - \mathbf{A}$, $q_+ = q_x + iq_y$, and $q_- = q_x - iq_y$. A is a complex number, while B and C are real numbers.

III. CHIRALITY OF EACH TDP

By using similar method that drives $k \cdot p$ effective Hamiltonian near A point, we can get the effective Hamiltonian near the TDPs. The 3×3 effective Hamiltonian around the TDP ($T1$) by Δ_4 and Δ_5 bands keeping only linear order terms for matrix elements is as follows:

$$H_{\Delta_4, \Delta_5} = \eta p_z + \begin{bmatrix} \alpha p_z & i\beta p_+ & \gamma p_- \\ -i\beta p_- & \alpha p_z & -\gamma p_+ \\ \gamma^* p_+ & -\gamma^* p_- & -\alpha p_z \end{bmatrix}, \quad (\text{S9})$$

where $\mathbf{p} = \mathbf{q} - \mathbf{q}_{T1} = (p_x, p_y, p_z)$, α , β and η are real numbers, and γ is a complex number.

The 2×2 Hamiltonian of the crossing point of $\Delta_{4\uparrow}$ and Δ_5 is:

$$H_{\Delta_{4\uparrow}, \Delta_5} = \eta p_z + \begin{bmatrix} \alpha p_z & \gamma p_- \\ \gamma^* p_+ & -\alpha p_z \end{bmatrix}, \quad (\text{S10})$$

and the effective Hamiltonian of the crossing point of $\Delta_{4\downarrow}$ and Δ_5 is:

$$H_{\Delta_{4\downarrow}, \Delta_5} = \eta p_z + \begin{bmatrix} \alpha p_z & -\gamma p_+ \\ -\gamma^* p_- & -\alpha p_z \end{bmatrix}. \quad (\text{S11})$$

Any 2×2 Hermitian matrix can be expanded by the Pauli matrices and the identity matrix:

$$H = \sum_i \sigma_0 v_{0i} p_i + \sum_{i,j} v_{ij} p_i \sigma_j, \quad (\text{S12})$$

where σ_0 is a 2×2 identity matrix, $i, j = x, y, z$, σ_j are one of three Pauli matrices. So Eq. S10 can be rewrote as:

$$H_{\Delta_{4\uparrow}, \Delta_5} = \eta p_z \sigma_0 + p_z \alpha \sigma_z + p_x \gamma_R \sigma_x + p_y \gamma_R \sigma_y - p_x \gamma_I \sigma_y + p_y \gamma_I \sigma_x, \quad (\text{S13})$$

and Eq. S11 can be rewrote as:

$$H_{\Delta_{4\downarrow}, \Delta_5} = \eta p_z \sigma_0 + p_z \alpha \sigma_z - p_x \gamma_R \sigma_x + p_y \gamma_R \sigma_y + p_x \gamma_I \sigma_y + p_y \gamma_I \sigma_x, \quad (\text{S14})$$

where γ_R and γ_I are the real and imaginary part of γ , respectively. So

$$V_{\Delta_{4\uparrow}, \Delta_5} \equiv [v_{ij}]_{\Delta_{4\uparrow}, \Delta_5} = \begin{bmatrix} \gamma_R & -\gamma_I & 0 \\ \gamma_I & \gamma_R & 0 \\ 0 & 0 & \alpha \end{bmatrix}, \quad (\text{S15})$$

$$V_{\Delta_{4\downarrow}, \Delta_5} \equiv [v_{ij}]_{\Delta_{4\downarrow}, \Delta_5} = \begin{bmatrix} -\gamma_R & \gamma_I & 0 \\ \gamma_I & \gamma_R & 0 \\ 0 & 0 & \alpha \end{bmatrix}. \quad (\text{S16})$$

According to Ref. 2, the chirality of a 2×2 Hamiltonian $\chi = \text{sgn}[\det(V)]$, so

$$\chi(\Delta_{4\uparrow}, \Delta_5) = -\chi(\Delta_{4\downarrow}, \Delta_5). \quad (\text{S17})$$

Similarly, the 3×3 effective Hamiltonian around the TDP by Δ_4 and Δ_6 bands is:

$$H_{\Delta_4, \Delta_6} = \eta' p_z + \begin{bmatrix} \alpha' p_z & i\beta' p_+ & \gamma' p_- \\ -i\beta' p_- & \alpha' p_z & \gamma' p_+ \\ \gamma'^* p_+ & \gamma'^* p_- & -\alpha' p_z \end{bmatrix}, \quad (\text{S18})$$

where α' , β' , and η' are real numbers and γ' is a complex number. Using similar method, we can prove that one TDP formed by Δ_4 and Δ_6 bands can also be viewed as two Weyl points with opposite chirality. Moreover, the results are invariant with the values of above $k \cdot p$ parameter. These results are similar with the TDPs induced by breaking time reversal symmetry in centrosymmetric system.³

* The authors contributed equally to this work.

† Electronic address: wjlu@issp.ac.cn

‡ Electronic address: ypsun@issp.ac.cn

¹ G. F. Koster, *Properties of the thirty-two point groups*, The MIT Press (1963).

² X. Wan, A. M. Turner, A. Vishwanath, and S. Y. Savrasov, *Phys. Rev. B* **83**, 205101 (2011).

³ C. H. Cheung, R. C. Xiao, M. C. Hsu, H. R. Fuh, Y. C. Lin, and C. R. Chang, arXiv:1709.07763.