

Supplementary Material: A new generation of effective core potentials from correlated calculations: 2nd row elements

M. Chandler Bennett^{1,2}, Guangming Wang¹, Abdulgani Annaberdiyev¹,
Cody A. Melton^{1,2}, Luke Shulenburger², and Lubos Mitas¹

*1) Department of Physics,
North Carolina State University, Raleigh,
North Carolina 27695-8202, USA*

and

*2) Sandia National Laboratories,
Albuquerque, New Mexico 87123, USA*

(Dated: September 7, 2018)

PACS numbers:

I. ATOMIC DATA

In Tables I-XVI, we show total energy components from various states of each of our ECPs from CCSD(T) calculations. All atomic calculations utilized the uncontracted aug-cc-pCVnZ basis sets where $n = \{T, Q, 5\}$. Each energy component, Hartree-Fock (SCF) and correlation, is extrapolated to the complete basis set (CBS) limit with the parameterized functions

$$E^{\text{SCF}}(n) = E_{\text{CBS}}^{\text{SCF}} + A \exp(-Bn) \quad (1)$$

and

$$E^{\text{Correlation}}(n) = E_{\text{CBS}}^{\text{Correlation}} + \frac{C}{(n + 3/8)^3} + \frac{D}{(n + 3/8)^5}, \quad (2)$$

respectively. The variables $A, B, C, D, E_{\text{CBS}}^{\text{SCF}}$ and $E_{\text{CBS}}^{\text{Correlation}}$ are fitting parameters.

TABLE I: Total energy components of various states of the Na atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^1$ 2S	-0.186167	-0.186202	-0.186202	-0.186203
[Ne] $3s^2$ 1S	-0.182241	-0.182308	-0.182308	-0.182313
Correlation	T	Q	5	CBS
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.024246	-0.024271	-0.02428	-0.02429

TABLE II: Total energy components of various states of the Na atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^1$ 2S	-47.355004	-47.356999	-47.357015	-47.357015
[Ne] 1S	-47.172935	-47.174928	-47.174944	-47.174944
[Ne] $3s^2$ 1S	-47.351159	-47.353181	-47.353197	-47.353197
Correlation	T	Q	5	CBS
[Ne] $3s^1$ 2S	-0.278836	-0.304079	-0.313965	-0.325551
[Ne] 1S	-0.273772	-0.298106	-0.307423	-0.318131
[Ne] $3s^2$ 1S	-0.302615	-0.327913	-0.337849	-0.349521

TABLE III: Total energy components of various states of the Mg atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^2$ 1S	-0.788370	-0.788345	-0.788370	-0.788370
[Ne] $3s^1$ 2S	-0.544766	-0.544744	-0.544763	-0.544763
Correlation	T	Q	5	CBS
[Ne] $3s^2$ 1S	-0.034907	-0.034994	-0.035031	-0.035077
[Ne] $3s^1$ 2S				

TABLE IV: Total energy components of various states of the Mg atom using our ccECP[He]

SCF		T	Q	5	CBS
[Ne] $3s^2$	1S	-62.927052	-62.927327	-62.927412	-62.927451
[Ne]	1S	-62.142427	-62.142705	-62.142790	-62.142828
[Ne] $3s^1$	2S	-62.684058	-62.684333	-62.684414	-62.684447
Correlation		T	Q	5	CBS
[Ne] $3s^2$	1S	-0.315920	-0.342027	-0.352077	-0.363681
[Ne]	1S	-0.270494	-0.294814	-0.304048	-0.314584
[Ne] $3s^1$	2S	-0.279223	-0.304733	-0.314498	-0.325718

TABLE V: Total energy components of various states of the Al atom using our ccECP[Ne]

SCF		T	Q	5	CBS
[Ne] $3s^23p^1$	2P	-1.876699	-1.876990	-1.876973	-1.876999
[Ne] $3s^1$	2S	-1.030941	-1.030946	-1.030933	-1.030949
[Ne] $3s^2$	1S	-1.674980	-1.675006	-1.674996	-1.675010
[Ne] $3s^23p^2$	3P	-1.878405	-1.878616	-1.878601	-1.878623
Correlation		T	Q	5	CBS
[Ne] $3s^23p^1$	2P	-0.058680	-0.059582	-0.059783	-0.059869
[Ne] $3s^1$	2S				
[Ne] $3s^2$	1S	-0.041715	-0.042057	-0.042137	-0.042175
[Ne] $3s^23p^2$	3P	-0.072470	-0.073840	-0.074155	-0.074305

TABLE VI: Total energy components of various states of the Al atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^23p^1$ 2P	-80.990584	-80.992910	-80.993046	-80.993054
[Ne] 1S	-79.112214	-79.114296	-79.114427	-79.114435
[Ne] $3s^1$ 2S	-80.144308	-80.146349	-80.146470	-80.146478
[Ne] $3s^2$ 1S	-80.789130	-80.791175	-80.791301	-80.791309
[Ne] $3s^23p^2$ 3P	-80.992087	-80.994343	-80.994480	-80.994489
Correlation	T	Q	5	CBS
[Ne] $3s^23p^1$ 2P	-0.305640	-0.355044	-0.373996	-0.395813
[Ne] 1S	-0.231923	-0.278793	-0.297044	-0.318327
[Ne] $3s^1$ 2S	-0.243946	-0.291640	-0.310130	-0.331614
[Ne] $3s^2$ 1S	-0.289386	-0.338035	-0.356793	-0.378485
[Ne] $3s^23p^2$ 3P	-0.319051	-0.368907	-0.387970	-0.409853

TABLE VII: Total energy components of various states of the Si atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^23p^2$ 3P	-3.671951	-3.672403	-3.672394	-3.672423
[Ne] $3s^1$ 2S	-1.639761	-1.639759	-1.639758	-1.639762
[Ne] $3s^2$ 1S	-2.813851	-2.813866	-2.813855	-2.813870
[Ne] $3s^23p^1$ 2P	-3.390158	-3.390719	-3.390718	-3.390749
[Ne] $3s^23p^3$ 4S	-3.707923	-3.708178	-3.708158	-3.708187
Correlation	T	Q	5	CBS
[Ne] $3s^23p^2$ 3P	-0.085857	-0.087646	-0.088188	-0.088666
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.047828	-0.048152	-0.048263	-0.048376
[Ne] $3s^23p^1$ 2P	-0.068862	-0.069816	-0.070130	-0.070437
[Ne] $3s^23p^3$ 4S	-0.100941	-0.103501	-0.104256	-0.104894

TABLE VIII: Total energy components of various states of the Si atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^2 3p^2$ 3P	-101.622367	-101.625367	-101.625788	-101.625857
[Ne] 1S	-97.946272	-97.948793	-97.949212	-97.949296
[Ne] $3s^1$ 2S	-99.589327	-99.591835	-99.592241	-99.592319
[Ne] $3s^2$ 1S	-100.765362	-100.767880	-100.768268	-100.768339
[Ne] $3s^2 3p^1$ 2P	-101.341340	-101.344430	-101.344845	-101.344910
[Ne] $3s^2 3p^3$ 4S	-101.657754	-101.660560	-101.660972	-101.661043
Correlation	T	Q	5	CBS
[Ne] $3s^2 3p^2$ 3P	-0.336020	-0.386298	-0.405726	-0.428232
[Ne] 1S	-0.232488	-0.278592	-0.296671	-0.317884
[Ne] $3s^1$ 2S	-0.246107	-0.293177	-0.311571	-0.333091
[Ne] $3s^2$ 1S	-0.298506	-0.346684	-0.365457	-0.387362
[Ne] $3s^2 3p^1$ 2P	-0.319161	-0.368422	-0.387552	-0.409811
[Ne] $3s^2 3p^3$ 4S	-0.351145	-0.402192	-0.421832	-0.444501

TABLE IX: Total energy components of various states of the P atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^2 3p^3$ 4S	-6.340560	-6.340906	-6.340875	-6.340917
[Ne] $3s^1$ 2S	-2.362967	-2.362961	-2.362981	-2.363013
[Ne] $3s^2$ 1S	-4.186007	-4.185996	-4.186002	-4.186015
[Ne] $3s^2 3p^1$ 2P	-5.265263	-5.265481	-5.265501	-5.265510
[Ne] $3s^2 3p^2$ 3P	-5.970169	-5.970641	-5.970646	-5.970670
[Ne] $3s^2 3p^4$ 3P	-6.320061	-6.320225	-6.320176	-6.320236
Correlation	T	Q	5	CBS
[Ne] $3s^2 3p^3$ 4S	-0.112732	-0.115948	-0.116922	-0.117781
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.053463	-0.053785	-0.053895	-0.054008
[Ne] $3s^2 3p^1$ 2P	-0.077685	-0.078756	-0.079108	-0.079454
[Ne] $3s^2 3p^2$ 3P	-0.097219	-0.099311	-0.099955	-0.100534
[Ne] $3s^2 3p^4$ 3P	-0.155101	-0.161173	-0.162879	-0.164211

TABLE X: Total energy components of various states of the P atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^23p^3$ 4S	-125.253282	-125.258161	-125.258394	-125.258405
[Ne] 1S	-118.899156	-118.903679	-118.903877	-118.903886
[Ne] $3s^1$ 2S	-121.271178	-121.275661	-121.275885	-121.275897
[Ne] $3s^2$ 1S	-123.100459	-123.104903	-123.105109	-123.105119
[Ne] $3s^23p^1$ 2P	-124.180512	-124.185217	-124.185449	-124.185461
[Ne] $3s^23p^2$ 3P	-124.884427	-124.889414	-124.889658	-124.889670
[Ne] $3s^23p^4$ 3P	-125.233338	-125.238037	-125.238254	-125.238264
Correlation	T	Q	5	CBS
[Ne] $3s^23p^3$ 4S	-0.365923	-0.418003	-0.438225	-0.461752
[Ne] 1S	-0.233066	-0.278627	-0.296756	-0.318285
[Ne] $3s^1$ 2S	-0.247871	-0.294566	-0.313051	-0.334912
[Ne] $3s^2$ 1S	-0.306293	-0.354270	-0.373190	-0.395491
[Ne] $3s^23p^1$ 2P	-0.330315	-0.379711	-0.399104	-0.421880
[Ne] $3s^23p^2$ 3P	-0.349999	-0.400799	-0.420623	-0.443787
[Ne] $3s^23p^4$ 3P	-0.408274	-0.463170	-0.484121	-0.508132

TABLE XI: Total energy components of various states of the S atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^2 3p^4$ 3P	-9.917772	-9.918088	-9.918140	-9.918156
[Ne] $3s^1$ 2S	-3.199330	-3.199320	-3.199373	-3.199413
[Ne] $3s^2$ 1S	-5.787262	-5.787214	-5.787263	-5.787312
[Ne] $3s^2 3p^1$ 2P	-7.490378	-7.490458	-7.490604	-7.490623
[Ne] $3s^2 3p^2$ 3P	-8.745383	-8.745651	-8.745767	-8.745853
[Ne] $3s^2 3p^3$ 4S	-9.587305	-9.587807	-9.587895	-9.587914
[Ne] $3s^2 3p^5$ 2P	-9.951496	-9.951654	-9.951678	-9.951687
Correlation	T	Q	5	CBS
[Ne] $3s^2 3p^4$ 3P	-0.165776	-0.173557	-0.175915	-0.177994
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.058888	-0.059247	-0.059376	-0.059518
[Ne] $3s^2 3p^1$ 2P	-0.085847	-0.087076	-0.087513	-0.087980
[Ne] $3s^2 3p^2$ 3P	-0.107509	-0.109937	-0.110748	-0.111559
[Ne] $3s^2 3p^3$ 4S	-0.124018	-0.127843	-0.129087	-0.130294
[Ne] $3s^2 3p^5$ 2P	-0.203222	-0.214387	-0.217619	-0.220276

TABLE XII: Total energy components of various states of the S atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^23p^4$ 3P	-151.909927	-151.916064	-151.916369	-151.916385
[Ne] 1S	-141.960834	-141.966569	-141.966790	-141.966799
[Ne] $3s^1$ 2S	-145.177455	-145.183135	-145.183423	-145.183438
[Ne] $3s^2$ 1S	-147.778004	-147.783595	-147.783878	-147.783893
[Ne] $3s^23p^1$ 2P	-149.483717	-149.489469	-149.489861	-149.489890
[Ne] $3s^23p^2$ 3P	-150.738321	-150.744306	-150.744672	-150.744696
[Ne] $3s^23p^3$ 4S	-151.578352	-151.584628	-151.584970	-151.584989
[Ne] $3s^23p^5$ 2P	-151.943481	-151.949466	-151.949744	-151.949757
Correlation	T	Q	5	CBS
[Ne] $3s^23p^4$ 3P	-0.420401	-0.477685	-0.499872	-0.525628
[Ne] 1S	-0.233065	-0.278340	-0.296665	-0.318728
[Ne] $3s^1$ 2S	-0.248757	-0.295307	-0.314039	-0.336489
[Ne] $3s^2$ 1S	-0.312562	-0.360586	-0.379826	-0.402804
[Ne] $3s^23p^1$ 2P	-0.339291	-0.389051	-0.408907	-0.432544
[Ne] $3s^23p^2$ 3P	-0.361203	-0.412731	-0.433178	-0.457406
[Ne] $3s^23p^3$ 4S	-0.378375	-0.431623	-0.452638	-0.477427
[Ne] $3s^23p^5$ 2P	-0.458354	-0.518979	-0.542040	-0.568391

TABLE XIII: Total energy components of various states of the Cl atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^2 3p^5$ 2P	-14.688897	-14.689151	-14.689385	-14.689386
[Ne] $3s^1$ 2S	-4.153414	-4.153407	-4.153487	-4.153506
[Ne] $3s^2$ 1S	-7.623920	-7.623836	-7.623936	-7.623950
[Ne] $3s^2 3p^1$ 2P	-10.068949	-10.069158	-10.069416	-10.069416
[Ne] $3s^2 3p^2$ 3P	-11.995324	-11.995572	-11.995854	-11.995856
[Ne] $3s^2 3p^3$ 4S	-13.435190	-13.435587	-13.435816	-13.435820
[Ne] $3s^2 3p^4$ 3P	-14.255149	-14.255690	-14.255918	-14.256085
[Ne] $3s^2 3p^6$ 1S	-14.784864	-14.784943	-14.785142	-14.785146
Correlation	T	Q	5	CBS
[Ne] $3s^2 3p^5$ 2P	-0.214761	-0.228169	-0.232225	-0.235794
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.064175	-0.064536	-0.064686	-0.064870
[Ne] $3s^2 3p^1$ 2P	-0.093905	-0.095211	-0.095684	-0.096199
[Ne] $3s^2 3p^2$ 3P	-0.117691	-0.120313	-0.121206	-0.122119
[Ne] $3s^2 3p^3$ 4S	-0.135384	-0.139587	-0.140982	-0.142366
[Ne] $3s^2 3p^4$ 3P	-0.179033	-0.187874	-0.190649	-0.193217
[Ne] $3s^2 3p^6$ 1S	-0.247496	-0.264870	-0.269980	-0.274288

TABLE XIV: Total energy components of various states of the Cl atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^23p^5$ 2P	-181.606009	-181.611999	-181.612377	-181.612402
[Ne] 1S	-166.876313	-166.881725	-166.882022	-166.882039
[Ne] $3s^1$ 2S	-171.052574	-171.057971	-171.058355	-171.058385
[Ne] $3s^2$ 1S	-174.540922	-174.546208	-174.546602	-174.546634
[Ne] $3s^23p^1$ 2P	-176.989287	-176.994921	-176.995449	-176.995504
[Ne] $3s^23p^2$ 3P	-178.914799	-178.920519	-178.921056	-178.921111
[Ne] $3s^23p^3$ 4S	-180.351450	-180.357396	-180.357852	-180.357890
[Ne] $3s^23p^4$ 3P	-181.172702	-181.178912	-181.179310	-181.179337
[Ne] $3s^23p^6$ 1S	-181.700830	-181.706653	-181.706998	-181.707020
Correlation	T	Q	5	CBS
[Ne] $3s^23p^5$ 2P	-0.472345	-0.535132	-0.559372	-0.587434
[Ne] 1S	-0.234937	-0.279240	-0.297555	-0.319976
[Ne] $3s^1$ 2S	-0.251472	-0.297158	-0.315912	-0.338747
[Ne] $3s^2$ 1S	-0.320306	-0.367597	-0.386907	-0.410321
[Ne] $3s^23p^1$ 2P	-0.349504	-0.398752	-0.418757	-0.442915
[Ne] $3s^23p^2$ 3P	-0.373428	-0.424710	-0.445416	-0.470303
[Ne] $3s^23p^3$ 4S	-0.391882	-0.445213	-0.466621	-0.492226
[Ne] $3s^23p^4$ 3P	-0.435816	-0.493986	-0.516894	-0.543868
[Ne] $3s^23p^6$ 1S	-0.506041	-0.572752	-0.598045	-0.626863

TABLE XV: Total energy components of various states of the Ar atom using our ccECP[Ne]

SCF	T	Q	5	CBS
[Ne] $3s^23p^6$ 1S	-20.778934	-20.779555	-20.779573	-20.779601
[Ne] $3s^1$ 2S	-5.207475	-5.207483	-5.207610	-5.207618
[Ne] $3s^2$ 1S	-9.665000	-9.664896	-9.665081	-9.665103
[Ne] $3s^23p^1$ 2P	-12.964117	-12.964630	-12.964915	-12.964920
[Ne] $3s^23p^2$ 3P	-15.676104	-15.676707	-15.677049	-15.677055
[Ne] $3s^23p^3$ 4S	-17.831518	-17.832105	-17.832395	-17.832680
[Ne] $3s^23p^4$ 3P	-19.266038	-19.266814	-19.266949	-19.266979
[Ne] $3s^23p^5$ 2P	-20.234894	-20.235814	-20.235893	-20.235923
Correlation	T	Q	5	CBS
[Ne] $3s^23p^6$ 1S	-0.259079	-0.279406	-0.285173	-0.289756
[Ne] $3s^1$ 2S				
[Ne] $3s^2$ 1S	-0.069503	-0.069933	-0.070038	-0.070099
[Ne] $3s^23p^1$ 2P	-0.101947	-0.103367	-0.103790	-0.104156
[Ne] $3s^23p^2$ 3P	-0.127850	-0.130675	-0.131528	-0.132275
[Ne] $3s^23p^3$ 4S	-0.146823	-0.151385	-0.152764	-0.153973
[Ne] $3s^23p^4$ 3P	-0.192939	-0.202715	-0.205588	-0.208007
[Ne] $3s^23p^5$ 2P	-0.229486	-0.244616	-0.248974	-0.252528

TABLE XVI: Total energy components of various states of the Ar atom using our ccECP[He]

SCF	T	Q	5	CBS
[Ne] $3s^23p^6$ 1S	-214.882779	-214.891228	-214.891473	-214.891481
[Ne] 1S	-194.015736	-194.023303	-194.023467	-194.023471
[Ne] $3s^1$ 2S	-199.262254	-199.269806	-199.270110	-199.270123
[Ne] $3s^2$ 1S	-203.750624	-203.758010	-203.758366	-203.758384
[Ne] $3s^23p^1$ 2P	-207.059126	-207.067196	-207.067647	-207.067674
[Ne] $3s^23p^2$ 3P	-209.774739	-209.782966	-209.783460	-209.783491
[Ne] $3s^23p^3$ 4S	-211.929923	-211.938199	-211.938633	-211.938657
[Ne] $3s^23p^4$ 3P	-213.368960	-213.377490	-213.377782	-213.377792
[Ne] $3s^23p^5$ 2P	-214.339213	-214.347933	-214.348210	-214.348219
Correlation	T	Q	5	CBS
[Ne] $3s^23p^6$ 1S	-0.516297	-0.586767	-0.613131	-0.642806
[Ne] 1S	-0.234846	-0.279060	-0.297392	-0.319883
[Ne] $3s^1$ 2S	-0.251903	-0.297609	-0.316402	-0.339315
[Ne] $3s^2$ 1S	-0.325612	-0.373081	-0.392439	-0.415887
[Ne] $3s^23p^1$ 2P	-0.356990	-0.406593	-0.426698	-0.450934
[Ne] $3s^23p^2$ 3P	-0.382642	-0.434514	-0.455381	-0.480384
[Ne] $3s^23p^3$ 4S	-0.402186	-0.456407	-0.478050	-0.503818
[Ne] $3s^23p^4$ 3P	-0.448215	-0.507975	-0.531290	-0.558527
[Ne] $3s^23p^5$ 2P	-0.485495	-0.550750	-0.575657	-0.604203

II. MOLECULAR DATA

TABLE XVII: All-electron (AE) UCCSD(T) Na2 ground state ($^1\Sigma_g$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	$0.737(13)^\dagger{}^a$	0.730(6)	-0.012(9)	-0.016(9)	-0.011(9)	-0.010(9)	-0.013(9)	0.015(9)	-0.002(10)
$\omega_e(\text{cm}^{-1})$	139.9^a	146(3)	-10(5)	-11(5)	-10(5)	-10(5)	-10(5)	-1(5)	-1(5)
$r_e(\text{\AA})$	3.079^b	3.069(18)	0.116(28)	0.103(28)	0.099(28)	0.094(28)	0.098(28)	-0.007(26)	0.005(29)
Potential Energy Surface									
U(2.20)		-0.0369	0.1300	0.0868	0.0887	0.0806	0.0843	-0.0460	0.00494
U(2.45)		-0.4296	0.0971	0.0773	0.0742	0.0686	0.0730	-0.0288	0.00487
U(2.70)		-0.6443	0.0628	0.0557	0.0500	0.0464	0.0510	-0.0201	0.00442
U(2.95)		-0.7324	0.0315	0.0322	0.0260	0.0241	0.0280	-0.0152	0.00528
U(3.20)		-0.7350	0.0056	0.0104	0.0047	0.0043	0.0070	-0.0133	-0.00323
U(3.45)		-0.6841	-0.0141	-0.0081	-0.0126	-0.0125	-0.0105	-0.0134	-0.00253

† Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XVIII: All-electron (AE) UCCSD(T) NaO ground state ($^2\Sigma$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	$2.67^\dagger{}^a$	2.52(2)	0.05(3)	0.06(2)	0.02(3)	-0.00(2)	0.01(2)	-0.02(3)	-0.01(3)
$\omega_e(\text{cm}^{-1})$	526^a	538(6)	-13(10)	-50(6)	-19(10)	-63(6)	-37(7)	-12(8)	0(8)
$r_e(\text{\AA})$	1.95^a	1.964(5)	0.044(9)	-0.012(5)	0.033(8)	-0.016(5)	0.007(6)	0.006(7)	0.001(6)
Potential Energy Surface									
U(1.475)		-0.0305	0.4054	-0.7805	0.1843	-0.8672	-0.3741	-0.0486	0.0252
U(1.725)		-2.1531	0.0936	-0.1597	0.0563	-0.1069	-0.0414	0.0124	0.0134
U(1.975)		-2.4941	-0.0390	-0.0732	-0.0169	-0.0127	-0.0181	0.0180	0.0090
U(2.225)		-2.2604	-0.0964	-0.0902	-0.0640	-0.0458	-0.0472	0.0047	0.0068

† Sum of experimental dissociation and zero-point energies

a Reference [2]

III. CL AND AR ALTERNATIVES

In Table XXXI, we share the alternative Ne-core ECP parameters for Cl and Ar which were generated with higher weights on the spectral component of the objective function. Additionally, the UCCSD(T) spectral errors and UCCSD(T) molecular errors from all-electron results are also provided. For Cl, the spectral errors are given in Table XXXII while the molecular errors are given in Figure 1 and Tables XXXIII and XXXIV. And for Ar, the spectral errors are given in Table XXXV while the molecular errors are given in Figure 2 and Table XXXVI. In the figures and tables, we have labeled these alternative ECPs as ccECP[Ne].S.

TABLE XIX: All-electron (AE) UCCSD(T) MgO ground state ($^1\Sigma^+$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	$2.5(2)^\dagger$ ^a	2.616(8)	0.01(1)	0.09(1)	0.14(1)	-0.02(1)	0.04(1)	-0.03(1)	0.00(1)
$\omega_e(\text{cm}^{-1})$	785.2^b	780(6)	-14(8)	-39(8)	-7(8)	-43(9)	-17(8)	-33(8)	-1(9)
$r_e(\text{\AA})$	1.749^c	1.738(2)	0.017(3)	0.005(3)	-0.020(3)	0.004(4)	-0.007(3)	0.007(3)	0.000(3)
Potential Energy Surface									
U(1.4)		-0.0876	0.2135	-0.2493	-0.6121	-0.2685	-0.3327	-0.1149	0.0013
U(1.5)		-1.5908	0.1230	-0.0830	-0.3654	-0.0738	-0.1695	-0.0054	0.0013
U(1.6)		-2.3414	0.0581	-0.0189	-0.2259	0.0012	-0.0844	0.0319	0.0016
U(1.7)		-2.6140	0.0113	-0.0037	-0.1484	0.0203	-0.0423	0.0368	0.0019
U(1.8)		-2.5929	-0.0230	-0.0132	-0.1096	0.0124	-0.0278	0.0239	0.0014

[†] Sum of experimental dissociation and zero-point energies

^a Reference [3]

^b Reference [4]

^c Reference [2]

TABLE XX: All-electron (AE) UCCSD(T) Al₂ ground state ($^3\Sigma_g$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs. All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	1.34(13) ^{† a}	1.423(3)	0.002(5)	0.014(5)	0.035(5)	0.010(5)	0.048(5)	0.009(5)	0.003(5)
$\omega_e(\text{cm}^{-1})$	285.8 ^b	285(1)	-1(2)	0(2)	2(2)	-1(2)	6(2)	-1(2)	0(2)
$r_e(\text{\AA})$	2.701 ^c	2.692(3)	0.014(4)	0.005(4)	-0.004(4)	0.004(4)	-0.021(4)	-0.005(4)	-0.000(4)
Potential Energy Surface									
U(2.1)		0.0717	0.0817	0.0101	-0.0473	-0.0009	-0.1253	0.0103	-0.0034
U(2.3)		-0.9124	0.0392	0.0002	-0.0409	-0.0007	-0.0900	0.0039	-0.0035
U(2.5)		-1.3265	0.0129	-0.0094	-0.0383	-0.0063	-0.0657	-0.0043	-0.0034
U(2.7)		-1.4265	-0.0023	-0.0154	-0.0359	-0.0113	-0.0477	-0.0102	-0.0033
U(2.9)		-1.3609	-0.0102	-0.0177	-0.0326	-0.0138	-0.0335	-0.0129	-0.0031

[†] Sum of experimental dissociation and zero-point energies

^a Reference [3]

^b Reference [4]

^c Reference [5]

TABLE XXI: All-electron (AE) UCCSD(T) AIO ground state ($^2\Sigma$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Potential Energy Surface									
$D_e(\text{eV})$	5.26(9) ^{† a}	5.28(1)	-0.02(1)	0.01(1)	0.06(1)	-0.24(1)	0.07(1)	0.04(1)	-0.01(1)
$\omega_e(\text{cm}^{-1})$	979.5 ^b	1028(5)	-4(7)	-36(6)	-26(6)	-90(6)	-12(6)	-36(6)	6(6)
$r_e(\text{\AA})$	1.618 ^c	1.611(2)	0.007(3)	0.004(2)	-0.002(2)	0.021(2)	-0.009(2)	0.004(2)	-0.003(2)
Potential Energy Surface									
U(1.22)		1.3650	0.2753	-0.4581	-0.6482	-0.3602	-0.7520	-0.4704	-0.0202
U(1.32)		-2.3179	0.1699	-0.1595	-0.3070	0.1097	-0.4055	-0.1652	-0.0153
U(1.42)		-4.2349	0.0980	-0.0432	-0.1495	0.2764	-0.2200	-0.0448	-0.0104
U(1.52)		-5.0781	0.0494	-0.0084	-0.0800	0.2913	-0.1202	-0.0064	-0.0064
U(1.62)		-5.2884	0.0179	-0.0056	-0.0508	0.2437	-0.0654	-0.0005	0.0209

[†] Sum of experimental dissociation and zero-point energies

^a Reference [3]

^b Reference [4]

^c Reference [2]

TABLE XXII: All-electron (AE) UCCSD(T) Si2 ground state ($^3\Sigma_g$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	3.239 ^{† a}	3.258(4)	-0.012(6)	-0.004(6)	0.015(6)	0.007(6)	0.085(6)	0.006(6)	0.011(6)
$\omega_e(\text{cm}^{-1})$	489.7 ^a	506(2)	-3(3)	-4(3)	-2(3)	-4(3)	9(3)	-4(3)	0(3)
$r_e(\text{\AA})$	2.246 ^b	2.242(2)	0.009(2)	0.004(3)	-0.004(3)	0.001(3)	-0.023(2)	0.003(2)	0.001(3)
Potential Energy Surface									
U(1.750)		0.1044	0.1444	-0.0001	-0.1319	-0.0554	-0.3778	-0.0134	-0.0011
U(1.875)		-1.7121	0.0935	0.0145	-0.0711	-0.0221	-0.2633	0.0072	-0.0052
U(2.000)		-2.7013	0.0563	0.0139	-0.0401	-0.0102	-0.1822	0.0112	-0.0060
U(2.125)		-3.1540	0.0298	0.0083	-0.0243	-0.0079	-0.1236	0.0087	-0.0076
U(2.250)		-3.2654	0.0113	0.0023	-0.0159	-0.0089	-0.0804	0.0047	-0.0142
U(2.375)		-3.1650	-0.0013	-0.0025	-0.0110	-0.0103	-0.0481	0.0010	-0.0138

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XXIII: All-electron (AE) UCCSD(T) SiO ground state ($^1\Sigma$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	8.313(87) ^{† a}	8.33(2)	-0.04(2)	-0.02(2)	0.02(2)	-0.16(2)	0.10(2)	0.01(2)	0.01(2)
$\omega_e(\text{cm}^{-1})$	1259 ^a	1278(7)	-6(10)	-24(8)	-25(8)	-52(8)	-1(9)	-21(9)	1(10)
$r_e(\text{\AA})$	1.510 ^b	1.509(2)	0.005(3)	0.000(2)	-0.004(2)	0.005(2)	-0.013(2)	-0.001(2)	0.000(2)
Potential Energy Surface									
U(1.15)		-0.1504	0.2614	-0.3946	-0.7175	-0.3682	-0.9623	-0.3928	-0.0212
U(1.25)		-4.9302	0.1693	-0.1252	-0.3053	0.0205	-0.5351	-0.1373	-0.0186
U(1.35)		-7.2762	0.1053	-0.0272	-0.1258	0.1447	-0.2942	-0.0390	-0.0146
U(1.45)		-8.1833	0.0609	0.0012	-0.0498	0.1551	-0.1536	-0.0055	-0.0111
U(1.55)		-8.2557	0.0300	0.0037	-0.0194	0.1264	-0.0695	0.0022	-0.0086
U(1.65)		-7.8612	0.0084	-0.0019	-0.0084	0.0913	-0.0184	0.0005	-0.0070

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [6]

TABLE XXIV: All-electron (AE) UCCSD(T) P2 ground state ($^1\Sigma_g$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	5.082(22) ^{† a}	4.96(2)	-0.03(2)	-0.06(2)	-0.03(2)	-0.03(2)	-0.13(2)	-0.05(2)	0.01(2)
$\omega_e(\text{cm}^{-1})$	769.5 ^a	769(8)	-7(10)	-11(11)	-7(11)	-8(11)	-15(11)	-9(11)	0(11)
$r_e(\text{\AA})$	1.893 ^b	1.887(3)	0.008(4)	0.005(4)	-0.004(4)	0.001(4)	0.002(4)	0.003(4)	0.001(4)
Potential Energy Surface									
U(1.55)		-1.2900	0.1760	0.0708	-0.1612	-0.0236	0.0331	0.0354	-0.0046
U(1.65)		-3.4331	0.1197	0.0846	-0.0555	0.0191	0.1027	0.0571	-0.0093
U(1.75)		-4.5340	0.0767	0.0774	-0.0006	0.0330	0.1281	0.0580	-0.0122
U(1.85)		-4.9504	0.0442	0.0629	0.0263	0.0336	0.1311	0.0505	-0.0122
U(1.95)		-4.9286	0.0197	0.0475	0.0379	0.0288	0.1234	0.0406	-0.0130

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XXV: All-electron (AE) UCCSD(T) PO ground state ($^2\Pi$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	6.16(13) ^{† a}	6.14(1)	-0.03(2)	-0.03(2)	-0.01(2)	-0.10(2)	-0.09(2)	-0.03(2)	0.01(2)
$\omega_e(\text{cm}^{-1})$	1233.3 ^b	1249(7)	-7(10)	-11(11)	-11(11)	-19(11)	-14(11)	-9(11)	0(10)
$r_e(\text{\AA})$	1.476 ^c	1.472(1)	0.004(2)	0.001(2)	-0.004(2)	0.003(2)	-0.002(2)	0.000(2)	0.000(2)
Potential Energy Surface									
U(1.2)		-1.9498	0.1467	-0.0306	-0.2437	0.0845	-0.1000	-0.0560	-0.0233
U(1.3)		-4.8025	0.0901	0.0229	-0.0790	0.1248	0.0246	0.0052	-0.0156
U(1.4)		-5.9524	0.0514	0.0319	-0.0108	0.1154	0.0732	0.0235	-0.0101
U(1.5)		-6.1369	0.0246	0.0270	0.0159	0.0944	0.0882	0.0258	-0.0068
U(1.6)		-5.7942	0.0059	0.0192	0.0248	0.0739	0.0881	0.0225	-0.0050

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [4]

^c Reference [2]

TABLE XXVI: All-electron (AE) UCCSD(T) S2 ground state ($^3\Sigma_g$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	4.408(3) ^{† a}	4.429(7)	-0.02(1)	0.00(1)	0.01(1)	0.01(1)	0.12(1)	-0.02(1)	0.02(1)
$\omega_e(\text{cm}^{-1})$	699.5 ^a	717(3)	-4(5)	-3(5)	-3(5)	-3(5)	17(5)	-5(5)	1(5)
$r_e(\text{\AA})$	1.889 ^b	1.886(2)	0.006(2)	0.001(2)	-0.006(2)	0.001(2)	-0.025(2)	0.003(2)	-0.001(2)
Potential Energy Surface									
U(1.5)		0.3808	0.1600	-0.0256	-0.2756	-0.0329	-0.7207	0.0272	-0.0263
U(1.6)		-2.2587	0.1076	-0.0055	-0.1444	-0.0117	-0.4797	0.0354	-0.0258
U(1.7)		-3.6725	0.0692	-0.0013	-0.0705	-0.0081	-0.3102	0.0306	-0.0241
U(1.8)		-4.3034	0.0412	-0.0023	-0.0284	-0.0100	-0.1899	0.0223	-0.0221
U(1.9)		-4.4421	0.0211	-0.0041	-0.0043	-0.0127	-0.1042	0.0143	-0.0199
U(2.0)		-4.2789	0.0066	-0.0056	0.0094	-0.0147	-0.0430	0.0079	-0.0177

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XXVII: All-electron (AE) UCCSD(T) SO ground state ($^3\Sigma$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	5.421(13) ^{† a}	5.412(7)	-0.021(9)	-0.01(1)	0.02(1)	-0.06(1)	0.03(1)	-0.02(1)	0.01(1)
$\omega_e(\text{cm}^{-1})$	1119 ^a	1165(3)	-5(4)	-6(5)	-7(5)	-13(5)	6(5)	-7(5)	0(5)
$r_e(\text{\AA})$	1.481 ^b	1.4772(8)	0.004(1)	0.001(1)	-0.005(1)	0.004(1)	-0.012(1)	0.001(1)	0.000(1)
Potential Energy Surface									
U(1.15)		0.7445	0.1464	-0.0520	-0.4014	0.1084	-0.5759	-0.0397	-0.0339
U(1.25)		-3.0835	0.0913	-0.0052	-0.1843	0.1147	-0.3000	0.0055	-0.0193
U(1.35)		-4.8278	0.0536	0.0053	-0.0830	0.0916	-0.1382	0.0164	-0.0103
U(1.45)		-5.3932	0.0277	0.0047	-0.0352	0.0667	-0.0429	0.0152	-0.0057
U(1.55)		-5.3060	0.0098	0.0014	-0.0128	0.0464	0.0128	0.0101	-0.0039

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XXVIII: All-electron (AE) UCCSD(T) Cl₂ ground state (¹Σ_g) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPS.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	2.5143(4) ^{† a}	2.526(9)	-0.01(1)	-0.01(1)	0.02(1)	-0.01(1)	0.04(1)	-0.01(1)	0.01(1)
$\omega_e(\text{cm}^{-1})$	559.6 ^a	533(5)	-3(7)	0(7)	1(7)	-2(7)	12(8)	-2(7)	1(7)
$r_e(\text{\AA})$	1.988 ^b	1.987(3)	0.005(5)	-0.001(5)	-0.009(5)	0.002(5)	-0.025(5)	0.002(5)	0.000(5)
Potential Energy Surface									
U(1.6)		0.5063	0.0825	-0.0122	-0.2120	0.0335	-0.4174	-0.0025	-0.0102
U(1.7)		-1.1542	0.0534	-0.0036	-0.1263	0.0279	-0.2635	0.0070	-0.0112
U(1.8)		-2.0502	0.0325	0.0011	-0.0736	0.0215	-0.1556	0.0110	-0.0117
U(1.9)		-2.4538	0.0174	0.0037	-0.0409	0.0162	-0.0796	0.0125	-0.0117
U(2.0)		-2.5435	0.0066	0.0053	-0.0208	0.0120	-0.0263	0.0127	-0.0114
U(2.1)		-2.4387	-0.0011	0.0059	-0.0085	0.0088	0.0102	0.0121	-0.0109

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [2]

TABLE XXIX: All-electron (AE) UCCSD(T) ClO ground state ($^2\Pi$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPS.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	2.802(1) ^{† a}	2.753(5)	-0.009(7)	-0.012(7)	0.035(7)	-0.060(7)	-0.026(8)	-0.016(7)	-0.001(7)
$\omega_e(\text{cm}^{-1})$	839.4 ^a	851(3)	-3(4)	-4(4)	0(4)	-13(4)	-4(4)	-6(4)	-2(4)
$r_e(\text{\AA})$	1.596 ^b	1.564(1)	0.004(1)	0.001(2)	-0.008(2)	0.007(2)	-0.007(2)	0.001(2)	0.001(2)
Potential Energy Surface									
U(1.17)		3.2728	0.1081	-0.1807	-0.5878	0.0054	-0.5093	-0.0260	-0.0119
U(1.27)		-0.1726	0.0674	-0.1125	-0.3375	0.0232	-0.2738	0.0076	-0.0002
U(1.37)		-1.8723	0.0400	-0.0694	-0.1919	0.0256	-0.1525	0.0172	0.0038
U(1.47)		-2.5904	0.0215	-0.0423	-0.1080	0.0244	-0.0766	0.0174	0.0034
U(1.57)		-2.7621	0.0089	-0.0259	-0.0592	0.0224	-0.0219	0.0146	0.0012
U(1.67)		-2.6326	0.0002	-0.0156	-0.0297	0.0205	0.0167	0.0114	-0.0010

[†] Sum of experimental dissociation and zero-point energies

^a Reference [1]

^b Reference [6]

TABLE XXX: All-electron (AE) UCCSD(T) ArH+ ground state ($^1\Sigma$) binding parameters and potential energy surface along with the errors from uncorrelated core (UC) and ECPs.

All energies in eV.

Qty.	Exp.	AE	Discrepancies from AE						
			UC	SBKJC	BFD	TN-DF	STU	ccECP[Ne]	ccECP[He]
Binding Parameters									
$D_e(\text{eV})$	4.059 ^{† a}	4.072(4)	0.004(6)	-0.001(6)	0.026(6)	0.001(6)	-0.015(6)	0.000(6)	0.001(6)
$\omega_e(\text{cm}^{-1})$	2723 ^a	2728(18)	-1(26)	2(25)	1(25)	0(26)	-8(25)	-1(25)	2(26)
$r_e(\text{\AA})$	1.286 ^a	1.282(1)	0.001(1)	0.000(1)	-0.004(1)	0.001(1)	-0.002(1)	0.000(1)	0.000(1)
Potential Energy Surface									
U(0.9)		0.4748	0.0307	0.0081	-0.1533	0.0213	-0.0770	-0.0033	0.0020
U(1.0)		-2.1495	0.0160	0.0067	-0.0913	0.0137	-0.0282	0.0014	0.0006
U(1.1)		-3.4435	0.0062	0.0041	-0.0562	0.0064	-0.0030	0.0014	0.0004
U(1.2)		-3.9723	-0.0003	0.0022	-0.0361	0.0012	0.0099	0.0004	0.0008
U(1.3)		-4.0653	-0.0045	0.0013	-0.0244	-0.0018	0.0161	-0.0006	0.0015
U(1.4)		-3.9164	-0.0072	0.0012	-0.0174	-0.0035	0.0186	-0.0013	0.0022

[†] Sum of experimental dissociation and zero-point energies


^a Reference [7]

TABLE XXXI: Parameter values for Cl and Ar ccECP[Ne].S potentials. The highest l value corresponds to the local channel.

Atom	Z_{eff}	l	$n_{l,k}$	$\alpha_{l,k}$	$\beta_{l,k}$
Cl	7	0	2	17.23708573	6.50888648
		0	2	4.31148447	46.76346700
		1	2	11.38275704	2.99464770
		1	2	3.83218762	28.01703410
		2	1	22.71655173	7.00000000
		2	3	78.57185685	159.01586213
		2	2	7.47352436	-15.65310650
Ar	8	0	2	26.06800734	7.67833187
		0	2	5.17516183	60.31303710
		1	2	17.09350095	3.47705723
		1	2	4.64108131	36.47236220
		2	1	8.98034652	8.00000000
		2	3	26.23635698	71.84277216
		2	2	7.10917092	-22.13000430

TABLE XXXII: All-electron (AE) UCCSD(T) ionization potentials and electron affinity of Cl along with the errors from ccECP[Ne].S. The uncontracted aug-cc-pCV5Z basis was used for all calculations. All values in eV. See Tab. V in main text for further description.

Qty.	AE	Discrepancy from AE
		ccECP[Ne].S
IP(I)	12.9388	-0.0427
IP(II)	23.7309	-0.0830
IP(III)	39.6980	0.0233
IP(IV)	53.1811	-0.0321
IP(V)	67.6102	-0.1362
IP(VI)	96.8961	-0.1292
IP(VII)	114.2079	-0.4249
EA	3.6210	-0.0184
AMAD		0.1112
LMAD		0.0480
MARE		0.0025



(a) Cl₂ binding curve discrepancies

(b) ClO binding curve discrepancies

FIG. 1: Binding energy discrepancies for (a) Cl₂ and (b) ClO molecules in their ground states $^1\Sigma_g$ and $^2\Pi$, respectively. The binding curves are relative to the AE UCCSD(T) binding curve. The shaded region indicates a discrepancy of chemical accuracy in either direction.

TABLE XXXIII: All-electron (AE) UCCSD(T) Cl₂ ground state ($^1\Sigma_g$) binding parameters and potential energy surface along with the errors from ccECP[Ne].S. All energies in eV.

Qty.	AE	Discrepancy from AE
		ccECP[Ne].S
Binding Parameters		
D_e (eV)	2.526(9)	-0.04(1)
ω_e (cm ⁻¹)	533(5)	-5(7)
r_e (Å)	1.987(3)	0.002(5)
Potential Energy Surface		
U(1.6)	0.5063	0.0254
U(1.7)	-1.1542	0.0377
U(1.8)	-2.0502	0.0406
U(1.9)	-2.4538	0.0391
U(2.0)	-2.5435	0.0358
U(2.1)	-2.4387	0.0317

TABLE XXXIV: All-electron (AE) UCCSD(T) ClO ground state ($^2\Pi$) binding parameters and potential energy surface along with the errors from ccECP[Ne].S. All energies in eV.

Qty.	Discrepancy from AE	
	AE	ccECP[Ne].S
Binding Parameters		
D_e (eV)	2.753(5)	-0.036(7)
ω_e (cm^{-1})	850(3)	-10(4)
r_e (\AA)	1.564(1)	0.003(2)
Potential Energy Surface		
U(1.17)	3.2728	0.0240
U(1.27)	-0.1726	0.0503
U(1.37)	-1.8723	0.0513
U(1.47)	-2.5904	0.0433
U(1.57)	-2.7621	0.0340
U(1.67)	-2.6326	0.0258

FIG. 2: Binding energy discrepancies for the ArH⁺ molecule in its ground state $^1\Sigma$. The binding curves are relative to the AE UCCSD(T) binding curve. The shaded region indicates a discrepancy of chemical accuracy in either direction.




TABLE XXXV: All-electron (AE) UCCSD(T) ionization potentials and electron affinity of Ar along with the errors from ccECP[Ne].S. The uncontracted aug-cc-pCV5Z basis was used for all calculations. All values in eV. See Tab. V in main text for further description.

Qty.	AE	Discrepancy from AE
		ccECP[Ne].S
IP(I)	15.7829	0.0202
IP(II)	27.6005	-0.0126
IP(III)	40.6121	-0.0862
IP(IV)	59.2804	0.0239
IP(V)	74.7437	-0.1283
IP(VI)	91.1085	-0.3698
IP(VII)	124.3356	-0.1401
IP(VIII)	143.4706	-0.5716
AMAD		0.1112
LMAD		0.0480
MARE		0.0025

TABLE XXXVI: All-electron (AE) UCCSD(T) ArH+ ground state ($^1\Sigma$) binding parameters and potential energy surface along with the errors from ccECP[Ne].S. All energies in eV.

Qty.	AE	Discrepancy from AE
		ccECP[Ne].S
Binding Parameters		
$D_e(\text{eV})$	4.072(4)	-0.002(6)
$\omega_e(\text{cm}^{-1})$	2728(18)	-2(25)
$r_e(\text{\AA})$	1.282(1)	-0.002(1)
Potential Energy Surface		
U(0.9)	0.4748	-0.0614
U(1.0)	-2.1495	-0.0297
U(1.1)	-3.4435	-0.0123
U(1.2)	-3.9723	-0.0027
U(1.3)	-4.0653	0.0026
U(1.4)	-3.9164	0.0054

-
- [1] D. Feller and K. A. Peterson, *The Journal of chemical physics* **110**, 8384 (1999).
- [2] K. Huber and G. Herzberg, Molecular spectra and molecular structure: IV. Constants of diatomic molecules (Van Nostrand Reinhold Co., 1979).
- [3] R. D. J. III, NIST Computational Chemistry Comparison and Benchmark Database NIST Standard Reference Database Number 101 Release 19, [Online]. Available: <http://cccbdb.nist.gov/> [2018, April 19]. National Institute of Standards and Technology, Gaithersburg, MD. (2018).
- [4] K. K. Irikura, *Journal of physical and chemical reference data* **36**, 389 (2007).
- [5] Z. Fu, G. W. Lemire, G. A. Bishea, and M. D. Morse, *The Journal of Chemical Physics* **93**, 8420 (1990).
- [6] F. J. Lovas, E. Tiemann, J. S. Coursey, S. A. Kotochigova, J. Chang, K. Olsen, and R. A. Dragoset, NIST Diatomic Spectral Database 114, [Online]. Available: www.physics.nist.gov/PhysRefData/MolSpec/Diatomic/index.html [2005, November]. Institut für Quantenoptik, Universität Hannover, Hannover, Germany (2005).
- [7] P. Rosmus, *Theoretica chimica acta* **51**, 359 (1979).