

# $\epsilon$ -greedy agents

Pull each arm once, then at step  $t$  pull random w/  $\epsilon_t$ , best w/  $1-\epsilon_t$  prob.

$$\begin{aligned} R_n &= \sum_{j=1}^m R_1^{j, \text{NAB}} + \sum_{i=1}^k \sum_{j=1}^m \sum_{t=m+1}^n \Delta_i \mathbb{P}(S_t=j, O_t^j=i) \\ &= \sum_{i=1}^k \sum_{j=1}^m \sum_{t=1}^n \Delta_i \mathbb{P}(S_t=j, O_t^j=i) \\ &= \sum_{i=1}^k \sum_{j=1}^m \Delta_i \mathbb{E}[T_i^j(n)] \end{aligned}$$

$$T_i^j(n) := \sum_{t=1}^n \sum_{i=1}^k \sum_{j=1}^m \mathbb{1}_{S_t=j} \mathbb{1}_{O_t^j=i}$$

$$\mathbb{P}(S_t=j) = \frac{\epsilon_t}{m} + (1-\epsilon_t) \mathbb{P}(\hat{\mu}_j^A(t-1) \geq \max_{j'} \hat{\mu}_{j'}^A(t-1))$$

$$\begin{aligned} \mathbb{P}(\hat{\mu}_j^A(t-1) \geq \max_{j'} \hat{\mu}_{j'}^A(t-1)) &\leq \mathbb{P}(\hat{\mu}_j^A(t-1) \geq \hat{\mu}_*^A(t-1)) \\ &= \mathbb{E} \left\{ \mathbb{E} \left[ \mathbb{1}_{\{\hat{\mu}_j^A(t-1) \geq \hat{\mu}_*^A(t-1)\}} \mid \mathcal{T}_\Sigma \right] \right\} \end{aligned}$$

$$T_\Sigma^j(n) := \sum_{i=1}^k T_i^j(n) \quad ; \quad \hat{\mu}_j^A(t-1) \in SG(\sqrt{\frac{1}{T_\Sigma^j(t-1)}})$$

$$\begin{aligned}
\mathbb{P}(\hat{\mu}_j^A(t-1) \geq \hat{\mu}_*^A(t-1) \mid T_\Sigma) &\leq e^{-\frac{(\Delta_j^A(t-1))^2}{2} \left( \frac{1}{T_\Sigma^*(t-1)} + \frac{1}{T_\Sigma^j(t-1)} \right)^{-1}} \\
&\leq e^{-\frac{(\Delta_j^A(t-1))^2}{4} T_\Sigma^*(t-1) \wedge T_\Sigma^j(t-1)} \\
&= e^{-\frac{\Delta^2}{4} T_\Sigma^*(t-1)} \vee e^{-\frac{\Delta^2}{4} T_\Sigma^j(t-1)} \\
&\leq e^{-\frac{\Delta^2}{4} \tilde{T}_\Sigma^*(t-1)} \vee e^{-\frac{\Delta^2}{4} \tilde{T}_\Sigma^j(t-1)} \\
&\leq e^{-\frac{\Delta^2}{4} \tilde{T}_\Sigma^*} + e^{-\frac{\Delta^2}{4} \tilde{T}_\Sigma^j}
\end{aligned}$$

where  $\tilde{T}_\Sigma^j(t)$  is the number of times the  $j^{\text{th}}$  recsys was selected during the random exploration rounds.

Note that  $\tilde{T}_\Sigma^j(m) = 1 + \sum_{t=m+1}^m b_t^j$  with  $b_t^j \sim \text{Bernoulli}(\frac{\varepsilon_t}{m})$  independent,  $\forall j$ .

$$\begin{aligned}
\Rightarrow \mathbb{E} \left[ e^{-\frac{(\Delta_j^A(t-1))^2}{4} \tilde{T}_\Sigma^j(t-1)} + e^{-\frac{(\Delta_j^A(t-1))^2}{4} \tilde{T}_\Sigma^*(t-1)} \right] \\
= 2 \mathbb{E} \left[ e^{-\frac{(\Delta_j^A(t-1))^2}{4} \tilde{T}_\Sigma^j(t-1)} \right] \\
= 2 e^{-\frac{(\Delta_j^A(t-1))^2}{4}} \prod_{k=m+1}^{t-1} \mathbb{E} e^{-\frac{(\Delta_j^A(t-1))^2}{4} b_k^j}
\end{aligned}$$

$$= 2 e^{-\frac{(\Delta_j^A(t-1))^2}{4}} \prod_{k=m+1}^{t-1} \left( \left(1 - \frac{\varepsilon_k}{m}\right) + \frac{\varepsilon_k}{m} e^{-\frac{(\Delta_j^A(t-1))^2}{4}} \right)$$

$$= 2 \exp \left\{ -\frac{\Delta^2}{4} + \sum_{k=m+1}^{t-1} \log \left( 1 - \frac{\varepsilon_k}{m} + \frac{\varepsilon_k}{m} e^{-\frac{\Delta^2}{4}} \right) \right\}$$

$$\stackrel{1+x \leq e^x}{\leq} 2 \exp \left\{ -\frac{\Delta^2}{4} + \sum_{k=m+1}^{t-1} \frac{\varepsilon_k}{m} (e^{-\frac{\Delta^2}{4}} - 1) \right\}$$

N: the  $1+x \leq e^x$  could have been taken

$$\dots = 2 \exp \left\{ -\frac{\Delta_j^A(t-1)^2}{4} - \frac{(1 - e^{-\frac{\Delta_j^A(t-1)^2}{4}})}{m} \sum_{k=m+1}^{t-1} \varepsilon_k \right\}$$

$\hookrightarrow$  at any time  $t$ , it's best to be best in  $\mathcal{R}_{t-1}^{j, \text{HAB}}$

• If  $\varepsilon_t = \varepsilon$ , then the 2nd term is

$\Omega(t)$ , if  $\varepsilon_t = \frac{\varepsilon}{t}$  then:

$$\log(t-1) + \frac{1}{t-1} - \log(m+1) - 1 \geq \log\left(\frac{t-1}{m+1}\right) - 1$$

$$\text{so} \geq \frac{\varepsilon}{m} (1 - e^{-\frac{\Delta^2}{4}}) \left( \log \frac{t-1}{m+1} - 1 \right)$$

! [assuming  $t$  suff. large] so

the speed of convergence for this

term is only a power not

exponential for

Crude bound on regret:

*We want to allow prior  
but not  $\Delta_i$   
(except  $ET \rightarrow \Delta_i$ )*

probably  $\frac{1}{k} \sum_{i=1}^k \Delta_i$  for any alg.

$$\sum_{j=1}^m R_{j, \text{DAS}} + \sum_{i=1}^k \sum_{j=1}^m \sum_{t=m+1}^k \Delta_i P(O_{tj}=i | S_t=j) P(S_t=j)$$

$$\text{2nd term: } \leq 2 \sum_{i=1}^k \Delta_i \sum_{j=1}^m \sum_{t=m+1}^m \exp \left\{ -\frac{\Delta_j^A (t-1)^2}{4} - \frac{(1 - e^{-\frac{\Delta_j^A (t-1)^2}{4}})}{m} \sum_{k=m+1}^{t-1} \epsilon_k \right\}$$

*add  $\frac{\epsilon_t}{m} + (1 - \epsilon_t)$*

$$\rightarrow \int_m^{m-1} \exp \left\{ -\frac{\Delta_j^A (t)^2}{4} - \frac{(1 - e^{-\frac{\Delta_j^A (t)^2}{4}})}{m} \sum_{k=m+2}^{t+1} \epsilon_k \right\} dt$$

$$\text{CRUDE-idea} \leq \int_m^{m-1} \exp \left\{ -\frac{(1 - e^{-\frac{\Delta_j^A (t)^2}{4}})}{m} \sum_{k=m+2}^{t+1} \epsilon_k \right\} dt$$

N: we'll have  $\sum_{t=m+1}^m \Delta_i \frac{\epsilon}{t} + \Delta_i (1 - \frac{\epsilon}{t}) \int \dots$

$$\leq \Delta_i \epsilon \log \frac{m}{m+1} + \Delta_i \int \dots$$

and the integral might go down

as  $e^{-\Omega(\log^2 n)} = O(\frac{1}{n})$  or faster

if the difference between